

Symbiotic *Bradyrhizobium japonicum* Reduces N₂O Surrounding the Soybean Root System via Nitrous Oxide Reductase

Reiko Sameshima-Saito,† Kaori Chiba, Junta Hirayama, Manabu Itakura, Hisayuki Mitsui, Shima Eda, and Kiwamu Minamisawa*

Graduate School of Life Sciences, Tohoku University, Katahira, Aoba-ku, Sendai 980-8577, Japan

Received 23 October 2005/Accepted 25 January 2006

N₂O reductase activity in soybean nodules formed with *Bradyrhizobium japonicum* was evaluated from N₂O uptake and conversion of ¹⁵N-N₂O into ¹⁵N-N₂. Free-living cells of USDA110 showed N₂O reductase activity, whereas a *nosZ* mutant did not. Complementation of the *nosZ* mutant with two cosmids containing the *nosRZDFYLX* genes of *B. japonicum* USDA110 restored the N₂O reductase activity. When detached soybean nodules formed with USDA110 were fed with ¹⁵N-N₂O, they rapidly emitted ¹⁵N-N₂ outside the nodules at a ratio of 98.5% of ¹⁵N-N₂O uptake, but nodules inoculated with the *nosZ* mutant did not. Surprisingly, N₂O uptake by soybean roots nodulated with USDA110 was observed even in ambient air containing a low concentration of N₂O (0.34 ppm). These results indicate that the conversion of N₂O to N₂ depends exclusively on the respiratory N₂O reductase and that soybean roots nodulated with *B. japonicum* carrying the *nos* genes are able to remove very low concentrations of N₂O.

Nitrous oxide (N₂O) is a key atmospheric greenhouse gas that contributes to global climate change through radiative warming and depletion of stratospheric ozone (1, 24). Agricultural land is a major source through denitrification and nitrification (14, 35) and contributes significantly to the net increase in atmospheric N₂O (1, 34). Several attempts have been made to reduce the emission of N₂O from agricultural systems (34, 35).

The complete denitrification of nitrate by bacteria to dinitrogen (N₂) is generally an anaerobic respiratory process, where the last step is mediated by N₂O reductase (54). The corresponding structural gene is *nosZ* and is assembled in the *nosRZDFYLX* gene operon (54). Several species capable of denitrification are also nitrogen-fixing bacteria, including rhizobia such as *Bradyrhizobium japonicum* (4, 6, 44, 49) and *Sinorhizobium meliloti* (7, 20). Indeed, genes responsible for denitrification have been found in rhizobial genomes (15, 23).

Earlier studies (19, 37) reported the evolution of ¹⁵N-N₂ from ¹⁵N-N₂O from sliced or detached soybean nodules. Recently, Velasco et al. (50) reported that *nosZ* and *nosR* insertion mutants of *B. japonicum* USDA110 accumulate N₂O when cultured microaerobically in the presence of nitrate. The *nosZ* gene was also expressed in soybean nodules (29). However, it has not yet been fully proved that N₂ evolution from N₂O by soybean nodules is mediated by N₂O reductase encoded by the *nosZ* gene in *B. japonicum* (4, 9, 38). The aims of this work were to confirm whether the *nos* gene cluster of *B. japonicum* is responsible for respiratory N₂O reduction to N₂ in nodules and to evaluate the capability of the nodulated roots to transform N₂O into N₂.

* Corresponding author. Mailing address: Graduate School of Life Sciences, Tohoku University, Katahira, Aoba-ku, Sendai 980-8577, Japan. Phone and fax: 81-22-217-5684. E-mail: kiwamu@ige.tohoku.ac.jp.

† Present address: Environmental Science for Human Life, Faculty of Agriculture, Shizuoka University, Ohya 836, Shizuoka 422-8529, Japan.

MATERIALS AND METHODS

Bacterial strains, plasmids, and media. The bacterial strains and plasmids used are listed in Table 1. *Bradyrhizobium* cells were grown at 30°C in HM salt medium (8) supplemented with 0.1% arabinose and 0.025% (wt/vol) yeast extract (Difco, Detroit, MI), which is termed HM medium here. HM medium was further supplemented with 0.55 μM Na₂MoO₄ · 2H₂O, 1 μM FeCl₃, and 1 μM CuSO₄ · 5H₂O (HMM medium) for the denitrification assay (44). *Escherichia coli* cells were grown at 37°C in Luria-Bertani medium (42). Antibiotics were added to the media at the following concentrations: for *B. japonicum*, 100 μg of tetracycline (Tc)/ml, 100 μg of spectinomycin (Sp)/ml, 100 μg of streptomycin (Sm)/ml, 100 μg of kanamycin (Km)/ml, and 50 μg of polymyxin B/ml; for *E. coli*, 15 μg of Tc/ml, 50 μg of Sp/ml, 50 μg of Sm/ml, 50 μg of Km/ml, and 100 μg of ampicillin/ml.

DNA manipulations. Isolation of plasmids, DNA ligation, and transformation of *E. coli* were performed as described by Sambrook et al. (42). DNA preparation and Southern hybridization were carried out as described previously (22, 30, 43).

Construction of a *B. japonicum* USDA110 *nosZ* mutant. A 4-kb BamHI DNA fragment identified from the genome sequence of *B. japonicum* USDA110 (23) was excised from BamHI-digested total DNA and inserted into the BamHI site of pTZ18R (Fig. 1A). The plasmid containing the fragment was selected by PCR amplification specific for the *nosZ* gene of *B. japonicum* USDA110 with primer 3 (5'-GAGGATCGTTTCGCATGAGCGACAGCGACAACAT-3') and primer 4 (5'-CGCTCCCGATCAGACCGATT-3'), in which the underlined nucleotides form the USDA110 *nosZ* sequence. From the cloned 4-kb DNA fragment, pK18mob-*nosZ*::Ω was constructed (Fig. 1A). Biparental mating was conducted on HM agar plates using *E. coli* S17-1 (47). Double crossover was verified by Southern hybridization with the 4-kb *nosRZD* fragment as a probe (Fig. 1A).

Genetic complementation by *nos* genes. Two cosmids, brc02856 and brc01733, carrying the *nos* gene cluster (*nosRZDFYLX*) were selected from the pKS800 cosmid library, which was constructed for the sequencing of *B. japonicum* USDA110 (23). The cosmids were introduced into *B. japonicum* strains by triparental mating via pRK2013 (39, 47, 52).

Gas chromatography. To determine N₂O concentrations, we used two ⁶³Ni electron capture detector gas chromatographs. In the quick N₂O analysis, 0.2 ml of sample gas was injected into a Shimadzu (Kyoto, Japan) GC-17A gas chromatograph with a CP-PoraBOND Q capillary column (internal diameter, 0.32 mm, length, 25 m; Varian, Palo Alto, CA). The temperatures of injection, the column, and the detector were 250, 65, and 300°C, respectively. In the sensitive N₂O analysis, 0.8 ml of sample gas was injected into a Shimadzu GC-14BPsE gas chromatograph equipped with tandem packed columns of a Porapak N (80/100 mesh; diameter, 0.3 mm; length, 1 m) and a Porapak Q (80/100 mesh; diameter, 0.3 mm; length, 3 m). The temperatures of injection, the column, and the detector were 80, 80, and 340°C, respectively. The carrier gas used for both N₂O analyses was composed of 5% (vol/vol) CH₄ in Ar. ¹⁵N-N₂ was determined with a thermal conductivity detector gas chromatograph as described previously (44).

TABLE 1. Bacterial strains and cosmids used in this study

Strain or plasmid	Characteristics ^a	Source or reference
Strains		
<i>Bradyrhizobium japonicum</i>		
USDA110	Wild type; <i>nosZ</i> ⁺	23
T7	Field isolate in Tokachi, Hokkaido, Japan; <i>nosZ</i> natural mutant	32, 43
USDA110Δ <i>nosZ</i>	USDA110 <i>nosZ</i> ::del/ins Ω cassette; Sm ^r Sp ^r	This study
<i>Escherichia coli</i>		
DH5α	<i>recA</i> ; cloning strain	Toyobo Inc.
JM109	<i>recA</i> ; cloning strain	Toyobo Inc.
HB101	<i>recA hsdR hsdM pro</i> Sm ^r	5
S17-1	<i>hsdR</i> RP4-2(kan::Tn7)(tet::Mu); Sm ^r Sp ^r	47
Plasmids		
pRK2013	ColE1 replicon carrying RK2 transfer genes; Km ^r	13
pTZ18R	Cloning vector; pMB1 <i>ori</i> ; Ap ^r	46
pTZ18R <i>nosZ</i>	pTZ18R carrying 4.0-kb <i>nosZ</i> fragment; Ap ^r	This study
pTZ18RΔ <i>SacI-nosZ</i>	<i>SacI</i> -deleted pTZ18R carrying 4.0-kb <i>nosZ</i> fragment; Ap ^r	This study
pHP45Ω	Plasmid carrying 2.1-kb Ω cassette; Sp ^r Sm ^r Ap ^r	40
pK18mob	Cloning vector; pMB1 <i>ori oriT</i> ; Km ^r	46
pK18mob- <i>nosZ</i> ::Ω	pK18mob carrying 4.0-kb <i>nosZ</i> fragment; Km ^r Sm ^r Sp ^r	This study
Cosmids		
pKS800	Derivative of broad-host-range cosmid pLAFR1; IncP Tc ^r	17, 23
brc02856	pKS800 carrying <i>nos</i> gene cluster ^b ; Tc ^r	23
brc01733	pKS800 carrying <i>nos</i> gene cluster ^b ; Tc ^r	23

^a Ap^r, ampicillin resistant; Tc^r, tetracycline resistant; Km^r, kanamycin resistant; Sm^r, streptomycin resistant; Sp^r, spectinomycin resistant.

^b See Fig. 1.

For the acetylene reduction assay, ethylene was measured by a flame ionization detection gas chromatograph, as described previously (12).

N₂O uptake by free-living cells. *B. japonicum* cells were washed with HMM medium by centrifugation (5,000 × *g*, 15 min, 4°C), and the cell density was adjusted to 6 × 10⁹ cells/ml (33). The cell suspension (4.6 ml, 6 × 10⁹ cells/ml) was introduced into a 123-ml airtight vial (V-100; Nichiden-Rika Glass, Kobe, Japan). After the gas phase was replaced with N₂, N₂O gas was introduced into the vial at a final concentration of approximately 0.1% (vol/vol).

¹⁵N-N₂ emission from ¹⁵N-labeled N₂O by free-living cells. *B. japonicum* cells (7 ml; 6 × 10⁹ cells/ml) were introduced into a 14-ml airtight specimen vial. After the N₂ gas replacement, ¹⁵N-N₂O (¹⁵N, >98 atom%; Cambridge Isotope Laboratories Inc., Andover, MA) was introduced. After incubation at 30°C for 67 h, the concentration of ¹⁵N-N₂ in the gas phase was determined by TCD gas chromatography (44).

Real-time PCR. Beacon Designer software (Premier Biosoft International, Palo Alto, CA) was used to design the *nosZ* primer set (*nosZ*2F, 5'-GTCGTC ATCTCAACCTCAAG; *nosZ*2R, 5'-TATTCATGCCATGCGGATTG). Total RNA was prepared as described previously (53). Quantitative reverse transcription-PCR analysis was carried out by the i-Cycler optical system (Bio-Rad Laboratories, Inc., Tokyo, Japan) as described previously (53).

Plant cultivation and inoculation. The surface-sterilized soybean seeds (*Glycine max* cv. Enrei) were germinated and transplanted to a Leonard jar (25, 48), which contained sterile vermiculite and nitrogen-free sterilized nutrient solution (31, 39). The cell suspension of *B. japonicum* (1 ml) was used for inoculation at 1 × 10⁷ cells per seed. For N₂O-feeding experiments, plants were grown in a growth chamber (LH200; Nippon Medical & Chemical Industries, Tokyo, Japan). For the nitrogen fixation experiment, soybeans were grown in pots (1 liter) filled with vermiculite in a phytotron (Koitoiron type KC; Koito Industries, Tokyo, Japan). A nitrogen-free sterilized nutrient solution was periodically supplied to the pots (31, 39).

N₂O uptake, ¹⁵N-N₂ emission, and ¹⁵N incorporation by nodules. After the nodulated roots of 45-day-old soybean plants were washed with water, the nodules were detached from the roots and weighed. The nodules were introduced into a 15-ml airtight vial. After the vials were sealed, ¹⁵N-N₂O was injected. After the gas measurements, the nodules were dried at 80°C for 3 days and then powdered with a mortar and pestle. The nodule samples were subjected to tracer ¹⁵N and total-N analyses by an automatic gas chromatograph-mass spectrometer

(EA1110-DELTA^{plus} Advantage ConFloIII system, Thermo Electron Co., Bremen, Germany).

The whole root system of 32-day-old soybean plants was inserted into a 51-ml test tube (19.4 mm in diameter by 176 mm in height) containing 20 ml nitrogen-free nutrient solution (31, 39). The test tube was sealed with a silicon rubber stopper with a gas sampling port and a hole in the center. This system enclosed the whole root system of intact soybeans in the test tube, leaving the aerial parts in air. Then N₂O gas was injected into the closed test tube.

Nitrogen fixation and plant growth. The nitrogen-fixing activity of nodules was examined by acetylene reduction assays in an incubation jar (300-ml capacity) containing the excised root system. Acetylene gas was injected into the jar at 15% (vol/vol), and the roots were incubated for 60 min at 25°C. Separated shoot, roots, and nodules were weighed.

Bacteroid antibiotic test. Nodules were removed from soybeans 18 days after inoculation with T7 or T7 with brc01733. Homogenates of surface-sterilized nodules were plated on HM agar medium with and without tetracycline.

RESULTS

Construction of the *nosZ* deletion mutant. To examine whether the *nos* gene cluster of *B. japonicum* was responsible for N₂O reductase activity, we constructed a *nosZ* deletion mutant of *B. japonicum* USDA110 (Table 1). After deletion of most of the *nosZ* gene from the cloned DNA fragment containing *nosRZD*, the Ω cassette was inserted into the position (Fig. 1). When free-living cells of USDA110 and USDA110Δ*nosZ* were anaerobically incubated in gas containing 0.1% (vol/vol) N₂O, the parent strain consumed N₂O, whereas the mutant did not show N₂O uptake (Fig. 2A). To further examine the N₂O reductase reaction, ¹⁵N-N₂O was supplied to cultures of *B. japonicum* USDA110 and USDA110Δ*nosZ*. Wild-type USDA110 stoichiometrically produced ¹⁵N-N₂ from ¹⁵N-N₂O, whereas USDA110Δ*nosZ* did not (data




FIG. 1. Construction of a *nosZ* mutant (A) and complementation cosmids inserts (B) of *Bradyrhizobium japonicum* USDA110. (A) Cloned fragments in pTZ18RΔSacI-*nosZ* and pK18mob-Δ*nosZ*::Ω are shown alongside the physical map of the *nos* gene cluster of *B. japonicum* USDA110. B, BamHI; S, SacI; X, XhoI; Sb, SacI blunt end; Bb, BamHI blunt end; Xb, XhoI blunt end. During blunt-end ligation, the Sb/Bb site produced a BamHI site in pK18mob-Δ*nosZ*::Ω. Upstream of *nosR* lies a consensus sequence of an FNR box (TTGAT-N₄-ATCAA) (50). The cloned 4-kb DNA fragment containing the *nosZ* gene in pTZ18R-*nosZ* was reconnected to pTZ18RΔSacI, which was produced by blunting the SacI site of pTZ18R, and pTZ18RΔSacI-*nosZ* was generated. The Ω cassette, which was excised from pHP45Ω (40), digested with BamHI, and blunt ended, was inserted into the blunt-ended SacI and XhoI sites of pTZ18RΔSacI-*nosZ* to generate pTZ18RΔSacI-*nosZ*::Ω. Finally, the fragment containing the *nosZ* and Ω cassette fusion was inserted between the SmaI and XbaI sites of pK18mob to generate pK18mob-*nosZ*::Ω. (B) Positions covered by two cosmids inserts, brc02856 and brc01733 (Table 1), in the genome of *B. japonicum* USDA110 (<http://www.kazusa.or.jp/rhizobase/>) (23).

not shown). These results suggest that *nosZ* of *B. japonicum* USDA110 is responsible for the N₂O reductase activity that converts N₂O to N₂.

Complementation of USDA110Δ*nosZ* with *nos* genes. To further examine the validity of *nosZ*-dependent N₂O reductase activity, two cosmids containing the *nos* gene cluster were introduced into USDA110Δ*nosZ* (Fig. 1B). The transconjugants were anaerobically incubated in the presence of 0.1%

(vol/vol) N₂O and tetracycline, and USDA110 and USDA110Δ*nosZ* carrying the pKS800 vector were used as controls (Fig. 2B). USDA110 and USDA110Δ*nosZ* carrying pKS800 (Fig. 2B) showed N₂O uptake profiles similar to those of wild-type USDA110 and USDA110Δ*nosZ*, respectively (Fig. 2A). However, the two cosmids in Fig. 1B restored strong N₂O uptake (Fig. 2B). The amino acid sequences of *B. japonicum nosRZDFYLX* (23, 50) have a high degree of similarity to those




FIG. 2. N₂O uptake by free-living cells of the *nosZ* mutant (A) and *nos* complementation of USDA110Δ*nosZ* (B) and T7 (C). pKS800 is the cosmid vector used for complementation (17). Two cosmid clones, brc02856 and brc01733, have inserts containing *nos* genes of USDA110 (Table 1; Fig. 1). N₂O reductase activities (means ± SDs) estimated from maximum rates of N₂O uptake were 4.6 ± 0.2 [USDA110(pKS800)], 22.9 ± 0.7 [USDA110Δ*nosZ*(brc02856)], and 23.6 ± 1.9 N₂O [USDA110Δ*nosZ*(brc01733)] μmol h⁻¹ 10⁷ cells⁻¹.

TABLE 2. Growth and N₂-fixing activity of soybean plants inoculated with *Bradyrhizobium japonicum* USDA110 or the *nosZ* mutant^a

Strain	Growth (g [fresh wt] plant ⁻¹)			N ₂ -fixing activity (C ₂ H ₂ -reducing activity) (μmol h ⁻¹ g of nodule [fresh wt] ⁻¹)
	Shoot	Root	Nodule	
USDA110	10.8 ± 0.7	4.8 ± 0.4	1.0 ± 0.1	10.3 ± 1.3
USDA110Δ <i>nosZ</i>	10.4 ± 0.9	4.5 ± 0.4	0.9 ± 0.1	9.0 ± 2.0
Uninoculated	6.9 ± 0.2	5.5 ± 0.2	0	0

^a Values for growth and N₂-fixing activity are shown as means ± SDs, with six replications.

of other N₂O reductases from *Pseudomonas stutzeri* (21). Both *B. japonicum* cosmids contain *nosRZDFYLX* and adjacent genes (Fig. 1B). Their addition to USDA110Δ*nosZ* produced similar activities of N₂O reductase, as indicated by N₂O uptake (Fig. 2B). Moreover, the introduction of a cosmid (brc02856 or brc01733) into USDA110Δ*nosZ* restored the strain's ability to convert ¹⁵N-N₂O into ¹⁵N-N₂ (data not shown). Thus, the complementation experiments confirmed that the *nos* gene cluster per se is responsible for N₂O reductase activity in *B. japonicum* USDA110.

Expression of *nosZ*. Strain USDA110Δ*nosZ* with the two cosmids containing the *nos* cluster rapidly and strongly absorbed N₂O, much more than the parent strain USDA110 containing the pKS800 vector (Fig. 2B). When the N₂O reductase activity was estimated from the maximum rate of N₂O uptake in Fig. 2B, the activities of USDA110Δ*nosZ* containing brc02856 and brc01733 (22.9 ± 0.7 and 23.6 ± 1.9 N₂O μmol h⁻¹ 10⁷ cells⁻¹) were five times that of wild-type USDA110 (4.6 ± 0.2 N₂O μmol h⁻¹ 10⁷ cells⁻¹). In addition, the lag time for the induction of N₂O reductase activity in the wild-type USDA110(pKS800) disappeared in the cosmid-complemented USDA110Δ*nosZ*.

Because we used pKS800, a derivative of cosmid pLAFR1 (RK2), as a vector for the *nos* complementation (17), it is possible that the cosmid introduction increases the copy number and thus results in higher expression of *nos* genes. USDA110Δ*nosZ* carrying brc01733 expressed the *nosZ* gene at 5-times-higher levels on the basis of total RNA than USDA110Δ*nosZ* carrying the pKS800 vector 1 h after anaer-

obic incubation in the presence of 0.1% (vol/vol) N₂O. This result suggests that the higher expression of the *nos* genes enhanced N₂O reductase activity and shortened the lag time for induction in USDA110Δ*nosZ* with the *nos* cosmids.

N₂O uptake and N₂ emission by nodules. Denitrification genes of *B. japonicum* are expressed in a microaerobic state in both legume nodules and free-living bacteria via the *fixLJ* and *fixK₂* regulatory cascades (3, 28). Mesa et al. (29) reported that the *nir*, *nor*, and *nos* denitrification genes in *B. japonicum* were expressed in soybean root nodules. We thus examined the nodulation phenotypes of the *nosZ* mutant. The nodule weight and nitrogen-fixing activity of USDAΔ*nosZ* showed no significant differences from those of USDA110 (Table 2), indicating that *nosZ* mutation does not alter symbiotic nitrogen fixation. This agrees with the results of a report on the nodulation kinetics (29) of a *nosZ* insertion mutant of *B. japonicum* (50).

To evaluate the extent of N₂O uptake and N₂ emission by N₂O reductase in soybean nodules, ¹⁵N-N₂O was supplied to detached nodules formed with the *nosZ* mutant or the parent strain (Fig. 3). Soybean nodules formed with USDA110 rapidly absorbed ¹⁵N-N₂O and emitted ¹⁵N-N₂ outside the nodules, whereas nodules formed with the *nosZ* mutant did not mediate the conversion of ¹⁵N-N₂O to ¹⁵N-N₂ at all (Fig. 3).

Because the transformation of ¹⁵N-N₂O into ¹⁵N-N₂ was likely stoichiometric (Fig. 3), we determined the amount of fixed ¹⁵N in the nodules and made a ¹⁵N balance sheet for ¹⁵N-N₂O exposure. ¹⁵N atoms of ¹⁵N-N₂O were distributed 98.5% ± 0.1% (mean ± standard deviation [SD]) outside the nodules as ¹⁵N-N₂ and 1.5% ± 0.1% (mean ± SD) fixed in the nodules. Thus, most N₂ molecules produced by N₂O reductase in bacteroids were released outside the nodules, and nitrogenase was not able to utilize most of them as a substrate for nitrogen fixation. This is probably due to the diffusion of N₂ into various compartments for N₂O reductase and nitrogenase. The apparent rate of N₂O uptake by USA110 nodules under about 3% N₂O (Fig. 3) was estimated to be 7.8 μmol h⁻¹ per g (fresh weight) of nodule; this was comparable with the acetylene-reducing activity (Table 2). These results indicate that the conversion of N₂O into N₂ in soybean nodules depends on the function of the *nos* gene cluster in *B. japonicum*.

N₂O uptake by intact soybeans nodulated with USDA110. To examine the extent of N₂O concentrations that the nodu-




FIG. 3. ¹⁵N-N₂O uptake and ¹⁵N-N₂ emission by detached soybean nodules formed with *Bradyrhizobium japonicum* USDA110 (A) and the *nosZ* mutant USDA110Δ*nosZ* (B). ¹⁵N-labeled N₂O was supplied to vials (15 ml) containing soybean nodules (1.6 to 1.8 g). Five independent experiments gave similar results.




FIG. 4. N₂O uptake from root systems of intact plants (A and B) and decapitated root systems (C) inoculated with the wild-type strain or the *nosZ* mutant of *Bradyrhizobium japonicum* USDA110. Initial gas phases were adjusted to approximately 800 ppm (A) (intact plant) and 0.34 ppm (B [intact plant] and C [decapitated root system]) of N₂O in air. Soybean plants that harbored 0.26 to 0.30 g (A), 0.15 to 0.22 g (B), or 0.18 to 0.36 g (C) of nodules (fresh weight per plant) were examined by a test tube system (see the text). Triplicate determinations were carried out, except that data for USDA110Δ*nosZ* inoculation in panel A show means of duplicate determinations. Error bars, SDs.

lated roots can take up, the whole root systems of intact plants nodulated by USDA110 or the *nosZ* mutant were enclosed. After the introduction of N₂O gas, N₂O concentrations around the whole root system were monitored (Fig. 4A and B). At an initial N₂O concentration of 800 ppm (Fig. 4A), the root systems nodulated by USDA110 absorbed N₂O, whereas the *nosZ* mutant-inoculated roots did not take up N₂O at all. After a 21-h incubation, the N₂O concentration reached 4 to 8 ppm

around the roots nodulated with USDA110 (Fig. 4A). Thus, the initial concentrations of N₂O were gradually decreased to 50, 5, 1, and 0.34 ppm. Surprisingly, the root systems nodulated by USDA110 were able to take up 0.34 ppm N₂O, reducing the concentration to 0.11 ± 0.02 ppm (mean \pm SD) 22 h after the start of incubation (Fig. 4B). When a shoot-decapitated root system was completely enclosed in the test tube with a rubber stopper to avoid gas leakage, the N₂O concentration around the root system nodulated by USDA110 decreased to below the detection limit (<0.02 ppm) by 17 h (Fig. 4C).

Generally, the N₂O concentration in soils (0.36 ppm to 8,300 ppm) is higher than that in ambient air (0.31 ppm) (1, 26, 34, 45). Therefore, these results strongly suggest that soybean nodules formed with *B. japonicum* carrying *nos* genes potentially scavenge N₂O gas in soil.

Introduction of *nos* genes into a naturally occurring *nosZ*-deficient strain. Strains of *B. japonicum* can apparently be classified into three denitrification types: full denitrifiers (up to N₂), truncated denitrifiers (up to N₂O), and nondenitrifiers (44, 49). Their incidence depends on the field site and the phylogeny (44). We tested whether introduction of the *nos* gene cluster into the naturally occurring *nosRZD*-deficient strain T7 would induce N₂O reductase activity. Introduction of *brc02856* or *brc01733* into strain T7 induced N₂O uptake, but T7 carrying the cosmid vector did not show such activity (Fig. 2C).

To test whether strain T7 carrying *brc01733* exhibited N₂O reductase activity in nodules, this strain was inoculated into soybean plants. However, the instability of cosmid *brc01733* interfered with this experiment: Bacteroid cells from the nodules lost cosmid *brc01733* during nodule development, as suggested by plate counts of tetracycline-resistant colonies (<10⁻⁶). Because similar instability of pLAFR1 derivatives has been reported (10), other strategies are needed for the symbiotic testing of strain T7 carrying *nos* genes.

DISCUSSION

The *nosZ* mutant of USDA110 was not able to convert ¹⁵N-N₂O to ¹⁵N-N₂ in nodules (Fig. 3) or as free-living cells (Fig. 2) and continued to produce nitrogen-fixing nodules (Table 2). Thus, the evolution of ¹⁵N-N₂ from ¹⁵N-N₂O in soybean nodules is due exclusively to N₂O reductase, which is encoded by *nosZ*.

An important finding of this work is that N₂O reductase expressed in *B. japonicum* can take up very low concentrations of N₂O from outside of the nodules, equivalent to the natural concentration of N₂O in air (approximately 0.31 ppm). Probably this indicates that N₂O reductase has a high affinity for N₂O as a substrate (54), and the gas movement through the tissues of the nodules does not severely interfere with the enzyme property.

Agricultural soil is a major source of the N₂O emissions that contribute to global warming (1, 34). The N₂O concentration in the soil gas phase fluctuates with water, organic matter, and nitrogen fertilizer but is generally higher than that of ambient air (1, 14, 18, 24, 26, 45). The results of N₂O uptake by soybean nodules (Fig. 3 and 4) suggest that soybean nodules formed with *B. japonicum* carrying *nos* genes scavenge N₂O in soil, thus lessening N₂O emission to the atmosphere from soybean fields.

However, several groups have reported that cultivation of legume crops often enhances N₂O emission from fields of alfalfa (11), soybean (51), white clover (36), and Bengal gram (16). Thus, we want to discuss the N₂O paradox of legume root nodules for those results and ours. One possible explanation for the N₂O paradox is that the nitrate absorbed via legume roots and nodule surfaces from the soil solution might be transported in nodules (27, 41), which would emit N₂O from nitrate in bacteroids that lack *nos* genes. Another explanation is that N₂O is produced by rhizosphere organisms from degrading nodules (51), where fixed nitrogen would become a source of N₂O emission irrespective of the *nos* genes in the bacteroids. If either assumption is correct, the presence of *nos* genes in rhizobia ought still to decrease N₂O levels in the rhizosphere and N₂O emission from field soils.

Rhizobial inoculation technology focuses on increasing legume yields. The *nos* gene-dependent transformation of N₂O into N₂ in soybean nodules prompted us to enhance this activity in bradyrhizobial inoculants—a technology that could be environmentally friendly. The introduction of cosmids carrying *nos* genes enhanced *nosZ* transcription and N₂O reductase activity (Fig. 2B), suggesting that up-regulation of *nos* gene expression would result in the enhancement of N₂O reductase activity in *B. japonicum*.

Natural populations of *B. japonicum* that lacks N₂O reductase activity were often dominant in soybean fields (44). To generate bradyrhizobial inoculants with high N₂O reductase activity in soybean nodules, one can adopt several strategies: (i) introduction of *nos* genes by using stable vectors or chromosome integration with strong promoters, (ii) selection of natural isolates with high N₂O reductase activity, and (iii) simple use of USDA110 and other wild-type strains carrying *nos* genes as inoculants. Denitrification steps, including N₂O reduction, are anaerobic respiration processes. It is thus important to examine the allocations of electrons to O₂ and N₂O in soybean nodules, because energy metabolism in bacteroids is supported by *cbb₃* terminal oxidase, which has a high affinity for O₂ and adapts to a microoxic environment in root nodules (2).

ACKNOWLEDGMENTS

This work was supported by grants 13007988-00 to R. Sameshima-Saito and 14360037 and 17380046 to K. Minamisawa from the Ministry of Education, Science, Sports and Culture of Japan.

We are grateful to M. J. Sadowsky (University of Minnesota) for providing *B. japonicum* strain USDA110; T. Kaneko and S. Tabata (Kazusa DNA Research Institute) for providing the cosmids carrying the *nos* gene cluster; and T. Ito (Tohoku University) for help in N₂O analysis by ⁶³Ni electron capture detector gas chromatography.

REFERENCES

- Banin, A., J. G. Lawless, and R. C. Whitten. 1984. Global N₂O cycles—terrestrial emissions, atmospheric accumulation and biospheric effects. *Adv. Space Res.* **4**:207–216.
- Batut, J. 2004. Genomics of the *ccoNOPQ*-encoded *cbb₃* oxidase complex in bacteria. *Arch. Microbiol.* **181**:89–96.
- Bedmar, E. J., E. F. Robles, and M. J. Delgado. 2005. The complete denitrification pathway of the symbiotic, nitrogen-fixing bacterium *Bradyrhizobium japonicum*. *Biochem. Soc. Trans.* **33**:141–144.
- Bhandari, B., and D. J. D. Nicholas. 1984. Denitrification of nitrate to nitrogen gas by washed cells of *Rhizobium japonicum* and by bacteroids from *Glycine max*. *Planta* **161**:81–85.
- Boyer, H. W., and D. Roulland-Dussoix. 1969. A complementation analysis of the restriction and modification of DNA in *Escherichia coli*. *J. Mol. Biol.* **41**:459–472.
- Breitenbeck, G. A., and J. M. Bremner. 1989. Ability of free-living cells of *Bradyrhizobium japonicum* to denitrify in soils. *Biol. Fertil. Soils* **7**:219–224.
- Chan, Y. K., L. Barran, and E. S. P. Bromfield. 1989. Denitrification activity of phage types representative of two populations of indigenous *Rhizobium meliloti*. *Can. J. Microbiol.* **35**:737–740.
- Cole, M. A., and G. H. Elkan. 1973. Transmissible resistance to penicillin G, neomycin, and chloramphenicol in *Rhizobium japonicum*. *Antimicrob. Agents Chemother.* **4**:248–253.
- Coyne, M. S., and D. D. Focht. 1987. Nitrous oxide reduction in nodules—denitrification or N₂ fixation? *Appl. Environ. Microbiol.* **53**:1167–1170.
- Dombrecht, B., J. Vanderleyden, and J. Michiels. 2001. Stable RK2-derived cloning vectors for the analysis of gene expression and gene function in Gram-negative bacteria. *Mol. Plant-Microbe Interact.* **14**:426–430.
- Duxbury, J. M., D. R. Bouldin, R. E. Terry, and R. L. Tate. 1982. Emission of nitrous oxide from soils. *Nature* **298**:462–464.
- Elbeltagy, A., K. Nishioka, T. Sato, H. Suzuki, B. Ye, T. Hamada, T. Isawa, H. Mitsui, and K. Minamisawa. 2001. Endophytic colonization and in planta nitrogen fixation by a *Herbaspirillum* sp. isolated from wild rice species. *Appl. Environ. Microbiol.* **67**:5285–5293.
- Figurski, D. H., and D. R. Helinski. 1979. Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in *trans*. *Proc. Natl. Acad. Sci. USA* **76**:1648–1652.
- Firestone, M. K., R. B. Firestone, and J. M. Tiedje. 1980. Nitrous oxide from soil denitrification: factors controlling its biological production. *Science* **208**:749–751.
- Galibert, F., T. M. Finan, S. R. Long, A. Puhler, P. Abola, F. Ampe, F. Barloy-Hubler, M. J. Barnett, A. Becker, P. Boistard, G. Bothe, M. Boutry, L. Bowser, J. Buhrmester, E. Cadieu, D. Capela, P. Chain, A. Cowie, R. W. Davis, S. Dreano, N. A. Federspiel, R. F. Fisher, S. Gloux, T. Godrie, A. Goffeau, B. Golding, J. Gouzy, M. Gurjal, I. Hernandez-Lucas, A. Hong, L. Huizar, R. W. Hyman, T. Jones, D. Kahn, M. L. Kahn, S. Kalman, D. H. Keating, E. Kiss, C. Komp, V. Lelaure, D. Masuy, C. Palm, M. C. Peck, T. M. Pohl, D. Portetelle, B. Purnelle, U. Ramsperger, R. Surzycki, P. Thebault, M. Vandenberg, F. J. Vorholter, S. Weidner, D. H. Wells, K. Wong, K. C. Yeh, and J. Batut. 2001. The composite genome of the legume symbiont *Sinorhizobium meliloti*. *Science* **293**:668–672.
- Ghosh, S., D. Majumdar, and M. C. Jain. 2002. Nitrous oxide emissions from kharif and rabi legumes grown on an alluvial soil. *Biol. Fertil. Soils* **35**:473–478.
- Hattori, Y., H. Omori, M. Hanyu, N. Kaseda, E. Mishima, T. Kaneko, S. Tabata, and K. Saeki. 2002. Ordered cosmid library of the *Mesorhizobium loti* MAFF303099 genome for systematic gene disruption and complementation analysis. *Plant Cell Physiol.* **43**:1542–1557.
- Heincke, M., and M. Kaupenjohann. 1999. Effects of soil solution on the dynamics of N₂O concentrations: a review. *Nutr. Cycl. Agroecosyst.* **55**:133–157.
- Hoch, G. E., K. C. Schneider, and R. H. Burris. 1960. Hydrogen evolution and exchange, and conversion of N₂O to N₂ by soybean root nodules. *Biochim. Biophys. Acta* **37**:273–279.
- Hollway, P., W. McCormick, R. J. Watson, and Y. K. Chan. 1996. Identification and analysis of the dissimilatory nitrous oxide reduction genes, *nosRZDFY*, of *Rhizobium meliloti*. *J. Bacteriol.* **178**:1505–1514.
- Honisch, U., and W. G. Zumft. 2003. Operon structure and regulation of the *nos* gene region of *Pseudomonas stutzeri*, encoding an ABC-type ATPase for maturation of nitrous oxide reductase. *J. Bacteriol.* **185**:1895–1902.
- Isawa, T., R. Sameshima, H. Mitsui, and K. Minamisawa. 1999. IS1631 occurrence in *Bradyrhizobium japonicum* highly reiterated sequence-possessing strains with high copy numbers of repeated sequences RSα and RSβ. *Appl. Environ. Microbiol.* **65**:3493–3501.
- Kaneko, T., Y. Nakamura, S. Sato, K. Minamisawa, T. Uchiumi, S. Sasamoto, A. Watanabe, K. Idesawa, M. Iriguchi, K. Kawashima, M. Kohara, M. Matsumoto, S. Shimpo, H. Tsuruoka, T. Wada, M. Yamada, and S. Tabata. 2002. Complete genomic sequence of nitrogen-fixing symbiotic bacterium *Bradyrhizobium japonicum* USDA110. *DNA Res.* **9**:189–197.
- Lashof, D. A., and D. R. Ahuja. 1990. Relative contribution of greenhouse gas emission to global warming. *Nature* **344**:529–531.
- Leonard, L. T. 1943. A simple assembly for use in the testing of cultures of rhizobia. *J. Bacteriol.* **45**:523–525.
- Li, X., K. Inubushi, and K. Sakamoto. 2002. Nitrous oxide concentration in an Andisol profile and emissions to the atmosphere as influenced by the application of nitrogen fertilizers and manure. *Biol. Fertil. Soils* **35**:108–113.
- Lucinski, R., W. Polcyn, and L. Ratajczak. 2002. Nitrate reduction and nitrogen fixation in symbiotic association *Rhizobium*-legumes. *Acta Biochim. Pol.* **49**:537–546.
- Mesa, S., E. J. Bedmar, A. Chanfon, H. Hennecke, and H. Fisher. 2003. *Bradyrhizobium japonicum* NnrR, a denitrification regulator, expands the FixLJ–FixK2 regulatory cascade. *J. Bacteriol.* **185**:3978–3982.
- Mesa, S., J. de Dios Alche, E. J. Bedmar, and M. J. Delgado. 2004. Expression of *nir*, *nor* and *nos* denitrification genes from *Bradyrhizobium japonicum* in soybean root nodules. *Physiol. Plant.* **120**:205–211.
- Minamisawa, K. 1990. Division of rhizobitoxine-producing and hydrogen-uptake positive strains of *Bradyrhizobium japonicum* by *nifDKE* sequence. *Plant Cell Physiol.* **31**:81–89.

31. Minamisawa, K., M. Itakura, M. Suzuki, K. Ichige, T. Isawa, K. Yuhashi, and H. Mitsui. 2002. Horizontal transfer of nodulation genes in soils and microcosms from *Bradyrhizobium japonicum* to *B. elkanii*. *Microbes Environ.* **17**:82–90.
32. Minamisawa, K., Y. Nakatsuka, and T. Isawa. 1999. Diversity and field site variation of indigenous populations of soybean bradyrhizobia in Japan by fingerprints with repeated sequences RS α and RS β . *FEMS Microbiol. Ecol.* **29**:171–178.
33. Miyamoto, T., M. Kawahara, and K. Minamisawa. 2004. Novel endophytic nitrogen-fixing clostridia from the grass *Miscanthus sinensis* as revealed by terminal restriction fragment length polymorphism analysis. *Appl. Environ. Microbiol.* **70**:6580–6586.
34. Mosier, A. R., and C. Kroeze. 2000. Potential impact on the global atmospheric N₂O budget of the increased nitrogen input required to meet future global food demands. *Chemosphere Global Change Sci.* **2**:465–473.
35. Mosier, A. R., C. Kroeze, C. Nevison, O. Oenema, S. Seitzinger, and O. van Cleemput. 1998. Closing the global N₂O budget: nitrous oxide emissions through the agricultural nitrogen cycle. *Nutr. Cycl. Agroecosyst.* **52**:225–248.
36. Mori, A., M. Honjito, H. Kondo, H. Matsunami, and D. Scholefield. 2005. Effects of plant species on CH₄ and N₂O fluxes from a volcanic grassland soil in Nasu, Japan. *Soil Sci. Plant Nutr.* **51**:19–27.
37. Mozen, M. M., and R. H. Burris. 1954. The incorporation of ¹⁵N-labelled nitrous oxide by nitrogen fixing agents. *Biochim. Biophys. Acta* **14**:577–578.
38. Murphy, S. G., and G. H. Elkan. 1965. Nitrogen metabolism of some strains of *R. japonicum* having different nodulating capabilities. *Can. J. Microbiol.* **11**:1039–1041.
39. Okazaki, S., M. Sugawara, and K. Minamisawa. 2004. *Bradyrhizobium elkanii rtxC* gene is required for expression of symbiotic phenotypes in the final step of rhizobitoxine biosynthesis. *Appl. Environ. Microbiol.* **70**:535–541.
40. Prentki, P., and H. M. Krisch. 1984. In vitro insertional mutagenesis with a selectable DNA fragment. *Gene* **29**:303–313.
41. Rigaud, J., F. J. Bergersen, G. L. Turner, and R. M. Daniel. 1973. Nitrate dependent anaerobic acetylene-reduction and nitrogen-fixation by soybean bacteroids. *J. Gen. Microbiol.* **77**:137–144.
42. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
43. Sameshima, R., T. Isawa, M. J. Sadowsky, T. Hamada, H. Kosai, A. Shutsrirung, H. Mitsui, and K. Minamisawa. 2003. Phylogeny and distribution of extra-slow-growing *Bradyrhizobium japonicum* harboring high copy numbers of RS α , RS β and IS1631. *FEMS Microbiol. Ecol.* **44**:191–202.
44. Sameshima-Saito, R., K. Chiba, and K. Minamisawa. 2004. New method of denitrification analysis of *Bradyrhizobium* field isolates by gas chromatographic determination of ¹⁵N-labeled N₂. *Appl. Environ. Microbiol.* **70**:2886–2891.
45. Sawamoto, T., K. Kusa, R. Hu, and R. Hatano. 2002. Dissolved N₂O, CH₄, and CO₂ in pipe drainage, seepage, and stream water in a livestock farm in Hokkaido, Japan. *Soil Sci. Plant Nutr.* **48**:433–439.
46. Schäfer, A., A. Tauch, W. Jäger, J. Kalinowski, G. Thierbach, and A. Pühler. 1994. Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* **145**:69–73.
47. Simon, R., U. Priefer, and A. Pühler. 1983. A broad host range mobilization system for *in vivo* genetic engineering: transposon mutagenesis in Gram negative bacteria. *Bio/Technology* **1**:784–791.
48. Trung, B. C., and S. Yoshida. 1983. Improvement of Leonard jar assembly for screening of effective *Rhizobium*. *Soil Sci. Plant Nutr.* **29**:97–100.
49. van Berkum, P., and H. H. Keyser. 1985. Anaerobic growth and denitrification among different serogroups of soybean rhizobia. *Appl. Environ. Microbiol.* **49**:772–777.
50. Velasco, L., S. Mesa, C. Xu, M. J. Delgado, and E. J. Bedmar. 2004. Molecular characterization of *nosRZDFLX* genes coding for denitrifying nitrous oxide reductase of *Bradyrhizobium japonicum*. *Antonie Leeuwenhoek* **85**:229–235.
51. Yang, L., and Z. Cai. 2005. The effect of growing soybean (*Glycine max.* L.) on N₂O emission from soil. *Soil Biol. Biochem.* **37**:1205–1209.
52. Yasuta, T., S. Okazaki, H. Mitsui, K. Yuhashi, H. Ezura, and K. Minamisawa. 2001. DNA sequence and mutational analysis of rhizobitoxine biosynthesis genes in *Bradyrhizobium elkanii*. *Appl. Environ. Microbiol.* **67**:4999–5009.
53. You, M., T. Nishiguchi, A. Saito, T. Isawa, H. Mitsui, and K. Minamisawa. 2005. Expression of the *nifH* gene of a *Herbaspirillum* endophyte in wild rice species: daily rhythm during the light-dark cycle. *Appl. Environ. Microbiol.* **71**:8183–8190.
54. Zumft, W. G. 1997. Cell biology and molecular basis of denitrification. *Microbiol. Mol. Biol. Rev.* **61**:533–616.