

# The *Dlx* Gene Complement of the Leopard Shark, *Triakis semifasciata*, Resembles That of Mammals: Implications for Genomic and Morphological Evolution of Jawed Vertebrates

David W. Stock<sup>1</sup>

Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, Colorado 80309-0334

Manuscript received May 31, 2004

Accepted for publication October 29, 2004

## ABSTRACT

Extensive gene duplication is thought to have occurred in the vertebrate lineage after it diverged from cephalochordates and before the divergence of lobe- and ray-finned fishes, but the exact timing remains obscure. This timing was investigated by analysis of the *Dlx* gene family of a representative cartilaginous fish, the leopard shark, *Triakis semifasciata*. *Dlx* genes encode homeodomain transcription factors and are arranged in mammals as three convergently transcribed bigene clusters. Six *Dlx* genes were cloned from *Triakis* and shown to be orthologous to single mammalian *Dlx* genes. At least four of these are arranged in bigene clusters. Phylogenetic analyses of *Dlx* genes were used to propose an evolutionary scenario in which two genome duplications led to four *Dlx* bigene clusters in a common ancestor of jawed vertebrates, one of which was lost prior to the diversification of the group. *Dlx* genes are known to be involved in jaw development, and changes in *Dlx* gene number are mapped to the same branch of the vertebrate tree as the origin of jaws.

COMPARISONS of vertebrate and invertebrate gene families have provided evidence for large-scale gene duplication in the vertebrate lineage after it diverged from its extant sister group, the cephalochordates (HOLLAND *et al.* 1994; SIDOW 1996; SPRING 1997; FURLONG and HOLLAND 2002; PANOPOULOU *et al.* 2003). One of the most extensively studied of these gene families is the Hox family of homeodomain transcription factors (PENDLETON *et al.* 1993; RUDDLE *et al.* 1994a; PRINCE 2002; WAGNER *et al.* 2003). The 39 Hox genes of mammals are arranged as four clusters of closely linked genes, each located on separate chromosomes. In contrast, the cephalochordate amphioxus has 14 Hox genes linked in a single cluster (FERRIER *et al.* 2000). The existence of a single Hox cluster in more distantly related invertebrate outgroups and the presence of multiple Hox clusters in actinopterygian (ray-finned) fishes suggest that a single Hox cluster gave rise to four prior to the divergence of actinopterygians and sarcopterygians (lobe-finned fishes and tetrapods; AMORES *et al.* 1998; FINNERTY and MARTINDALE 1998; BALAVOINE *et al.* 2002; PRINCE 2002; WAGNER *et al.* 2003). Additional Hox cluster duplications are likely to have occurred in the actinopterygian lineage after it diverged from sarcopterygi-

ans (AMORES *et al.* 1998, 2004; PRINCE 2002; WAGNER *et al.* 2003; CHIU *et al.* 2004).

Less information is available for assessing whether four Hox clusters were present prior to the divergence of osteichthyans (actinopterygians and sarcopterygians) from chondrichthyans (cartilaginous fishes) or the divergence of lampreys and hagfishes from gnathostomes (jawed vertebrates, which include osteichthyans and chondrichthyans). Recent analyses of lamprey Hox genes have suggested that not all of the duplications leading to the four-cluster condition antedated the lamprey-gnathostome divergence and, furthermore, that independent Hox duplications occurred in the lamprey lineage (FORCE *et al.* 2002; IRVINE *et al.* 2002; FRIED *et al.* 2003; WAGNER *et al.* 2003). While only two Hox clusters have been described to date from a chondrichthyan (KIM *et al.* 2000), it has been speculated that the four-cluster condition characterized the ancestral gnathostome (WAGNER *et al.* 2003).

Hox cluster duplications are commonly described as the result of complete genome duplications (SIDOW 1996; FURLONG and HOLLAND 2002; LARHAMMAR *et al.* 2002), but this interpretation remains controversial (HUGHES *et al.* 2001; MARTIN 2001; HORTON *et al.* 2003; HUGHES and FRIEDMAN 2003). Much attention in this controversy has focused on the linkage of additional families of paralogous genes to the Hox clusters of vertebrates and more specifically on the question of whether their duplication history parallels that of the Hox genes (LUNDIN 1993; RUDDLE *et al.* 1994b; BAILEY *et al.* 1997; HUGHES *et al.* 2001; LARHAMMAR *et al.* 2002; HUGHES

Sequence data from this article have been deposited with the EMBL/GenBank Data Libraries under accession nos. AY738118–AY738123.

<sup>1</sup>Address for correspondence: Department of Ecology and Evolutionary Biology, N122 Ramaley, Campus Box 334, University of Colorado, Boulder, CO 80309-0334. E-mail: david.stock@colorado.edu

and FRIEDMAN 2003; LUNDIN *et al.* 2003). As pointed out by LARHAMMAR *et al.* (2002), many of the analyses of these gene families have not included a sample of vertebrate lineages adequately representing the diversity of the group. A partial exception to this generalization is the Dlx family of homeodomain transcription factors related to the *Drosophila* gene *Distal-less*.

Dlx genes have been shown to play a role in the development of a variety of vertebrate structures, including the brain, ears, nose, jaws, teeth, and limbs (PANGANIBAN and RUBENSTEIN 2002). Mammals possess six Dlx genes arranged as three convergently transcribed bigene clusters, with each cluster located on a different Hox-containing chromosome (PANGANIBAN and RUBENSTEIN 2002; SUMIYAMA *et al.* 2003). Analyses of the eight described zebrafish Dlx genes (arranged as three bigenes clusters and two unlinked single genes) have suggested a duplication history resembling that of Hox genes (STOCK *et al.* 1996; AMORES *et al.* 1998; NEIDERT *et al.* 2001; SUMIYAMA *et al.* 2003). Specifically, duplication of an ancestral bigene cluster led to four clusters in a common ancestor of osteichthyans, followed by loss of a cluster before the divergence of actinopterygians and sarcopterygians. These events were followed by independent duplication and losses of Dlx genes in the actinopterygian lineage. As in the Hox gene family, it has been suggested that independent Dlx duplications have occurred in the lamprey lineage (NEIDERT *et al.* 2001).

No data have been available to indicate whether four Dlx bigene clusters arose before the divergence of the main lineages of gnathostomes. Such data are of interest for a number of reasons. Jaws, the possession of which gives gnathostomes their name, are believed to be patterned by Dlx genes (DEPEW *et al.* 2002) and expression differences exist between lamprey and tetrapod Dlx genes (NEIDERT *et al.* 2001; SHIGETANI *et al.* 2002). The possibility that Dlx genes were involved in the evolution of gnathostome jaws raises the question of whether gene duplication events were important in addition to changes in expression. The linkage of Hox and Dlx genes in osteichthyans allows interpretation of an identical duplication history of these genes as evidence for duplication of at least large chromosomal regions. In addition, information on Dlx genes of chondrichthyans may aid the interpretation of the incompletely characterized Hox clusters of this group. Finally, if it is assumed that Hox and Dlx duplications are the result of whole-genome duplications, information on the Dlx genes of chondrichthyans would provide insight on the organization of the genome of the common ancestor of gnathostomes.

This study seeks to address the above issues through a characterization of the Dlx gene complement of a representative chondrichthyan, the leopard shark, *Triakis semifasciata*. Two main alternative hypotheses tested are (1) that the four bigene-cluster-condition postdated

the divergence of osteichthyans and chondrichthyans and (2) that the four-cluster condition antedated this divergence. The first scenario predicts that shark genes will be orthologous to multiple osteichthyan genes, while the second predicts that shark genes will be orthologous to single osteichthyan genes. Under either scenario, sharks may have lost orthologs of gnathostome genes or undergone independent gene duplication. The cloning and analyses of *Triakis* Dlx genes presented herein are interpreted as support for the origin of four bigene clusters and the subsequent loss of one of these prior to the divergence of the main gnathostome lineages. No evidence was found for additional Dlx duplications or losses in the chondrichthyan lineage. The acquisition of the three-cluster condition of sharks and mammals therefore maps to the same branch of the phylogenetic tree of extant vertebrates as the origin of jaws.

## MATERIALS AND METHODS

**Specimens:** Gravid females of the Leopard shark, *T. semifasciata*, were provided by fisherman participating in the Pajaro Valley Rod and Gun Club Shark Derby of July 10, 1994. The specimens were captured in Elkhorn Slough, Monterey County, California. Embryos were collected from the females by dissection and were dechorionated, separated from the yolk, and staged according to the series constructed by BALLARD *et al.* (1993) for *Scyliorhinus canicula*, a member of the same order. The embryos were then frozen on dry ice and stored at  $-70^{\circ}$ .

**RNA isolation:** A single embryo 30 mm in total length (approximate stage 30) was ground to a fine powder in liquid  $N_2$  using a mortar and pestle. Total cellular RNA was extracted from the powder using the RNazol B reagent (Biotecx Laboratories, Houston) according to the manufacturer's instructions.

**Reverse transcriptase-mediated PCR of Dlx homeoboxes:** Reverse transcription of total RNA was carried out at  $42^{\circ}$  using an RNase H<sup>-</sup> Moloney murine leukemia virus reverse transcriptase (Superscript II; Invitrogen, Carlsbad, CA) and random hexamer primers. Single-stranded cDNA from this reaction was used as the template for PCR amplification with degenerate Dlx primers described by STOCK *et al.* (1996). These included the primers 5'-GCCGGGATCCCAARCCNMG NACNATHAYTC-3' (where the underlined sequence represents a restriction site added to the 5'-end) and 5'-TTYT GRAACCADATYTTNAC-3', which bind codons near the 5'- and 3'-ends, respectively, of the homeoboxes of all previously known Dlx genes. The additional primers used (5'-GCCGGGATCCCGNAARAARATHMGNAARCC-3' and 5'-GCCGGGATCCATNGTNAAYGGNAARCCNAA-3') bind to regions conserved in a subset of vertebrate Dlx genes and located just upstream of the homeobox. The thermocycling profile used was 2 min of denaturation at  $94^{\circ}$  followed by 40 cycles of 1 min at  $94^{\circ}$ , 1 min at  $45^{\circ}$ , and 1 min at  $72^{\circ}$ .

**PCR amplification of 5' and 3' cDNA ends:** To obtain complete cDNA sequences corresponding to the reverse transcriptase-mediated PCR (RT-PCR) products described above, the homeobox sequences determined were used to design gene-specific primers for the rapid amplification of cDNA ends (RACE) procedure (FROHMAN 1990). 3'-RACE was carried out on poly(A)<sup>+</sup> RNA using the Marathon RACE method (CHENCHIK *et al.* 1996) and 5'-RACE was carried out on total RNA by the SMART RACE method (CHENCHIK *et al.* 1998).

In both cases, the RNA came from the same individual used for RT-PCR and the protocol followed that provided by the manufacturer (BD CLONTECH, Palo Alto, CA). RACE products corresponding to the gene of interest were identified by Southern blotting with oligonucleotide probes or by restriction enzyme digestion.

**PCR amplification of exons and introns from genomic DNA:**

The exon/intron structure of Dlx genes has been described for human (NAKAMURA *et al.* 1996; SUMIYAMA *et al.* 2002), mouse (MCGUINNESS *et al.* 1996; NAKAMURA *et al.* 1996; LIU *et al.* 1997; SUMIYAMA *et al.* 2002), and zebrafish (ELLIES *et al.* 1997). Intron location is conserved in all described cases, with one intron 5' of the homeobox and a second intron within. To determine whether this genomic structure is also conserved with Dlx genes of Triakis, PCR primers were designed to amplify from genomic DNA the entire coding region of each gene in two overlapping fragments. These two fragments extended from the 5'-UTR to the predicted second exon and from this exon to the 3'-UTR. This strategy identifies all introns located within the coding region of the genes, but cannot exclude the possibility of introns at the extreme ends of the 5'- and 3'-UTRs.

Genomic DNA (extracted from individual, unstaged embryos by overnight proteinase K digestion at 55° followed by phenol/chloroform extraction and isopropanol precipitation) was used as a template for long PCR (BARNES 1994). These PCR reactions employed the Expand DNA polymerase mixture and associated buffer (Roche, Indianapolis) and the TaqStart Antibody (CLONTECH), with the latter allowing a "hot start" (KELLOGG *et al.* 1994). Thermocycling consisted of 2 min of denaturation at 94°, followed by 35 cycles of 30 sec at 94° and 5 min at 68°. PCR products were subcloned and intron positions were determined by sequencing. The sizes of the introns were estimated by a combination of partial sequencing and agarose gel electrophoresis of PCR products.

**Genomic long PCR:** The presence of Dlx bigene clusters in Triakis was investigated by long PCR, using conditions as described above (with the exception that the annealing/extension step at 68° was increased to 10 or 15 min). PCR products were characterized by restriction mapping, subcloning, and partial sequencing.

**Cloning and sequencing:** PCR products were subcloned into pBluescript II SK+ (Stratagene, La Jolla, CA) or pCR4-TOPO (Invitrogen) by standard procedures (SAMBROOK *et al.* 1989). Single- or double-stranded plasmid DNA was subjected to automated sequencing. With the exception of a few nucleotides in the 5'-untranslated regions, each nucleotide of the six Triakis Dlx cDNAs was determined from at least five different clones, representing each strand at least twice. Sequences were manipulated and translated using programs in the DNASTAR package (DNASTar, Madison, WI).

**Phylogenetic analyses:** Amino acid sequences of *Distal-less*-related genes were obtained from the GenBank database with the following accession numbers: nematode (*Caenorhabditis elegans*) C28A5.4 (Z32680); fruit fly (*Drosophila melanogaster*) Dll (S47947); acorn worm (*Ptychodera flava*) Pf-Dlx (AB028221); tunicate (*Ciona intestinalis*) Dll-A (AJ278696), Dll-B (AJ278697), and Dll-C (AJ278698); amphioxus (*Branchiostoma floridae*) *AmphiDll* (U47058); lamprey (*Petromyzon marinus*) DlxA (AY010116), DlxB (AY010117), DlxC (AY010118), and DlxD (AY010119); lamprey (*Lethenteron japonicum*) LjDLX1/6 (AB048759); zebrafish (*Danio rerio*) *dlx1* (U67842), *dlx2* (U03875), *dlx3* (X65060), *dlx4* (U03876), *dlx5* (U67843), *dlx6* (U67844), *dlx7* (U67845), and *dlx8* (U67846); frog (*Xenopus laevis*) *Xdll* (D10259), *X-DLL1* (A56570), *Xdll-2* (S74210), *X-dll2* (L09730), *X-dll3* (L09729), and *X-dll4* (L09728); newt (*Notophthalmus viridescens*) *NvHBox-4* (X63531) and *NvHBox-5* (X63532); chicken (*Gallus gallus*) *DLX3* (AJ243432) and *Dlx5* (U25274);

mouse (*Mus musculus*) *Dlx1* (NM\_010053), *Dlx2* (NM\_010054), *Dlx3* (NM\_010055), *Dlx5* (U67840), and *Dlx7* (AF452637); and human (*Homo sapiens*) *DLX1* (XM\_087198), *DLX2* (NM\_004405), *DLX3* (NM\_005220), *DLX5* (NM\_005221), *DLX6* (NM\_005222), and *DLX7* (AF452638). The nomenclature of human, mouse, and zebrafish Dlx genes was adjusted from the above GenBank records where necessary to follow PANGANIBAN and RUBENSTEIN (2002). Related genes outside of the Distal-less family were obtained for use as outgroups (STOCK *et al.* 1996). These sequences were mouse *Barx1* (Y07960) and *Msx1* (NM\_010835) and *D. melanogaster* *Msh* (U33319).

Upon inspection of initial alignments, a few of the above sequences were interpreted as containing errors or truncations of their 5'-ends. Adjustments to these sequences were made as follows. The lamprey sequences *DlxD* (*P. marinus*) and *LjDLX1/6* (*L. japonicum*) are clearly orthologous on the basis of a high degree of nucleotide identity. However, comparison of these genes with each other and with other vertebrate Dlx genes suggests that *DlxD* is truncated at its 5'-end, while *LjDLX1/6* has a frameshift error in the region 3' of the homeobox (which itself has a six-nucleotide deletion found in no other animals). A composite lamprey *DlxD* sequence was therefore constructed with the missing amino terminus of *P. marinus* *DlxD* replaced by the corresponding region from *L. japonicum* *LjDLX1/6*. The reported human *DLX6* sequence similarly appears to have an internal methionine designated as the start codon. Translation of the genomic sequence of a BAC clone (accession AC004774) revealed the likely true start codon in frame and part of the same exon; this region was added to the reported amino acid sequence. *X. laevis* *Xdll-2* was reported to have a truncated 5'-end. The missing nucleotides were added to this sequence from an *X. laevis* EST that appeared to be derived from the same gene (accession BF613971). Finally, a single amino acid adjustment was made to the homeodomain of zebrafish *dlx2a* on the basis of unpublished sequence data, which exhibited a better match to all other sequenced animal *Distal-less*-related genes. After these modifications, the only incomplete coding region remaining among the chordate sequences analyzed was that of *P. marinus* *DlxB* (reported as such by NEIDERT *et al.* 2001).

The 61-amino-acid homeodomain of all Dll/Dlx genes may be aligned reliably without gaps. However, alignment outside of the homeodomain of the most distantly related genes is problematic (STOCK *et al.* 1996; NEIDERT *et al.* 2001). To identify the location of the root of the tree of vertebrate Dlx genes, an initial phylogenetic analysis of homeodomain sequences was attempted. Pairwise distances between amino acid sequences were calculated as percentage of similarities [*P*-values, as recommended by ZHANG and NEI (1996) in their analysis of Hox gene homeodomains] and used to construct neighbor-joining trees with the program MEGA, version 2.1 (KUMAR *et al.* 2001). To reduce the number of identical sequences in the analyses, gnathostome sequences were restricted to those of human, zebrafish, and Triakis.

To provide increased resolution of the phylogeny of vertebrate Dlx sequences, an alignment of complete vertebrate Dlx sequences was constructed using Clustal X (THOMPSON *et al.* 1997). The profile alignment feature of this program was used to align sequences in clades identified from the analysis of homeodomain sequences, followed by sequential alignment of these subalignments into a global alignment. Adjustments to alignments at all stages of the process were made either by changing alignment parameters for specific regions or by manual editing. The alignment is available as supplementary material at <http://www.genetics.org/supplemental/>.

Maximum likelihood and neighbor-joining analyses of complete Dlx protein sequences were conducted with the PROML,




FIGURE 1.—Genomic organization of Triakis Dlx genes. Phylogenetically related genes are aligned in columns. Boxes indicate protein-coding portions of exons (homeodomain in black), angled lines indicate introns, and horizontal lines indicate intergenic regions. Transcriptional orientation is denoted by arrows. All distances between the start and stop codons of an individual gene are drawn to scale, while intergenic distances (listed) are not. Question mark indicates lack of data from genomic PCR to support or refute linkage predicted from phylogenetic analyses.

PROTDIST, and NEIGHBOR programs of the PHYLIP package (version 3.6; FELSENSTEIN 1989). Both types of analyses employed the Jones-Taylor-Thornton model of amino acid change (JONES *et al.* 1992) and a gamma distribution of rates among sites (coefficient of variation is 0.707, corresponding to an  $\alpha$ -value of 2.0, as recommended by KUMAR *et al.* 2001). For maximum likelihood analysis, six rate categories, 40 random addition sequences, and global rearrangements were used.

## RESULTS

**Six Dlx genes detected in *T. semifasciata*:** Sixty-five cloned Dlx homeobox fragments obtained by RT-PCR were sequenced. These fragments arose from two independent reactions performed with each of three different primer combinations (using a common antisense primer). The sequences obtained fell into six classes and were used to design primers for 5'- and 3'-RACE PCR reactions. Cloning and sequencing of the latter PCR products resulted in complete cDNA sequences that were divergent enough to confirm that the six classes represented different loci rather than multiple alleles. The loci were named according to their orthology to other vertebrate Dlx genes (PANGANIBAN and RUBENSTEIN 2002), as inferred from the phylogenetic analyses described below. These loci and the number of homeobox clones obtained by RT-PCR are *Dlx1* (5), *Dlx2* (2), *Dlx3* (35), *Dlx4* (10), *Dlx5* (5), and *Dlx6* (8).

**Genomic organization of Triakis Dlx genes:** The exon/intron structures and linkage arrangements of the six *T. fasciata* Dlx genes are shown in Figure 1. All six genes possess two introns and three exons. The second intron occupies an identical location in the homeobox of all genes and is phase 0 (between codons). The first intron is phase 1 (between the first and second nucleotides of a codon) in all genes and is located 50–68 nucleotides 5' of the homeobox. Intron 1 ranges in size from 128 bp (*Dlx2*) to  $\sim$ 1.8 kb (*Dlx3*), while intron 2 ranges in size from 221 bp (*Dlx2*) to  $\sim$ 3.9 kb (*Dlx4*).

Analysis of genomic long PCR products revealed that Triakis possesses at least two Dlx bigene clusters (*Dlx1*-

*Dlx2* and *Dlx5*-*Dlx6*), with the genes oriented in the convergently transcribed arrangement characteristic of chordate Dlx genes (Figure 1). Furthermore, the linkage of these specific orthologs (identified by phylogenetic analysis of amino acid sequences—see below) is the same as that described in other gnathostome vertebrates (PANGANIBAN and RUBENSTEIN 2002). The stop codons of *Dlx1* and *Dlx2* are separated by  $\sim$ 7.9 kb, while those of *Dlx5* and *Dlx6* are separated by  $\sim$ 10.2 kb. Despite attempts with multiple sets of primers, no genomic PCR product was obtained that included *Dlx3* and *Dlx4*, genes whose presumed orthologs form a bigene cluster in other gnathostomes.

**Phylogenetic analysis of bilaterian Dlx homeodomain sequences:** A neighbor-joining tree constructed from homeodomain sequences is shown in Figure 2. Gnathostome Dlx sequences in this tree are found in two main clades. One of these (35% bootstrap support) contains *Dlx1*, *Dlx4*, *Dlx6*, and one lamprey Dlx sequence, while the other contains *Dlx2*, *Dlx3*, *Dlx5*, and three lamprey sequences (63% bootstrap support). One tunicate Dll gene forms the sister group to the *Dlx1*/*Dlx4*/*Dlx6* clade, while the single reported Dll/Dlx genes of *Drosophila*, the hemichordate *Ptychodera*, and the cephalochordate amphioxus, as well as one tunicate Dll gene form a clade outside of both gnathostome-containing Dlx clades. An additional tunicate Dll gene and the single reported nematode (*C. elegans*) Dll/Dlx gene group outside of other animal Dll/Dlx genes.

This analysis of homeodomain sequences provides little phylogenetic resolution (as judged by low bootstrap values). Nevertheless, it reveals that Triakis possesses three members of each of the main gnathostome Dlx clades and suggests that the root of the vertebrate Dlx tree lies on the internal branch connecting these clades.

**Phylogenetic analysis of complete vertebrate Dlx sequences:** The phylogenetic analyses of Dlx genes reported by STOCK *et al.* (1996) and NEIDERT *et al.* (2001) provided evidence for the existence of six clades of




FIGURE 2.—Neighbor-joining analysis of Dll/Dlx homeodomain sequences. Numbers indicate the percentage of 1000 bootstrap replicates in which the indicated node was present. The tree is rooted at the midpoint of the longest path between two taxa. Brackets indicate the two main clades of vertebrate Dlx genes named after the mammalian members.

osteichthyan Dlx genes, each possessing a single mammalian Dlx gene and one or two zebrafish genes. The results of maximum likelihood and neighbor-joining analysis of complete vertebrate Dlx genes shown in Figure 3 indicate that each of these clades also contains a single Triakis gene. The monophyly of five of these six clades is supported by bootstrap values of at least 99% with one or both methods of analysis. Somewhat lower bootstrap values for the monophyly of the Dlx4 clade (74 and 76%) may be due to the rapid evolution of mammalian Dlx4 apparent in the tree in Figure 3.

If the six Dlx genes isolated from Triakis are in fact the orthologs of the osteichthyan genes, then they would be expected to exhibit a sister group relationship to all the other members of a given clade. This relationship is found for all clades other than Dlx2 in which the expected relationship is contradicted by an internal branch with relatively weak bootstrap support (64%; Figure 3).

The phylogenetic analysis in Figure 3 reveals 100% bootstrap support for an internal branch separating lamprey *DlxA*, *DlxB*, and *DlxC* from lamprey *DlxD*. It is likely that this branch also contains the root of the tree (on the basis of midpoint rooting in Figure 3 and

outgroup analysis in Figure 2). If this is the true location of the root of the tree, then *DlxA*, *DlxB*, and *DlxC* are members of the vertebrate Dlx2/Dlx3/Dlx5 clade and *DlxD* is a member of the Dlx1/Dlx4/Dlx6 clade. The analysis shown in Figure 3 provides little support for the phylogenetic position of the lamprey genes within these major clades, with the exception of strong bootstrap support (93%) for a sister group relationship between lamprey *DlxA* and *DlxC*.

## DISCUSSION

**Orthology of the Triakis Dlx genes:** The number of Dlx genes isolated from Triakis is the same as that reported from mammals and all are supported as members of clades containing a single human gene (Figure 3). It has been argued that the six-Dlx-gene condition of the osteichthyan common ancestor arose by the loss of a Dlx bigene cluster linked to the HoxC complex (STOCK *et al.* 1996; NEIDERT *et al.* 2001; SUMIYAMA *et al.* 2003). Because of the phylogenetic position of chondrichthyans as the sister group of all other gnathostomes in the analyses, it is not possible to rule out the orthology of one or two of the Dlx genes of Triakis to such HoxC-




FIGURE 3.—Maximum likelihood and neighbor-joining analysis of vertebrate *Dlx* genes. Tree topology and branch lengths are from a maximum likelihood analysis, with midpoint rooting. Nodes with single numbers indicate the results of a bootstrap maximum likelihood analysis with 100 replicates and options as described in MATERIALS AND METHODS, except for the use of two random addition sequences for each replicate. Where two numbers appear, the upper is from the bootstrap maximum likelihood analysis and the lower is from a bootstrap neighbor-joining analysis with 1000 replications. Brackets indicate the six clades of jawed vertebrate *Dlx* genes, each named after and containing a single Triakis gene.

linked *Dlx* genes. The failure to amplify a PCR product containing Triakis *Dlx3* and *Dlx4* is in fact consistent with the possibility that one of these genes is orthologous to a *Dlx* gene lost in the ancestry of gnathostomes. This result does not constitute positive evidence for the non-orthology of one of these genes to osteichthyan *Dlx* genes, however. Not all primer combinations attempted amplified the other bigene clusters and it is possible that *Dlx3* and *Dlx4* are too far apart to be reliably amplified by the PCR methods employed. Given the high degree of divergence between osteichthyan *Dlx* clades relative to that within, orthology of a Triakis *Dlx* gene to one lost from osteichthyans would be expected to be associated with extensive divergence of the Triakis sequence from its closest relatives in the tree. No clear evidence exists for this (Figure 3), and it is concluded on the basis of parsimony that each Triakis *Dlx* gene is orthologous to

the members of the osteichthyan *Dlx* clade of the same name.

**Conservation of *Dlx* intron structure:** Comparison of the locations of introns in Triakis *Dlx* genes with those of zebrafish (*dlx1a*, *dlx2a*, *dlx3b*, *dlx5a*, and *dlx6a*; ELLIES *et al.* 1997), mouse (*Dlx1*, *Dlx2*, *Dlx3*, and *Dlx4*; MCGUINNNESS *et al.* 1996; SUMIYAMA *et al.* 2002), and human (*DLX3*, *DLX4*, *DLX5*, and *DLX6*; SUMIYAMA *et al.* 2002; GenBank accession no. AC004774) revealed the exact correspondence of intron location among *Dlx* orthologs, with the exception of the first intron in zebrafish *dlx3b*, which was displaced by two amino acids from the corresponding alignment position in shark and mammal orthologs. All vertebrate *Dlx* genes exhibited an identical location for the second intron (within the homeobox), while the first intron varied in location (but not phase) by up to six alignment positions.

The location of the intron in the homeobox of vertebrate Dlx genes is identical to that of *Drosophila*. Five additional introns are present in the *Drosophila* gene (VACHON *et al.* 1992), however, and in the absence of outgroup information, it is not possible to determine whether the differences in intron number arose through loss in vertebrates or gain in *Drosophila*. Of the three reported Dll genes in the tunicate *C. intestinalis*, two have been reported to have a homeobox intron in the same location as that of vertebrates (DI GREGORIO *et al.* 1995). The structure of the region upstream of the homeobox has not been reported for any of the genes, but *Ci-DllB* has two introns 3' of the homeobox (CARACCILO *et al.* 2000). As discussed below, it is likely that Dlx bigene clusters arose by a tandem duplication before the divergence of vertebrates and tunicates. The virtually identical intron/exon structure of members of bigene clusters (more distantly related than orthologous tunicate and vertebrate Dlx genes) suggests that *Ci-DllB* gained introns after the divergence of vertebrates and tunicates. Such gain of introns over this time scale is not unusual (LYNCH and RICHARDSON 2002).

**Dlx arrangement as bigene clusters:** A convergently transcribed bigene-cluster arrangement has been described for two of the three Dll genes of the tunicate *C. intestinalis* (CARACCILO *et al.* 2000), six of the eight zebrafish Dlx genes (ELLIES *et al.* 1997; GHANEM *et al.* 2003), four Dlx genes of the pufferfishes *Takifugu rubripes* and *Spheroides nephelus* (GHANEM *et al.* 2003), and the six mammalian Dlx genes (reviewed by SUMIYAMA *et al.* 2003). The *Dlx1/Dlx2* and *Dlx5/Dlx6* bigene clusters identified for Triakis are consistent with the organization of Dlx genes in the other vertebrates examined. Furthermore, the intergenic distances (between stop codons) of 7.9 and 10.2 kb, respectively, fall within the range of those reported for ray-finned fish and mammals (5.3–12.3 kb and ~3–10.7 kb, respectively; GHANEM *et al.* 2003; SUMIYAMA *et al.* 2003). The bigene cluster for which no evidence was obtained in *T. semifasciata*, *Dlx3/Dlx4*, contains an intergenic region of ~17.6 kb in humans (SUMIYAMA *et al.* 2003). Such a distance between these genes in Triakis may explain the failure to obtain an intergenic PCR product. In the absence of evidence of Dlx genomic organization in the lamprey, the existence of Dlx bigene clusters in Triakis provides further support for the hypothesis that this arrangement is primitive for chordates (STOCK *et al.* 1996; NEIDERT *et al.* 2001; SUMIYAMA *et al.* 2003).

**Evolutionary history of vertebrate Dlx genes:** A scenario for the evolution of chordate Dlx genes is shown in Figure 4. The closer relationship of Dlx genes among rather than within bigene clusters provides the evidence for the postulated initial tandem duplication of Dlx genes shown on branch 1. That this event antedates the divergence of vertebrates and tunicates is hypothesized on the basis of the presence of bigene clusters in both groups. It should be noted, however, that phylogenetic




FIGURE 4.—Scenario for the evolution of Dlx genes showing selected deuterostome taxa. Boxes indicate individual Dlx genes arranged in columns according to phylogenetic relationship. Horizontal lines joining boxes indicate demonstrated linkage relationships. See text for details of events postulated to have occurred along numbered branches.

analysis of homeodomain sequences (Figure 2) did not yield the predicted clustering of all tunicate Dll genes as members of one or the other of the two main clades of vertebrate Dlx genes. Such a relationship was found for *Dll-C*, but was weakly supported, while the linked *Dll-A* and *Dll-B* genes clustered outside of all vertebrate Dlx genes. Rather than postulating the complex pattern of gain and loss implied by a literal interpretation of the tree in Figure 2, the scenario in Figure 4 follows STOCK *et al.* (1996) in interpreting the bigene clusters of tunicates and vertebrates as unlikely to have evolved independently.

Placing the initial tandem duplication on branch 1 of Figure 4 implies that the single Dll gene isolated from amphioxus was ancestrally (if not still) a member of a bigene cluster. The position of the amphioxus sequence in Figure 2 does not support this assertion, but the expected clustering in one of the two major clades of vertebrate Dlx genes is not strongly contradicted. Whether the initial tandem duplication occurred before protostomes and/or hemichordates diverged from tunicates and vertebrates remains equivocal; *Drosophila* and hemichordate sequences cluster outside of all vertebrate Dlx genes but are members of a clade containing amphioxus and tunicate sequences. A literal interpretation of the tree would require loss of a Dll-like gene from vertebrates and loss of a Dlx-like gene from protostomes, hemichordates, and amphioxus.

The events on branch 2 of Figure 4 that led to three tunicate Dll genes after an initial tandem duplication remain unclear because of the contradiction between the results of phylogenetic analyses and gene linkage described above. The scenario in Figure 4 postulates

independent duplications on branches 2 (tunicate lineage) and 3 (vertebrate lineage) on the basis of the common view that duplication of Hox clusters (linked to Dlx genes) all occurred after the divergence of amphioxus from vertebrates (HOLLAND *et al.* 1994; SIDOW 1996; AMORES *et al.* 1998; FINNERTY and MARTINDALE 1998; FERRIER *et al.* 2000; FORCE *et al.* 2002; FURLONG and HOLLAND 2002; IRVINE *et al.* 2002; PANOPOULOU *et al.* 2003; WAGNER *et al.* 2003).

The general pattern of Dlx bigene cluster duplication from one (branch 1) to two (branch 3) to four (branch 5) shown in Figure 4 is based on the hypothesis that the four Hox clusters of mammals arose by two rounds of genome duplication (SIDOW 1996; FURLONG and HOLLAND 2002; LARHAMMAR *et al.* 2002). It is important to note that this hypothesis remains controversial (HUGHES *et al.* 2001; MARTIN 2001; HORTON *et al.* 2003; HUGHES and FRIEDMAN 2003) and that Dlx genes cannot in themselves refute or support such a means of duplication because only three bigene clusters are found in jawed vertebrates that have not undergone relatively recent genome duplication. The exact placement of genome duplications shown in Figure 4 (one on branch 3 before lampreys diverged from gnathostomes and one on branch 5 afterward) is based on the analyses of Hox genes by FORCE *et al.* (2002) and FRIED *et al.* (2003) and remains tentative because of lack of phylogenetic resolution of lamprey Dlx sequences (this study and NEIDERT *et al.* 2001). Independent Dlx duplication in the lamprey lineage relative to that in gnathostomes is required by the scenario in Figure 4 and is supported by clustering of lamprey *DlxA* and *DlxC* in Figure 3 and in the analysis of NEIDERT *et al.* (2001).

The most significant finding of this study bearing on the evolutionary history of vertebrate Dlx genes is evidence that the common ancestor of extant gnathostomes had three Dlx bigene clusters (branch 5 of Figure 4). This is suggested by the analyses described above that identify all six Triakis Dlx genes as orthologous to specific osteichthyan Dlx genes. That this condition evolved by the loss of a bigene cluster is postulated on the basis of the presence of such clusters on only three of the four mammalian Hox-containing chromosomes (STOCK *et al.* 1996; SUMIYAMA *et al.* 2003). While the relationship of lamprey Dlx genes shown in Figure 3 differs somewhat from that obtained in the analysis of NEIDERT *et al.* (2001), both suggest that the acquisition of four bigene clusters and loss of one occurred after gnathostomes diverged from lampreys.

No evidence was found for independent Dlx gene duplication in the lineage leading to Triakis (branch 6 of Figure 4), in contrast to those leading to the lamprey (branch 4, discussed above), zebrafish, and *X. laevis*. The latter two species (separate derivatives of branch 7, not shown in Figure 4) are thought to have undergone genome duplications after diverging from the other vertebrates examined (KOBEL and DU PASQUIER 1986; AMORES *et al.* 1998, 2004; FRIED *et al.* 2003) and these

duplications have been postulated to have given rise to additional Dlx genes (STOCK *et al.* 1996; AMORES *et al.* 1998). Because of potential biases in the ability of degenerate PCR primers to amplify multiple Dlx genes and the fact that cDNA-based amplification is limited to genes expressed at the embryonic stage examined, it is not possible to rule out the presence of additional, uncharacterized Dlx in the Triakis genome, however.

#### **Dlx genes and the origin of gnathostome characters:**

The earliest divergence among extant jawed vertebrates separates chondrichthyans from osteichthyans (MAISEY 1986; JANVIER 1996, 2001). As argued above, the presence of three Dlx bigene clusters characterized the common ancestor of both lineages and arose by duplication and loss after the divergence of lampreys and gnathostomes (branch 5 of Figure 4). This same internal branch, the stem gnathostome lineage (JANVIER 2001), was undoubtedly the location of the origin of jaws, the defining feature of gnathostomes. Dlx genes have been shown to be involved in patterning the jaws of mice (DEPEW *et al.* 2002; PANGANIBAN and RUBENSTEIN 2002), and the expression patterns of Dlx genes differ between the oral regions of lampreys and gnathostomes (MYOJIN *et al.* 2001; NEIDERT *et al.* 2001; COHN 2002; SHIGETANI *et al.* 2002). In addition to changes in the expression of existing Dlx genes playing a role in the origin of jaws, the scenario for Dlx evolution illustrated in Figure 4 raises the possibility that Dlx gene duplication (and loss) were involved as well.

JANVIER (2001) lists 22 characters in addition to jaws that are unique to gnathostomes among living vertebrates and that are likely to be preserved in fossils. Determining whether these originated along the stem gnathostome lineage (and hence may have been related to the acquisition of three Dlx bigene clusters) rather than representing reversals in extant jawless vertebrates depends critically on the relationship of fossil jawless fishes ("ostracoderms") branch from the stem gnathostome lineage rather than clustering with extant jawless fishes (JANVIER 1996, 2001; DONOGHUE *et al.* 2000) increase the number of characters that may have arisen after the three Dlx bigene-cluster condition. Those whose development requires Dlx genes include paired nasal openings, three semicircular canals, teeth, and pectoral fins (see JANVIER 2001 and DONOGHUE *et al.* 2000 for character evolution and PANGANIBAN and RUBENSTEIN 2002 for a review of Dlx function).

#### **The structure of the ancestral gnathostome genome:**

Recent proponents of the hypothesis of two rounds of genome duplication in vertebrates have argued that both occurred before the divergence of actinopterygians and sarcopterygians (AMORES *et al.* 1998, 2004; WAGNER *et al.* 2003). The presence of three Dlx bigene clusters in Triakis and mammals is consistent with both having also occurred before the divergence of extant gnathostome taxa. While it has been proposed that the ancestral gnathostome had four Hox clusters (AMORES



*et al.* 1998, 2004; WAGNER *et al.* 2003), only two Hox clusters have been reported from chondrichthyans to date (KIM *et al.* 2000). The linkage of Dlx genes to Hox genes in mammals and zebrafish (AMORES *et al.* 1998; SUMIYAMA *et al.* 2003), argued to be a common feature of the bilaterian ancestor (POLLARD and HOLLAND 2000), suggests that at least three Hox clusters were present in chondrichthyan ancestors, and the hypothesis of multiple Hox and Dlx clusters arising by whole-genome duplication suggests that there were four. Consistent with this prediction, the two identified chondrichthyan Hox clusters are each orthologous to single osteichthyan clusters (KIM *et al.* 2000; WAGNER *et al.* 2003).

If Dlx bigene clusters arose by whole-genome duplication, other families of osteichthyan gene duplicates with single orthologs in invertebrates should have multiple orthologs in chondrichthyans. Although chondrichthyan sequences are underrepresented in studies of vertebrate gene and genome duplication, evidence does exist from a number of gene families supporting this prediction. Early studies of gene number by isozyme electrophoresis suggested gene duplication after the divergence of vertebrates and invertebrates and before the divergence of osteichthyans and chondrichthyans for the lactate dehydrogenase (LDH), malate dehydrogenase, creatine kinase, and glucose phosphate isomerase gene families (MARKERT *et al.* 1975; FISHER *et al.* 1980). Similar duplication histories have been proposed for several Wnt paralogs (SIDOW 1992), proteasome Z subunits (KASAHARA *et al.* 1996), LDH genes [the sequence analysis of STOCK *et al.* (1997) and STOCK and POWERS (1998) confirmed the results of early isozyme studies], dioxin receptors (HAHN *et al.* 1997), enolases (TRACY and HEDGES 2000), Otx genes (GERMOT *et al.* 2001), Tbx genes (TANAKA *et al.* 2002), Emx genes (DEROBERTE *et al.* 2002), thyroid hormone receptors (ESCRIVA *et al.* 2002), dopamine receptors (LE CROM *et al.* 2003), and neuropeptide Y receptors (SALANECK *et al.* 2003). The Dlx gene data presented in this study are therefore consistent with data from other gene families in suggesting that the distinctive morphological differences between osteichthyans and chondrichthyans arose in organisms with a similar overall gene content.

The author thanks Gregor Caillet, Giacomo Bernardi, Stephanie Clendennen, and anonymous participants in the Elkhorn Slough Shark Derby for assistance in obtaining *Triakis* specimens; Kenneth Weiss for support during the early stages of this project; R. Travis Merritt and Anne Buchanan for technical assistance; and William Jackman, Jason Pardo, Josh Trapani, Sarah Wise, and two reviewers for helpful comments. Funding was provided by National Science Foundation grants SBR 9408402 to K. Weiss and IBN 0092487 to the author.

#### LITERATURE CITED

- AMORES, A., A. FORCE, Y.-L. YAN, L. JOLY, C. AMEMIYA *et al.*, 1998 Zebrafish *hox* clusters and vertebrate genome evolution. *Science* **282**: 1711–1714.
- AMORES, A., T. SUZUKI, Y.-L. YAN, J. POMEROY, A. SINGER *et al.*, 2004 Developmental roles of pufferfish *Hox* clusters and genome evolution in ray-finned fish. *Genome Res.* **14**: 1–10.
- BAILEY, W. J., J. KIM, G. P. WAGNER and F. H. RUDDLE, 1997 Phylogenetic reconstruction of vertebrate Hox cluster duplications. *Mol. Biol. Evol.* **14**: 843–853.
- BALAVOINE, G., R. DE ROSA and A. ADOUTTE, 2002 Hox clusters and bilaterian phylogeny. *Mol. Phylogenet. Evol.* **24**: 366–373.
- BALLARD, W. W., J. MELLINGER and H. LECHENAULT, 1993 A series of normal stages for development of *Scyliorhinus canicula*, the Lesser spotted dogfish (*Chondrichthyes: Scyliorhinidae*). *J. Exp. Zool.* **267**: 318–336.
- BARNES, W. M., 1994 PCR amplification of up to 35-kb DNA with high fidelity and high yield from  $\lambda$  bacteriophage templates. *Proc. Natl. Acad. Sci. USA* **91**: 2216–2220.
- CARACCILO, A., A. DI GREGORIO, F. ANIELLO, R. DI LAURO and M. BRANNO, 2000 Identification and developmental expression of three *Distal-less* homeobox containing genes in the ascidian *Ciona intestinalis*. *Mech. Dev.* **99**: 173–176.
- CHENCHIK, A., F. MOQADAM and P. SIEBERT, 1996 A new method for full-length cDNA cloning by PCR, pp. 273–321 in *A Laboratory Guide to RNA: Isolation, Analysis and Synthesis*, edited by P. A. KRIEG. Wiley-Liss, New York.
- CHENCHIK, A., Y. ZHU, L. DIATCHENKO, R. LI, J. HILL *et al.*, 1998 Generation and use of high-quality cDNA from small amounts of total RNA by SMART PCR, pp. 305–319 in *Gene Cloning and Analysis by RT-PCR*, edited by P. SIEBERT and J. LARRICK. BioTechniques Press, Westborough, MA.
- CHIU, C.-H., K. DEWAR, G. P. WAGNER, K. TAKAHASHI, F. RUDDLE *et al.*, 2004 Bichir *HoxA* cluster sequence reveals surprising trends in ray-finned fish genomic evolution. *Genome Res.* **14**: 11–17.
- COHN, M. J., 2002 Lamprey *Hox* genes and the origin of jaws. *Nature* **416**: 386–387.
- DEPEW, M. J., T. LUFKIN and J. L. R. RUBENSTEIN, 2002 Specification of jaw subdivisions by *Dlx* genes. *Science* **298**: 381–385.
- DEROBERTE, Y., J. L. PLOUHINEC, T. SAUKA-SPENGLER, C. LE MENTEC, B. BARATTE *et al.*, 2002 Structure and expression of three *Emx* genes in the dogfish *Scyliorhinus canicula*: functional and evolutionary implications. *Dev. Biol.* **247**: 390–404.
- DI GREGORIO, A., A. SPAGNUOLO, F. RISTORATORE, M. PISCHETOLA, F. ANIELLO *et al.*, 1995 Cloning of ascidian homeobox genes provides evidence for a primordial chordate cluster. *Gene* **156**: 253–257.
- DONOGHUE, P. C., P. L. FOREY and R. J. ALDRIDGE, 2000 Conodont affinity and chordate phylogeny. *Biol. Rev. Camb. Philos. Soc.* **75**: 191–251.
- ELLIES, D. L., D. W. STOCK, G. HATCH, G. GIROUX, K. M. WEISS *et al.*, 1997 Relationship between the genomic organization and the overlapping embryonic expression patterns of the zebrafish *dlx* genes. *Genomics* **45**: 580–590.
- ESCRIVA, H., L. MANZON, J. YOUSON and V. LAUDET, 2002 Analysis of lamprey and hagfish genes reveals a complex history of gene duplications during early vertebrate evolution. *Mol. Biol. Evol.* **19**: 1440–1450.
- FELSENSTEIN, J., 1989 PHYLIP—Phylogeny Inference Package, Version 3.2. *Cladistics* **5**: 164–166.
- FERRIER, D. E. K., C. MINGUILLÓN, P. W. H. HOLLAND and J. GARCIA-FERNÁNDEZ, 2000 The amphioxus Hox cluster: deuterostome posterior flexibility and *Hox14*. *Evol. Dev.* **2**: 284–293.
- FINNERTY, J. R., and M. Q. MARTINDALE, 1998 The evolution of the Hox cluster: insights from outgroups. *Curr. Opin. Genet. Dev.* **8**: 681–687.
- FISHER, S. E., J. B. SHAKLEE, S. D. FERRIS and G. S. WHITT, 1980 Evolution of five multilocus isozyme systems in the chordates. *Genetica* **52/53**: 73–85.
- FORCE, A., A. AMORES and J. H. POSTLETHWAIT, 2002 *Hox* cluster organization in the jawless vertebrate *Petromyzon marinus*. *J. Exp. Zool. (Mol. Dev. Evol.)* **294**: 30–46.
- FRIED, C., S. J. PROHASKA and P. F. STADLER, 2003 Independent Hox-cluster duplications in lampreys. *J. Exp. Zool. Mol. Dev. Evol.* **299B**: 18–25.
- FROHMAN, M. A., 1990 RACE: rapid amplification of complementary DNA ends, pp. 28–38 in *PCR Protocols: A Guide to Methods and Applications*, edited by M. A. INNIS, D. H. GELFAND, J. J. SNINSKY and T. J. WHITE. Academic Press, New York.
- FURLONG, R. F., and P. W. H. HOLLAND, 2002 Were vertebrates octoploid? *Philos. Trans. R. Soc. Lond. B* **357**: 531–544.
- GERMOT, A., G. LECOINTRE, J.-L. PLOUHINEC, C. LE MENTEC, F. GIRA-

- DOT *et al.*, 2001 Structural evolution of *Otx* genes in craniates. *Mol. Biol. Evol.* **18**: 1668–1678.
- GHANEM, N., O. JARINOVA, A. AMORES, Q. LONG, G. HATCH *et al.*, 2003 Regulatory roles of conserved intergenic domains in vertebrate *Dlx* bigene clusters. *Genome Res.* **13**: 533–541.
- HAHN, M. E., S. I. KARCHNER, M. A. SHAPIRO and S. A. PERERA, 1997 Molecular evolution of two vertebrate aryl hydrocarbon (dioxin) receptors (AHR1 and AHR2) and the PAS family. *Proc. Natl. Acad. Sci. USA* **94**: 13743–13748.
- HOLLAND, P. W., J. GARCIA-FERNÁNDEZ, N. A. WILLIAMS and A. SIDOW, 1994 Gene duplications and the origins of vertebrate development. *Dev. Suppl.*, 125–133.
- HORTON, A. C., N. R. MAHADEVAN, I. RUVINSKY and J. J. GIBSON-BROWN, 2003 Phylogenetic analyses alone are insufficient to determine whether genome duplication(s) occurred during early vertebrate evolution. *J. Exp. Zool. Mol. Dev. Evol.* **299B**: 41–53.
- HUGHES, A. L., and R. FRIEDMAN, 2003 2R or not 2R: testing hypotheses of genome duplication in early vertebrates. *J. Struct. Funct. Genomics* **3**: 85–93.
- HUGHES, A. L., J. DA SILVA and R. FRIEDMAN, 2001 Ancient genome duplications did not structure the human *Hox*-bearing chromosomes. *Genome Res.* **11**: 771–780.
- IRVINE, S. Q., J. L. CARR, W. J. BAILEY, K. KAWASAKI, N. SHIMIZU *et al.*, 2002 Genomic analysis of *Hox* clusters in the sea lamprey *Petromyzon marinus*. *J. Exp. Zool. Mol. Dev. Evol.* **294**: 47–62.
- JANVIER, P., 1996 *Early Vertebrates*. Clarendon Press, Oxford.
- JANVIER, P., 2001 Ostracoderms and the shaping of the gnathostome characters, pp. 172–186 in *Major Events in Early Vertebrate Evolution: Paleontology, Phylogeny, Genetics and Development*, edited by P. E. AHLBERG. Taylor & Francis, New York.
- JONES, D. T., W. R. TAYLOR and J. M. THORNTON, 1992 The rapid generation of mutation data matrices from protein sequences. *Comput. Appl. Biosci.* **8**: 275–282.
- KASAHARA, M., M. HAYASHI, K. TANAKA, H. INOKO, K. SUGAYA *et al.*, 1996 Chromosomal localization of the proteasome Z subunit gene reveals an ancient chromosomal duplication involving the major histocompatibility complex. *Proc. Natl. Acad. Sci. USA* **93**: 9096–9101.
- KELLOGG, D. E., I. RYBALKIN, S. CHEN, N. MUKHAMEDOVA, T. VLASIK *et al.*, 1994 TaqStart antibody: hot start PCR facilitated by a neutralizing monoclonal antibody directed against *Taq* DNA polymerase. *BioTechniques* **16**: 1134–1137.
- KIM, C.-B., C. AMEMIYA, W. BAILEY, K. KAWASAKI, J. MEZEY *et al.*, 2000 *Hox* cluster genomics in the horn shark, *Heterodontus francisci*. *Proc. Natl. Acad. Sci. USA* **97**: 1655–1660.
- KOBEL, H. R., and L. DU PASQUIER, 1986 Genetics of polyploid *Xenopus*. *Trends Genet.* **2**: 310–315.
- KUMAR, S., K. TAMURA, I. B. JAKOBSEN and M. NEI, 2001 MEGA2: molecular evolutionary genetics analysis software. *Bioinformatics* **17**: 1244–1245.
- LARHAMMAR, D., L.-G. LUNDIN and F. HALLBÖÖK, 2002 The human *Hox*-bearing chromosome regions did arise by block or chromosome (or even genome) duplications. *Genome Res.* **12**: 1910–1920.
- LE CROM, S., M. KAPSIMALI, P.-O. BARÔME and P. VERNIER, 2003 Dopamine receptors for every species: gene duplications and functional diversification in Craniates. *J. Struct. Funct. Genomics* **3**: 161–176.
- LIU, J. K., I. GHATTAS, S. LIU, S. CHEN and J. L. R. RUBENSTEIN, 1997 *Dlx* genes encode DNA-binding proteins that are expressed in an overlapping and sequential pattern during basal ganglia differentiation. *Dev. Dyn.* **210**: 498–512.
- LUNDIN, L. G., 1993 Evolution of the vertebrate genome as reflected in paralogous chromosomal regions in man and the house mouse. *Genomics* **16**: 1–19.
- LUNDIN, L.-G., D. LARHAMMAR and F. HALLBÖÖK, 2003 Numerous groups of chromosomal regional paralogies strongly indicate two genome doublings at the root of vertebrates. *J. Struct. Funct. Genomics* **3**: 53–63.
- LYNCH, M., and A. O. RICHARDSON, 2002 The evolution of spliceosomal introns. *Curr. Opin. Genet. Dev.* **12**: 701–710.
- MAISEY, J. G., 1986 Heads and tails: a chordate phylogeny. *Cladistics* **2**: 201–256.
- MARKERT, C. L., J. B. SHAKLEE and G. S. WHITT, 1975 Evolution of a gene. Multiple genes for LDH isozymes provide a model of the evolution of gene structure, function and regulation. *Science* **189**: 102–114.
- MARTIN, A., 2001 Is tetralogy true? Lack of support for the “one-to-four” rule. *Mol. Biol. Evol.* **18**: 89–93.
- MCGUINNESS, T., M. H. PORTEUS, S. SMIGA, A. BULFONE, C. KINGSLEY *et al.*, 1996 Sequence, organization, and transcription of the *Dlx-1* and *Dlx-2* locus. *Genomics* **35**: 473–485.
- MYOJIN, M., T. UEKI, F. SUGAHARA, Y. MURAKAMI, Y. SHIGETANI *et al.*, 2001 Isolation of *Dlx* and *Emx* gene cognates in an agnathan species, *Lampetra japonica*, and their expression patterns during embryonic and larval development: conserved and diversified regulatory patterns of homeobox genes in vertebrate head evolution. *J. Exp. Zool. Mol. Dev. Evol.* **291**: 68–84.
- NAKAMURA, S., D. W. STOCK, K. L. WYDNER, J. A. BOLLEKENS, K. TAKESHITA *et al.*, 1996 Genomic analysis of a new mammalian *Distal-less* gene: *Dlx7*. *Genomics* **38**: 314–324.
- NEIDERT, A. H., V. VIRUPANNAVAR, G. W. HOOKER and J. A. LANGE-LAND, 2001 Lamprey *Dlx* genes and early vertebrate evolution. *Proc. Natl. Acad. Sci. USA* **98**: 1665–1670.
- PANGANIBAN, G., and J. L. R. RUBENSTEIN, 2002 Developmental functions of the *Distal-less/Dlx* homeobox genes. *Development* **129**: 4371–4386.
- PANOPOULOU, G., S. HENNIG, D. GROTH, A. KRAUSE, A. J. POUSTKA *et al.*, 2003 New evidence for genome-wide duplications at the origin of vertebrates using an amphioxus gene set and completed animal genomes. *Genome Res.* **13**: 1056–1066.
- PENDLETON, J. W., B. K. NAGAI, M. T. MURTHA and F. H. RUDDLE, 1993 Expansion of the *Hox* gene family and the evolution of chordates. *Proc. Natl. Acad. Sci. USA* **90**: 6300–6304.
- POLLARD, S. L., and P. W. H. HOLLAND, 2000 Evidence for 14 homeobox gene clusters in human genome ancestry. *Curr. Biol.* **10**: 1059–1062.
- PRINCE, V. E., 2002 The *Hox* paradox: more complex(es) than imagined. *Dev. Biol.* **249**: 1–15.
- RUDDLE, F. H., J. L. BARTELS, K. L. BENTLEY, C. KAPPEN, M. T. MURTHA *et al.*, 1994a Evolution of *Hox* genes. *Annu. Rev. Genet.* **28**: 423–442.
- RUDDLE, F. H., K. L. BENTLEY, M. T. MURTHA and N. RISCH, 1994b Gene loss and gain in the evolution of vertebrates. *Dev. Suppl.*, 155–161.
- SALANECK, E., D. H. ARDELL, E. T. LARSON and D. LARHAMMAR, 2003 Three neuropeptide Y receptor genes in the spiny dogfish, *Squalus acanthias*, support en bloc duplications in early vertebrate evolution. *Mol. Biol. Evol.* **20**: 1271–1280.
- SAMBROOK, J., E. F. FRITSCH and T. MANIATIS, 1989 *Molecular Cloning: A Laboratory Manual*, Ed. 2. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- SHIGETANI, Y., F. SUGAHARA, Y. KAWAKAMI, Y. MURAKAMI, S. HIRANO *et al.*, 2002 Heterotopic shift of epithelial-mesenchymal interactions in vertebrate jaw evolution. *Science* **296**: 1316–1319.
- SIDOW, A., 1992 Diversification of the *Wnt* gene family on the ancestral lineage of vertebrates. *Proc. Natl. Acad. Sci. USA* **89**: 5098–5102.
- SIDOW, A., 1996 Gen(om)e duplications in the evolution of early vertebrates. *Curr. Opin. Genet. Dev.* **6**: 715–722.
- SPRING, J., 1997 Vertebrate evolution by interspecific hybridization—Are we polyploid? *FEBS Lett.* **400**: 2–8.
- STOCK, D. W., and D. A. POWERS, 1998 A monophyletic origin of heart-predominant lactate dehydrogenase (LDH) isozymes of gnathostome vertebrates: evidence from the cDNA sequence of the spiny dogfish (*Squalus acanthias*) LDH-B. *Mol. Mar. Biol. Biotechnol.* **7**: 160–164.
- STOCK, D. W., D. L. ELLIES, Z. ZHAO, M. EKKER, F. H. RUDDLE *et al.*, 1996 The evolution of the vertebrate *Dlx* gene family. *Proc. Natl. Acad. Sci. USA* **93**: 10858–10863.
- STOCK, D. W., J. M. QUATTRO, G. S. WHITT and D. A. POWERS, 1997 Lactate dehydrogenase (LDH) gene duplication during chordate evolution: the cDNA sequence of the LDH of the tunicate *Styela plicata*. *Mol. Biol. Evol.* **14**: 1273–1284.
- SUMIYAMA, K., S. Q. IRVINE, D. W. STOCK, K. M. WEISS, K. KAWASKI *et al.*, 2002 Genomic structure and functional control of the *Dlx3-7* bigene cluster. *Proc. Natl. Acad. Sci. USA* **99**: 780–785.
- SUMIYAMA, K., S. Q. IRVINE and F. H. RUDDLE, 2003 The role of gene duplication in the evolution and function of the vertebrate *Dlx/distal-less* bigene clusters. *J. Struct. Funct. Genomics* **3**: 151–159.

- TANAKA, M., A. MÜNSTERBERG, W. G. ANDERSON, A. R. PRESCOTT, N. HAZON *et al.*, 2002 Fin development in a cartilaginous fish and the origin of vertebrate limbs. *Nature* **416**: 527–531.
- THOMPSON, J. D., T. J. GIBSON, F. PLEWNIAK, F. JEANMOUGIN and D. G. HIGGINS, 1997 The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **25**: 4876–4882.
- TRACY, M. R., and S. B. HEDGES, 2000 Evolutionary history of the enolase gene family. *Gene* **259**: 129–138.
- VACHON, G., B. COHEN, C. PFEIFLE, M. E. MCGUFFIN, J. BOTAS *et al.*, 1992 Homeotic genes of the Bithorax complex repress limb development in the abdomen of the *Drosophila* embryo through the target gene *Distal-less*. *Cell* **71**: 437–450.
- WAGNER, G. P., C. AMEMIYA and F. RUDDLE, 2003 Hox cluster duplications and the opportunity for evolutionary novelties. *Proc. Natl. Acad. Sci. USA* **100**: 14603–14606.
- ZHANG, J., and M. NEI, 1996 Evolution of Antennapedia-class homeobox genes. *Genetics* **142**: 295–303.

Communicating editor: S. YOKOYAMA

