

Carbonic Anhydrase and Its Influence on Carbon Isotope Discrimination during C₄ Photosynthesis. Insights from Antisense RNA in *Flaveria bidentis*¹

Asaph B. Cousins*, Murray R. Badger, and Susanne von Caemmerer

Molecular Plant Physiology Group (A.B.C., M.R.B., S.V.C.) and Australian Research Council Centre of Excellence in Plant Energy Biology (M.R.B.), Research School of Biological Sciences, Australian National University, Canberra, Australian Capital Territory 2601, Australia

In C₄ plants, carbonic anhydrase (CA) facilitates both the chemical and isotopic equilibration of atmospheric CO₂ and bicarbonate (HCO₃⁻) in the mesophyll cytoplasm. The CA-catalyzed reaction is essential for C₄ photosynthesis, and the model of carbon isotope discrimination ($\Delta^{13}\text{C}$) in C₄ plants predicts that changes in CA activity will influence $\Delta^{13}\text{C}$. However, experimentally, the influence of CA on $\Delta^{13}\text{C}$ has not been demonstrated in C₄ plants. Here, we compared measurements of $\Delta^{13}\text{C}$ during C₄ photosynthesis in *Flaveria bidentis* wild-type plants with *F. bidentis* plants with reduced levels of CA due to the expression of antisense constructs targeted to a putative mesophyll cytosolic CA. Plants with reduced CA activity had greater $\Delta^{13}\text{C}$, which was also evident in the leaf dry matter carbon isotope composition ($\delta^{13}\text{C}$). Contrary to the isotope measurements, photosynthetic rates were not affected until CA activity was less than 20% of wild type. Measurements of $\Delta^{13}\text{C}$, $\delta^{13}\text{C}$ of leaf dry matter, and rates of net CO₂ assimilation were all dramatically altered when CA activity was less than 5% of wild type. CA activity in wild-type *F. bidentis* is sufficient to maintain net CO₂ assimilation; however, reducing leaf CA activity has a relatively large influence on $\Delta^{13}\text{C}$, often without changes in net CO₂ assimilation. Our data indicate that the extent of CA activity in C₄ leaves needs to be taken into account when using $\Delta^{13}\text{C}$ and/or $\delta^{13}\text{C}$ to model the response of C₄ photosynthesis to changing environmental conditions.

Isotope analysis of atmospheric CO₂ is an important tool for monitoring changes in the global exchange of CO₂ (Flanagan and Ehleringer, 1998; Yakir and Sternberg, 2000). However, to interpret the atmospheric CO₂ isotopic signature requires an understanding of the isotopic fractionation steps associated with specific processes during leaf gas exchange (Yakir and Sternberg, 2000). Leaf level models of carbon isotope exchange ($\Delta^{13}\text{C}$) in C₄ plants have been used for many years to help interpret the response of C₄ plants to changing environmental conditions. However, only recently has the genetic manipulation of the C₄ photosynthetic apparatus provided an opportunity to reexamine the C₄ leaf level models of $\Delta^{13}\text{C}$ (von Caemmerer et al., 1997a, 1997b).

Most C₄ plants utilize a compartmentalized CO₂-concentrating mechanism between the mesophyll and bundle sheath cells (BSC) to increase the CO₂ partial pressure ($p\text{CO}_2$) around the site of Rubisco in the BSC.

The first enzymatic step in C₄ photosynthesis is the reversible hydration reaction catalyzed by carbonic anhydrase (CA), which converts CO₂ to bicarbonate (HCO₃⁻) in the mesophyll cytoplasm. Subsequently, HCO₃⁻ is fixed via phosphoenolpyruvate carboxylase (PEPC) into a four-carbon acid that diffuses to the BSC for decarboxylation (Kanai and Edwards, 1999). The specialized biochemistry and leaf anatomy of C₄ plants results in a $p\text{CO}_2$ around the site of Rubisco severalfold higher than current atmospheric levels, significantly reducing the rates of photorespiration (Hatch, 1987; Kanai and Edwards, 1999).

The carbon isotope discrimination during C₄ photosynthesis is determined by the fractionation that occurs during diffusion of CO₂ into the leaf, its conversion to HCO₃⁻ via CA, and the subsequent carboxylation reactions catalyzed by PEPC and Rubisco (Peisker, 1982; Farquhar, 1983; Peisker and Henderson, 1992; von Caemmerer et al., 1997a). The extent to which Rubisco can fractionate against CO₂ is determined by the amount of leakiness (ϕ), defined as the fraction of CO₂ fixed by PEPC that subsequently leaks out of the BSC. If the BSC were gas tight, then all of the CO₂ released into the BSC would be fixed by Rubisco and no fractionation would occur at this step. However, CO₂ can leak out of the BSC, allowing Rubisco to influence the overall discrimination during C₄ photosynthesis (Farquhar, 1983; Peisker and Henderson, 1992).

Differences in the ratio of CO₂ partial pressures between the intercellular airspace and the atmosphere (p_i/p_a) along with ϕ are the main factors attributed to

¹ This work was supported by a National Science Foundation international postdoctoral fellowship (to A.B.C.).

* Corresponding author; e-mail asaph.cousins@anu.edu.au; fax 61-2-61255075.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Susanne von Caemmerer (susanne.caemmerer@anu.edu.au).

Article, publication date, and citation information can be found at www.plantphysiol.org/cgi/doi/10.1104/pp.106.077776.

variation in $\Delta^{13}\text{C}$ in C₄ plants (Farquhar, 1983). The ratio p_i/p_a is primarily determined by stomatal conductance, whereas ϕ depends on the physical conductance of the BSC walls and the balance between the C₄ and C₃ cycles. Little change in ϕ was determined with gas exchange and $\Delta^{13}\text{C}$ measurements in various C₄ plants under a variety of environmental conditions (Henderson et al., 1992). However, with the use of antisense technologies, it has been shown that $\Delta^{13}\text{C}$ and ϕ increase when the capacity of the C₃ cycle is reduced relative to the C₄ cycle (von Caemmerer et al., 1997a, 1997b). Growth conditions (e.g. elevated CO₂ and water stress) have also been reported to influence the balance of the C₄ and C₃ cycles, leading to an altered isotopic composition of dry matter (Watling et al., 2000; Williams et al., 2001), although the influence on CA was not addressed in these studies.

There is limited research concerning the influence of CA activity on $\Delta^{13}\text{C}$ in C₄ plants. Recent work indicates that CA activity in wild-type *Flaveria bidentis* is in excess and does not limit CO₂ assimilation under normal conditions (von Caemmerer et al., 2004). *F. bidentis* lines with reduced levels of CA, due to the expression of antisense constructs targeted to a putative mesophyll cytosolic CA, showed that rates of CO₂ assimilation were unaffected by a decrease in CA activity until activity was less than 20% of wild type (von Caemmerer et al., 2004). Although large changes in CA activity had little effect on photosynthetic rates, according to the $\Delta^{13}\text{C}$ theory developed by Farquhar in 1983 (see "Materials and Methods"), the decrease in the hydration reaction of CO₂ (V_h) relative to the rate of PEPC carboxylation (V_p) should increase $\Delta^{13}\text{C}$ potentially without a corresponding change in the rate of net CO₂ assimilation (Farquhar, 1983).

In this article, we use *F. bidentis* plants with low CA activity to examine the influence of the hydration reaction of CO₂ on $\Delta^{13}\text{C}$ during C₄ photosynthesis. These results are discussed in relation to measurements of $\Delta^{13}\text{C}$ made in *F. bidentis* under various irradiances, as well as plants with reduced levels of Rubisco.

RESULTS

Carbon Isotope Discrimination

Light Response Curves

In the mass spectrometric gas-exchange system used here for online $\Delta^{13}\text{C}$ measurements, the leaf chamber gas outlet of a LI-6400 gas-exchange system (LI-COR) was directly coupled to a mass spectrometer (micro-mass ISOPRIME; Micromass Ltd.) via a gas-permeable silicone membrane (Fig. 1). This allowed the measurement of the $^{13}\text{C}/^{12}\text{C}$ ratio of the CO₂ in the airstream without prior purification of that CO₂. We measured rates of net CO₂ assimilation and $\Delta^{13}\text{C}$ in *F. bidentis* wild-type plants in response to photon flux density (PFD) to test our online systems with previously published values of $\Delta^{13}\text{C}$ from C₄ plants (Henderson et al., 1992). A summary of the symbols used in the text are shown in Table I. Net CO₂ assimilation increased with PFD to near-saturating rates (Fig. 2a). However, there was little change in $\Delta^{13}\text{C}$, p_i/p_a , and BSC CO₂ leakiness (ϕ), except at the two lowest light levels (Fig. 2, b–d). There was more uncertainty in the $\Delta^{13}\text{C}$ measurements made at low light because of the higher ratio of the rate of CO₂ entry into the chamber to the rate of net CO₂ assimilation by the leaf (ξ ; see Fig. 2, legend). Leakiness was calculated by rearranging Equation 2 (see equations in "Materials and Methods") and substituting b_4 with Equation 3, with the assumption that the initial CO₂ carboxylation reaction catalyzed by PEPC to the rate of CO₂ hydration by CA (V_p/V_h) was zero. These gas-exchange and $\Delta^{13}\text{C}$ measurements are similar to those previously reported for *Amaranthus edulis* and *Zea mays* under similar measurement conditions (Henderson et al., 1992).

Rubisco Small Subunit Plants

Net CO₂ assimilation in *F. bidentis* plants with reduced levels of Rubisco caused by antisense RNA constructs targeted to the nuclear-encoded gene for




Figure 1. Arrangement of the gas flow controllers, the LI-6400 gas exchange system, and the mass spectrometer system used for simultaneous measurements of leaf gas exchange and carbon isotope discrimination. Switching between gas samples was controlled by a manual four-way valve. The zero and reference readings were made before and after each leaf measurement and averaged during the calculations.

Table I. Symbols used in the text

Symbol	Description
A	Net CO ₂ assimilation
a	Fractionation during diffusion of CO ₂ from the chloroplast to the atmosphere (4.4‰)
a_l	Fractionation of CO ₂ diffusion through a liquid (0.7‰)
BSC	Bundle sheath cells
b_3	Combined discrimination of Rubisco, respiration, and photorespiration (see Eq. 4)
b_4	Combined discrimination of PEPC, respiration, and hydration/dehydration of CO ₂ (see Eq. 3 and Fig. 4)
b_{PEPC}	Discrimination by PEPC (2.2‰)
$\Delta^{13}C$	Carbon isotope discrimination
CA	Carbon anhydrase
e	Fractionation during respiration (3‰ or -6‰)
e_s	Fractionation as CO ₂ dissolves (1.1‰)
e_b	Equilibrium fractionation factor for the catalyzed hydration/dehydration of CO ₂ (-9‰)
f	Discrimination during photorespiration (10‰ or -6.8‰)
ϕ	The fraction of CO ₂ fixed by PEPC that subsequently leaks out of the BSC
g_w	The internal conductance to the diffusion of CO ₂ between the intercellular air space and the site of carboxylation in the mesophyll cytoplasm
h	Catalyzed fractionation during CO ₂ hydration (1.1‰)
k_{CA}	Rate constant of carbonic anhydrase
K_c	Michaelis constant of Rubisco for CO ₂
K_o	Michaelis constant of Rubisco for O ₂
K_p	Michaelis constant of PEPC for CO ₂
M_d	Rate of mitochondrial respiration
M_m	Rate of mitochondrial respiration in the mesophyll cells
M_s	Rate of mitochondrial respiration in the BSC
p_{CO_2}	Partial pressure of CO ₂
p_e	p_{CO_2} of dry air entering the leaf chamber
p_i	p_{CO_2} of the intercellular airspace
p_m	p_{CO_2} of the mesophyll cytoplasm
p_o	p_{CO_2} of dry air leaving the leaf chamber
R_e	¹³ C/ ¹² C of the air entering the leaf chamber
R_o	¹³ C/ ¹² C of the air leaving the leaf chamber
PFD	Photon flux density
PSII	PSII
ξ	$p_e/(p_e - p_o)$
s	Fractionation during the leakage of CO ₂ from the BSC (1.8‰)
V_c	Rate of Rubisco carboxylation
V_{cmax}	Maximal rate of Rubisco carboxylation
V_h	Rate of CO ₂ hydration
V_o	Rate of photorespiration
V_p	Rate of PEP carboxylation $(A + M_d)/(1 - \phi)$ or $(p_m V_{pmax})/(p_m + K_p)$
V_{pmax}	Maximal rate of PEPC carboxylation

the small subunit of Rubisco (anti-SSu plants) had rates between 40% to 80% of wild-type plants (Table II). Additionally, the ratio of p_i/p_a , $\Delta^{13}C$, and ϕ were higher in the anti-SSu-plants as compared with wild-type plants (Table II). The parameter ϕ was determined from simultaneous gas-exchange and isotope measurements and solving for ϕ in Equation 2. Our measurements of $\Delta^{13}C$ and leaf gas exchange

are similar to previously published values by von Caemmerer et al. (1997b). The comparison of our results to previously published $\Delta^{13}C$ values shows that our system can accurately and consistently monitor the influence of both environmental conditions and perturbations to the C₄ photosynthetic apparatus on instantaneous carbon isotope discrimination.

CA Plants

Carbon isotope discrimination ($\Delta^{13}C$) increased as CA activity decreased in the *F. bidentis* plants containing the antisense RNA constructs targeted to the putative cytosolic CA (anti-CA plants; Fig. 3). CA activity, reported here as a rate constant (k_{CA} $\mu\text{mol m}^{-2} \text{s}^{-1} \text{Pa}^{-1}$), was determined on leaf extracts using mass spectrometry to measure the rates of ¹⁸O₂ exchange from doubly labeled ¹³C¹⁸O₂ to H₂¹⁶O (see "Materials and Methods"). Interestingly, $\Delta^{13}C$ was more sensitive than net CO₂ assimilation to changes in CA activity as $\Delta^{13}C$ increased in some anti-CA plants, whereas net CO₂ assimilation remained similar to wild-type plants (Fig. 3). In these anti-CA plants with reduced CA activity and wild-type rates of net CO₂ assimilation, $\Delta^{13}C$ increased 1‰ to 2‰, which is a large shift for C₄ photosynthesis (Fig. 3c, inset; Table III). In the anti-CA plants, net CO₂ assimilation rates and p_i/p_a were similar to wild-type plants, except when CA activities were less than 20% of wild type (Fig. 3, a and b). Anti-CA plants with extremely low levels of CA activity (<5% of wild type) and low rates of net CO₂ assimilation had extremely high values of $\Delta^{13}C$ (Fig. 3c).

Nearly all of the measured values of $\Delta^{13}C$ fall within the theoretical relationship of $\Delta^{13}C$ to p_i/p_a as predicted from the model of C₄ carbon isotope discrimination developed by Farquhar (1983; Fig. 4; see "Materials and Methods"). Only in the anti-CA plants with extremely low CA activity do the measured values of $\Delta^{13}C$ fall outside the predicted values (Fig. 4). The theoretical relationship of $\Delta^{13}C$ and p_i/p_a was calculated with a ϕ value of 0.24, and the initial CO₂ carboxylation reaction catalyzed by PEPC relative to the CO₂ hydration by CA (V_p/V_h) was assumed to be either zero or 1 (as indicated in Fig. 4). The b_4 parameter, which is the combined fractionation associated with PEPC, respiration, and the isotopic equilibrium during the dissolution of CO₂ and conversion to HCO₃⁻, used in these calculations was determined with either the CA catalyzed (solid lines) or the spontaneous uncatalyzed (dotted lines) CO₂ and HCO₃⁻ hydration and dehydration fractionation factors (see "Materials and Methods").

To characterize the influence of CA activity on $\Delta^{13}C$, independent of changes in net CO₂ assimilation, we pooled the data of anti-CA plants with reduced CA activity and wild-type-like photosynthetic rates. $\Delta^{13}C$ was higher in the anti-CA plants compared to the wild-type plants, whereas p_i/p_a was unchanged (Table III). The in vivo CA activity (CA_{leaf}), which is the product of k_{CA} and the p_{CO_2} in the mesophyll cytoplasm (p_m),




Figure 2. a, Net CO₂ assimilation rate. b, Ratio of intercellular to ambient CO₂ partial pressures (p_i/p_a). c, Carbon isotope discrimination ($\Delta^{13}C$). d, Bundle sheath leakiness to CO₂ (ϕ) as a function of PFD ($\mu\text{mol quanta m}^{-2} \text{s}^{-1}$). Measurements were made at a $p\text{CO}_2$ of 52 Pa, a $p\text{O}_2$ of 4.8 kPa, and a leaf temperature of 30°C. Shown are the means \pm the SE of measurements made on three to five leaves from two *F. bidentis* wild-type plants. Values for ξ (Eq. 1) were 29.9 ± 0.75 , 15.7 ± 0.54 , 7.11 ± 0.24 , 5.5 ± 0.23 , and 4.9 ± 0.17 at PFDs of 150, 300, 800, 1,400, and 2,000 $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$. ϕ was calculated from Equation 5, assuming $V_p/V_h = 0$.

was significantly less in the anti-CA relative to the wild-type plants (Table III). The value of p_m was calculated with an internal conductance to the diffusion of CO₂ between the intercellular airspace and the site of carboxylation in the mesophyll cytoplasm (g_w) of $10 \text{ mol m}^{-2} \text{s}^{-1} \text{ Pa}^{-1}$. The ratio of V_p/V_h determined from the online measurements of $\Delta^{13}C$ was approximately 6 times greater in the anti-CA plants than in the wild-type plants (Table III). It appears that a rather large decrease in leaf CA activity in *F. bidentis* can maintain the chemical equilibrium between CO₂ and HCO₃⁻ needed to sustain photosynthesis, but limits the isotopic equilibrium causing $\Delta^{13}C$ to increase without changes in ϕ .

It should be noted that the absolute value of V_p/V_h determined this way is largely influenced by ϕ and slightly by g_w . For example, changing g_w from 6 to $10 \text{ mol m}^{-2} \text{s}^{-1} \text{ Pa}^{-1}$ shifts calculations of V_p/V_h from 0.08 to 0.07 and 0.55 to 0.46 for wild-type and anti-CA plants, respectively. However, changing ϕ

from 0.24 to 0.10, assuming a constant g_w of $10 \text{ mol m}^{-2} \text{s}^{-1} \text{ Pa}^{-1}$, causes V_p/V_h to increase from 0.07 to 0.61 in the wild-type plants and from 0.46 to 0.99 in the anti-CA plants (Table III). In the anti-CA plants, which have a reduced capacity to concentrate CO₂ within the BSC, it is predicted from the C₄ photosynthetic model that ϕ will decrease relative to the wild-type plants (see below), which would increase the difference of V_p/V_h between the wild-type and the anti-CA plants. V_p/V_h can also be approximated from gas exchange and in vitro CA activity as V_p/CA_{leaf} where CA_{leaf} is calculated as $k_{CA}p_m$ and V_p is calculated as $(A + M_d)/(1 - \phi)$ (von Caemmerer, 2000). The parameter M_d is the daytime rate of mitochondrial respiration assumed to be $2 \mu\text{mol m}^{-2} \text{s}^{-1}$. The ratio of V_p/V_h determined from the in vitro assays of CA activity was approximately 4 times greater in the anti-CA plants than in the wild-type plants (Table III). The absolute value of V_p/V_h calculated in this manner is also influenced by changes in g_w and ϕ , although neither

Table II. CA rate constant (k_{CA}), net CO₂ assimilation rate (A), ratio of intercellular to atmospheric CO₂ partial pressure (p_i/p_a), online $\Delta^{13}C$ discrimination, and leakiness of CO₂ out of the BSCs (ϕ) in the anti-SSu plants from the primary transformant 136-13

For calculation of ϕ , the V_p/V_h ratio was assumed to be zero. Measurements were made at a $p\text{CO}_2$ of 52 Pa, a $p\text{O}_2$ of 4.8 kPa, a PFD of $2,000 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$, and a leaf temperature of 30°C. $n = 4$ for the wild-type plants.

	k_{CA}	A	p_i/p_a	$\Delta^{13}C$	Leakiness ϕ
	$\text{mol m}^{-2} \text{s}^{-1} \text{ Pa}^{-1}$	$\mu\text{mol m}^{-2} \text{s}^{-1}$		‰	
136-13-11#1	52	17.0	0.67	6.1	0.43
136-13-12#4	88	19.5	0.64	6.5	0.45
136-13-12#3	62	27.9	0.72	6.0	0.42
136-13-11#2	81	29.9	0.71	5.4	0.39
Wild type	69 ± 2	37.6 ± 2.6	0.55 ± 0.02	2.5 ± 0.4	$0.25 \pm .02$




Figure 3. Net CO₂ assimilation rate, the ratio of intercellular to ambient *p*CO₂ (p_i/p_a), and carbon isotope discrimination ($\Delta^{13}\text{C}$) as a function of the rate constant of leaf CA (k_{CA} $\mu\text{mol m}^{-2} \text{s}^{-1} \text{Pa}^{-1}$). The inset in c shows the expanded scale of $\Delta^{13}\text{C}$ where net CO₂ assimilation is relatively constant. Each point represents a measurement made on a different plant grown in a glasshouse at ambient CO₂ or in a growth cabinet at 1% CO₂: wild-type plants grown at ambient CO₂ (\square); anti-CA plants grown at ambient CO₂ (\blacksquare); wild-type grown at 1% CO₂ (\circ , \triangle); and anti-CA plants grown at 1% CO₂ (\bullet , \blacktriangle). Measurements were made at 2,000 $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$, leaf temperature of 30°C, and an inlet CO₂ concentration of either 38 Pa in air (\triangle , \blacktriangle) or 52 Pa of CO₂ in a 90.5 kPa of N₂ and 4.8 kPa of O₂ gas mixture (\square , \blacksquare). The lines represent the best fit for all measurements (wild-type and anti-CA plants) made at either 38 or 52 Pa of CO₂.

parameter has a large influence on the relative changes of V_p/V_h between wild-type and anti-CA plants.

Dry Matter $\delta^{13}\text{C}$

Leaf dry matter $\delta^{13}\text{C}$, the ratio of $^{13}\text{C}/^{12}\text{C}$ of the sample relative to the standard Vienna Pee Dee Belemnite (VPDB), was lower in plants with low levels of CA and correlated with increases in $\Delta^{13}\text{C}$ (Fig. 5). Leaf $\delta^{13}\text{C}$ was determined on plants germinated and grown in a glasshouse. After collecting an entire leaf for $\delta^{13}\text{C}$, the three plants with very low CA and photosynthetic rates were transferred after several weeks to the 1% CO₂ growth cabinets before leaf gas-exchange measurements were made. Otherwise, the leaf opposite to the one used for gas exchange was sampled for $\delta^{13}\text{C}$.

Photosynthetic and Carbon Isotope Discrimination Models

It has recently been shown that low leaf CA activity in *F. bidentis* reduces the capacity of the C₄ cycle by limiting the rate of PEPC carboxylation of HCO₃⁻ (V_p) (von Caemmerer et al., 2004). Here, we use the C₄ photosynthetic model developed by Berry and Farquhar (1978) and von Caemmerer (2000) to predict the response of net CO₂ assimilation, bundle sheath *p*CO₂, photorespiration (V_o), and ϕ to changes in the activity of PEPC due to a limitation in CA activity. In the C₄ photosynthetic model, the CA-mediated hydration/dehydration reaction of CO₂ within the mesophyll cytoplasm has not been incorporated. However, manipulating V_p within the model simulates the effect of changing CA activity and leads to a diminished ability to concentrate CO₂ within the BSC, which decreases both the photosynthetic rate and ϕ (Fig. 6, a and b).

The outputs from the C₄ photosynthetic model, specifically the rates of Rubisco carboxylation (V_c), V_o , V_p , ϕ , and the *p*CO₂ in the BSC, were then incorporated into the model of C₄ carbon isotope discrimination ($\Delta^{13}\text{C}$) developed by Farquhar (1983). The $\Delta^{13}\text{C}$ model was used to determine which photosynthetic parameters would influence $\Delta^{13}\text{C}$ consistent with our experimental data and to demonstrate the influence of ϕ on $\Delta^{13}\text{C}$ independent of changes in V_p/V_h . The model in Figure 6 included sufficient CA activity to keep V_p/V_h close to zero as V_p changes and p_i/p_a were held constant at 0.4. As shown in Figure 6c, when ϕ and the *p*CO₂ in the BSC are low, $\Delta^{13}\text{C}$ decreases as the ability of Rubisco to fractionate is reduced. Additionally, the $\Delta^{13}\text{C}$ model accounts for the effects of fractionation during respiration (e) and photorespiration (f); however, there is uncertainty in the specific values of factors e and f in the model (Gillon and Griffiths, 1997; Ghashghaie et al., 2003). Therefore, to test the influence of these parameters on the $\Delta^{13}\text{C}$ model, various values of e (3‰ versus -6‰) and f (-6.8‰ versus 10‰) were used. Even at low CO₂ assimilation rates, relatively large changes in e and f had only a small influence on $\Delta^{13}\text{C}$ (Fig. 6c).

Table III. Net CO₂ assimilation rate (*A*), the ratio of intercellular to ambient pCO₂ (*p_i/p_a*), CA_{leaf} calculated as (*k_{CA}p_m*), Δ¹³C, and *V_p/V_h* for *F. bidentis* wild-type plants and anti-CA plants with low CA_{leaf} activity and wild-type-like net CO₂ assimilation rates

Measurements were made at a pCO₂ of 52 Pa, a pO₂ of 4.8 kPa, PFD of 2,000 μmol quanta m⁻² s⁻¹, and a leaf temperature of 30°C. *V_p/V_h* was estimated either by online Δ¹³C measurements* using Eqs. 3 and 5 ("Materials and Methods") or estimated from *V_p/CA_{leaf}*** where *V_p* was calculated as (*A* + *M_d*)/(1 - φ) and CA_{leaf} *g_w* was assumed to be either 10 or 6 μmol m⁻² s⁻¹ Pa⁻¹, and φ was set at either 0.24 or 0.1. *M_d* is the daytime rate of respiration assumed to be 2 μmol m⁻² s⁻¹. *n* = 7 and 5 for anti-CA and wild-type plants, respectively.

	<i>A</i>	<i>p_i/p_a</i>	CA _{leaf}	Δ ¹³ C	<i>V_p/V_h</i>				
					<i>g_w</i> = 10 φ = 0.24	<i>g_w</i> = 6 φ = 0.24	<i>g_w</i> = 10 φ = 0.1	<i>g_w</i> = 6 φ = 0.1	
	μmol m ⁻² s ⁻¹		μmol m ⁻² s ⁻¹	‰					
Anti-CA	42 ± 0.6	0.48 ± 0.02	228 ± 37	4.4 ± 0.2	Δ ¹³ C*	0.46 ± 0.06	0.55 ± 0.07	0.99 ± 0.08	1.1 ± 0.08
					In vitro**	0.29 ± 0.04	0.40 ± 0.04	0.24 ± 0.03	0.30 ± 0.04
Wild type	40 ± 0.1	0.47 ± 0.02	774 ± 48	3.3 ± 0.2	Δ ¹³ C*	0.07 ± 0.07	0.08 ± 0.08	0.61 ± 0.06	0.62 ± 0.08
					In vitro**	0.07 ± 0.01	0.09 ± 0.01	0.06 ± 0.01	0.07 ± 0.01

DISCUSSION

Carbon isotope discrimination increased in the short term during leaf gas exchange (Δ¹³C) and the carbon isotope composition of leaf dry matter (δ¹³C) decreased in transformants containing reduced levels of leaf CA. As was previously reported (von Caemmerer et al., 2004), leaf CA activity appears to be in excess to maintain steady-state rates of net CO₂ assimilation in *F. bidentis* at high light. A nearly 80% decrease in leaf CA activity was needed before net CO₂ assimilation was affected when measured at a CO₂ partial pressure (pCO₂) of 52 Pa (Fig. 3a). The CA activity required to maintain wild-type-like photosynthetic rates increased when measurements were conducted at a lower pCO₂ of 38 Pa (Fig. 3a). This is in agreement with previously published work where reduced leaf cytosolic CA activity affected the initial slope of the CO₂ response curve in *F. bidentis* when the rate of CO₂ hydration limited the supply of HCO₃⁻ for PEPC carboxylation (von Caemmerer et al., 2004).

Carbon Isotope Discrimination and CA Activity

According to the C₄ photosynthetic model (von Caemmerer, 2000), a limitation in the supply of cytosolic HCO₃⁻ will lead to a decrease in the initial CO₂ carboxylation reaction catalyzed by PEPC and reduce the capacity of the C₄ pump to concentrate CO₂ within the BSC. A reduced pCO₂ in the BSC leads to a decrease in the rate of net CO₂ assimilation as well as a lower BSC CO₂ leakiness (φ). In the model of C₄ carbon isotope discrimination, the main factors that influence Δ¹³C are changes in the intercellular to ambient CO₂ partial pressures (*p_i/p_a*) and φ (Farquhar, 1983). When the ratio of PEPC carboxylation to the hydration reaction of CO₂ (*V_p/V_h*) is near zero (i.e. CA activity is high relative to PEPC carboxylation), the C₄ carbon isotope model predicts that Δ¹³C will decrease as φ decreases (Fig. 6). The Δ¹³C modeling illustrates that when the front end of the C₄ cycle is diminished (either by reduced CA and/or PEPC activity or anything else), φ decreases and Δ¹³C associated with φ also decreases (Fig. 6). However, in the anti-CA plants,

which potentially reduced the ability to concentrate CO₂ in the BSC, Δ¹³C increased, which cannot be explained in the model by decreases in φ, but can be explained by changes in *V_p/V_h*.

Due to the high levels of mesophyll cytoplasmic CA activity in C₄ plants, it is generally assumed that CO₂ and HCO₃⁻ are in close chemical equilibrium. Under such conditions, the ratio of *V_p/V_h* in Equation 3 approaches zero and can be omitted from the calculation of *b₄*. However, when *V_p/V_h* tends away from zero, Equation 3 can be expressed with the fractionation factors provided in "Materials and Methods" as *b₄* = -5.7 + 7.9 *V_p/V_h* at 25°C. The *b₄* fractionation factor becomes more positive as *V_p/V_h* increases and Δ¹³C increases even without changes in *p_i/p_a* and φ. As shown in Figure 4, varying *V_p/V_h* between 0 and 1 can




Figure 4. Carbon isotope discrimination (Δ¹³C) as a function of the ratio of intercellular to ambient pCO₂ (*p_i/p_a*). The white symbols are wild-type plants and the black symbols are anti-CA plants. Other symbols and measurement conditions are as described in Figure 3. The lines represent the theoretical relationship of Δ¹³C and *p_i/p_a*, where φ = 0.24, the ratio of the PEPC carboxylation to the CO₂ hydration reaction (*V_p/V_h*) is either 0 or 1, and the *b₄* parameter is calculated with the catalyzed (solid lines, *b₄* = -5.7 + 7.9 *V_p/V_h*) and uncatalyzed (dotted lines, *b₄* = -4.5 + 12.5 *V_p/V_h*) CO₂ and HCO₃⁻ hydration and dehydration fractionation factors. *g_w* was assumed to be large such that *p_i* = *p_m*.




Figure 5. Leaf dry matter $\delta^{13}\text{C}$, determined on the entire leaf opposite to the one used for gas exchange and enzyme analysis, plotted against changes in online carbon isotope discrimination ($\Delta^{13}\text{C}$). The white symbols are wild-type plants and the black symbols are anti-CA plants. All plants were germinated and grown in a glasshouse under ambient atmospheric CO_2 conditions. The plants with extremely low $\delta^{13}\text{C}$ values (●) were transferred to the 1% CO_2 growth cabinets after tissue was collected for $\Delta^{13}\text{C}$ analysis.

have a large impact on $\Delta^{13}\text{C}$, especially when p_i/p_a is high. Nearly all the variation in $\Delta^{13}\text{C}$ in the anti-CA plants can be explained by changes in V_p/V_h (Fig. 4). Only when CA activity and photosynthetic rates decline dramatically do changes in V_p/V_h and p_i/p_a not accurately predict $\Delta^{13}\text{C}$ (see below for further discussion).

Variation in the Ratio of PEPC Carboxylation to CO_2 Hydration by CA

The large change in V_p/V_h without changes in photosynthesis in the anti-CA plants (Table III) indicates that $\Delta^{13}\text{C}$ is more sensitive to a reduction in CA activity than net CO_2 assimilation. The influence of V_p/V_h on $\Delta^{13}\text{C}$ is predicted by the C_4 photosynthetic model for carbon isotope discrimination developed by Farquhar (1983), and here we demonstrate the influence of CA activity on $\Delta^{13}\text{C}$ in a C_4 plant. In wild-type *F. bidensis* plants, CA appears to be in excess for supporting photosynthesis and V_p/V_h approaches zero. However, it has been reported that CA activity in most C_4 species is only just sufficient to support photosynthetic rates, especially in C_4 monocots (Hatch and Burnell, 1990; Gillon and Yakir, 2000, 2001), and the influence of V_p/V_h on $\Delta^{13}\text{C}$ may be greater in these C_4 species. For example, we measured (A.B. Cousins, M. R. Badger, and S. von Caemmerer, unpublished data) CA activity in *Z. mays* as $266 \pm 22 \mu\text{mol m}^{-2} \text{s}^{-1}$, which is similar to our anti-CA plants with wild-type

photosynthetic rate (see Fig. 3c; Table III). Gillon and Yakir (2001) reported even lower CA activity for a number of C_4 grasses, which correspond to the low CA activities we measured in anti-CA plants shown in Figure 3c, inset. The values of V_p/V_h estimated from the CA activity and net CO_2 assimilation from this article indicate that $\Delta^{13}\text{C}$ will differ in C_4 plants that have been reported to contain a range of CA_{leaf} activity (2–529 $\mu\text{mol m}^{-2} \text{s}^{-1}$) with generally similar photosynthetic rates (Gillon and Yakir, 2001). However, a




Figure 6. Modeling the response of net CO_2 assimilation (a); the $p\text{CO}_2$ in the BSC and BSC CO_2 leakiness, ϕ (b); and $\Delta^{13}\text{C}$ (c) in response to changes in PEPC activity (V_p). The C_4 model used a V_{cmax} of $60 \mu\text{mol m}^{-2} \text{s}^{-1}$, a bundle sheath conductance to CO_2 per leaf area of $0.03 \mu\text{mol m}^{-2} \text{s}^{-1} \text{Pa}^{-1}$, K_m of PEPC for CO_2 (K_p) of 8 Pa, K_m of Rubisco for CO_2 (K_c) and O_2 (K_o) of 65 Pa and 45 kPa, fraction of PSII in the BSC 0.2 and Rubisco specificity of 2,590 Pa/Pa in the gas phase, and mitochondrial respiration was $2 \mu\text{mol m}^{-2} \text{s}^{-1}$, one-half of which was assumed to occur in the mesophyll. The $p\text{O}_2$ in the mesophyll was assumed to be 20 kPa. Carbon isotope discrimination was calculated using the C_4 photosynthetic model output and a constant p_i/p_a of 0.4. V_h was set at $2,000 \mu\text{mol m}^{-2} \text{s}^{-1}$ and V_p/V_h varied between 0.001 and 0.1, causing only a 0.3‰ shift in $\Delta^{13}\text{C}$ at a constant ϕ . The lines for $\Delta^{13}\text{C}$ represent models determined with Equation 2 by substituting the b_4 and b_3 factors with Equations 3 and 4, respectively. The lines for $\Delta^{13}\text{C}$ represent models using different fractionation factors for respiration (3‰ dotted line and -6% solid and dashed lines) and photorespiration (10‰ dashed line and -6.8% dotted and solid lines).

systematic investigation of the influence of V_p/V_h on $\Delta^{13}\text{C}$ in a range of C₄ species needs to be conducted.

Changes in V_p/V_h and its influence on $\Delta^{13}\text{C}$ also have important implications for interpreting physiological processes responsible for changes in $\Delta^{13}\text{C}$ and $\delta^{13}\text{C}$ during C₄ photosynthesis, particularly in response to changing environmental conditions. Water stress, reduced nitrogen availability, and atmospheric CO₂ availability have all been reported to increase $\Delta^{13}\text{C}$ in C₄ plants by 1‰ to 3‰ (Meinzer et al., 1994; Ranjith et al., 1995; Buchmann et al., 1996; Saliendra et al., 1996; Meinzer and Saliendra, 1997; Meinzer and Zhu, 1998; Watling et al., 2000). Variation in $\Delta^{13}\text{C}$ in these reports has been interpreted as changes in either p_i/p_a and/or ϕ , with the apparent assumption that V_p/V_h remains close to zero in all treatments. However, there are a few reports in the literature that suggest CA activity in C₄ plants is also influenced by environmental conditions, including nitrogen status, atmospheric CO₂ availability, and salt stress (Cervigni et al., 1971; Burnell et al., 1990; Brownell et al., 1991), implying that environmental conditions may also alter V_p/V_h and thus influence measured $\Delta^{13}\text{C}$. Because leaf CA activity in C₄ plants is largely dependent on the internal CO₂ partial pressures, conditions that influence CO₂ availability, such as water stress and growth under elevated atmospheric CO₂, will also alter V_p/V_h . C₄ photosynthesis generally operates near CO₂-saturating conditions at current atmospheric $p\text{CO}_2$ such that a reduction in p_i due to stomatal closure will cause V_p/V_h to increase. However, under such conditions, the ratio of p_i/p_a also decreases and the influence of V_p/V_h on $\Delta^{13}\text{C}$ decreases as shown in Figure 4. Alternatively, it has been shown that, under well-watered conditions, C₄ photosynthesis generally does not respond to increases in atmospheric $p\text{CO}_2$ (McLeod and Long, 1999; Ghannoum et al., 2000; Wall et al., 2001; Ainsworth and Long, 2005; Leakey et al., 2006). However, because p_i/p_a is generally constant with changing atmospheric $p\text{CO}_2$ and CA has a very high K_m for HCO₃⁻, an increase in CO₂ availability will increase V_{hr} , whereas PEPC is generally saturated around ambient $p\text{CO}_2$ and V_p will not change. This raises the possibility that growth under future atmospheric CO₂ conditions will alter $\Delta^{13}\text{C}$ regardless of other environmental changes if CA is limiting.

Both online and in vitro measurements of V_p/V_h indicated that changes in CA activity have a significant influence on $\Delta^{13}\text{C}$ without changes in p_i/p_a and ϕ . It must be noted that, although changes in g_w have a subtle effect on estimates of V_p/V_h , variation in ϕ can lead to large shifts in the absolute values of V_p/V_h when determined from the $\Delta^{13}\text{C}$ measurements. There are no direct means of measuring ϕ , but it can be estimated using $\Delta^{13}\text{C}$ measurements when V_p/V_h is assumed to be close to zero. The use of antisense technology targeted toward the C₄ PEPC enzyme would provide a range of V_p/V_h values and would allow an estimate of ϕ when V_p/V_h was known to be close to zero.

Low CA and Photosynthetic Mutants

In the majority of CA plants, the increase in $\Delta^{13}\text{C}$ can be explained by changes in the ratio of V_p/V_h and p_i/p_a (Fig. 4). However, this explanation does not hold true for plants with very low leaf CA activity and photosynthetic rates. Potentially, the amount of direct fixation of atmospheric CO₂ in the BSC, leakage of HCO₃⁻ from the BSC, as well as photorespiration and respiration would influence $\Delta^{13}\text{C}$ especially when net CO₂ assimilation is inhibited. Theoretically, CO₂ assimilation by direct diffusion of CO₂ from the atmosphere into the BSC would increase the $\Delta^{13}\text{C}$ as the exchange of CO₂ between the atmosphere and the BSC would allow Rubisco to fractionate against the heavier carbon isotope. However, a low conductance of CO₂ diffusion across the BSC (g_w mmol m⁻² s⁻¹ Pa⁻¹) is an essential component of the C₄ CO₂-concentrating mechanism and limits the amount of direct fixation of CO₂ under ambient CO₂ concentrations (Jenkins et al., 1989; Brown and Byrd, 1993; He and Edwards, 1996; von Caemmerer, 2000; Kiirats et al., 2002). Therefore, even when the initial carboxylation reaction of the C₄ pump is limited by low CA activity or low light, there would be little, if any, direct fixation of CO₂ in the BSC and minimal influence on $\Delta^{13}\text{C}$.

Alternatively, because ¹³C concentrates in HCO₃⁻ and Rubisco preferentially fixes ¹²C, leakage of HCO₃⁻ out of the BSC would change the fractionation factor associated with CO₂ leakage from the BSC (s from Eq. 2). However, with the relatively low CA activity in the BSC, it is unlikely that CO₂ and HCO₃⁻ would be in full isotopic equilibrium and there would be little influence on s (Farquhar, 1983; von Caemmerer et al., 1997a; Ludwig et al., 1998). Additionally, the influence of respiration and photorespiration on the modeled value of $\Delta^{13}\text{C}$ will increase as the rates of net CO₂ assimilation decrease. However, changing the fractionation effect of respiration (e in Eqs. 3 and 4) and photorespiration (f in Eq. 4) to a range of values reported in the literature (Ghashghaie et al., 2003) had only a slight influence on the modeled $\Delta^{13}\text{C}$ even at low photosynthetic rates (Fig. 6c).

As previously mentioned, Equation 3 simplifies to $b_4 = -5.7 + 7.9 V_p/V_h$ at 25°C when the catalyzed fractionation values for e_b of -9.0‰ and h of 1.1‰ are used. However, if the interconversion of CO₂ and HCO₃⁻ occurs via the spontaneous uncatalyzed reaction, e_b and h become -7.8‰ and 6.9‰, respectively, and Equation 3 is $b_4 = -4.5 + 12.5 V_p/V_{hr}$, causing the b_4 value to become larger, leading to an increase in $\Delta^{13}\text{C}$ (Fig. 4). The catalyzed and uncatalyzed values of e_b and h are taken from previously published work on the hydration and dehydration of CO₂ and HCO₃⁻ (Mook et al., 1974; Marlier and O'Leary, 1984; Paneth and O'Leary, 1985). The proportion of catalyzed to uncatalyzed hydration/dehydration reactions may have an influence on the $\Delta^{13}\text{C}$ when the photosynthetic rates are extremely low, such as in the anti-CA plants with extremely low CA activity, but it would have little, if any, influence in wild-type plants.

Carbon Isotope Discrimination Increases at Low Light

The response of $\Delta^{13}\text{C}$ in C_4 plants to various light levels has not been well characterized, but is an important factor to consider when interpreting dry matter $\delta^{13}\text{C}$ of plants exposed to different light environments or leaves within a canopy. The increase in $\Delta^{13}\text{C}$ and estimated values of ϕ in *F. bidentis* (Fig. 2) are similar to earlier reports that showed that $\Delta^{13}\text{C}$ generally increases as the PFD decreases (Henderson et al., 1992; Peisker and Henderson, 1992; Tazoe et al., 2005). Buchmann et al. (1996) also showed that $\Delta^{13}\text{C}$, calculated from leaf $\delta^{13}\text{C}$ values, in a number of C_4 plants was greater at low PFD. The low conductance of CO_2 diffusion across the BSC needed for C_4 photosynthesis would limit the direct fixation of CO_2 by Rubisco, even under low light, and its influence on $\Delta^{13}\text{C}$ should be minimal. However, it has been demonstrated with the C_4 photosynthetic model that ϕ increases at low PFD as more electron transport is needed for recycling of photorespired CO_2 (von Caemmerer, 2000). The predicted change in ϕ at low PFD by the C_4 photosynthetic model is consistent with our current experimental evidence, as well as earlier published results (Henderson et al., 1992). The evidence from both online and dry matter isotope measurements indicates that growth light conditions need to be considered when interpreting carbon isotope discrimination in C_4 plants.

CONCLUSION

CA activity in wild-type *F. bidentis* appears to be in excess to maintain net CO_2 assimilation; however, reducing leaf CA activity had a relatively large influence on $\Delta^{13}\text{C}$, often without changes in net CO_2 assimilation. The influence of CA activity on $\Delta^{13}\text{C}$ was also evident in the leaf dry matter $\delta^{13}\text{C}$. The model of $\Delta^{13}\text{C}$ developed by Farquhar (1983) predicted the influence of changes in PEPC carboxylation relative to the hydration reaction of CO_2 (V_p/V_h) on $\Delta^{13}\text{C}$, except when photosynthetic rates and CA activity were dramatically reduced. It will be important to take the extent of CA activity in C_4 leaves into account when using $\Delta^{13}\text{C}$ and/or $\delta^{13}\text{C}$ to model leaf level and global C_4 photosynthesis in response to changing environmental influences. The influence of environmental conditions on leaf CA activity, V_p/V_h , and thus on $\Delta^{13}\text{C}$ warrants further investigation.

Additionally, the amount of CA activity in a leaf plays an important role in determining C^{18}O discrimination during C_4 photosynthesis because CA enhances the rate of oxygen exchange between CO_2 and leaf H_2O and thus determines the extent of isotopic equilibrium. The anti-CA plants will be used to test whether changing leaf CA activity influences C^{18}O discrimination under similar environmental conditions and whether high CA activity, relative to photosynthetic rates, corresponds to complete isotopic equilibrium between CO_2 and leaf H_2O as predicted.

MATERIALS AND METHODS

Growth Conditions

Flaveria bidentis plants were previously transformed with antisense RNA constructs targeted to either the nuclear-encoded gene for the small subunit of Rubisco (anti-SSu plants) or a putative cytosolic CA (anti-CA plants; Furbank et al., 1996; von Caemmerer et al., 1997b, 2004). The segregating T_1 generations of anti-CA primary transformants with photosynthetic rates similar to wild type were grown during the summer months in a glasshouse under natural light conditions (27°C d/ 18°C night temperatures). Anti-CA and anti-SSu plants (segregating T_2 generation from primary transformant 136-13) with low photosynthetic capacities and wild-type plants were grown under 1% CO_2 in a controlled environment growth cabinet at a photosynthetic PFD of $400 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ at plant height and air temperature of 27°C during the day and 18°C at night with a 14-h daylength. Three plants with very low CA and photosynthetic rates were germinated and grown for several weeks in the glasshouse. Subsequently, these plants were transferred to the 1% CO_2 growth cabinets before leaf gas-exchange measurements were made. Plants were grown in 5-L pots in garden mix with 2.4 to 4 g Osmocote/L soil (15/4.8/10.8/1.2 N/P/K/Mg + trace elements: B, Cu, Fe, Mn, Mo, Zn; Scotts Australia Pty Ltd.) and watered daily.

Gas-Exchange Measurements

Plants from either the glasshouse or growth cabinet were transferred to the gas-exchange system, where one of the uppermost fully expanded leaves was placed into the leaf chamber of the LI-6400 and allowed to equilibrate at a leaf temperature of 30°C and $2,000 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ for a minimum of 1.5 h. Air entering the leaf chamber was prepared by using mass flow controllers (MKS Instruments) to obtain a gas mix of 90.5 kPa of dry nitrogen and 4.8 kPa oxygen (Fig. 1). A portion of the nitrogen/oxygen mixture was used to zero the mass spectrometer to correct for N_2O and other contaminants contributing to the 44 and 45 peaks. Pure CO_2 ($\delta^{13}\text{C} = -29\text{‰}$; VPDB) was added to the remaining airstream to obtain a CO_2 partial pressure of approximately 52 Pa. Alternatively, some measurements were made by mixing pure CO_2 with CO_2 -free air and using the CO_2 -free air as a zero.

The different gas mixtures had no apparent influence on leaf gas exchange or ^{13}C isotope discrimination. Low oxygen (4.8 kPa) was used to minimize contamination of the 46 peak caused by the interaction of O_2 and N_2 to produce NO_2 with the mass spectrometer source element. This was important when looking at C^{18}O discrimination (A.B. Cousins, M.R. Badger, and S. von Caemmerer, unpublished data). The CO_2 used during the gas-exchange measurements had a similar isotopic signature to the CO_2 in the high CO_2 growth cabinet. This minimized the influence of respired CO_2 on the $\Delta^{13}\text{C}$ measurements in plants with low photosynthetic rates.

The gas mixtures were fed to the inlet of the LI-6400 console and a flow rate of $200 \mu\text{mol s}^{-1}$ was maintained over the leaf. The remaining airstream was vented or used to determine the isotopic composition of air entering the leaf chamber (Fig. 1). The efflux from the leaf chamber was measured by either replacing the match valve line with a line connected directly to the mass spectrometer or by placing a tee in the match valve line, allowing flow to both the mass spectrometer and the match valve simultaneously. Gas-exchange parameters were determined by the LI-6400, and $p\text{CO}_2$ leaving the chamber was subsequently corrected for the dilution of CO_2 by water vapor (von Caemmerer and Farquhar, 1981).

Isotopic Measurements

The efflux from the leaf chamber and the gas mix supplied to the LI-6400 system was linked to a mass spectrometer through an ethanol/dry ice water trap and a thin, gas-permeable silicone membrane, which was housed in a temperature-controlled cuvette. Masses 44 and 45 were monitored continuously and the carbon isotope discrimination during CO_2 exchange, $\Delta^{13}\text{C}$, was calculated from the ratio of mass 45 to 44 in the reference air, determined before and after each sample measurement, entering the chamber (R_e), and the composition of the sample air leaving the leaf chamber (R_s) as described by Evans et al. (1986):

$$\Delta = \frac{-\xi(R_e/R_o - 1)}{1 + \xi(R_e/R_o - 1)}, \quad (1)$$

where $\xi = p_e/(p_e - p_o)$, and p_e and p_o are the CO_2 partial pressures of dry air entering and leaving the leaf chamber, respectively. A summary of the

symbols used in the text is listed in Table I. Zero values for the 44 and 45 peaks were determined before and after the sample measurements were subtracted from both the sample and reference measurements prior to determining the mass ratios. The zero values were typically 1% of the 44 and 45 peaks at 4.8 kPa oxygen and 2% at 20 kPa oxygen.

Calculations of Carbon Isotope Discrimination

The model of C₄ carbon isotope discrimination ($\Delta^{13}\text{C}$) of Farquhar (1983) was used to determine which factors in the model would influence $\Delta^{13}\text{C}$ consistent with our experimental data. The simplified model predicts that:

$$\Delta^{13}\text{C} = a + (b_4 + (b_3 - s)\phi - a)p_i/p_a, \quad (2)$$

where a (4.4‰) is the fractionation during diffusion of CO₂ and s (1.8‰) is the fractionation during CO₂ leakage from the BSCs. The combined fractionation of PEPC, respiration, and the isotopic equilibrium during dissolution of CO₂ and conversion to HCO₃⁻ (b_4) is calculated as (Farquhar, 1983):

$$b_4 = (b_p + e_s + e_b)(1 - V_p/V_h) + (e_s + h)V_p/V_h - eM_m/V_p, \quad (3)$$

where b_p (2.2‰) is the fractionation by PEPC (O'Leary, 1981), e_s (1.1‰) is the fractionation as CO₂ dissolves (O'Leary, 1984), and e_b (-9‰) is the equilibrium fractionation factor of the catalyzed hydration/dehydration reactions of CO₂ and HCO₃⁻ (Mook et al., 1974). Alternatively, during the hydration/dehydration reactions, the uncatalyzed equilibrium fractionation factor $e_b = -7.8‰$ (Marlier and O'Leary, 1984). The fractionation when CO₂ and HCO₃⁻ are not at equilibrium is dependent on the rate of CO₂ hydration (V_h), the rate of PEPC (V_p), e_s , and the catalyzed fractionation during CO₂ hydration (h). The catalyzed hydration reaction has a fractionation factor of 1.1‰ (calculated by summing the catalyzed CO₂ and HCO₃⁻ equilibrium fractionation factor -9.0‰ and the catalyzed dehydration fractionation factor 10.1‰; Mook et al., 1974; Paneth and O'Leary, 1985), whereas the uncatalyzed reaction has a 6.9‰ fractionation factor (Marlier and O'Leary, 1984). The fractionation attributed to mitochondrial respiration is e at a rate of mesophyll CO₂ release of M_m .

The combined fractionation of Rubisco (30‰), respiration, and photorespiration (b_3) can be calculated as:

$$b_3 = 30 - s - e(M_m + M_s)/V_c - fV_o/V_c, \quad (4)$$

where V_c is the rate of Rubisco carboxylation reaction, M_s is the rate of BSC mitochondrial respiration, V_o is the rate of photorespiration, and f is the discrimination of photorespiration (Farquhar, 1983).

Equation 2 assumes that the internal conductance to the diffusion of CO₂ between the intercellular airspace and the site of carboxylation in the mesophyll cytoplasm (g_w) is large, such that p_i is equal to the $p\text{CO}_2$ at the site of PEPC carboxylation (p_c). If g_w is low, then Equation 2 can be modified to:

$$\Delta^{13}\text{C} = a + (b_4 + (b_3 - s)\phi - a)p_i/p_a + A/(g_w p_a)(e_s + a_1 - b_4 - (b_3 - s)\phi), \quad (5)$$

where A is the net rate of CO₂ assimilation and a_1 (0.7‰) is the fractionation of CO₂ diffusion through a liquid (O'Leary, 1984).

CA Activity Measurements

CA activity was measured on leaf extracts using mass spectrometry to measure the rates of ¹⁸O₂ exchange on doubly labeled ¹³C¹⁸O₂ to H₂¹⁶O (Badger and Price, 1989; von Caemmerer et al., 2004). Measurements of leaf extracts were made at 25°C with a subsaturating total carbon concentration of 1 mM. The hydration rates were calculated from the enhancement in the rate of ¹⁸O loss over the uncatalyzed rate. We then applied this factor to the nonenzymatic first-order rate constant calculated at pH 7.4 appropriate for the mesophyll cytosol (Furbank et al., 1989) and report the CA activity as a first-order rate constant k_{CA} (mol m⁻² s⁻¹ Pa⁻¹). $k_{\text{CA}}p_m$ then gives the in vivo CA activity at that particular cytosolic $p\text{CO}_2$. Leaf samples were collected after the gas-exchange measurements on the same leaf material and subsequently frozen in liquid nitrogen and stored at -80°C.

Dry Matter $\delta^{13}\text{C}$

The opposite leaf to the one used during gas exchange was collected and oven dried at 70°C, and ground with a mortar and pestle. A subsample of

ground tissue was weighed and the isotopic composition determined by combustion in a Carlo Erba elemental analyzer; the CO₂ was analyzed by mass spectrometry. δ was calculated as $[(R_{\text{sample}} - R_{\text{standard}})/R_{\text{standard}}] \times 1,000$, where R_{sample} and R_{standard} are the ¹³C/¹²C of the sample and the standard VPDB, respectively. Dry matter $\delta^{13}\text{C}$ was determined on glasshouse-grown plants only because there were large fluctuations in the carbon isotopic composition of the air in the growth cabinets.

Photosynthetic Model

The C₄ photosynthetic model developed by Berry and Farquhar (1978) and von Caemmerer (2000) was used to predict the response of net CO₂ assimilation, bundle sheath $p\text{CO}_2$, p_i/p_a , photorespiration, and ϕ to changes in the amount of PEPC activity (V_p). Manipulating V_p within the photosynthesis model was used to simulate the effect of changes in CO₂ hydration rates (V_h). The outputs from the C₄ photosynthetic model, specifically the rates of Rubisco carboxylation (V_c), V_o , V_p , ϕ , and the $p\text{CO}_2$ in the BSC, were incorporated into the model of C₄ carbon isotope discrimination ($\Delta^{13}\text{C}$) developed by Farquhar (1983). The $\Delta^{13}\text{C}$ model was used to determine which photosynthetic parameters would influence $\Delta^{13}\text{C}$ consistent with our experimental data.

ACKNOWLEDGMENTS

We thank Sue Wood for the carbon isotope analysis of dry matter samples and Howard Griffiths for his helpful comments on earlier versions of this manuscript.

Received January 25, 2006; revised March 12, 2006; accepted March 13, 2006; published March 16, 2006.

LITERATURE CITED

- Ainsworth EA, Long SP (2005) What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy. *New Phytol* **165**: 351–371
- Badger MR, Price GD (1989) Carbonic-anhydrase activity associated with the *Cyanobacterium Synechococcus* Pcc7942. *Plant Physiol* **89**: 51–60
- Berry JA, Farquhar GD (1978) The CO₂-concentrating function of C₄ photosynthesis. A biochemical model. In DO Hall, J Coombs, TW Goodwin, eds, *Proceedings of the Fourth International Congress on Photosynthesis*. Biochemical Society, London, pp 119–131
- Brown RH, Byrd GT (1993) Estimation of bundle-sheath cell conductance in C₄ species and O₂ insensitivity of photosynthesis. *Plant Physiol* **103**: 1183–1188
- Brownell P, Bielig L, Grof C (1991) Increased carbonic anhydrase activity in leaves of sodium-deficient C₄ plants. *Aust J Plant Physiol* **18**: 589–592
- Buchmann N, Brooks JR, Rapp KD, Ehleringer JR (1996) Carbon isotope composition of C₄ grasses is influenced by light and water supply. *Plant Cell Environ* **19**: 392–402
- Burnell JN, Suzuki I, Sugiyama T (1990) Light induction and the effect of nitrogen status upon the activity of carbonic anhydrase maize leaves. *Plant Physiol* **94**: 384–387
- Cervigni T, Teofani F, Bassanelli C (1971) Effect of CO₂ on carbonic anhydrase in *Avena sativa* and *Zea mays*. *Phytochemistry* **10**: 2991–2994
- Evans JR, Sharkey TD, Berry JA, Farquhar GD (1986) Carbon isotope discrimination measured concurrently with gas-exchange to investigate CO₂ diffusion in leaves of higher-plants. *Aust J Plant Physiol* **13**: 281–292
- Farquhar GD (1983) On the nature of carbon isotope discrimination in C₄ species. *Aust J Plant Physiol* **10**: 205–226
- Flanagan LB, Ehleringer JR (1998) Ecosystem-atmosphere CO₂ exchange: interpreting signals of change using stable isotope ratios. *Trends Ecol Evol* **13**: 10–14
- Furbank RT, Chitty JA, von Caemmerer S, Jenkins CLD (1996) Antisense RNA inhibition of RbcS gene expression reduces Rubisco level and photosynthesis in the C₄ plant *Flaveria bidentis*. *Plant Physiol* **111**: 725–734
- Furbank RT, Jenkins CLD, Hatch MD (1989) CO₂ concentrating mechanism of C₄ photosynthesis—permeability of isolated bundle sheath-cells to inorganic carbon. *Plant Physiol* **91**: 1364–1371

- Ghannoum O, von Caemmerer S, Ziska LH, Conroy JP** (2000) The growth response of C_4 plants to rising atmospheric CO_2 partial pressure: a reassessment. *Plant Cell Environ* **23**: 931–942
- Ghoshgaie J, Badeck F-W, Lanigan G, Nogue S, Tcherkez G, Deleens E, Cornic G, Griffiths H** (2003) Carbon isotope fractionation during dark respiration and photorespiration in C_3 plants. *Phytochemistry Reviews* **2**: 145–161
- Gillon J, Yakir D** (2001) Influence of carbonic anhydrase activity in terrestrial vegetation on the O-18 content of atmospheric CO_2 . *Science* **291**: 2584–2587
- Gillon JS, Griffiths H** (1997) The influence of (photo)respiration on carbon isotope discrimination in plants. *Plant Cell Environ* **20**: 1217–1230
- Gillon JS, Yakir D** (2000) Naturally low carbonic anhydrase activity in C_4 and C_3 plants limits discrimination against (COO)-O-18 during photosynthesis. *Plant Cell Environ* **23**: 903–915
- Hatch MD** (1987) C_4 photosynthesis—a unique blend of modified biochemistry, anatomy and ultrastructure. *Biochim Biophys Acta* **895**: 81–106
- Hatch MD, Burnell JN** (1990) Carbonic anhydrase activity in leaves and its role in the first step of C_4 photosynthesis. *Plant Physiol* **93**: 825–828
- He DX, Edwards GE** (1996) Estimation of diffusive resistance of bundle sheath cells to CO_2 from modeling of C_4 photosynthesis. *Photosynth Res* **49**: 195–208
- Henderson SA, von Caemmerer S, Farquhar GD** (1992) Short-term measurements of carbon isotope discrimination in several C_4 species. *Aust J Plant Physiol* **19**: 263–285
- Jenkins CLD, Furbank RT, Hatch MD** (1989) Inorganic carbon diffusion between C_4 mesophyll and bundle sheath-cells—direct bundle sheath CO_2 assimilation in intact leaves in the presence of an inhibitor of the C_4 pathway. *Plant Physiol* **91**: 1356–1363
- Kanai R, Edwards GE** (1999) The biochemistry of C_4 photosynthesis. In R Sage, R Monson, eds, *C_4 Plant Biology*. Academic Press, San Diego, pp 49–87
- Kiirats O, Lea PJ, Franceschi VR, Edwards GE** (2002) Bundle sheath diffusive resistance to CO_2 and effectiveness of C_4 photosynthesis and refixation of photorespired CO_2 in a C_4 cycle mutant and wild-type *Amaranthus edulis*. *Plant Physiol* **130**: 964–976
- Leakey ADB, Uribelarrea M, Ainsworth EA, Naidu SL, Rogers A, Ort DR, Long SP** (2006) Photosynthesis, productivity, and yield of maize are not affected by open-air elevation of CO_2 concentration in the absence of drought. *Plant Physiol* **140**: 779–790
- Ludwig M, von Caemmerer S, Price GD, Badger MR, Furbank RT** (1998) Expression of tobacco carbonic anhydrase in the C_4 dicot *Flaveria bidentis* leads to increased leakiness of the bundle sheath and a defective CO_2 -concentrating mechanism. *Plant Physiol* **117**: 1071–1081
- Marlier J, O'Leary M** (1984) Carbon kinetic isotope effects on the hydration of carbon dioxide and dehydration of bicarbonate ion. *J Am Chem Soc* **106**: 5054–5057
- McLeod AR, Long SP** (1999) Free-air carbon dioxide enrichment (FACE) in global change research: a review. *Adv Ecol Res* **28**: 1–56
- Meinzer FC, Plaut Z, Saliendra NZ** (1994) Carbon isotope discrimination, gas-exchange, and growth of sugarcane cultivars under salinity. *Plant Physiol* **104**: 521–526
- Meinzer FC, Saliendra NZ** (1997) Spatial patterns of carbon isotope discrimination and allocation of photosynthetic activity in sugarcane leaves. *Aust J Plant Physiol* **24**: 769–775
- Meinzer FC, Zhu J** (1998) Nitrogen stress reduces the efficiency of the C_4 CO_2 concentrating system, and therefore quantum yield, in Saccharum (sugarcane) species. *J Exp Bot* **49**: 1227–1234
- Mook W, Bommerson J, Staverman W** (1974) Carbon isotope fractionation between dissolved bicarbonate and gaseous carbon dioxide. *Earth Planet Sci Lett* **22**: 169–176
- O'Leary M** (1981) Carbon isotope fractionation in plants. *Phytochemistry* **20**: 553–567
- O'Leary M** (1984) Measurement of the isotope fractionation associated with the diffusion of carbon dioxide in aqueous solution. *J Phys Chem* **88**: 823–825
- Paneth P, O'Leary M** (1985) Carbon isotope effect on dehydration of bicarbonate ion catalyzed by carbonic anhydrase. *Biochemistry* **24**: 5143–5147
- Peisker M** (1982) The effect of CO_2 leakage from bundle sheath-cells on carbon isotope discrimination in C_4 plants. *Photosynthetica* **16**: 533–541
- Peisker M, Henderson SA** (1992) Carbon—terrestrial C_4 plants. *Plant Cell Environ* **15**: 987–1004
- Ranjith SA, Meinzer FC, Perry MH, Thom M** (1995) Partitioning of carboxylase activity in nitrogen-stressed sugarcane and its relationship to bundle sheath leakiness to CO_2 , photosynthesis and carbon isotope discrimination. *Aust J Plant Physiol* **22**: 903–911
- Saliendra NZ, Meinzer FC, Perry M, Thom M** (1996) Associations between partitioning of carboxylase activity and bundle sheath leakiness to CO_2 , carbon isotope discrimination, photosynthesis, and growth in sugarcane. *J Exp Bot* **47**: 907–914
- Tazoe Y, Noguchi K, Terashima I** (2005) Effects of growth light and nitrogen nutrition on the organization of the photosynthetic apparatus in leaves of a C_4 plant, *Amaranthus cruentus*. *Plant Cell Environ* doi/10.1111/j.1365-3040.2005.01453.x
- von Caemmerer S** (2000) *Biochemical Models of Leaf Photosynthesis*. CSIRO Publishing, Collingwood, Australia
- von Caemmerer S, Farquhar GD** (1981) Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. *Planta* **153**: 376–387
- von Caemmerer S, Ludwig M, Millgate A, Farquhar GD, Price D, Badger M, Furbank RT** (1997a) Carbon isotope discrimination during C_4 photosynthesis: insights from transgenic plants. *Aust J Plant Physiol* **24**: 487–494
- von Caemmerer S, Millgate A, Farquhar GD, Furbank RT** (1997b) Reduction of ribulose-1,5-bisphosphate carboxylase/oxygenase by antisense RNA in the C_4 plant *Flaveria bidentis* leads to reduced assimilation rates and increased carbon isotope discrimination. *Plant Physiol* **113**: 469–477
- von Caemmerer S, Quinn V, Hancock NC, Price GD, Furbank RT, Ludwig M** (2004) Carbonic anhydrase and C_4 photosynthesis: a transgenic analysis. *Plant Cell Environ* **27**: 697–703
- Wall GW, Brooks TJ, Adam R, Cousins AB, Kimball BA, Pinter PJ, LaMorte RL, Triggs L, Ottman MJ, Leavitt SW, et al** (2001) Elevated atmospheric CO_2 improved Sorghum plant water status by ameliorating the adverse effects of drought. *New Phytol* **152**: 231–248
- Watling J, Press M, Quick W** (2000) Elevated CO_2 induces biochemical and ultrastructural changes in leaves of the C_4 cereal sorghum. *Plant Physiol* **123**: 1143–1152
- Williams DG, Gempko V, Fravolini A, Leavitt SW, Wall GW, Kimball PA, Pinter PJ, LaMorte R** (2001) Carbon isotope discrimination by Sorghum bicolor under CO_2 enrichment and drought. *New Phytol* **150**: 285–293
- Yakir D, da SL Sternberg L** (2000) The use of stable isotopes to study ecosystem gas exchange. *Oecologia* **123**: 297–311