

Expression of the thyroid hormone receptor gene, $erbA\alpha$, in B lymphocytes: alternative mRNA processing is independent of differentiation but correlates with antisense RNA levels

Michelle L. Hastings, Christine Milcarek¹, Kathlee Martincic¹, Martha L. Peterson² and Stephen H. Munroe*

Department of Biology, Marquette University, Milwaukee, WI 53233, USA, ¹Department of Molecular Genetics and Biochemistry, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261, USA and ²Department of Pathology and Laboratory Medicine, University of Kentucky College of Medicine, Lexington, KY 40536, USA

Received July 8, 1997; Revised and Accepted September 10, 1997

ABSTRACT

The $erbA\alpha$ gene encodes two α -thyroid hormone receptor isoforms, TR α 1 and TR α 2, which arise from alternatively processed mRNAs, $erbA\alpha$ 1 (α 1) and $erbA\alpha$ 2 (α 2). The splicing and alternative polyadenylation patterns of these mRNAs resemble that of mRNAs encoding different forms of immunoglobulin heavy chains, which are regulated at the level of alternative processing during B cell differentiation. This study examines the levels of $erbA\alpha$ mRNA in eight B cell lines representing four stages of differentiation in order to determine whether regulation of the alternatively processed α 1 and α 2 mRNAs parallels the processing of immunoglobulin heavy chain mRNAs. Results show that the pattern of α 1 and α 2 mRNA expression is clearly different from that observed for immunoglobulin heavy chain mRNAs. B cell lines display characteristic ratios of α 1/ α 2 mRNA at distinct stages of differentiation. Furthermore, expression of an overlapping gene, Rev-Erb α (RevErb), was found to correlate strongly with an increase in the ratio of α 1/ α 2 mRNA. These results suggest that alternative processing of $erbA\alpha$ mRNAs is regulated by a mechanism which is distinct from that regulating immunoglobulin mRNA. The correlation between RevErb and $erbA\alpha$ mRNA is consistent with negative regulation of α 2 via antisense interactions with the complementary RevErb mRNA.

INTRODUCTION

Thyroid hormone receptors (TRs) mediate the cellular response to thyroid hormone (T3) by regulating target gene transcription (1–3). In all vertebrates TRs are the products of two different genes, $erbA\alpha$ and $erbA\beta$ (4). The mammalian $erbA\alpha$ gene produces two mRNAs, $erbA\alpha$ 1 (α 1) and $erbA\alpha$ 2 (α 2), through

alternative processing of the 3'-end of its pre-mRNA transcript (5–7). These mRNAs give rise to receptor isoforms with antagonistic functions. α 1 codes for the α -thyroid hormone receptor (TR α 1), whereas α 2 codes for an orphan nuclear receptor (TR α 2) which does not bind T3 (6,7). TR α 2 competes with TR α 1 and TR β for specific DNA binding sites, thereby antagonizing T3 action (8). Because the $erbA\alpha$ gene produces both a transcriptional activator (TR α 1) and its specific inhibitor (TR α 2), regulation of the alternative processing of α 1 and α 2 mRNA may provide an important mechanism for determining the cellular response to thyroid hormone.

Alternative processing of the 3'-end of $erbA\alpha$ RNA transcripts involves competition between splicing and polyadenylation. Polyadenylation at an upstream site yields α 1 mRNA, whereas processing from a 5' splice site (ss) within the final exon of α 1 to a downstream 3' ss produces α 2 mRNA. The levels of α 1 and α 2 vary in different tissues and at different developmental stages (9,10). However, the mechanisms which regulate expression of the alternatively processed mRNAs are not well understood.

Two general models have been described for the regulation of alternative splicing and polyadenylation (11). In some cases alternative processing is regulated by transcript-specific factors which alter the efficiency of either splicing or polyadenylation. In other instances, the activity of one or more constitutive components of the mRNA processing apparatus is altered. In the latter case, the processing of many unrelated transcripts may be affected.

The organization of the $erbA\alpha$ gene and the alternative mRNA processing of its transcripts appear similar to that of the immunoglobulin (Ig) heavy chain genes. The Ig heavy chain gene produces two functionally distinct mRNAs which encode heavy chains for the membrane-bound (mb) and secreted (sec) forms of Ig. Like α 1, Ig sec mRNA processing utilizes an upstream polyadenylation site. Similar to α 2, Ig mb mRNA processing utilizes a 5' ss in the last common exon to splice to a downstream exon. Regulation of Ig heavy chain mRNA processing in B lymphocytes has been well characterized. In early stages of B cell

*To whom correspondence should be addressed. Tel: +1 414 288 1485; Fax: +1 414 288 7357; Email: munroes@vms.csd.mu.edu

development downstream splicing of Ig mb mRNA is equal to or greater than that of Ig sec mRNA processing. At later stages upstream polyadenylation of Ig sec mRNA predominates (12). Several studies suggest that the balance between splicing and polyadenylation required for processing of Ig mb and sec mRNAs is regulated by a change in the level of a general polyadenylation factor (13–16). In view of these findings, it is possible that erbA α mRNA processing may parallel that of Ig heavy chain mRNA during B cell development.

A distinguishing feature of the mammalian erbA α gene locus is the presence of a third gene, Rev-ErbA α (RevErb), encoded on the DNA strand opposite erbA α (17,18). The 3' exon of RevErb overlaps with the α 2-specific 3' exon but not with α 1 sequence. The unusual organization of these genes results in α 2 and RevErb mRNAs which are complementary at their 3'-ends allowing the possible formation of antisense/sense RNA duplexes. Such basepairing interactions between the RevErb and α 2 mRNAs could negatively regulate α 2 mRNA levels, and therefore offer a transcript-specific mechanism by which α 1 and α 2 mRNA levels are regulated.

In this study, we examine the expression of α 1 and α 2 mRNA from B cells representing different stages of differentiation. We find that the thyroid hormone receptor, TR α 1, and the orphan receptors TR α 2 and RevErb are expressed at all stages of differentiation but at varying levels in the different cell lines. Our results indicate that the regulation of alternative RNA processing of α 1 and α 2 is distinct from that of Ig sec and mb RNA processing regulation. However, changes in the relative levels of α 1 and α 2 correlate strongly with variations in levels of RevErb mRNA.

MATERIALS AND METHODS

Cell lines

All cell lines are from the mouse B cell lineage and were grown as previously described (13,14,19,20). 70Z/3.12 represents a pre-B cell stage line with equal amounts of sec and mb IgM mRNA (20). WEHI-231 is an early B cell with equal amounts of sec and mb IgM (20). The M12 cell line represents an early B cell which has lost its endogenous heavy chain but expresses approximately equal amounts of sec and mb IgM mRNA when transfected with an IgM gene (13). The A20 and 2PK3 cell lines are memory B cells and produce about equal quantities of sec and mb IgG heavy chain mRNA (14). 4T001 is a plasmacytoma cell line which secretes large amounts of γ 2b, κ molecules of IgG (14). S194 is a plasmacytoma cell line which has lost its endogenous heavy chain but produces a large excess of sec over mb mRNA when transfected with an IgM gene (13). J558L is a plasmacytoma cell line which has lost its endogenous α heavy chain but when transfected with an IgG gene sec mRNA is expressed in excess over mb mRNA (19).

Recombinant plasmids and RNA probes

The erbA α probe used for northern analysis was prepared from a 600 nt *Xba*I–*Sac*I DNA fragment excised from plasmid p α 2HN. p α 2HN contains an *Eco*RI–*Hinc*II fragment which is common to both α 1 and α 2. This fragment was isolated from plasmid p α 2 Δ C–*Sac*/stop (21) and subcloned in pBluescript KS⁺ (Stratagene). The probe was uniformly labeled with [α -³²P]dCTP by random oligonucleotide-primed synthesis (22). For RNase protection assays, a single-stranded antisense riboprobe of α 1 and

α 2 common sequence was prepared from pB3EOP which contains the 162 nt *Pst*I–*Eco*0109I erbA α DNA fragment cloned between the *Apa*I–*Pst*I sites of pBluescript KS⁺. This fragment of the erbA α gene spans the α 2-specific 5' ss within the 3'-most exon of α 1. pB3EOP was linearized with *Xba*I and transcribed with T3 RNA polymerase to produce a 209 nt probe. Of this probe, 162 nt are complementary to α 1 mRNA and 135 nt are complementary to α 2 mRNA. The RevErb riboprobe was made from pB4E6 which contains 303 nt of a *Bgl*III–*Bsu*36I fragment from pB4-1 (17) cloned into pBluescript KS⁺. pB4E6 was linearized with *Xba*I and transcribed with T3 RNA polymerase to produce a 360 nt probe, 130 nt of which are complementary to RevErb exon 6. RNA probes were uniformly labeled with [α -³²P]UTP and purified by electrophoresis (22). Plasmids p α 2 Δ C–*Sac*/stop and pB4-1 were generously provided by M.A. Lazar, University of Pennsylvania.

RNA isolation and northern analysis

Cytoplasmic RNAs were isolated by detergent lysis and phenol/chloroform extraction as previously described (19). The RNA was passed over oligo(dT) columns to isolate poly(A)⁺ RNA. Electrophoresis of 1–3 μ g of poly(A)⁺ RNA on a 1.0% agarose–0.22 M formaldehyde gel was run in buffer containing 0.22 M formaldehyde. The RNA was transferred by capillary action to Nytran (Schleicher and Schuell, Inc.) after which the RNA was UV-irradiated and hybridized to labeled probes (23). RNA size markers (GIBCO-BRL) were included on the gel and stained with methylene blue after transfer to Nytran (22).

RNase protection assays

Labeled probes were hybridized to 10–30 μ g target RNAs as previously described (22). The resulting hybrids were digested with 1.5 μ l RNase (1 mg/ml RNase A, 20 000 U/ml RNase T1, Ambion, Inc.) at 30°C for 1 hr. Protected RNAs were denatured in formamide and resolved on 5.5% polyacrylamide–urea gels. Band intensities were quantified by radioanalytic scanning with an AMBIS 100 Image Analyzer. Background was subtracted by using regions of identical size located immediately above each experimental band. RevErb/ α 2 ratios were computed from experiments in which probes were prepared in parallel using identical mixes of labeled and unlabeled nucleotides to ensure identical specific activities. The specific activity of the probes ranged from 9×10^8 to 2×10^9 c.p.m./nmol between experiments. Molar ratios were calculated after correcting for length and composition of the protected RNA fragments.

RESULTS

Structural organization and expression of erbA α mRNAs in B cell lines

In mammals, a single locus codes for the two alternatively processed erbA α mRNAs and the overlapping RevErb mRNA (Fig. 1A). To determine if the erbA α mRNAs are expressed in B lymphocytes, we assayed α 1 and α 2 mRNA levels in two mouse B cell lines representing different stages of differentiation. The cell line 70Z/3 is a tumor line arrested in the pre-B cell stage (20) and J558L is a myeloma cell line representing a late-stage B cell or plasma cell line (19). The C6 rat astrocytoma cell line, which is well-characterized for erbA α mRNA levels (9), was also




Figure 1. Expression of *erbA α* and *RevErb* in B cell lines. (A) Structural organization of the overlapping *c-erbA α* locus. Arrows indicate the direction of transcription, boxes represent exons, lines represent introns. Shaded boxes represent 5' and 3' untranslated regions. Vertical, dotted lines demarcate *RevErb* and $\alpha 2$ complementary sequence. The $\alpha 1$, $\alpha 2$ and *RevErb* poly(A) sites are noted. The alternative splicing of the $\alpha 2$ -specific exon is represented by the angled line above the map. Probes used in RNase protection assays are indicated as bars beneath the sequence with length of the protected fragments indicated in nt. (B) Northern blot analysis of 1 μ g 70Z/3 (lane 1), 2 μ g J558L (lane 2) and 3 μ g C6 (lane 3) polyA⁺ cytoplasmic RNA. Blots were hybridized with a probe from the region common to $\alpha 1$ and $\alpha 2$.

analyzed. Northern blot experiments using an *erbA α* specific probe show the expected 5.5 kb $\alpha 1$ mRNA and 2.6 kb $\alpha 2$ mRNA which correspond to the documented sizes of C6 cell $\alpha 1$ and $\alpha 2$ mRNA (Fig. 1B) (17). These results demonstrate that both *erbA α* mRNAs are expressed at early and late stages of B cell differentiation.

Alternative mRNA processing of *erbA α* mRNA

We next asked whether the balance between alternative processing of $\alpha 1$ and $\alpha 2$ mRNA parallels that of Ig *sec* and *mb* mRNA at different stages of B cell differentiation. Cytoplasmic mRNA was analyzed from eight mouse cell lines representing four different stages of B cell development. In addition to the pre-B and plasmacytoma cell lines described above, mRNA from two mature B cell lines [WEHI-231 (20) and M12 (13)], two cell lines representing memory B cells [2PK3 and A20 (14)], and two additional plasma cell lines [4T001 (14) and S194 (13)] were examined to determine the levels of expression of $\alpha 1$ and $\alpha 2$ mRNA. RNase protection assays of mRNA from these eight cell lines show that the pre-B cell line, 70Z/3, has the highest ratio of $\alpha 1/\alpha 2$ mRNA, 3.6, indicating a predominance of upstream polyadenylation over downstream splicing (Fig. 2A and B). The lowest $\alpha 1/\alpha 2$ ratios are seen in the plasma cell lines, S194 and J558L and the mature B cell line, M12, which demonstrates an excess of $\alpha 2$ splicing over polyadenylation. The ratios of $\alpha 1/\alpha 2$ mRNA in the different B cell lines are clearly distinct from those




Figure 2. Differential *erbA α* alternative RNA processing in B cell lines representing various stages of differentiation. (A) RNase protection assays of B cell cytoplasmic mRNA using the *erbA α* and *RevErb*-specific probes shown in Figure 1A. Assays were carried out with 10 μ g J558L RNA, 20 μ g 70Z/3, M12, 4T001, S194 RNA and 30 μ g WEHI-231, 2PK3, A20 RNA. Assays were carried out in parallel except for those shown in lanes 3 and 7 which were analyzed in a separate experiment. Lanes 9 and 10 show results of the same experiment in lanes 4 and 5 except for a 2-fold increase in exposure time. The specific B cell lines and their representative stage in differentiation is indicated above each lane. The protected fragments corresponding to $\alpha 1$, $\alpha 2$ and *RevErb* mRNAs are indicated at left. (B) Comparison of the ratio of $\alpha 1/\alpha 2$ mRNA and Ig *sec/mb* mRNAs (13,14,19,20) is shown schematically.

seen for *sec* and *mb* mRNA from the Ig heavy chain gene (Fig. 2B). For example, the levels of Ig *sec* (polyadenylated) mRNA levels are nearly equal to those of Ig *mb* (spliced) mRNA at early stages of B cell differentiation but increase sharply as cells switch almost exclusively to production of Ig *sec* mRNA in plasma cells. Although changes in the levels of *erbA α* mRNAs during B cell differentiation are opposite from the Ig mRNA alternative processing pattern, B cell lines representing distinct developmental stages display characteristic ratios of $\alpha 1/\alpha 2$ mRNA (Fig. 2B, Table 1).

Expression of *RevErbA α* mRNA in B cells

The complementary, overlapping organization of the *erbA α* and *RevErb* genes may have important implications for the coordinate regulation of these genes. To determine whether the differential expression of *erbA α* RNA processing during B cell differentiation correlates with changes in expression of *RevErb*, we assessed the levels of *RevErb* mRNA in parallel with $\alpha 1$ and $\alpha 2$ mRNA in the same eight B cell lines. *RevErb* mRNA is expressed

in all of the cell lines but at variable levels (Fig. 2A). The highest expression of RevErb is seen in the 70Z/3 mRNA, which shows an 8-fold excess of RevErb over $\alpha 2$. Conversely, S194, J558L and M12 mRNA, the cell lines with the lowest $\alpha 1/\alpha 2$ (1.0–0.8), also have the lowest RevErb/ $\alpha 2$ ratio (0.5–0.9) though their absolute levels of RevErb extend over a broad range. In general, increases in the ratio of RevErb/ $\alpha 2$ correlate strongly with increases in the ratio of $\alpha 1/\alpha 2$ (Fig. 3, Table 1).

DISCUSSION

B lymphocyte cell lines provide a unique system for investigating the molecular basis of developmentally regulated alternative processing. Such processing has been intensively studied for Ig heavy chain genes during B cell differentiation. In this study we show that *erbA α* mRNAs are expressed at different levels in B cell lines representing specific stages of development. The ratio of $\alpha 1/\alpha 2$ mRNA varies in a manner which suggests stage-specific regulation of mRNA levels, but the pattern of expression is clearly different from the variations in *sec/mb* mRNA levels. Characterization of mRNA levels from the overlapping RevErb gene shows that expression of RevErb mRNA correlates with changes in the $\alpha 1/\alpha 2$ ratio. These observations have important implications both for regulation of alternative processing and for the role of thyroid hormone during B cell development.

Molecular analysis of Ig heavy chain mRNA processing has shown that regulation of polyadenylation and splicing depends on changes in the levels of essential processing factors (14,16). Of particular interest to this study is the observation that unrelated pre-mRNAs which undergo similar 3'-end processing are differentially processed throughout B cell differentiation in the same manner as Ig *sec* and *mb* heavy chain mRNAs (13,15). On the basis of these results, *erbA α* mRNA processing would also be predicted to parallel that of Ig heavy chain mRNAs, with the $\alpha 1/\alpha 2$ ratio lowest in pre-B cells, highest in plasmacytoma cells and at intermediate levels in mature and memory B cells. Our results show a different and more complex pattern of variation in the $\alpha 1/\alpha 2$ expression than predicted. Mature B cells and plasma cells both have relatively low $\alpha 1/\alpha 2$ ratios while higher ratios are observed for memory cells and the pre-B cell line. These results suggest that the balance between alternative splicing and




Figure 3. RevErb mRNA expression in B cell lines related to changes in $\alpha 1/\alpha 2$ mRNA. The ratio of RevErb/ $\alpha 2$ mRNA was quantified and compared to $\alpha 1/\alpha 2$ mRNA. Data is presented in a scatter plot with each point representing a specific cell line. The representative developmental stage for each cell line is indicated by specific symbols.

polyadenylation of *erbA α* mRNAs is regulated in a manner independent of Ig heavy chain alternative mRNA processing.

One possible mechanism for regulating expression of *erbA α* mRNAs involves interactions with the RevErb mRNA. Since this overlapping mRNA is complementary to $\alpha 2$, but not $\alpha 1$, variations in RevErb expression may differentially affect $\alpha 1$ and $\alpha 2$ expression. The results presented here are consistent with such a relationship. Despite wide variations in the levels at which these mRNAs are expressed in the eight lymphocyte cell lines analyzed (Table 1), a strong correlation is evident between the ratio of RevErb/ $\alpha 2$ and the ratio of $\alpha 1/\alpha 2$ (Fig. 3). Specifically, when the ratio of RevErb/ $\alpha 2$ is between 0.4 and 1.2, the $\alpha 1/\alpha 2$ ratio is <1.5 , and when the RevErb/ $\alpha 2$ is >2.0 the $\alpha 1/\alpha 2$ ratio is ≥ 2.5 . Thus, an increase from 1.5 to 2.0 in the RevErb/ $\alpha 2$ ratio may represent a threshold associated with a 2- to 3-fold increase in $\alpha 1/\alpha 2$. Since the ratio of RevErb/ $\alpha 2$ varies over a wider range than the ratio of $\alpha 1/\alpha 2$ and there is no evident correlation in the variation of RevErb and $\alpha 2$ levels, interactions between these genes or their mRNA products may be modulated by additional factors.

Table 1. Quantitation of $\alpha 1$, $\alpha 2$ and RevErb mRNA in B cell lines

Cell line	$\alpha 1$ amol/ μ g	$\alpha 2$ amol/ μ g	RevErb amol/ μ g	$\alpha 1/\alpha 2$ (S.D.)	RevErb/ $\alpha 2$ (S.D.)
70Z/3	5.5	1.7	12.9	3.6 (0.3)	7.9 (0.4)
WEHI-231	0.9	0.75	0.65	1.3 (0.2)	1.0 (0.3)
M12	2.3	3.2	1.4	0.82 (0.16)	0.46 (0.04)
2PK3	0.85	0.22	0.5	3.3 (1.0)	2.4 (0.4)
A20	0.6	0.3	0.5	2.5	2.0
4T001	2.4	1.6	1.8	1.5 (0.0)	1.2 (0.5)
S194	2.9	3.1	2.8	1.0 (0.2)	0.91 (0.00)
J558L	8.1	9.6	5.3	0.94 (0.12)	0.63 (0.18)

Levels of *erbA α* and RevErb mRNAs determined by RNase protection assays as shown in Figure 2. A quantitative determination of the $\alpha 1$ and $\alpha 2$ mRNA levels in each of the cell lines was obtained after correction for mRNA loading and specific activity of the protected fragments. The abundance of both *erbA α* mRNAs is $\sim 0.1\%$ that of mRNA from the housekeeping gene GAPDH (data not shown). Ratios of $\alpha 1/\alpha 2$ mRNA represent the average of three or four experiments. Data from the A20 cell line represents a single experiment. Absolute values and RevErb/ $\alpha 2$ ratios are the average of two experiments carried out in parallel with probes of identical specific activity.

The relationship between RevErb expression and erbA α mRNA processing, described above, may reflect either direct or indirect interactions between these genes or their products. Several other studies have suggested that increased expression of RevErb mRNA may result in a decrease in α 2 mRNA relative to α 1 (24–26). Cycloheximide treatment of a particular pituitary tumor cell line results in a coordinate increase in both RevErb and the α 1/ α 2 ratio (24). Similarly, when adipocyte cells are induced to differentiate *in vitro*, an increase in RevErb mRNA and the α 1/ α 2 ratio is observed (25). The increase in α 1/ α 2 observed in both these studies clearly reflects alterations in the processing of erbA α pre-mRNA, since neither the rate of transcription nor the relative stability of α 1 and α 2 are altered (24,25). However, it should be noted that in both cases the treatments used cause multiple, widespread alterations in cellular metabolism, in addition to altering levels of RevErb and erbA α mRNA.

In a third study antisense RNA corresponding to the overlapping portion of RevErb mRNA was shown to efficiently block splicing of erbA α pre-mRNA *in vitro* (26). In this case the increasing levels of the antisense RNA clearly block mRNA splicing. However, the relevance of this *in vitro* system as a model for physiological regulation is not clear. Antisense regulation at the level of alternative mRNA processing may provide a novel mechanism for the regulation of functionally antagonistic nuclear receptors. Although artificial antisense transcripts and oligonucleotides have been employed to repress expression of many genes, naturally occurring antisense regulation has not been well characterized in higher eukaryotes (27–29). The clear correlation between RevErb expression and α 1/ α 2 mRNA levels described here for eight lymphocyte cell lines, and the results of previous studies (24–26), support a direct role for RevErb in the regulation of relative levels of erbA α mRNAs. Rigorous demonstration of a physiologically relevant antisense mechanism will require evidence that base-pairing interactions between α 2 and RevErb transcripts are essential for the negative regulation of α 2-specific splicing *in vivo*.

The differential expression of α 1 and α 2 mRNAs may have important physiological consequences for B cell differentiation. However, little is known regarding the role of the receptor protein during lymphocyte regulation. This study is the first to report the expression of mRNAs for specific TRs in B cells. Earlier studies have shown that T3 stimulates B cell proliferation and differentiation (30,31), which appears to be a result of changes in T3 reception during differentiation (32). One specific role for TRs in B cell differentiation may involve direct interactions of TRs with the transcription factor AP-1, which regulates stage-specific functions required for B cell proliferation and differentiation (33). The activity of AP-1 is modulated by a number of agents including ligand-activated transcription factors which belong to the TR-related subfamily (34). Both TR α and TR β have been shown to interfere with AP-1 activated transcription in a T3-dependent manner via direct protein–protein interactions (35,36). Thus, the activity of AP-1 during B cell differentiation may be modulated by alterations in the levels of TR α 1 and its antagonist, TR α 2.

ACKNOWLEDGEMENTS

We thank Mitch A. Lazar for providing recombinant materials, for helpful discussions and for a critical reading of the manuscript.

This research was supported by National Institutes of Health grants #DK48034 to S.H.M. and #GM50145 to C.M. and a National Science Foundation grant #MCB-9507513 to M.L.P. M.L.H. was supported by a pre-doctoral fellowship from the Arthur J. Schmitt Foundation.

REFERENCES

- Sap, J., Munoz, A., Damm, K., Goldberg, Y., Ghysdael, J., Leutz, A., Beug, H. and Vennstrom, B. (1986) *Nature*, **324**, 635–640.
- Weinberger, C., Thompson, C.C., Ong, E.S., Lebo, R., Gruol, D.J. and Evans, R.M. (1986) *Nature*, **324**, 641–646.
- Evans, R.M. (1988) *Science*, **240**, 889–895.
- Lazar, M.A. (1993) *Endocr. Rev.*, **14**, 184–193.
- Benbrook, D. and Pfahl, M. (1987) *Science*, **238**, 788–791.
- Izumo, S. and Mahdavi, V. (1988) *Nature*, **334**, 539–542.
- Mitsuhashi, T.G., Tennyson, G.E. and Nikodem, V.M. (1988) *Proc. Natl. Acad. Sci. USA*, **85**, 5804–5808.
- Katz, D. and Lazar, M.A. (1993) *J. Biol. Chem.*, **268**, 20904–20910.
- Jannini, E.A., Mitsuhashi, T. and Nikodem, V.M. (1992) *Biochem. Biophys. Res. Comm.*, **184**, 739–745.
- Leonard, J.L., Farwell, A.P., Yen, P.M., Chin, W.W. and Stula, M. (1994) *Endocrinology*, **135**, 548–555.
- Edwards-Gilbert, G., Veraldi, K.L. and Milcarek, C. (1997) *Nucleic Acids Res.*, **25**, 2547–2561.
- Peterson, M.L. (1994) In Snow, E.C. (ed.), *Handbook of B and T Lymphocytes*. Academic Press, San Diego, CA., pp. 321–342.
- Peterson, M.L. (1994) *Mol. Cell. Biol.*, **14**, 7891–7898.
- Edwards-Gilbert, G. and Milcarek, C. (1995) *Mol. Cell. Biol.*, **15**, 6420–6429.
- Matis, S.A., Martincic, K. and Milcarek, C. (1996) *Nucleic Acids Res.*, **24**, 4684–4692.
- Takagaki, Y., Siepelt, R.L., Peterson, M.L. and Manley, J.L. (1996) *Cell*, **87**, 941–952.
- Lazar, M.A., Hodin, R.A., Darling, D.S. and Chin, W.W. (1989) *Mol. Cell. Biol.*, **9**, 1128–1136.
- Miyajima, N., Horiuchi, R., Shibuya, Y., Fukushige, S., Matsubara, K., Toyoshima, K. and Yamamoto, T. (1989) *Cell*, **57**, 31–39.
- Kobrin, B.J., Milcarek, C. and Morrison, S.L. (1986) *Mol. Cell. Biol.*, **6**, 1687–1697.
- Genovese, C. and Milcarek, C. (1990) *Mol. Immunol.*, **17**, 69–81.
- Katz, D., Berrodin, T.J. and Lazar, M.A. (1992) *Mol. Endocrinol.*, **6**, 805–814.
- Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor University Press, Cold Spring Harbor, NY.
- Peterson, M.L., Bryman, M.B., Peiter, M. and Cowan, C. (1994) *Mol. Cell. Biol.*, **14**, 77–86.
- Lazar, M.A., Hodin, R.A., Guemalli, C. and Chin, W.W. (1990) *J. Biol. Chem.*, **265**, 12859–12863.
- Chawla, A. and Lazar, M.A. (1993) *J. Biol. Chem.*, **268**, 16265–16269.
- Munroe, S.H. and Lazar, M.A. (1991) *J. Biol. Chem.*, **266**, 22083–22086.
- Krystal, G.W. (1992) In Erickson, R.P. and Izant, J.G. (eds.), *Gene Regulation: Biology of Antisense RNA and DNA*. Raven Press, Ltd., NY, pp. 11–20.
- Hildebrandt, M. and Nellen, W. (1992) *Cell*, **69**, 197–204.
- Lee, R.C., Feinbaum, R.L. and Ambros, V. (1993) *Cell*, **75**, 843–854.
- Paavonen, T. (1982) *Scand. J. Immunol.*, **15**, 211–215.
- Chatterjee, S. and Chandel, A.S. (1983) *Acta Endocrinol.*, **103**, 95–100.
- Csaba, G. and Suder, F. (1978) *Horm. Metab. Res.*, **14**, 455–456.
- Grant, P.A., Thompson, C.B. and Pettersson, S. (1995) *EMBO J.*, **14**, 4501–4513.
- Pfahl, M. (1993) *Endocr. Rev.*, **14**, 651–658.
- Saatcioglu, F., Bartunek, P., Deng, T., Zenke, M. and Karin, M. (1993) *Mol. Cell. Biol.*, **13**, 3675–3685.
- Zhang, X.-K., Wills, K.N., Husmann, M., Hermann, T. and Pfahl, M. (1991) *Mol. Cell. Biol.*, **11**, 6016–6025.