

Hsk1 kinase is required for induction of meiotic dsDNA breaks without involving checkpoint kinases in fission yeast

Keiko Ogino*, Kouji Hirota[†], Seiji Matsumoto*, Tadayuki Takeda[‡], Kunihiro Ohta[†], Ken-ichi Arai^{§¶}, and Hisao Masai*^{||}

*Genome Dynamics Project and [§]Department of Integrated Life Science, Tokyo Metropolitan Institute of Medical Science, Bunkyo-ku, Tokyo 113-8613, Japan; [†]Genetic System Regulation Laboratory, The Institute of Physical and Chemical Research (RIKEN), Wako, Saitama 351-01, Japan; and [‡]Computational and Experimental Systems Biology Group, The Institute of Physical and Chemical Research (RIKEN), Genomic Sciences Center, Yokohama, Kanagawa 230-0045, Japan

Communicated by Masayasu Nomura, University of California, Irvine, CA, March 30, 2006 (received for review October 8, 2005)

Cdc7 kinase, conserved through evolution, is known to be essential for mitotic DNA replication. The role of Cdc7 in meiotic recombination was suggested in *Saccharomyces cerevisiae*, but its precise role has not been addressed. Here, we report that Hsk1, the Cdc7-related kinase in *Schizosaccharomyces pombe*, plays a crucial role during meiosis. In a *hsk1* temperature-sensitive strain (*hsk1-89*), meiosis is arrested with one nucleus state before meiosis I in most of the cells and meiotic recombination frequency is reduced by one order of magnitude, whereas premeiotic DNA replication is delayed but is apparently completed. Strikingly, formation of meiotic dsDNA breaks (DSBs) are largely impaired in the mutant, and Hsk1 kinase activity is essential for these processes. Deletion of all three checkpoint kinases, namely Cds1, Chk1, and Mek1, does not restore DSB formation, meiosis, or Cdc2 activation, which is suppressed in *hsk1-89*, suggesting that these aberrations are not caused by known checkpoint pathways but that Hsk1 may regulate DSB formation and meiosis. Whereas transcriptional induction of some *rec* genes and horsetail movement are normal, chromatin remodeling at *ade6-M26*, a recombination hotspot, which is prerequisite for subsequent DSB formation at this locus, is not observed in *hsk1-89*. These results indicate unique and essential roles of Hsk1 kinase in the initiation of meiotic recombination and meiosis.

Cdc7-related kinase | checkpoint control | chromatin remodeling | meiosis | meiotic recombination

Although recent studies revealed the presence of a number of conserved factors involved in mitotic DNA replication (1), it is largely unknown whether they play roles in meiotic replication and recombination (2, 3). Studies in fission and budding yeasts indicated essential roles of Cdc2 kinase (1) and Clb5- and Clb6-dependent Cdc28 kinase (4), respectively, in premeiotic DNA replication. Orc, Cdc18, and MCM are also required for premeiotic DNA synthesis (3, 5), although their requirement may not be identical to that during mitotic replication (2). It is not known whether initiation is regulated by a common mechanism or the same set of replication origins is used during premeiotic S phase. In budding yeast, strong coupling of DNA replication and meiotic recombination has been shown (6–9).

Cdc7 kinase plays an essential role in firing replication origins during mitotic growth (10, 11). It was reported previously that meiotic recombination frequency is reduced, whereas premeiotic DNA replication proceeds, in the budding yeast *cdc7^{ts}* mutants and that Cdc7 may play a role in synaptonemal complex formation during meiosis, a step stabilizing homologous pairing during meiotic recombination (12, 13), but the precise meiotic roles of Cdc7 have remained elusive. We report here that Hsk1, the fission yeast homologue of Cdc7 kinase, is required for efficient meiotic recombination and meiotic nuclear division. During meiosis, Hsk1 is required specifically for formation of dsDNA breaks (DSBs), an early event for meiotic recombination. The

loss of DSB and meiosis are not corrected by loss of three major checkpoint kinases of fission yeast, suggesting that known checkpoint pathways are not involved. Our results also indicate requirement of *hsk1*⁺ for chromatin remodeling at the site of meiosis-specific DSBs. These results show unique and essential roles for Hsk1 kinase during meiotic processes.

Results

Defective Meiosis in *hsk1-89* Mutant. We previously reported *hsk1-89*, a temperature-sensitive allele of *hsk1*⁺ encoding the fission yeast Cdc7 homologue, and showed that *hsk1*⁺ function is essential for mitotic S phase (14). We noticed that the *hsk1-89/hsk1-89* homozygous diploid cells frequently failed to complete meiosis, and most cells were arrested with one nucleus (Fig. 1A), indicative of the requirement of Hsk1 kinase for early stage(s) of meiosis, including premeiotic DNA synthesis, meiotic recombination, or meiosis I. When vegetatively growing diploid cells were shifted to nitrogen-free media at 25°C, DNA content in the majority of the WT cells initially shifted to 2C (C = haploid genome equivalent) (4 h after shift), and then reshifted to 4C by 8 h, consistent with successful progression of premeiotic DNA replication (Fig. 1B Left). In contrast, the *hsk1-89* diploid cells exhibited slow transition to 2C DNA content and 40–50% population kept being arrested in G₁ phase until 16 h after nitrogen deprivation. They shifted to 4C DNA content by 25 h (Fig. 1B Right). These results show that initial transfer to 2C DNA content after nitrogen starvation and subsequent premeiotic DNA replication are delayed in the *hsk1-89/hsk1-89* homozygous diploid.

pat1⁺ encodes a protein kinase that phosphorylates Mei2 protein and negatively regulates meiosis at both the stage of premeiotic DNA replication and that of meiosis I (15, 16). In *pat1-114* cells, meiosis can be induced in a haploid state at a nonpermissive temperature in a more synchronous manner. After nitrogen starvation for 16 h at 25°C, *pat1-114* was shifted to 32°C (nonpermissive for mitotic growth) in medium containing nitrogen sources. Analyses of DNA content at various times after release indicate that premeiotic DNA replication occurs at 1.5–3 h after shift to a nonpermissive temperature in *pat1-114* (Fig. 1C Left). By 8 h after shift, four nuclei compartments similar to spores were observed (Fig. 1D Lower). In the *hsk1-89* background, transition to 2C DNA, detectable on FACS, did not start until 2.5 h after temperature shift, and DNA content slowly increased until it became a 2C amount by 4–5 h after the

Conflict of interest statement: No conflicts declared.

Abbreviations: DSB, dsDNA break; TK, thymidine kinase; MM, minimal medium.

[¶]Present address: Laboratory for Systems Biology and Medicine, Research Center for Advanced Science and Technology, University of Tokyo, Meguro-ku, Tokyo 153-8904, Japan.

^{||}To whom correspondence should be addressed. E-mail: hmasai@rinshoken.or.jp.

© 2006 by The National Academy of Sciences of the USA




Fig. 1. Meiotic defect in *hsk1-89* cells. (A and B) Diploid cells, NI298 (*hsk1⁺/hsk1⁺*) or NI325 (*hsk1-89/hsk1-89*), grown in minimal medium to 5×10^6 cells per ml, were starved for nitrogen at 25°C for 24 h and examined under a microscope after staining with DAPI (A) or analyzed by FACS at the indicated times after starvation (B). (C) JZ767 (*pat1-114*, WT) and NI394 (*pat1-114 hsk1-89*) cells were starved for nitrogen sources for 16 h at 25°C and released into media containing nitrogen at 32°C, nonpermissive for *pat1-114* (0 h), to permit induction of meiosis. Cells were collected at various times after release, and DNA contents were analyzed by FACS. The difference of timing for premeiotic DNA replication among the two strains is most obvious at 3 h after induction (highlighted by a box). (D) Morphology of JZ767 (*pat1-114*, WT) and NI394 (*pat1-114 hsk1-89*) cells induced into meiosis. The photos were taken at 8 h after release at 32°C. Spore-like compartments containing three to four nuclei were observed in *pat1-114*, whereas most cells were arrested with one nucleus in *pat1-114 hsk1-89*. The graphs indicate the fractions of cells with one, two, or three to four nuclei at various time points after release. (E) JZ767 (*pat1-114*, WT) or NI394 (*pat1-114 hsk1-89*) cells were induced into meiosis, as described in *Materials and Methods*, except that release from nitrogen starvation was conducted at 34°C, semipermissive for *hsk1-89*. DNA content was analyzed by FACS at each time point after release. At this temperature, DNA replication proceeds in a kinetics similar to that of *hsk1⁺* cells, apparently completing the process at 3 h after induction (highlighted by a box). (Magnifications: $\times 250$, A; $\times 290$, D.)

induction (Fig. 1C Right). Most cells were with one nucleus even at 8 h after shift (Fig. 1D). Therefore, meiosis induced in haploid cells by inactivation of Pat1 is also arrested in *hsk1-89* cells before meiosis I, and premeiotic DNA replication is delayed.

hsk1-89 cells do not form colonies at 30–32°C, but can grow at 37°C, albeit at a reduced rate (17). This growth property may be caused by induction of chaperone protein(s) at 37°C, which may partially reactivate the kinase-compromised Hsk1-89 protein. When meiosis was induced in *pat1-114 hsk1-89* at 34°C, at which *hsk1-89* cells could form small colonies after prolonged incubation, very little delay of premeiotic S phase was observed, and DNA replication was mostly completed within 3 h after temperature shift, suggesting that the partially restored mutant Hsk1-89 protein could support premeiotic DNA replication in nearly normal timing (Fig. 1E). However, most cells still arrested with one nucleus under this condition (data not shown), indicating that meiosis was blocked.




Fig. 2. Induction of meiotic DSBs in *hsk1-89* cells. (A and B) JZ767 (*pat1-114*, WT), NI394 (*pat1-114 hsk1-89*), and KO272 (*pat1-114 rec12Δ*) cells were induced into meiosis by temperature shift to 32°C (A) or 34°C (B). At the times indicated, DNAs were analyzed on pulsed-field gel electrophoresis. DNA fragments derived from DSBs induced during meiosis are indicated by brackets. Chr 1, Chr 2, and Chr 3 indicate the positions of chromosomes 1, 2, and 3, respectively. DNA plugs were prepared as described (3). The size of chromosome 3 in *hsk1-89* cells is shorter than that in *hsk1⁺* cells. A similar observation was also made in another *hsk1ts* cell (*hsk1-1312*; ref. 44) and other replication mutant cells. The alteration of the size of chromosome 3 is most likely to be caused by the expansion or shrinkage of the rRNA-encoding DNA repeats on this chromosome (data not shown). (C) DNA from *hsk1⁺* (WT) or *hsk1-89* cells was analyzed on a pulsed-field gel under an altered electrophoresis condition. (Left) Ethidium bromide staining of the gel. (Right) Southern blot hybridization of the same gel with a radioactive probe containing the *ura1* gene located ≈ 0.75 Mb from the left end of the chromosome I. Arrowheads indicate three major fragments generated during meiotic DSB, which were detected by this probe (19). M indicates *Saccharomyces cerevisiae* chromosome DNA markers (BioWhittaker). (D) DSBs induced at the *ade6-M26* locus during *pat1*-induced meiosis were examined under K342 (WT) and KO162 (*hsk1-89*) backgrounds. DNA plugs were prepared as described (25). Digestion of DNA plugs and preparation of the probe for Southern blotting were conducted as described (20). M indicates EcoT14I-digested DNA size markers (TAKARA). The arrowhead indicates the position of M26. The bracket indicates the position of the breaks.

Meiosis-Induced DSBs Are Not Generated in *hsk1-89* Cells. Initiation of meiotic recombination is marked by induction of multiple DSBs on the chromosomes (7, 18). DSBs generate broken chromosomes that can be detected in pulsed-field gel electrophoresis as fragments migrating faster in the gel. In *hsk1⁺* cells, DSBs appear between 2 and 4 h after temperature shift of *pat1-114* cells, and their intensity reaches maximum at 3 h (Fig. 2A), roughly correlating with the completion of premeiotic DNA synthesis. These broken chromosomes are not detected in a mutant lacking *rec12⁺* encoding the protein that makes DSBs (ref. 19 and Fig. 2A Right), suggesting that they are indeed intermediates of meiotic recombination. In contrast, broken chromosomes are not detected in *hsk1-89* even at later times when the bulk of DNA has been synthesized (5–8 h after temperature shift; Fig. 2A Left). DSBs were not detected in *hsk1-89* cells at 34°C either, where premeiotic DNA replication proceeded in normal timing (Figs. 1E and 2B).

DSBs can be detected in a more sensitive manner by analyzing the broken chromosomes near one end of chromosome I by hybridization using a radioactive *ura1* probe (19). Chromosome DNAs were separated on pulsed-field gel electrophoresis under a different condition. Broken chromosomes were visible in the ethidium bromide-stained gel in *hsk1⁺* cells at 3–6 h after induction, whereas they were not obvious in *hsk1-89* cells (Fig. 2C Left). When this gel was hybridized with the *ura1* probe, multiple bands, including three major bands previously identified (19), were detected in the *hsk1⁺* cells, whereas the intensities of these bands were significantly reduced in the *hsk1-89* background (Fig. 2C Right). We next examined DSBs at *ade6-M26*, a recombination hot spot, where specific meiotic DSBs have




Fig. 3. BrdU labeling of the replicating chromosomes and analyses on pulsed-field gel electrophoresis. JZ767 (*pat1-114*, WT) or NI394 (*pat1-114 hsk1-89*) cells harboring the plasmid indicated were grown in the absence of thiamine, and meiosis was induced as described in *Materials and Methods*. At the times indicated, genomic DNAs were analyzed on pulsed-field gel electrophoresis. (Right) After the run, the gel was blotted with anti-BrdU antibody. (Left) Ethidium bromide staining of the same gel. The bracket indicates the positions of meiotic DSBs.

been detected (20). These DSBs can be detected at 4–6 h after induction of meiosis in *pat1-114* cells with the *rad50S* background that accumulates unprocessed meiotic DSBs (21, 22). However, these cleavages were not detected in the *hsk1-89* background (Fig. 2D). These data indicate an essential role of the *hsk1*⁺ function in the induction of meiotic DSBs.

Premeiotic DNA Replication Is Likely to Be Completed in *hsk1-89* Cells Under a Nonpermissive Condition for Mitotic DNA Replication. Strong coupling of premeiotic DNA replication and recombination in budding yeast (6–9) suggests a possibility that DNA replication may not be completed in *hsk1-89* cells, which may be affecting the initiation of meiotic recombination. Therefore, we have examined the extent of premeiotic DNA replication in *hsk1-89* cells. It is known that the replicating DNA does not enter agarose in pulsed-field gel electrophoresis. Only after completion and decatenation of the replicated chromosomes can the chromosomes enter agarose and become separated as distinct chromosome bands. We labeled the DNA with BrdU and monitored the progression of premeiotic DNA replication by Western blotting of the chromosomes fractionated by pulsed-field gel electrophoresis using anti-BrdU antibody (23). For this purpose, we have introduced a thymidine kinase (TK)-expressing plasmid in yeast cells. Under this condition, the progression of *pat1*-induced meiosis at 32°C was delayed for some unknown reason. Complete transition to 2C DNA content required 4 h in WT cells and 7 h in *hsk1-89* cells (Fig. 7, which is published as supporting information on the PNAS web site). In WT cells, the chromosome bands incorporating BrdU were visible at 3 h after the induction of meiosis and increased at later hours (Fig. 3 Upper), whereas they were detectable at 5 h after the induction and reached maximum at 8 h in *hsk1-89* cells (Fig. 3 Lower). Delayed appearance of the replicated chromosomes in *hsk1-89* cells is consistent with the results of FACS analyses of DNA contents. Nevertheless, this result strongly indicates that the three chromosomes are replicated to completion during the meiotic process in the mutant cells. The DSBs were not observed even at 12 h after induction, the timing long after completion of the premeiotic DNA replication in *hsk1-89* cells (Fig. 8, which is published as supporting information on the PNAS web site), indicating that the formation of DSBs is not simply delayed but

is indeed impaired. We also confirmed the replication of the specific segments of the chromosome by directly analyzing the replicated molecules (Fig. 9, which is published as supporting information on the PNAS web site). These results mostly rule out the possibility that loss of DSB is caused by the incomplete replication of the genome. However, we cannot completely rule out the possibility that minor defects in DNA replication that cannot be detected by the methods used in this study are responsible for the loss of DSBs. In budding yeast, it was previously reported that the overall extent of chromosome replication is not significantly affected in *cdc7^{ts}* mutants during the course of meiosis (12, 13), and our results in fission yeast are consistent with these reports.

Hsk1 Kinase Activity Is Required for Timely Premeiotic DNA Synthesis and DSB Formation. Both premeiotic DNA synthesis and DSBs were restored in the mutant by introduction of a plasmid expressing WT Hsk1, whereas kinase-negative (K129D) or kinase-attenuated (K129R,K130S) forms of Hsk1 (24) did not restore them under the same conditions (Fig. 10A and B, which is published as supporting information on the PNAS web site, and data not shown), indicating that the kinase activity of Hsk1 is required for timely DNA synthesis and induction of DSBs during meiosis. The K129R,K130S mutant can complement the temperature-sensitive growth of *hsk1-89* when overexpressed (24). However, this mutant did not restore defective meiosis, including timing of premeiotic DNA synthesis (Fig. 10A) and DSB formation even after overproduction (Fig. 10B). These results show that the requirement of Hsk1 kinase activity during meiosis may be more stringent than that for mitotic growth. The size of chromosome 3 is shorter in *hsk1-89* than in WT cells, but is partially corrected by introduction of a plasmid expressing the WT Hsk1 but not the kinase-attenuated one. The shortening of chromosome 3 is observed not only in nitrogen-starved cells but also in vegetatively growing cells, and thus is not caused by the problems in the process of meiosis. We have shown that it is caused by reduction of the copy numbers of the rRNA-encoding DNA on chromosome 3 (data not shown).

Transcriptional Induction of Some *rec* Genes and Horsetail Movement in *hsk1-89* Are Normal. A series of *rec* genes are transcriptionally activated during meiosis. The induction of these *rec* genes, which depends on the Cdc10–Rep1 transcriptional complex (7, 25), is needed for meiotic recombination. We have examined whether this transcriptional activation is affected in *hsk1-89* cells. Under the *hsk1*⁺ background, *rec6*⁺, *rec7*⁺, *rec8*⁺, and *rec12*⁺ genes were induced at 2 h after temperature shift in *pat1-114* cells and disappeared by 5–6 h. Under the *hsk1-89* background, all of these genes were induced at identical timings. The transcripts were detected until 5–7 h after induction and disappeared by 8 h (Fig. 4A and Fig. 11A, which is published as supporting information on the PNAS web site). The *hsk1*⁺ transcript, on the other hand, displayed biphasic induction, peaking at 1 and 5 h after induction of meiosis, whereas the *dfp1*⁺/*him1*⁺ transcript, encoding an activation subunit for Hsk1 (10, 24, 26, 27), increased during the first 4 h and decreased after that in *hsk1*⁺ cells, suggesting that transcriptions of *hsk1*⁺ and *dfp1*⁺/*him1*⁺ may be under regulation distinct from that for *rec* genes. They were induced with similar timing in *hsk1-89* cells, and their levels were maintained until later hours. On the protein level, Dfp1/Him1 protein starts to increase at 1.5 h after temperature shift, around the time of premeiotic DNA synthesis, whereas the Hsk1 protein level transiently increased at 3–3.5 h after the shift, roughly the time of DSB formation, and diminished again until it started to increase again at 5.5 h after induction. In *hsk1-89* cells, Hsk1 protein slightly increased at 2.5–3 h after the shift and increased again only at 7–8 h, whereas Dfp1/Him1 protein showed delayed induction at 3.5–5.0 h after the shift (Figs. 4B and 11B). These




Fig. 4. Expression of various proteins and horsetail movement during meiosis in *hsk1-89* cells. (A and B) JZ767 (*pat1-114*, WT) and NI394 (*pat1-114 hsk1-89*) cells were induced into meiosis, and RNA and whole-cell extracts were prepared at the times indicated after temperature shift. They were analyzed, respectively, by Northern analyses using the probes for the genes indicated (A) or by Western analyses with the various antibodies indicated (B). AS, culture growing asynchronously at 25°C. The band intensities were quantified, and relative intensities are presented in Fig. 11. (C). Time-lapse observation of horsetail movement of *hsk1-89/hsk1-89*. To examine the effect of *hsk1-89* mutation on horsetail movement, NI322 (*hsk1-89 h⁺*) and NI324 (*hsk1-89 h⁻*) cells were conjugated on a sporulation agar (SPA) plate. After incubation at 30°C for 17 h, cells were stained with Hoechst 33342 (4 μg/ml in SPA) and mounted on a thin SPA layer on a glass slide, and pictures were taken every 30 s with a Zeiss Axiophoto equipped with an AQUACOSMOS imaging system (Hamamatsu Photonics, Hamamatsu City, Japan). Photos of representative cells at selected time points are shown.

results indicate that at least the transcriptional induction of some meiotic recombination genes and replication factors are intact in *hsk1-89* cells. Similarly, telomere clustering and horsetail movement, required for efficient homologue pairing and meiotic recombination (28), were detected in the *hsk1-89/hsk1-89* diploid cells as actively as in the WT diploid cells (Fig. 4C).

Known Checkpoint Kinases Are Not Involved in Suppression of DSB and Meiosis in *hsk1-89*. We then examined whether the meiotic defects in *hsk1-89* result from checkpoint-mediated inhibition caused by perturbation of the preceding events. We therefore investigated the effects of mutations in checkpoint kinases known to be involved in DNA damage and replication checkpoint responses. Delay of premeiotic DNA synthesis and generation of one nucleus cells were not suppressed by either *chk1* or *cds1* mutations (Fig. 12, which is published as supporting information on the PNAS web site, and data not shown). Similarly, DSBs were not generated even in the *chk1* or *cds1* null background (Fig. 5). Furthermore, neither replication timing nor DSB formation was restored in *cds1Δ chk1Δ* double mutant. We also examined the effect of *mek1* mutation, which encodes a meiosis-specific checkpoint kinase related to Cds1 (29, 30). Introduction of *mek1Δ* in *hsk1-89* did not restore timely premeiotic DNA replication, DSBs, or meiosis. Furthermore, they were not restored even in the presence of *cds1Δ mek1Δ* double or *cds1Δ mek1Δ chk1Δ* triple mutations. The inability of checkpoint mutations to rescue the defects of *hsk1-89* is in a sharp contrast to hydroxyurea-induced inhibition of DSB formation, which is restored by mutations in the Rad3–Cds1 checkpoint pathway (31, 32).

Cdc2 kinase needs to be activated before cell division, and this activation is partly regulated by phosphorylation of Tyr-15 residue (1, 33). Cdc2 protein was present at more or less constant levels throughout meiosis, except for later stages when it signif-




Fig. 5. Effect of disruption of Cds1, Chk1, Mek1, and combinations of these checkpoint kinase mutations on DSB formation. *pat1-114* (WT) or *pat1-114 hsk1-89* cells in combination with checkpoint mutations indicated were induced into meiosis, and DSB formation was analyzed by pulsed-field gel electrophoresis. The brackets indicate the positions of the meiotic breaks. Chr 1, Chr 2, and Chr 3 indicate the positions of chromosomes 1, 2, and 3, respectively.

icantly increased in its abundance (Fig. 4B). In contrast, phosphorylation of Tyr-15 of Cdc2 increased at 2–3.5 h after the shift and then disappeared in *hsk1⁺* cells, whereas this phosphorylation was maintained until later stages in *hsk1-89* cells (Figs. 4B and 11). This sustained phosphorylation of Tyr-15 of Cdc2 may be at least partially responsible for inhibition of meiosis I and II in the *hsk1* mutant. We examined whether the Tyr-15 phosphorylation in *hsk1-89* is regulated by checkpoint kinases. Under *cds1Δ*, *chk1Δ*, *mek1Δ*, *cds1Δ mek1Δ*, *cds1Δ chk1Δ*, or *cds1Δ chk1Δ mek1Δ* backgrounds, sustained phosphorylation of Cdc2 Tyr-15 was still observed (Fig. 13, which is published as supporting information on the PNAS web site), suggesting that these checkpoint kinases are not involved in inhibition of dephosphorylation of this tyrosine residue.

These results show that delay of premeiotic DNA replication, inhibition of meiotic DSB, or meiotic divisions in *hsk1-89* is not under the regulation of known checkpoint kinases.

Table 1. *hsk1-89* mutant is defective in meiotic recombination

Cell type	Intragenic recombination frequencies (Ade ⁺ recombinants/total colonies counted), M26 × 469	Intergenic recombination frequencies, %		
		Leu ⁺ Mat1-M*	Leu ⁺ His ⁺⁺	Leu ⁻ His ⁻ †
WT	82/10572 (0.78%)	12	13.5	12.5
<i>hsk1-89</i>	8/7080 (0.11%)	0.92	2.08	1.56

The frequencies of meiotic intragenic recombination were measured in the WT (K37 × K41, WT) and *hsk1-89* (KO185 × KO461) backgrounds by random spore analyses. Ade⁺ frequencies were determined by red (Ade⁻) or white (Ade⁺) colony color selection on yeast extract plates. The frequencies of meiotic intergenic recombination was measured on the *leu1-mat1* and *leu1-his5* intervals on the chromosome 2 by random spore analyses of crosses in the following: *, WT (NT145 × K27, WT) and *hsk1-89* (KO147 × KO159) backgrounds or †, diploid WT (D20) and *hsk1-89* (D21) cells. Glucanase-treated cells were further treated by 30% ethanol for 5 min to kill remaining diploid cells. The surviving colonies were examined for the markers indicated. All the Leu⁺ His⁺ colonies were confirmed to be haploids.




Fig. 6. Chromatin remodeling during meiosis in *hsk1-89* cells. Chromatin remodeling was examined in the *ade6-M26* region during meiosis induced in diploid cells D20 (*M26 hsk1⁺/M26 hsk1⁺*, Left) and D21 (*M26 hsk1-89/M26 hsk1-89*, Right). The arrowheads indicate the bands derived from the chromatin remodeling observed in the WT cells. Micrococcal nuclease was titrated (0, 5, 10, or 20 units per ml for *hsk1⁺/hsk1⁺* and 0, 10, or 20 units per ml for *hsk1-89/hsk1-89*) at each time point, as indicated by the filled triangles. The open arrow indicates the coding region and direction of the *ade6* gene. The positions of the XhoI site, ATG initiation codon, and M26 DSB sites are also indicated. N, MNase digestion of naked DNA.

Meiotic Recombination Is Impaired in *hsk1-89* Cells. Reduced levels of DSBs in *hsk1-89* predict the reduced recombination frequency. We have therefore measured recombination frequency in spores generated in *hsk1-89* cells. Approximately 10% of the *hsk1-89* cells undergo sporulation and generate four-spored asci. We measured the intragenic recombination frequency between *ade6-469* and *ade6-M26* and intergenic recombination frequency in the *leu1-mat1* and *leu1-his5* loci (Table 1). We found that the intragenic recombination frequencies decreased by ≈ 7 -fold in *hsk1-89* cells (0.78% and 0.11% in WT and *hsk1-89*, respectively). The intergenic recombination frequencies in the WT cells were 12.0% for *leu1/mat1* and 13.5% (*leu⁺ his⁺*)/12.5% (*leu⁻ his⁻*) for *leu1/his5*, respectively, which were roughly consistent with the values reported (29). In contrast, they were 0.92% and 2.08% (*leu⁺ his⁺*)/1.56% (*leu⁻ his⁻*), respectively, in the *hsk1-89* cells, which are about the same as the values observed in some *rec* mutants (34). These results show that recombination is indeed impaired in *hsk1-89* cells, as was reported in *cdc7^{ts}* cells in budding yeast (12, 13).

Chromatin Remodeling at the Site of DSB Is Impaired in *hsk1-89* Cells. Chromatin remodeling is known to play an important role in the generation of DSBs at *ade6-M26* (20, 35, 36). We have examined whether the *hsk1-89* mutation affects chromatin remodeling at this locus. In the *hsk1⁺* background, the alteration of micrococcal nuclease cleavage patterns started to appear at 3–4 h after induction of meiosis in diploid cells and continued to increase until 6 h (Fig. 6 Left and data not shown), indicative of alteration of chromatin structures. In contrast, in *hsk1-89* cells, very little change in the cleavage pattern was detected even at 16 h after induction of meiosis (Fig. 6 Right). Strong chromatin remodeling observed in the WT cells was not obvious in *hsk1-89* cells during *pat1*-induced meiosis (Fig. 14, which is published as supporting information on the PNAS web site). These results show that the *hsk1* mutation abrogates chromatin remodeling at the site of meiotic DSBs, which may cause the loss of DSBs.

Discussion

The data presented in this study demonstrate essential roles of *hsk1⁺*, the fission yeast homologue of Cdc7 kinase known to be essential for mitotic S phase (10, 26, 27), during meiosis. Our results indicate that *hsk1⁺* is required for both meiotic recombination and meiotic nuclear and cell divisions. Most strikingly,

our data show that *hsk1⁺* is required for induction of DSBs, one of the initial events for meiotic recombination. These meiotic defects in the *hsk1* mutant are not caused by known checkpoint pathways.

Strong coupling between premeiotic DNA replication and meiotic recombination has been reported in budding yeast (6–9). Although premeiotic DNA replication is slightly delayed in *hsk1-89* cells, evidence indicates that recombination and meiotic defects may not be consequences of incomplete DNA replication. (i) Analyses of replicated chromosomes on pulsed-field gel electrophoresis strongly indicate the completion and proper decatenation of all of the chromosomes (Figs. 3 and 7). (ii) Analyses of BrdU-labeled replicated molecules strongly suggest the replication of the entire genome in *hsk1-89* cells (Fig. 9). (iii) At 34°C, a temperature that permits a limited growth of *hsk1-89*, premeiotic DNA replication is induced apparently in normal timing, and yet no DSB is induced and cells arrest with one nucleus (Figs. 1 and 2). (iv) Both intragenic and intergenic recombination frequency is significantly reduced in the mutant spores, which may have completed meiotic processes, including premeiotic DNA replication and meiotic division (Table 1). Although the deteriorative effect of minor defects in premeiotic DNA replication on DSB formation cannot be formally ruled out, these findings strongly indicate that Hsk1 either directly or indirectly regulates meiotic recombination. They are in keeping with previous reports showing that the bulk amount of DNA synthesis during meiosis is not affected in *cdc7^{ts}* mutants (12, 13). Apparent completion of premeiotic DNA replication in *hsk1-89* under the condition nonpermissive for mitotic DNA replication suggests differential roles of Cdc7 kinase in DNA replication during mitotic and meiotic phases.

Chromatin remodeling at a recombination hot spot is not induced in the *hsk1-89* cells, suggesting a possibility that Hsk1 may regulate the processes of chromatin remodeling. We previously reported that sister chromatid cohesion during mitosis is partially impaired in *hsk1-89* cells (14). It is an intriguing possibility that defective meiotic sister chromatid cohesion in *hsk1-89* may lead to impaired chromatin remodeling, a prerequisite for meiotic recombination.

The requirement of Hsk1 for meiotic nuclear division was unexpected. It was reported that meiosis I can proceed in mutants lacking recombination functions, including *rec12⁺*, essential for DSB induction (37). Therefore, the block of meiosis I in *hsk1-89* cells may not be simply caused by the absence of recombination. In mammalian cells, Cdc7 has been implicated in mitosis as well (38). Therefore, we speculate that Hsk1 may play some unknown essential functions during meiotic nuclear and cell divisions independent from its S-phase functions.

We recently reported severe defects in the development of testes and ovaries in genetically engineered mice with an attenuated Cdc7 kinase level (39). Analyses of defective testes indicate arrest of spermatogenesis at premeiotic phases or early prophase I of meiosis I, which is roughly consistent with premeiotic arrest in *hsk1-89* cells. Thus, the essential roles of Cdc7 kinase during the processes of meiosis may be conserved through evolution.

Materials and Methods

Yeast Strains, Media, and Genetics. *Schizosaccharomyces pombe* strains used in this study are listed in Table 2, which is published as supporting information on the PNAS web site. Cells were grown in rich medium or minimal medium (MM) containing the required supplements. Sporulation agar medium containing 10 g/liter of glucose, 1 g/liter of KH_2PO_4 , 1 ml of $\times 1,000$ vitamin stock solution per liter⁻¹, or malt extract agar medium was used for genetic crosses and sporulation. Nitrogen-free derivative MM-N was used for nitrogen starvation experiments as de-

scribed (40). General genetic methods (41) and the procedure for transformation (42) were as described.

Induction of Meiosis in *pat1-114* Cells. Cells, grown at 25°C in MM (1% glucose) supplemented with 100 µg/ml of leucine and 100 µg/ml of adenine per liter⁻¹ to 5 × 10⁶ cells/ml, were washed with MM-NH₄Cl (1% glucose) four times and resuspended in MM-NH₄Cl (1% glucose) supplemented with 25 µg/ml of leucine, followed by incubation for 16 h at 25°C. After addition of an equal volume of prewarmed MM (containing 1 g/liter of NH₄Cl and 1% glucose) supplemented with 50 µg/ml of leucine and 70 µg/ml of adenine, meiosis was induced by shifting the temperature to 32°C or 34°C. Equal volumes of samples were withdrawn at each time point from each culture for FACS analyses and DNA analyses on pulsed-field gel electrophoresis.

Detection of Meiotic DSBs by Pulsed-Field Gel Electrophoresis. *pat1-114* cells released from nitrogen starvation were harvested every hour and treated as described (3). Pulsed-field gel electrophoresis was carried out in 0.6% chromosomal grade agarose gel (Bio-Rad) on a Bio-Rad CHEF-DRIII apparatus and recirculated at 14°C. Electrophoresis was performed for 48–72 h at 1.5 V/cm in 0.5× TAE buffer (Tris-acetate-EDTA), with a switch time of 30 min at an included angle of 106°. For analyses by Southern hybridization, electrophoresis was conducted in 0.8% chromosomal grade agarose gel (Bio-Rad) for 24 h at 6 V/cm in 0.5× TBE buffer (89 mM Tris/89 mM boric acid/2 mM EDTA, pH 8.0), with a linearly ramped 60- to 120-s switch time at an included angle of 120°, as described (19).

Incorporation of BrdU. JZ767 and NI394 harboring pREP41-DP-hENT-TK (expressing human TK gene; a gift from Hisao Masukata, Osaka University, Osaka, Japan) or pREP1, grown in the absence of thiamine, were starved for nitrogen source for 16 h and transferred to 32°C to induce meiosis. BrdU (400 µg/ml) was added to the culture 30 min before temperature shift, and cells were harvested every hour (23). Genomic DNA was prepared, and pulsed-field gel electrophoresis was carried out as described above. After electrophoresis, the gel was stained with ethidium bromide

and soaked in 0.25 M HCl for 20 min, followed by incubation in a solution of 0.5 M NaOH and 1.5 M NaCl for 30 min and transfer to nylon membrane (Hybond N+, Amersham Pharmacia Biosciences). The membrane was washed in 2× SSC briefly and dried. Immunodetection was conducted with anti-BrdU antibody (a gift from Katsuyuki Tamai, Cyclex, Nagano, Japan).

Antibodies. Anti-Hsk1 and anti-Dfp1/Him1 antibodies were affinity-purified and used as described (26, 43). B-5-1-2 (Sigma), anti-Cdc2 (sc-53, PSTAIRE; Santa Cruz Biotechnology), and anti-phospho-Cdc2 (Tyr-15; Cell Signaling Technology, Beverly, MA) antibodies were used to detect α-tubulin, Cdc2, and tyrosine-phosphorylated Cdc2, respectively.

Analyses of Chromatin Remodeling at the *ade6-M26* Locus. Analyses of chromatin structures by indirect end labeling were done as reported (35). To analyze chromatin structures around the *ade6*⁺ promoter, genomic DNA was digested with XhoI and separated by electrophoresis in 1.5% agarose gel (40 cm long) in TAE buffer (Tris-acetate-EDTA). The separated DNA fragments were alkali-transferred to charged nylon membranes (Biodyne B membrane, Pall). The probe used for detection of the *ade6*⁺ locus was as described (35).

See *Supporting Text*, which is published as supporting information on the PNAS web site, for more details.

We thank Dr. Yuichi Iino (University of Tokyo, Tokyo) for the *pat1-114* strain; Dr. Yoshinori Watanabe (University of Tokyo, Tokyo) for various strains, valuable suggestions for many aspects of this work, and advice for recombination frequency measurement; Dr. Takehiko Shibata (RIKEN, Saitama, Japan) for valuable suggestions; Dr. Hisao Masukata for the gift of pREP41-DP-hENT-TK; Dr. Pedro San-Segundo (Universidad de Salamanca, Salamanca, Spain) for the *mek1Δ* strain; Dr. Katsuyuki Tamai for anti-BrdU antibody; and Ai Ishii (Tokyo Metropolitan Institute of Medical Science) for suggestions in conducting FACS analyses. This work was supported in part by grants-in-aid for scientific research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (to H.M.) and grants for basic research from Bio-oriented Technology Research Advancement Institution (to K. Ohta and T. Shibata).

- Iino, Y., Hiramine, Y. & Yamamoto, M. (1995) *Genetics* **140**, 1235–1245.
- Forsburg, S. L. & Hodson, J. A. (2000) *Nat. Genet.* **25**, 263–268.
- Murakami, H. & Nurse, P. (2001) *Nat. Genet.* **28**, 290–293.
- Stuart, D. & Wittenberg, C. (1998) *Genes Dev.* **12**, 2698–2710.
- Lindoner, K., Gregan, J., Montgomery, S. & Kearsley, S. E. (2002) *Mol. Biol. Cell.* **13**, 435–444.
- Borde, V., Goldman, A. S. H. & Lichten, M. (2000) *Science* **290**, 806–809.
- Davis, L. & Smith, G. R. (2001) *Proc. Natl. Acad. Sci. USA* **98**, 8395–8402.
- Smith, K. N., Penkner, A., Ohta, K., Klein, F. & Nicolas, A. (2001) *Curr. Biol.* **11**, 88–97.
- Murakami, H., Botde, V., Shibata, T., Lichten, M. & Ohta, K. (2003) *Nucleic Acids Res.* **31**, 4085–4090.
- Masai, H. & Arai, K. (2002) *J. Cell. Physiol.* **190**, 287–296.
- Sclafani, R. A. (2000) *J. Cell Sci.* **113**, 2111–2117.
- Schild, D. & Byers, D. (1978) *Chromosoma* **70**, 109–130.
- Hollingsworth, E. R. & Sclafani, A. R. (1993) *Chromosoma* **102**, 415–420.
- Takeda, T., Ogino, K., Tatebayashi, K., Ikeda, H., Arai, K. & Masai, H. (2001) *Mol. Biol. Cell* **12**, 1257–1274.
- Iino, Y. & Yamamoto, M. (1985) *Proc. Natl. Acad. Sci. USA* **82**, 2447–2451.
- Watanabe, Y. & Yamamoto, M. (1994) *Cell* **78**, 487–498.
- Matsumoto, S., Ogino, K., Noguchi, E., Russell, P. & Masai, H. (2005) *J. Biol. Chem.* **280**, 42536–42542.
- Fox, M. E. & Smith, G. R. (1998) *Prog. Nucleic Acid Res. Mol. Biol.* **61**, 345–378.
- Cervantes, M. D., Farah, J. A. & Smith, G. R. (2000) *Mol. Cell* **5**, 883–888.
- Steiner, W. W., Schreckhise, R. W. & Smith, G. R. (2002) *Mol. Cell* **9**, 847–855.
- Cao, L., Alani, E. & Kleckner, N. (1990) *Cell* **61**, 1089–1101.
- Young, J. A., Schreckhise, R. W., Steiner, W. W. & Smith, G. R. (2002) *Mol. Cell* **9**, 253–263.
- Lengronne, A., Pasero, P., Bensimon, A. & Schwob, E. (2001) *Nucleic Acid Res.* **29**, 1433–1442.
- Ogino, K., Takeda, T., Matsui, E., Iiyama, H., Taniyama, C., Arai, K. & Masai, H. (2001) *J. Biol. Chem.* **276**, 31376–31387.
- Lin, Y. & Smith, G. R. (1994) *Genetics* **136**, 769–779.
- Masai, H., Miyake, T. & Arai, K. (1995) *EMBO J.* **14**, 3094–3104.
- Brown, G. W. & Kelly, T. J. (1998) *J. Biol. Chem.* **273**, 22083–22090.
- Chikashige, Y., Ding, D. Q., Funabiki, H., Haraguchi, T., Mashiko, S., Yanagida, M. & Hiraoka, Y. (1994) *Science* **264**, 270–273.
- Pérez-Hidalgo, L., Moreno, S. & San-Segundo, P. A. (2003) *J. Cell Sci.* **116**, 259–271.
- Shimada, M., Nabeshima, K., Tougan, T. & Nojima, H. (2002) *EMBO J.* **21**, 2807–2818.
- Tonami, Y., Murakami, H., Shirahige, K. & Nakanishi, M. (2005) *Proc. Natl. Acad. Sci. USA* **102**, 5797–5801.
- Ogino, K. & Masai, H. (2006) *J. Biol. Chem.* **281**, 1338–1344.
- Gold, K. L. & Nurse, P. (1989) *Nature* **342**, 39–45.
- DeVaux, L. C., Hoagland, N. A. & Smith, G. R. (1992) *Genetics* **130**, 251–262.
- Mizuno, K., Emura, Y., Baur, M., Kohli, J., Ohta, K. & Shibata, T. (1997) *Genes Dev.* **11**, 876–886.
- Yamada, T., Mizuno, K., Hirota, K., Kon, N., Wahls, W. P., Hartsuiker, E., Murofushi, H., Shibata, T. & Ohta, K. (2004) *EMBO J.* **23**, 1792–1803.
- Davis, L. & Smith, G. R. (2003) *Genetics* **163**, 857–874.
- Yoshizawa-Sugata, N., Ishii, A., Taniyama, C., Matsui, E., Arai, K. & Masai, H. (2004) *J. Biol. Chem.* **280**, 13062–13070.
- Kim, J. M., Takemoto, N., Arai, K. & Masai, H. (2003) *EMBO J.* **22**, 5260–5272.
- Ishiki, T., Mochizuki, N., Maeda, T. & Yamamoto, M. (1992) *Gen. Dev.* **6**, 2455–2462.
- Moreno, S., Klar, A. & Nurse, P. (1991) *Methods Enzymol.* **194**, 795–823.
- Okazaki, K., Okazaki, N., Kume, K., Jinno, S., Tanaka, K. & Okayama, H. (1990) *Nucleic Acids Res.* **18**, 6485–6489.
- Takeda, T., Ogino, K., Matsui, E., Cho, M. K., Kumagai, H., Miyake, T., Arai, K. & Masai, H. (1999) *Mol. Cell Biol.* **19**, 5535–5547.
- Snaith, H. A., Brown, G. W. & Forsburg, S. L. (2000) *Mol. Cell Biol.* **21**, 7922–7932.