

Simpson and Ballard set a good example.⁵ Their study highlights how tests do not always measure what they set out to measure; in this case decision making in the orals. Swanwick and Chana raise interesting issues on the high validity of evidential, locally-based assessment, suggesting ways of enhancing the reliability.⁶ If we believe the published literature,¹¹ the task in hand is challenging.

The danger of current change is the impetus with which it is taking place. The new approach may not lead us to the Holy Grail for a test that accurately predicts future unobserved practice; but we do face a major change in philosophy. Assessment programmes designed to ensure education is efficiently and appropriately delivered are replacing the examinations that are known to be reliable but are at times lacking in validity. We need well-designed research to support or refute this new rationale.

Val Wass

Professor of Community-based Education,
Manchester

REFERENCES

1. Flexner A. *Medical education in the United States and Canada*. Bethesda, MD: Science and Health Publications, 1910.
2. Miller GE. The assessment of clinical skills/competence/performance. *Acad Med* 1990; **65**: S63–S67.
3. Schuwirth LWT, van der Vleuten CPM. The use of clinical simulations in assessment. *Med Educ* 2003; **37** Suppl 1: 65–71.
4. Wass V, van der Vleuten CPM. The long case. *Med Educ* 2004; **38**(11): 1176–1180.
5. Simpson RG, Ballard KD. What is being assessed in the MRCGP oral examination? A qualitative study. *Br J Gen Pract* 2005; **55**: 430–436.
6. Swanwick T, Chana N. Workplace assessment for licensing in general practice. *Br J Gen Pract* 2005; **55**: 461–467.
7. van der Vleuten CPM, Schuwirth LWT. Assessing professional competence: from methods to programmes. *Med Educ* 2005; **39**: 309–317.
8. Wass V, van der Vleuten CPM, Shatzer J, Jones R. Assessment of clinical competence. *Lancet* 2001; **357**: 945–949.
9. Regehr G. Trends in medical education research. *Acad*

Med 2004; **79**(10): 939–947.

10. Talbot M. Monkey see, monkey do: a critique of the competency model in graduate medical education. *Med Educ* 2004; **38**: 587–592.
11. Norcini JJ, Blank LL, Duffy FD, Fortuna GS. The mini-CEX: a method for assessing clinical skills. *Ann Intern Med* 2003; **138**(6): 476–481.
12. Williams RG, Klamen DK, McGaghie WC. Cognitive, social and environmental sources of bias in clinical performance ratings. *Teach Learn Med* 2003; **15**(4): 270–292.

ADDRESS FOR CORRESPONDENCE

Val Wass

Professor of Community-based Education,
First Floor, Rusholme Health Centre,
Walmer St, Manchester M14 5NP
E-mail: valerie.wass@manchester.ac.uk

Making general practice fit for the 21st century

For some while it has been customary to argue that good primary care is an essential part of an effective, comprehensive system of health care. Barbara Starfield's work comparing the systems in different countries has supplied convincing objective evidence to support the contention, where countries with poor orientation towards primary care have worse health outcomes.¹ The same relationship has been found when comparing the different US states,¹ and on a more local scale, mortality rates in English hospitals have been found to be closely related to the supply of GPs in the area.² Policy experts observing from countries lacking such systems, most obviously the US, bemoan their deprivation in this respect.³ Older doctors and patients lament the passing of the familiar family doctor, who through long knowledge of the patients became a family friend. The difficulty with this vision is that such long-term commitment has become unusual, and erodes through the mobility of both patients and the needs of doctors for career development and adherence to the

European Working Time Directive. So unusual that it is now unrealistic to plan a system on the basis of long-term relationships. Worse is the risk that constructing a system on the basis of such relationships sets it up to fail. It is worth taking stock to define what the task should be now, and how we can set about achieving it.

First and foremost, doctors working in primary care have to practise medicine to a high degree of technical competence. It is easy to dismiss this, or regard it as a truism. But truisms are often true. It has been customary to take technical excellence for granted, either by assuming its universal existence among all primary care doctors, or through fear of what we might find if we looked a bit harder. However, we know that there have always been doctors working in primary care whose practice would not, if tested, reach an acceptable standard. One of the benefits of a robust system of revalidation, when it is in place, is that it will tell us whether such doctors comprise 0.1%, 1% or 10% of the total. Technical

competence demands two added dimensions. First, doctors all have to keep abreast of a discipline that changes constantly, with the appearance of new diseases (and the disappearance of some old ones), development of new diagnostic tests and techniques, and a constant supply of new therapies. Second, doctors working in primary care should continue to aspire to be good generalists. One of the strengths of primary care is to have enough expertise across all areas of medicine to relieve patients of any anxiety that they need to decide the appropriate specialist to deal with their problem. Also the last thing patients want is a primary care doctor who can say as specialists can (and sometimes do), 'Sorry but this problem is outside my area of expertise'. Because the idea of excellence which turns its face against specialism is so alien in our culture, the insistence that primary care has to remain rooted in generalism is a vital message to get into the heads of policy makers. This is especially true right now in the UK, with developments pulling in the opposite

direction: the creation of GPs with special interests, a contract offering doctors the opportunity to opt out of particular clinical areas, and the trend towards part-time working.

Modern general practice implies expertise in areas beyond purely medical competence. GPs have to practise within a defensible ethical framework. There is the principle of justice, implying respect for the wide range of cultural sensitivities encountered, and attention to the cost-effectiveness of decisions and how they impinge on the rest of the system. There is also the requirement to respect patients' autonomy, with the need to involve them, as far as they wish, in decisions affecting their health. In passing, it's worth pointing out that the need to promote shared decision making is both more high-minded, more truly respecting of patients' autonomy and more complex than the meretricious choice of provider so assiduously promoted by policy makers in the UK. These extra dimensions can feel as if there are numerous invisible observers in the consulting room looking over our shoulders. Not just the medical experts but also a panoply of ethicists, anthropologists, economists, and management consultants.

Providing care of high technical quality has become easier with electronic access to up-to-date information, in the form of primary research, systematic reviews and guidelines. Decision support, both electronic and delivered by other means is following close behind. Standard GP systems already provide prompts and alerts, and such aids will become more sophisticated, delivering individualised data on prognosis, and risks and benefits of different treatments.⁴ What's more, either because of such assistance or despite it, there is convincing evidence that standards of technical care are improving, and have been doing so for some time before the introduction of the 2004 contract.⁵

Which leaves a question mark hanging over personal care. Because in the past it was felt to be such a permanent feature of primary care we may have taken it for granted. Patients have always valued personal continuity of care,⁶⁻⁹ but others, most notably policy makers, have assumed that immediate access is more important. However, recent research on changes to

access,¹⁰⁻¹¹ together with the more dramatic press coverage shortly before the recent UK election has confirmed its importance. The more that the value of trust is recognised to lie at the heart of effective health care,¹² the more important personal care becomes. If patient-centred care is to become anything other than an empty slogan, it will require much better personal care, with doctors being more sensitive to the personal agendas that patients bring to shared decision making. So, with long-term personal continuity slowly eroding, and no longer to be assumed, the central task of primary care in the coming years will be to find ways of delivering personal care in the context of much shorter, more intense, relationships between doctors and patients. Different models are being proposed for personal care, and the evidence for what works in routine primary care is, as yet, scanty. For future researchers the vital criteria follow from the arguments put forward here: it has to be an approach that is not based on long-term relationships, that can be deployed in relatively short encounters and that facilitates shared decision making.

Accomplishing this task will involve everyone with any influence on primary care. Clinicians themselves will want to assess their own abilities more than they do at present, and it may become a regular part of appraisal. Those responsible for postgraduate education (and assessment) will wish to develop methods to help doctors learn the necessary skills. The work to date on shared decision making has so far indicated that the skills can be learnt and deployed,¹³ but that does not guarantee that they will be applied in all consultations.¹⁴ It will certainly be a long learning process for most of us, patients as well as doctors. There is, as has been shown in the past, coherence in teaching, assessing and maintaining professional attitudes, and the vital first step is to accept that such attitudes are essential. This approach will also demand longer times for consultation than the 10 minutes that is now the UK norm, and planners will have to be convinced that patient-centred medicine requires long term investment.

David Jewell
Editor

Acknowledgements

This article draws on ideas shared at a seminar held by the Nuffield Trust on 26 April 2005. The seminar is held under 'Chatham House Rules', which means that it is not permitted to attribute views to individuals. Thanks, therefore to all the unnamed people whose views have contributed, and to Roger Jones and Nigel Mathers who commented on an early draft.

REFERENCES

1. Starfield B. New paradigms for quality in primary care. *Br J Gen Pract* 2001; **51**: 303-309.
2. Jarman B, Gault S, Alves B, *et al*. Explaining differences in English hospital death rates using routinely collected data. *BMJ* 1999; **318**: 1515-1520.
3. Ferrer RL, Hambidge SJ, Maly RC. The essential role of generalists in health care systems. *Ann Intern Med* 2005; **142**: 691-699.
4. Montgomery AA, Fahey T, Peters TJ. A factorial randomised controlled trial of decision analysis and an information video plus leaflet for newly diagnosed hypertensive patients. *Br J Gen Pract* 2003; **53**: 446-453.
5. Campbell S, Steiner A, Robison J, *et al*. Is the quality of care in general medical practice improving? Results of a longitudinal observational study. *Br J Gen Pract* 2003; **53**: 298-304.
6. Kearley KE, Freeman GK, Heath A. An exploration of the value of the personal doctor-patient relationship in general practice. *Br J Gen Pract* 2001; **51**: 712-718.
7. Schers H, Webster S, van den Hoogen H, *et al*. Continuity of care in general practice: a survey of patients' views. *Br J Gen Pract* 2002; **52**: 459-462.
8. Schers H, van den Hoogen H, Bor H, *et al*. Preference for a general practitioner and patients' evaluations of care: a cross-sectional study. *Br J Gen Pract* 2004; **54**: 693-694.
9. Mainous AG, Goodwin MA, Stange KC. Patient-physician shared experiences and value patients place on continuity of care. *Ann Fam Med* 2004; **2**: 452-454.
10. Pickin DM, O' Cathain A, Sampson FC, Dixon S. Evaluation of Advanced Access in the National Primary Care Collaborative. *Br J Gen Pract* 2004; **54**: 334-340.
11. Pascoe SW, Neal RD, Allgar VL. Open-access versus bookable appointment systems: survey of patients attending appointments with general practitioners. *Br J Gen Pract* 2004; **54**: 367-369.
12. Maynard A, Bloor K. Trust, performance management and the new GP contract. *Br J Gen Pract* 2003; **53**: 754-755.
13. Elwyn G, Edwards A, Hood K, *et al*. Achieving involvement: process outcomes from a cluster randomized trial of shared decision making skill development and use of risk communication aids in general practice. *Fam Pract* 2004; **21**: 337-346.
14. Edwards A, Elwyn G, Wood F, *et al*. Shared decision making and risk communication in practice: a qualitative study of GPs' experiences. *Br J Gen Pract* 2005; **55**: 6-13.

ADDRESS FOR CORRESPONDENCE

David Jewell

British Journal of General Practice
Royal College of General Practitioners
14 Princes Gate, Hyde Park
London SW7 1PU
E-mail: david.jewell@bristol.ac.uk