

Structure-Activity Relationships for Ulcerogenic and Adrenocorticolytic Effects of Alkyl Nitriles, Amines, and Thiols

by Sandor Szabo* and Edward S. Reynolds*

In rats, a single administration of acrylonitrile (vinyl cyanide) produces a rapidly occurring bilateral adrenal apoplexy. Structure-activity studies have shown that a close derivative, propionitrile (ethyl cyanide), causes duodenal ulcer without markedly affecting the adrenal glands. Prolonging the two-carbon chain of propionitrile by a methyl group (*n*-butyronitrile) enhances, replacing the methyl by bromide or nitrile decreases, while substitution by an amino group abolishes the ulcerogenic potency and variably affects the adrenocorticolytic action. On assaying a large number of nonnitrile compounds as well for ulcerogenic effect, such as thiols and amines, this effect was found to be related to a two-carbon structure bearing electronegative radicals on one or both ends of the chain.

Alkyl nitriles (e.g., acrylonitrile, propionitrile) are used extensively in the manufacture of acrylic fibers, plastics, and surface coatings and as intermediates in the synthesis of antioxidants, pharmaceuticals, and dyes. Many people are exposed to these compounds, since 5-30% of acrylonitrile may remain as monomers in various synthetic products or may become liberated with the aging of polymers (1). Thus, health implications of these compounds are of considerable interest.

Previous Studies

While studying the effect of hormonal and nonhormonal steroids as well as various hepatic drug-metabolizing enzyme inducers on the toxicity of alkyl nitriles, we recently found (2) that in rats acrylonitrile produces bilateral adrenal apoplexy. Hemorrhage and necrosis develop rapidly (1-2 hr) and occur mostly in the outer layers of the adrenal cortex.

Frequently, subcapsular bleeding is seen as well (Fig. 1). Pretreatment of rats with ACTH (3) or phenobarbital, unlike PCN (pregnenolone-16 α -carbonitrile) or spironolactone (4) offers complete protection against the adrenal lesions and mortality induced by acrylonitrile.

Experiments on structure-activity relationships of similar compounds revealed that propionitrile (ethyl cyanide) had no adrenocorticolytic action but produced solitary, often perforating duodenal ulcers in rats (5). These lesions occurred 3-5 mm from the pylorus of the stomach, mostly on the antimesenteric site of the duodenum and usually penetrated into the liver (Fig. 2). Microscopic examination revealed erosion of the mucosa or superficial or deep necrosis of the duodenal wall infiltrated by varying numbers of polymorphonuclear leukocytes (Fig. 3). The duodenal ulcer invariably developed 24-48 hr after the initial administration of propionitrile and was not accompanied by gastric ulcers.

Cysteamine also produced duodenal ulcers (6). Similarities in structure or biologic action between acrylonitrile, propionitrile, and cysteamine were not apparent. Hence, the ulcerogenic

* Department of Pathology, Peter Bent Brigham Hospital and Harvard Medical School, Boston, Massachusetts 02115.




FIGURE 1. Gross and microscopic appearance of adrenals of two rats both given acrylonitrile: (A,*a*) pretreatment with ACTH protects against adrenal necrosis, gland appears normal; (A) gross, adrenal at arrow ($\times 2$); (*a*) microscopic ($\times 49$); (B,*b*) massive cortical and subcapsular hemorrhagic necrosis in animals not protected by ACTH, (B) gross, adrenal is hemorrhagic at arrow ($\times 2.5$); (*b*) micro, extensive hemorrhagic necrosis in outer layers of adrenal cortex and in subcapsular area ($\times 49$).

effect could not be ascribed to a common property (Table 1). However, more recently, these drugs were noted to have a common "two-carbon" or vicinal carbon (2-C) skeleton, bearing in some cases an electron-negative, high bond energy and/or reactive radical(s).

Based on these assumptions, we have now tested numerous compounds having a two

carbon atom skeleton for possible ulcer-producing effects. Preliminary results have been reported elsewhere (7).

Current Research

Throughout the experiments, female Sprague-Dawley rats with an initial body weight of




FIGURE 2. Perforated duodenal ulcer produced by propionitrile (mucosal surface): stomach at top, duodenum at bottom; ulcer at arrows, $\times 45$.

100 g or, if so stated, 200 g were maintained on Purina Lab Chow and tap water *ad libitum*. The compounds were given by mouth (PO) and/or subcutaneously (SC) thrice daily for 4 days, unless otherwise stated at the dose level shown in Table 2. Autopsy was performed soon after the animal died or on the fifth day, when survivors were killed by chloroform. The criterion for "positive ulcer" was a lesion identifiable by gross examination.

For every compound, detailed dose-response and toxicity studies were performed (in some cases up to 50 rats were used per drug). Hence, only representative findings of a large series of investigations are reported.

It can be seen (Table 2) that close derivatives of propionitrile, e.g., *n*-butyronitrile, 3-bromopropionitrile, ethylene dicyanide, unlike 3-aminopropionitrile, showed prominent ulcerogenic effect. However, all of these compounds exerted slight adrenocorticolytic action.

Effects of alkylamines and thiols were compared with those of alkyl nitriles. Of the alkyl amines, ethylamine was the most active ulcerogenic compound and also produced adrenal

necrosis. Methylamine and propylamine, having structures with one and three carbon atoms, respectively, exert weaker ulcerogenic effects; *n*-butylamine was completely devoid of ulcerogenic action. A hydroxyl attached to the β carbon of ethylamine (resulting in ethanolaamine) abolished ulcerogenic potency.

Alkyl aminothiols, close derivatives of cysteamine (cystamine and 2-*n*-butylaminoethanethiol, containing the cysteamine molecule "hidden" in a long carbon chain) produced a high incidence of duodenal ulcer and adrenal necrosis.

In contrast, ethanethiol was not ulcerogenic. However, a derivative of this drug, 2-diethylaminoethanethiol, like 1,2-propanedithiol and less actively 1,3-propanedithiol, showed marked ulcerogenic activity. The four-carbon compound (1-butanethiol) was inactive, while 1,4-butanedithiol had very mild effect.

Discussion

The results presented here demonstrate that certain alkyl nitriles readily produce adrenal necrosis and/or duodenal ulcer in rats. The mechanism of action and the high specificity of localization of the lesion (adrenal and/or duodenum) are poorly understood. No clear structure-activity relationship can be established for the adrenocorticolytic action, although it is striking that the nonsaturated acrylonitrile (vinyl cyanide) causes adrenal apoplexy, the saturated derivative propionitrile (ethyl cyanide) produces duodenal ulcer, and cysteamine (β -mercaptoethylamine) is very active in affecting both of these organs.

There are other data on the ulcerogenic effect of these compounds. It was recently found that propionitrile or cysteamine stimulates gastric secretion and acid output (8). Duodenal ulcers following the administration of cysteamine and propionitrile can be prevented by antacids and ligation of the pylorus (9). Addition at the terminal carbon of propionitrile with a methyl group (*n*-butyronitrile) enhances the ulcerogenic effect. Bromine or nitrile addition decreases and an amino group completely abolishes the ulcerogenic action. Thus, in this case, the ulcerogenic potency of these radicals added to the terminal carbon of propionitrile may be presented as follows: $-\text{CH}_3 > -\text{CN} \cong -\text{Br} >> -\text{NH}_2$.




FIGURE 3. Stages of duodenal ulcer development (stomach at top, serosa on left, mucosa on right): (A) early, mucosal erosion in *n*-butyronitrile-treated rat (arrows) showing villous amputation (H&E, $\times 39$); (B) transmural necrosis (arrows) and perforation (at bottom) of duodenum from rats given propionitrile (PAS, $\times 39$).

Table 1. Effect of acrylonitrile, propionitrile, and cysteamine on duodenum and adrenal gland.

Compound	Duodenal ulcer ^a	Adrenal necrosis ^a	Mortality ^b	Dose, mg/100 g
Acrylonitrile $\text{CH}_2=\text{CH}-\text{CN}$	0/10	10/10	10/10	20, (IV, $\times 1$)
Propionitrile $\text{CH}_3-\text{CH}_2-\text{CN}$	8/10	0/10	10/10	6 (1st day), 8 (2nd day), 10 (3rd, 4th day); SC $\times 3$
Cysteamine $\text{H}_2\text{N}-\text{CH}_2-\text{CH}_2-\text{SH}$	9/10	8/10	2/10	30, PO $\times 3$ or 100, PO $\times 1$

^a Ratio positive/total.

^b Ratio dead/total.

It can also be assumed that the ulcerogenic effect of drugs is mostly, but not exclusively, related to a two-carbon group bearing radical(s), e.g., $-\text{CH}_3$, $-\text{CN}$, $-\text{SH}$, $-\text{NH}_2$. One of the most important of these observations is the fact that ethylamine, a part of the histamine molecule 4-(2-ethylamine)imidazole, which has

been known for a long time to be a potent stimulant of gastric secretion, easily produces duodenal ulcer. It has also been described that 24-hr infusion of histamine in rats results in duodenal ulcer (10,11). Our findings, however, demonstrate that the entire histamine molecule is not necessary for this action! The two-carbon

Table 2. Production of duodenal ulcer and adrenal necrosis by alkyl compounds.

Compound	Duodenal ulcer	Adrenal necrosis	Mortality	Dose, mg/100 g
Nitriles				
Propionitrile CH ₃ -CH ₂ -CN	8/10	0/10	10/10	6 (1st day), 8 (2nd day), 10 (3rd, 4th day); SC (200 g)
<i>n</i> -Butyronitrile CH ₃ -CH ₂ -CH ₂ -CN	8/10	2/10	4/10	10 (1st, 2nd day), 20 (3rd, 4th day); SC (200 g)
3-Bromopropionitrile Br-CH ₂ -CH ₂ -CN	2/5	3/5	5/5	10; PO
Ethylenedicyanide NC-CH ₂ -CH ₂ -CN	2/5	2/5	2/5	4 (1st day), 8 (2nd day), 10 (3rd, 4th day); SC
3-Aminopropionitrile H ₂ N-CH ₂ -CH ₂ -CN	0/5	2/5	5/5	50; SC
Amines				
Methylamine CH ₃ -NH ₂	1/5	0/5	5/5	20; PO
Ethylamine CH ₃ -CH ₂ -NH ₂	4/8	3/8	8/8	120; SC (200 g)
Propylamine CH ₃ -CH ₂ -CH ₂ -NH ₂	1/5	0/5	4/5	40; PO
Butylamine CH ₃ -CH ₂ -CH ₂ -CH ₂ -NH ₂	0/5	0/5	5/5	30; PO
Ethanolamine HO-CH ₂ -CH ₂ -NH ₂	0/5	0/5	1/5	50; PO or SC
Aminothiols				
Cystamine·2 HCl (H ₂ N-CH ₂ -CH ₂) ₂ S ₂	4/5	5/5	5/5	20; PO
2- <i>n</i> -Butylaminoethanethiol CH ₃ -(CH ₂) ₃ -NH-CH ₂ -CH ₂ -SH	5/10	4/10	5/10	20; PO
Thiols				
Ethanethiol CH ₃ -CH ₂ -SH	0/5	0/5	0/5	20 (1st, 2nd day), 40 (3rd day); PO
2-Diethylaminoethanethiol (CH ₃ -CH ₂ -NH) ₂ -CH ₂ -CH ₂ -SH	2/5	0/5	4/5	25; PO
1,2-Propanedithiol HS-CH ₂ -CH ₂ -CH ₃	4/5	0/5	2/5	0.5 (1st day), 2 (2nd day), 5 (3rd, 4th day); SC
1,3-Propanedithiol HS-CH ₂ -CH ₂ -CH ₂ -SH	2/5	0/5	5/5	4 (1st day), 8 (2nd day), 10 (3rd, 4th day); SC
1-Butanethiol HS-CH ₂ -CH ₂ -CH ₂ -CH ₃	0/5	2/5	0/5	20 (1st, 2nd day), 40 (3rd, 4th day); PO
1,4-Butanedithiol HS-CH ₂ -CH ₂ -CH ₂ -CH ₂ -SH	1/5	2/5	5/5	15; PO

ethylamine (fragment of histamine) is capable of producing perforating duodenal ulcers.

Adrenal necrosis and duodenal ulcer produced by alkyl derivatives and some other compounds may be useful animal models to study the mechanism of development of these lesions. Until recently, only a few drugs, e.g., 7,12-dimethylbenzo[*a*]anthracene (12), hexadimethrine bromide (13,14), thioguanine (15), thioacetamide (16), and basic polyglutamic acids (17) were known to produce adrenal necrosis in rats. Among these compounds acrylonitrile acts the most rapidly; adrenal hemorrhage and necrosis (resembling the Waterhouse-Friderichsen syndrome in man) occur

within 1-2 hr, and most of the animals die within 3-4 hr.

Similarly, there are no readily available models for study of duodenal ulcer. It has long been known that stress and various compounds cause gastric but not duodenal ulcers in rats (18). Only chronic deficiency of pantothenic acid in certain strains of rats (19) and infusion of histamine and other secretagogues (e.g., carbachol, pentagastrin) in fasted rats (10,11) are known to produce duodenal ulcers. Most of the compounds presented here cause similar lesion after a single SC or PO dose or after multiple injections at short intervals.

Although there is no overt evidence that these chemicals may in fact be responsible for the production of duodenal ulcers or adrenal apoplexy or atrophy in man, this possibility must be considered in the future.

Acknowledgement

Most of the experiments reported were performed at the Institute of Experimental Medicine and Surgery at the University of Montreal. Hence, special thanks are due to Dr. Hans Selye.

REFERENCES

1. Allara, D. L. Aging of polymers. *Environ. Health Perspect.* 11: 29 (1975).
2. Szabo, S., and Selye, H. Adrenal apoplexy and necrosis produced by acrylonitrile. *Endokrinologie* 57: 405 (1971).
3. Szabo, S., and Selye, H. Adrenal apoplexy produced by acrylonitrile and its prevention by ACTH and hypophysectomy. *Fed. Proc.* 30: 307 (1971).
4. Szabo, S., and Selye, H. Effect of phenobarbital and steroids on the adrenal apoplexy produced by acrylonitrile in rats. *Endocrinol. Exp.* 6: 141 (1972).
5. Szabo, S., and Selye, H. Duodenal ulcers produced by propionitril in rats. *Arch. Pathol.*, 93, 390 (1972).
6. Selye, H., and Szabo, S. An experimental model for the production of perforating duodenal ulcers by cysteamine in the rat. *Nature* 244: 458 (1973).
7. Szabo, S. New approach to experimental ulcerogenesis: Production of duodenal ulcer by derivatives of propionitrile and cysteamine. *Proc. 5th World Congr. Gastroenterol., Mexico City* (1974).
8. Szabo, S., et al. Effect of propionitrile and cysteamine on gastric secretion in rats. *Clin. Res.*, 22: 370A (1974).
9. Robert, A., et al. A new model to test antiulcer agents, cysteamine-induced duodenal ulcers. *Fed. Proc.* 33: 310 (1974).
10. Robert, A., Stout, D. J., and Dale, J. E. Production by secretagogues of duodenal ulcers in the rat. *Gastroenterology* 59: 95 (1970).
11. Robert, A., Stowe, D. F., and Nezamis, J. E. Possible relation between smoking and peptic ulcer. *Nature* 433: 497 (1971).
12. Huggins, C., and Morii, S. Selective adrenal necrosis and apoplexy induced by 7,12-dimethylbenz[α]anthracene. *J. Exp. H. Med.*, 114: 741 (1961).
13. Selye, H., Gabbiani, G., and Tuchweber, B. Organ lesions produced by hexadimethrine and their modification by various agents. *Med. Exp.* 8: 74 (1966).
14. Kovacs, K., Carroll, R., and Tapp, E. The pathogenesis of hexadimethrine necrosis of the pituitary and adrenal. *Arzneimittel-Forsch.* 16: 516 (1966).
15. Szabo, S., et al. Production of hemorrhagic necrosis in the rat adrenal by thioguanine or thioacetamide and its prevention by hypophysectomy. *Proc. Can. Fed. Biol. Soc.* 15: 341 (1972).
16. Lazar, G., and Szabo, S. Effect of hypophysectomy and catatoxic steroids on thioacetamide-induced adrenal lesions in rats. *Endokrinologie* 61: 97 (1973).
17. Lazar, G., Szabo, S., and Husztik, E. The experimental production of hemorrhagic lesions in the rat adrenal, liver and lung by basic polyglutamic acid derivatives. *Res. Exp. Med. (Z. Ges. Exp. Med.)* 159: 58 (1972).
18. Selye, H. *Hormones and Resistance*. Springer Verlag, Berlin, 1971.
19. Berg, B. N., Zucker, T. F., and Zucker, L. M. Duodenal ulcers produced on a diet deficient in pantothenic acid. *Proc. Soc. Exp. Biol. Med.*, 71: 374 (1949).