

A LuxS-Dependent Cell-to-Cell Language Regulates Social Behavior and Development in *Bacillus subtilis*

Esteban Lombardía, Adrián J. Rovetto, Ana L. Arabolaza, and Roberto R. Grau*

Departamento de Microbiología, Facultad de Ciencias Bioquímicas y Farmacéuticas, FCByF-UNR e Instituto de Biología Molecular y Celular de Rosario, IBR-CONICET, Suipacha 531, Rosario (2000), Argentina

Received 30 January 2006/Accepted 4 April 2006

Cell-to-cell communication in bacteria is mediated by quorum-sensing systems (QSS) that produce chemical signal molecules called autoinducers (AI). In particular, LuxS/AI-2-dependent QSS has been proposed to act as a universal lexicon that mediates intra- and interspecific bacterial behavior. Here we report that the model organism *Bacillus subtilis* operates a luxS-dependent QSS that regulates its morphogenesis and social behavior. We demonstrated that *B. subtilis* luxS is a growth-phase-regulated gene that produces active AI-2 able to mediate the interspecific activation of light production in *Vibrio harveyi*. We demonstrated that in *B. subtilis*, luxS expression was under the control of a novel AI-2-dependent negative regulatory feedback loop that indicated an important role for AI-2 as a signaling molecule. Even though luxS did not affect spore development, AI-2 production was negatively regulated by the master regulatory proteins of pluricellular behavior, SinR and Spo0A. Interestingly, wild *B. subtilis* cells, from the undomesticated and probiotic *B. subtilis* natto strain, required the LuxS-dependent QSS to form robust and differentiated biofilms and also to swarm on solid surfaces. Furthermore, LuxS activity was required for the formation of sophisticated aerial colonies that behaved as giant fruiting bodies where AI-2 production and spore morphogenesis were spatially regulated at different sites of the developing colony. We proposed that LuxS/AI-2 constitutes a novel form of quorum-sensing regulation where AI-2 behaves as a morphogen-like molecule that coordinates the social and pluricellular behavior of *B. subtilis*.

Bacteria not only behave as self-sufficient individuals but also act as communities capable of cell-cell communication (7, 15, 20, 42, 53). This social interaction leads to the coordination of communitarian activities that resemble, in their complexity, the behaviors observed in multicellular organisms (20, 42, 49, 53, 57). This microbial phenomenon is known as quorum sensing, a process by which bacteria monitor their cell population density by measuring the concentration of small secreted signal molecules called autoinducers (57). Even though a vast number of diverse quorum-sensing systems exist (41, 46, 57), they can be divided into two established paradigms that regulate the intraspecific behavior in many bacteria: (i) LuxI/LuxR-type quorum-sensing systems in gram-negative bacteria responsible for the production of *N*-acyl-L-homoserine lactone autoinducers or type I autoinducers and (ii) oligopeptide/two component-type quorum-sensing circuits in gram-positive bacteria responsible for the production of autoinducer peptides (4, 17, 26, 32, 44).

However, in recent years, a considerable amount of information has been gained on the existence of another type of quorum-sensing system that not only seems to participate in intraspecific bacterial behavior but also seems to regulate the interspecific interactions among bacteria of different genera. This system involves the production of autoinducer-2 (AI-2) signal molecules, mediated by the activity of the LuxS enzyme, in response to cell density (15, 46, 59, 62). It has been dem-

onstrated in numerous studies that the AI-2 molecules present in the supernatant of luxS-positive bacteria have the capacity to induce the AI-2 reporter species *Vibrio harveyi*, the model organism originally selected for the study of AI-2-dependent quorum sensing (57). LuxS/AI-2 have been shown to control a variety of cellular processes, such as production of pathogenicity factors, toxin production, biofilm formation, and swarming motility (14, 16, 28, 31, 34, 37, 38, 50, 58). More recently, it has been shown under laboratory conditions that interspecific luxS-dependent quorum sensing from *E. coli* modulated luxS-dependent quorum sensing in *V. harveyi* and pathogenic *V. cholerae* (63). Therefore, LuxS and its product AI-2 have been proposed to serve as a universal language (bacterial Esperanto or universal bacterial lexicon) used by bacteria to mediate their intra- and interspecific behaviors (59, 62).

Alternatively, it has been proposed that LuxS, the synthase for AI-2, has another role in the cell, in which it functions as an integral component of the activated methyl cycle (AMC) to avoid (jointly with the activity of the Pfs enzyme; see below) the intracellular accumulation of the toxic metabolite *S*-adenosyl-L-homocysteine (SAH). In fact, AI-2 is a by-product of the AMC, which recycles *S*-adenosyl-L-methionine (SAM), the main methyl donor in cells (35, 51, 55, 60, 61). As part of the AMC, SAM is converted to *S*-adenosyl-L-homocysteine (SAH), which is subsequently detoxified by the Pfs enzyme to generate adenosine and the sole intracellular source of substrate for LuxS, *S*-ribosyl-homocysteine (SRH). LuxS then produces the precursor of AI-2, 4,5-dihydroxy-2,3-pentanedione (DPD) during the conversion of SRH to homocysteine. DPD is able to undergo spontaneous cyclization to either of the two forms (R or S) of 2,4-dihydroxy-2-methylhydro-3-furanone (DHMF). Hydration of *S*-DHMF yields *S*-TMHF that subse-

* Corresponding author. Mailing address: Departamento de Microbiología, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531, Rosario (2000), Argentina. Phone: (54) 341-4353377. Fax: (54) 341-4350661. E-mail: robertograu@fulbrightweb.org.

TABLE 1. *Bacillus subtilis* strains

Strain	Relevant genotype	Comments and/or source (reference) ^a
RG1337 (JH642)	Wild-type domesticated	Laboratory stock (2)
RG1338	<i>amyE::luxS-lacZ cat</i>	This study
RG1339	<i>luxS::pJM103 cat</i>	This study
RG1340	<i>luxS::pJM103 cat::spc</i>	pCm::Spc (BGSC) ^b →RG1339
RG1341	<i>amyE::luxS-lacZ cat luxS::pJM103 cat::spc</i>	RG1338→RG1339
RG4365	Wild-type undomesticated natto strain	Laboratory stock (A. Nakamura)
RG4366	<i>amyE::luxS-lacZ cat</i>	This study
RG4367	<i>luxS::pJM103 cat</i>	This study
RG4368	<i>amyE::sspB-lacZ cat</i>	RG348 (36)→RG4365
RG4369	<i>amyE::luxS-lacZ cat luxS::pJM103 cat::spc</i>	This study
RG12604	<i>amyE::abrB-lacZ cat</i>	Laboratory stock (2)
RG1342	<i>amyE::abrB-lacZ cat luxS::pJM103 cat::spc</i>	RG12604→RG1340
RG438	<i>amyE::sinR-lacZ cat</i>	Laboratory stock (18)
RG1343	<i>amyE::sinR-lacZ cat luxS::pJM103 cat::spc</i>	RG438→RG1340
RG19005	<i>amyE::spo0A-lacZ cat</i>	Laboratory stock (36)
RG1344	<i>amyE::spo0A-lacZ cat luxS::pJM103 cat::spc</i>	RG19005→RG1340
RG1345	<i>amyE::luxS-lacZ cat ΔabrB::spc</i>	RG1338→RG12607 (2)
RG1346	<i>amyE::luxS-lacZ cat ΔsinR::phea</i>	RG1338→RG432 (18)
RG1347	<i>amyE::luxS-lacZ cat Δspo0A::ery</i>	RG955 (36)→RG1338
RG4380	<i>amyE::luxS-lacZ cat Δspo0A::ery</i>	RG955 (36)→RG4366
RG4381	<i>amyE::luxS-lacZ cat ΔsinR::phea</i>	RG432 (18)→RG4366

^a Strain construction is indicated by an arrow. Chromosomal DNA or plasmid DNA listed at the tail of the arrow was used to transform the strains listed at the head of the arrow.

^b BGSC, *Bacillus* Genetic Stock Center, Ohio State University, Columbus, Ohio.

quently forms a diester with boric acid to generate the first characterized active form of AI-2 with a key role in the regulation of quorum sensing of marine *V. harveyi* (11, 47). On the other hand, *R*-DHMF hydrates to form *R*-THMF, the active form of enteric AI-2 that has recently been cocrystallized with its putative cellular transporter/receptor LsrB (39, 54). Furthermore, *R*-THMF, once internalized, is phosphorylated to sequester it into the cytoplasm, where *R*-THMF-phosphate might function as an active intracellular autoinducer (39, 54). The findings that more than one type of furanone-derived molecule, depending on the bacterial species and the surrounding environment, can behave as the AI-2 in quorum sensing made it necessary to use the expression “AI-2” as a generic term to indicate a family of interconverting molecules with autoinducer activity (*R*-THM-furanone, *S*-THM-furanone-borate, and *R*-THM-furanone-phosphate) (see the legend to Fig. 2A for details and references 55 and 57).

So far, LuxS/AI-2-dependent quorum sensing has been studied in diverse gram-positive and gram-negative bacteria (55, 57), but it is far from clear how widely this AI-2 family of autoinducers is used in bacterial signaling during the expression of wild behaviors in nature (33, 35, 51, 55, 57, 61). One microorganism that serves as an excellent model to study the regulation of cell development and social behavior is the gram-positive spore-forming soil-bacterium *Bacillus subtilis* (5, 10, 12, 21, 56). At present, there are two well-studied cell density-dependent signaling systems that have been described in this bacterium. These two QSS depend on the production of specific autoinducer peptides that are encoded by *com* and *phr* genes that control sporulation and competence development (21, 26, 32, 44). However, it has been recently reported that the detection of a LuxS-dependent signaling in *Bacillus anthracis* and *Bacillus cereus* with implicitness in density-dependent gene expression and pathogenesis for both bacteria (3, 27). Both

luxS orthologs, from *B. cereus* and *B. anthracis*, show a high degree of identity with a putative *luxS* ortholog located on the chromosome of the sequenced reference strain 168 of *B. subtilis*. Therefore, we were intrigued to know whether or not a LuxS/AI-2-dependent signaling system operates in *B. subtilis* and, if this is the case, what consequences it would have on the social behavior and developmental potential of this model organism.

MATERIALS AND METHODS

Bacterial strains and growth conditions. The two wild-type *B. subtilis* strains used in this study were the domesticated, laboratory-reference strain RG1337 (JH642 laboratory stock; James A. Hoch, The Scripps Research Institute, La Jolla, Calif.) and the undomesticated and wild *B. subtilis* natto strain RG4365 (laboratory stock; Akira Nakamura, University of Tsukuba, Japan). These two strains, as well as their isogenic derivatives (Table 1), were grown in Luria-Bertani broth (LB), Schaeffer's sporulation medium (SM), or Spizizen minimal salts medium supplemented with glucose (0.5%) plus the supplements as indicated. The antibiotics, when required, were used at the following final concentrations: 5 μg/ml chloramphenicol, 100 μg/ml spectinomycin, 1 μg/ml erythromycin, and 2.5 μg/ml pleuromycin. For sporulation efficiency, *B. subtilis* strains were grown in SM for 20 h and then treated with 10% CHCl₃ for 15 min before being plated as previously described (2). Transformation of *B. subtilis*, to obtain isogenic derivatives of the parental strains, was carried out as previously described (18). β-Galactosidase assays from *B. subtilis* strains harboring *lacZ* fusions were assayed as described previously and the specific activity was expressed in Miller units (2). The β-galactosidase experiments described in the figure legends were independently repeated three to five times, and a representative set of results is shown in each figure. All cultures were grown at 37°C. *V. harveyi* BB170 was kindly provided by B. Bassler (Princeton University) and grown in AB medium overnight at 30°C as described previously (6, 19). AB medium consists of 10 mM K₂HPO₄ (pH 7.0), 0.3 M NaCl, 0.05 M MgSO₄, 0.2% vitamin-free Casamino Acids (Difco), 2% glycerol, and 1 mM L-arginine.

For the swarming analysis, we used B medium (1), which contains 15 mM (NH₄)₂SO₄, 8 mM MgSO₄ · 7H₂O, 27 mM KCl, 7 mM sodium citrate · 2H₂O, and 50 mM Tris-HCl (pH 7.5) supplemented with 0.6 mM KH₂PO₄, 2 mM CaCl₂ · 2H₂O, 1 mM FeSO₄ · 7H₂O, 10 mM MnSO₄ · 4H₂O, 4.5 mM glutamic acid, 780 μM tryptophan, 860 μM lysine, and 0.2% (wt/vol) glucose. All plates




FIG. 1. Organization and biological activity of *luxS* of *B. subtilis*. (A) Representation of the DNA region containing *luxS* and its adjacent genes. In addition, we showed the location and size of the DNA segments amplified and used for plasmid and strain constructions (see Materials and Methods for details). Rho-independent transcriptional terminators and the relevant restriction sites used for cloning are shown on the diagram. (B) The wild-type reporter strain RG1338 (*luxS-lacZ::amyE*) was grown in SM and growth was measured as the increase in optical density at 525 nm; samples were collected at the indicated times and assayed for β -galactosidase activity expressed in Miller units (2, 36). T_0 represents the transition from vegetative to stationary phase. Essentially, the same pattern of *luxS* expression (with a maximum of 30 to 45 Miller units at T_{-1}) was observed with the *B. subtilis* natto strain RG4366 (data not shown). (C) Induction of bioluminescence in *V. harveyi* strain BB170 by CFCM prepared from wild-type RG1337 cells and its isogenic *luxS*-negative derivative RG1338 (Table 1). At time zero, CFCM from the different strains was added to the *V. harveyi* culture at a final concentration of 10% (vol/vol) and light production was recorded. Symbols for the addition of CFCM at each experimental condition are as follows: ■, *V. harveyi* plus CFCM from RG1337 cells; ○, *V. harveyi* plus CFCM from RG1338 cells; ◆, *V. harveyi* plus heated (80°C, 20 min) CFCM from RG1337 cells; ×, *V. harveyi* without CFCM addition. Data for a representative set of experiments are shown. Essentially, the same results as the ones obtained with CFCM from RG1337 and RG1338 cells were obtained using CFCM from the *B. subtilis* natto strains RG4365 and RG4367, respectively, and the *V. harveyi* BB170 strain (data not shown).

were prepared by supplementing the medium with 0.7% of agar and used during the day.

Detection of SAM and SAH. For the detection of SAM and SAH, *B. subtilis* strains were grown in Spizizen minimal salts medium until 1 h after the end of the exponential phase. Standard reagents, SAM iodide salt, SAH, and 1-heptanesulfonic acid sodium salt were all obtained from Sigma Chemical Company. SAM and SAH levels were determined by reverse-phase ion-pair high-performance liquid chromatography (Δ KTA basic high-performance liquid chromatograph [Amersham]). To prepare the high-performance liquid chromatography extracts, frozen pellets from cultures were resuspended in chilled 0.01 N hydrochloric acid (HCl) and sonicated on ice. After centrifugation, supernatants were filtered through a 0.2- μ m filter (Orange Scientific) at 4°C. The mobile phase consisted of 87% 5 mM heptanesulfonic acid (Sigma Co.), with the pH adjusted to 3.2 with HCl and 13% acetonitrile (EM Science). The flow rate was 1 ml/min through an Altech C_{18} column (4.6 by 250 mm), and the absorbance was monitored at 254 nm. The amounts of SAM and SAH in the samples were determined by comparison with a standard solution of 10 μ M each of SAM and SAH in 0.01 N HCl. All runs, including standards, were done three times. SAM and SAH levels were expressed as picomoles per 1×10^9 CFU of the corresponding *B. subtilis* culture.

Plasmids and strains constructions. Strains RG1338 and RG4366 are isogenic derivatives from RG1337 and RG4365, respectively, that carry a transcriptional reporter fusion to the *luxS* promoter region (Table 1). To obtain the promoter region of *luxS*, we proceeded to a PCR amplification of chromosomal DNA from strain RG1337 using the oligonucleotides 5' TTCTTCAGGTACCTTTCTGAT GCAAG 3' and 5' TTATTAGGATCCCGTCTGTCCAC 3' (the introduced restriction sites for KpnI and BamHI are underlined; boldface indicates substitutions used to create the restriction site). The PCR product (230 bp) containing the promoter region of *luxS* was cloned into the *mcs* of the vector pJM116 to make a transcriptional fusion to the *lacZ* gene of *E. coli* (2, 56). The resulting plasmid (pER1) (Fig. 1A) was linearized and integrated into the chromosome of wild-type competent cells via double crossover at the *amyE* locus by transformation and selection for resistance to chloramphenicol (2, 56). For the construction of *B. subtilis* *luxS*-negative mutant strains (RG1339 and RG4367), we amplified a 420-bp internal fragment of *luxS* by using chromosomal DNA from strain RG1337 as a template and the oligonucleotides 5' ggTTgTggATCCATATgTA AgACATTgCg 3' and 5' TTCTTCTggATCCTgTgAAAgCCAgAAACgC 3'. The amplified PCR fragment was cloned into the integrative vector pJM103 (2, 57) at the unique BamHI site, generating pER2 (Fig. 1A). This plasmid was used to transform competent cells of RG1337 and RG4365 and to select, by a single-

crossover event, chloramphenicol-resistant colonies that were screened by Southern blotting and PCR to corroborate the integration of pER2 into the *luxS* locus. To change the antibiotic resistance marker (chloramphenicol resistance) of the strains to one of a spectinomycin resistance, we used the plasmid pCm::Spc isolated from the *E. coli* ECE74 strain (*Bacillus* Genetic Stock Center). For the overexpression of *luxS* in the multicopy vector pHT315 (25 to 50 copies/cells), we amplified from strain RG1337 a truncated version of *luxS* lacking its promoter region but containing its own ribosomal binding and ATG sites (plasmid pER3) (Fig. 1A). To this end, we used oligonucleotides 5' gAAATTACATTTCTgCag AAggggAgAg 3' and 5' CCTAAATggTACCAAACgCTTACAgC 3' to clone the amplified PCR fragment between the PstI and KpnI restriction sites of pHT315, giving rise to the multicopy plasmid pER3 (Fig. 1A). This plasmid and the vector pHT315 were introduced into competent cells of the *luxS*-negative mutant reporter strain RG1341 (Table 1) and selected for resistance to erythromycin (15 μ g/ml).

AI-2 bioassay. *B. subtilis* strains were cultured in LB for 12 h (to stationary phase). The stationary-phase cultures were then inoculated into LB at a 1:20 dilution and grown until mid-exponential phase. The culture supernatants were collected by centrifugation at $8,000 \times g$ and passed through a 0.45- μ m filter (Orange Scientific) to remove cells. This cell-free conditioned medium (CFCM) was stored at -20°C until studied. The bioassay was performed essentially as described previously (63). Briefly, the *V. harveyi* BB170 *luxN::Tn5*, AI-1 sensor-negative AI-2 sensor-positive (6, 63) reporter strain was grown for 16 h with aeration at 30°C in AB medium, and then diluted 1:10,000 in fresh AB medium. Cell-free culture supernatants of *B. subtilis* strains to be tested for AI-2 activity were then added to the diluted *V. harveyi* culture at a 10% (vol/vol) final concentration. The cultures were shaken at 30°C and light production was measured every hour.

Biofilm (pellicle) formation. Overnight cultures of *B. subtilis* were grown in LB to stationary phase. Then, 50 μ l of these cultures were diluted in 2 ml of fresh LB supplemented with 10% yeast extract and incubated at 37°C for 48 h.

Swarming motility. B medium (1) fortified with 0.7% agar was inoculated with 1 μ l of 2×10^8 cells/ml grown to mid-log phase at 37°C in LB. The inoculated petri dishes were then incubated at 37°C for 24 h.

Fruiting body formation. For colony architecture analysis, the *luxS*⁺ and *luxS*-negative natto strains were grown for 12 h in LB, reaching a cell density of 5×10^7 viable cells/ml, and then 5 μ l of each culture were placed onto SM-agar plates supplemented with 10% yeast extract plus (when indicated) 5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside (X-Gal, 80 μ g/ml) and incubated at 37°C .

Microscope observations. The developed swarm, biofilm and fruiting-body plates were visualized with a Stemi 2000 (Carl Zeiss, Germany) stereomicroscope using a KL1500LCD (Carl Zeiss, Germany) illumination system. A PowerShot A80 system (Canon) was used to capture the photographs for swarm, fruiting bodies, and biofilm images.

RESULTS

***B. subtilis luxS* gene is actively transcribed, and its product, AI-2, mediates interspecific signaling with *V. harveyi*.** AI-2-dependent quorum sensing signaling is currently found in more than 55 gram-positive and gram-negative bacterial species, leading to the suggestion that AI-2 constitutes a universal language for interspecies communication (15, 46, 50, 53, 57, 59). To uncover the situation in *B. subtilis*, we characterized the pattern of expression of the *B. subtilis luxS* ortholog and the supposed biological activity of its product (AI-2) using the *V. harveyi* bioassay. An examination of the annotation of the complete genome sequence of the *B. subtilis* European strain 168 (<http://genolist.pasteur.fr/SubtilList>) showed only one ortholog (formerly named *yjtB*) of the *V. harveyi luxS* gene (data not shown). *luxS* of *B. subtilis* seems to be organized as a monocistronic unit of 471 bp flanked by rho-independent transcriptional terminators and not linked to any specific gene class (Fig. 1A and data not shown). To examine the *in vivo* activity of *luxS* in *B. subtilis*, we cloned the 5' end of this gene as a *lacZ* transcriptional fusion (strain RG1338) (Table 1). As shown in Fig. 1B, the use of this construct permitted demonstration that *luxS* was actively transcribed. *luxS* expression showed a continuous augmentation of its activity during the exponential growth of *B. subtilis* in LB, reaching a maximum value of 20 to 25 Miller units of β -galactosidase activity one hour before the onset of the stationary phase (T_0), and decreased upon entry into stationary phase (Fig. 1B).

The temporally regulated activity of the *luxS* promoter opened the possibility that *B. subtilis* operates a previously undetected LuxS-dependent quorum-sensing signaling system apart from the well-described autoinducer peptide-dependent quorum-sensing systems that regulate sporulation and competence development (21). Therefore, to confirm that *B. subtilis* produced active AI-2, we recur to the use of the *V. harveyi* AI-2 reporter assay (6, 63). This AI-2 assay takes advantage of a deficiency in AI-1 sensor signaling in the *V. harveyi* strain BB170. Without the *luxN* AI-1-encoded sensor, strain BB170 can exhibit only bioluminescence in response to AI-2 (6, 63). When an overnight culture of the *V. harveyi* strain BB170 is diluted 1:10,000 (to yield low cell density), the level of endogenous AI-2 is reduced below the threshold required for luminescence. Under these experimental conditions, the addition of exogenous cell-free supernatants from bacteria possessing LuxS activity (and, hence, producing AI-2) can restore the bioluminescence phenotype of the BB170 cells (6, 57, 63). Therefore, we prepared CFCM from the wild-type strain RG1337 grown in LB until 1 h before T_0 . This CFCM was added to the BB170 cells prepared as indicated above, and the luminescence induced by *B. subtilis* culture supernatants was measured (Fig. 1C). As observed, the cell-free supernatant from the wild-type strain RG1337 significantly induced the luminescence of BB170 (Fig. 1C). In contrast, the luminescence of BB170 was induced neither by heat-treated CFCM (to destroy AI-2-dependent activity) of wild-type RG1337 cells nor

by supernatant of the RG1339 isogenic *luxS*-negative mutant strain (absence of AI-2 activity). These results indicated that *B. subtilis* harbors an active *luxS* gene (Fig. 1B) that codes for an AI-2-mediated activity able to be recognized as the AI-2-specific signaling pathway of *V. harveyi* (Fig. 1C).

Transcription of *luxS* is regulated by a negative autoregulatory feedback loop. The LuxS enzyme is responsible for the last enzymatic step of AI-2 synthesis (43, 45, 47, 55, 57), but at the same time, it has an important function within central metabolism as part of the activated methyl cycle (AMC) of the cell for recycling (jointly with Pfs) of the toxic intermediate SAH (51, 55, 57, 61, 62). In contrast to what has been reported in *Salmonella enterica* serovar Typhimurium and other bacteria for which *luxS* expression is constitutive (8, 55, 57), we demonstrated that in *B. subtilis*, *luxS* activity was temporally regulated (Fig. 1B). Therefore, it was interesting to analyze whether the synchronized expression of *luxS* was growth-phase regulated as a reflection of cell density (quorum sensing) or, in contrast, if *luxS* was regulated by central metabolism in order to maintain appropriate nontoxic levels of the intermediates of the AMC (Fig. 2A). To this end, we recurred to the use of an RG1337-isogenic *luxS*-negative mutant strain that harbored a *luxS-lacZ* reporter fusion at the nonessential *amyE* locus (strain RG1341) (Table 1). This construct permitted analysis of the expression of *luxS* in the absence of AI-2-dependent quorum sensing. First, we noticed that the absence of LuxS activity did not produce any detrimental effect on growth or final cellular yield of the *luxS*-negative mutant strain, in comparison to the wild-type strain (Fig. 2B and data not shown). More interesting, as is shown in Fig. 2B, the *luxS*-dependent β -galactosidase levels of the AI-2-deficient RG1341 culture far exceeded the levels of β -galactosidase activity accumulated by a culture of AI-2-proficient cells throughout the growth period (strain RG1338) (Table 1). Since the growth of the *luxS*-negative mutant strain was indistinguishable from the growth of the wild-type strain (Fig. 2B) and the two cultures reached similar cellular yields (data not shown), it can be assumed that the accumulation of intermediates of the AMC did not reach growth-toxic levels. To support this conclusion, we measured the intracellular levels of SAM and SAH accumulated by LuxS-proficient and LuxS-deficient cells. As shown in Fig. 2C, even though the intracellular levels of both intermediates of the AMC were a little higher in LuxS-deficient cells, their concentrations (less than nM) were far lower than the concentrations reported to be toxic in bacteria (33, 40, 43, 51, 55). Therefore, the dramatic difference in activity of the *luxS* gene in favor of the AI-2-deficient mutant strain (Fig. 2B) was not due to intracellular toxicity.

In contrast, we hypothesized that the regulation of *luxS* transcription was (i) metabolically regulated by metabolites of the AMC and/or (ii) cell density-regulated by AI-2 (Fig. 2A). In order to distinguish between these possibilities, we measured the activity of the *luxS-lacZ* reporter fusion in LuxS-proficient and LuxS-deficient cells grown in the presence of intermediates of the AMC or in the presence of CFCM prepared from wild-type (presence of AI-2) or *luxS*-negative (absence of AI-2) cultures. As shown in Fig. 2D, the activity of the *luxS* promoter in the wild-type strain was not affected by the addition of SAH (or other intermediates of the AMC, such as SAM and Met [data not shown]) or by the addition of CFCM




FIG. 2. LuxS activity regulates *luxS* expression in *B. subtilis*. (A) A diagram of the activated methyl cycle (AMC) of the cell pointing out the nature of the products with AI-2 activity derived from the activity of the LuxS enzyme. Also shown are two possibilities, discussed in the text, for the regulation of *luxS* expression. One is by metabolic regulation mediated by metabolites such as SAH (possibility 1), and the other is by quorum-sensing regulation by AI-2 (possibility 2). (B) Up-regulation of *luxS* expression in AI-2-deficient *B. subtilis* cells. Growth and β -galactosidase activity were measured as indicated. Growth (\square) and *luxS* activity (\blacksquare) of the RG1338 culture (wild-type strain, AI-2-positive) and growth (\circ) and *luxS* activity (\bullet) of the RG1341 culture (*luxS*-negative mutant strain, AI-2 deficient) are shown. Essentially the same results were obtained using the *B. subtilis* natto strains RG4366 and RG4369 instead of RG1338 and RG1341, respectively (data not shown). (C) Intracellular levels of SAM and SAH accumulated by AI-2-proficient (strain RG1337) and AI-2-deficient (strain RG1339) cells. A representative set of experiments made by triplicate is shown. Essentially the same levels of SAH and SAM were detected in cultures of the *B. subtilis* natto strains RG4365 and RG4367 (data not shown). (D) Regulation of *luxS* expression by CFCM containing (or not containing) AI-2 activity. The figure shows the β -galactosidase activity accumulated until 1 h after T_0 by cultures of the strains RG1338 (*luxS*⁺, AI-2-proficient) and RG1341 (*luxS* negative, AI-2 deficient) with each of them harboring a reporter *luxS-lacZ* fusion at the nonessential *amyE* locus (Table 1). Each culture of the strains RG1338 and RG1341 was grown in LB in the presence of the additions indicated in the figure. SAH was added at a final concentration of 1 mM; CFCM were added at a final concentration ranging from 1 to 5% (vol/vol) in each set of experiments repeated in triplicate (a representative set of data is shown). AI-2-positive CFCM and AI-2-deficient CFCM were derived from RG1337 and RG1339 cultures, respectively, as indicated in Materials and Methods. Essentially the same results were obtained using CFCM from cultures of the *B. subtilis* natto strains RG4365 (instead of CFCM from RG1337 cells) and RG4367 (instead of CFCM from RG1339 cells), and the results were corroborated by using the reporter natto strains RG4366 and RG4369 (data not shown). (E) In vivo down-regulation of *luxS* expression by up-regulation of AI-2 production. Growth and β -galactosidase activities expressed by the strain RG1341 harboring the multicopy plasmid pER3 (\circ and \bullet , respectively) overexpressing AI-2 and the control plasmid pHT315 (\square and \blacksquare , respectively).




FIG. 3. Spo0A and SinR are negative regulators of AI-2 quorum sensing. (A to C) Activity of the regulatory genes *abrB*, *sinR*, and *spo0A* in the presence and absence of LuxS activity. (A) RG12604 and RG1342; (B) RG438 and RG1343; (C) RG19005 and RG1344. Symbols: ■, LuxS-proficient cells; ●, LuxS-deficient cells. (D to F) Effects of AbrB, SinR, and Spo0A on *luxS* expression. (D) RG1338 (■) and RG1345 (●); (E) RG1338 (■) and RG1346 (●); (F) RG1338 (■) and RG1347 (●). Cells were grown in LB and samples were collected at the indicated times and assayed for β -galactosidase activity expressed in Miller units (18).

prepared from a *luxS*-negative culture. In contrast, the addition of wild-type-derived CFCM (presence of AI-2) to a wild-type culture produced a slight, but reproducible, inhibition of *luxS* activity (Fig. 2D). Moreover, the addition of CFCM containing AI-2, but not CFCM prepared from a *luxS*-negative culture (absence of AI-2), to a LuxS-deficient culture produced a significant reduction of the activity of the *luxS* promoter. These results strongly indicated that it was not the accumulation of intermediates of the AMC (possibility 1 in Fig. 2A) that resulted in the higher activity of the *luxS* promoter and suggested the existence of a negative AI-2-dependent autoregulatory loop operating on *luxS* expression. Supporting this view, the heat treatment (to destroy the activity of AI-2 molecules) of CFCM derived from wild-type cells resulted in the complete loss of the original inhibitory effect on the activity of the *luxS* promoter (Fig. 2D).

If, as we suggested, AI-2 has an inhibitory effect on *luxS* expression (possibility 2 in Fig. 2A), it can be postulated that the overproduction or accumulation of AI-2 should decrease *luxS* transcription. To test this hypothesis, we overexpressed *luxS* without its native promoter region (to avoid intrinsic regulation and/or titer effects) in the multicopy vector pHT315 (see Fig. 1A and Materials and Methods). This multicopy plasmid has been shown to overexpress cloned genes from constitutive internal promoters (2, 26, 56). As is shown in Fig. 2E, the

growth of the culture harboring the multicopy plasmid overexpressing *luxS* (pER3) (Fig. 1A) was not affected in comparison with the growth of the culture harboring pHT315 without insert. Confirming the original hypothesis of the existence of an AI-2-dependent negative feedback loop on *luxS*, the expression of *luxS* driven from its chromosomal copy with its native promoter showed a dramatic reduction throughout growth in the presence of the plasmid overproducing AI-2 (Fig. 2E).

AI-2 production is under the control of the master regulatory proteins of social behavior, Spo0A and SinR. The observation that *luxS* expression reached a maximum level at the end of the exponential phase of growth and declined after the commencement of the stationary phase (Fig. 1B) led us to further examine if key *B. subtilis* transition state regulators are involved in the regulation of *luxS* expression and vice versa (9, 10, 21). In particular, we investigated the *B. subtilis* transcription factors (Spo0A, AbrB, and SinR) that participate in quorum-sensing regulation of biofilm, swarming, and spore development (9, 10, 22, 29, 30). As shown in Fig. 3, the levels of expression of the developmental genes *abrB*, *sinR*, and *spo0A* were essentially the same in LuxS-proficient and LuxS-deficient cells, discarding the possibility that AI-2 would be affecting the expression of those master regulatory proteins (Fig. 3A to C). On the other hand, while *luxS* activity was not affected in AbrB-deficient cells (Fig. 3D), *luxS* expression was dramati-




FIG. 4. LuxS activity is required for biofilm formation in *B. subtilis*. Top-down images of pellicle (biofilms) made by LuxS-proficient (left panel) cells (strain RG4365) and LuxS-deficient (right panel) cells (strain RG4367) in LB fortified with 10% yeast extract after 2 days at 37°C without shaking. Note the differences in rugosity and architecture between both biofilms. Scale bar, 1 cm.

cally induced in Spo0A- and SinR-deficient cultures (Fig. 3E and F), showing the existence of a Spo0A/SinR-dependent negative-regulatory circuit decreasing *luxS* expression. Moreover, these findings suggested the possibility of a physiologic connection between AI-2-dependent quorum sensing and pluricellular social behaviors regulated by Spo0A and SinR (see below).

LuxS activity is required for biofilm formation and swarming motility in *B. subtilis*. LuxS and/or AI-2 have been shown to affect normal biofilm formation and swarming motility in different pathogenic bacteria (13, 57). The domesticated *B. subtilis* strains routinely used in research laboratories (i.e., strains 168 and JH642) do not develop a high degree of spatial organization and do not form highly structured or surface-associated communities. However, it has been recently reported that wild, undomesticated strains of *B. subtilis*, which were not subjected to random mutagenesis or to repeated rounds of selections for fast growth in rich culture media, were capable of social and multicellular behaviors (9, 10, 12, 22, 29, 30). At the moment, only the undomesticated *B. subtilis* NCIB 3610 isolate (or Marburg strain, from which the strains 168 and JH642 are derived) has been selected for the studies of multicellular behavior in bacilli (9, 10, 29). However, wild behaviors should be common and well-distributed attributes of most bacteria in nature (20, 42, 49). One type of undomesticated *B. subtilis* isolate whose social behavior is unreported and is a model of study in our laboratory because of its probiotic properties, is the Japanese *B. subtilis* natto strain RG4365 (Table 1 and references 5, 24, and 48). We confirmed that the pattern of expression of *luxS* in the *B. subtilis* natto strain RG4366 (Table 1) and its up-regulation in an AI-2-deficient background (strain RG4369) were indistinguishable from the results obtained with the domesticated strains RG1338 and RG1341, respectively (data not shown). In light of these findings, we were motivated to know to what extent LuxS would be required for the full manifestation of two previously described social behaviors of *B. subtilis* NCIB 3610 that were under Spo0A and SinR regulation: biofilm formation and swarming migration (10, 12, 29, 30). First, we noted that the wild-type *B. subtilis* natto strain (RG4365) formed, in standing liquid cultures incubated at 37°C for 48 h, robust and highly struc-




FIG. 5. LuxS activity is required for swarming motility in *B. subtilis*. Swarming ability of AI-2-proficient (strain RG4365) and AI-2-deficient (RG4367) cells after 24 h of incubation in B medium at 37°C. See Materials and Methods for details. Scale bar, 1 cm.

tured biofilms that were very similar to the previously reported thick and vein-like biofilms formed by strain NCIB 3610 (9, 10). In contrast, biofilm formation of the *luxS*-negative RG4367 natto strain was delayed and morphologically altered showing a clear final defect on pellicle formation and biofilm maturation (Fig. 4).

When we analyzed the ability of the LuxS-proficient and LuxS-deficient *B. subtilis* natto cells (strains RG4365 and RG4367, respectively) to swarm on solid surfaces, we found that LuxS activity was also essential for this cell density-dependent social motility behavior (Fig. 5). These results strongly argue for a novel level of regulation and complexity (LuxS-dependent quorum sensing) apart from the previously reported effects of Spo0A and SinR (12, 22), acting on the manifestation of the social and pluricellular behaviors of wild *B. subtilis* isolates.

LuxS activity produces a morphogen-like developmental signal crucial for pluricellular aerial architecture in *B. subtilis*. While colonies of the domesticated *B. subtilis* strains (i.e., 168 and JH642) are flat, smooth, and poorly differentiated (9, 10), colonies of the wild-type *B. subtilis* natto strain form thick and very sophisticated aerial structures. In Fig. 6, we show the time course of formation and maturation of the aerial colony architecture produced by the wild-type *B. subtilis* natto strain RG4365. Early in morphogenesis, the borders of the developing colony acquired a vein-like architecture with avenues of cells centrally converging to give rise, at later times (48 h to 72 h of development), to a palisade of cells that formed a central cylinder emerging from the base of the colony (Fig. 6). During the development of this elegant arrangement of differentiating cells, the cavity of the elevating cylinder (or cellular tower) remained open, full of a viscous fluid containing few trapped cells (data not shown). Later in development (96 h of development), the crest of the aerial tower was covered and closed with several layers of cells (Fig. 6) that finally differentiated into mature spores (see below and Fig. 7A).




FIG. 6. LuxS activity is required for aerial colony architecture in *B. subtilis*. Time course development of colony morphogenesis of AI-2-proficient (strain RG4365) and AI-2-deficient (strain RG4367) cells on solid SM fortified with 10% yeast extract. Scale bar, 1 mm.

Microbial aerial structures are typically sites for sporulation in fungi and filamentous bacteria. This is also the case for the microscopic fruiting bodies that were shown to project from the colonies of the wild strain NCIB 3610 (10). To prove that wild-type *B. subtilis* natto cells would also differentiate into spores in special regions of the developing aerial colony (Fig. 6), we used a construct for which the promoter region of the sporulation-specific gene *sspB*, which is expressed late in sporulation, was fused to the *lacZ* gene and integrated in a single copy into the chromosome of the wild-type *B. subtilis*

natto strain RG4365 (strain RG4368) (Table 1). This reporter strain was grown on solid medium that contained the X-Gal indicator of LacZ activity to point out the sites in the aerial colony where the spores were formed. As observed in Fig. 7A, spore morphogenesis was restricted to the pinnacle of the colony, a finding that indicated that spore development took place in a temporal and spatially regulated process that was highly coordinated with the formation of a sophisticated aerial colony. In fact, the extremely differentiated natto colony resembled a giant fruiting body (Fig. 7A). In the undomesticated NCIB 3610 (and also in myxobacteria), fruiting bodies are microscopic aerial projections of the colony (10), but in the case of *B. subtilis* natto, the whole aerial colony behaved as a unique and enormous fruiting body visible at first glance (Fig. 6 and 7A). Interestingly, when the behavior of the *luxS*-negative *B. subtilis* natto mutant strain was analyzed on solid medium, a tremendous defect in colony morphogenesis was immediately observed (Fig. 6). It took longer for the LuxS-deficient cells to make a structured colony (72 h or more) that finally failed to fully differentiate. In fact, LuxS-deficient cells were unsuccessful in their commitment to aerial-colony development because, at longer times of incubation (96 h or more), the immature AI-2-deficient aerial structure collapsed (Fig. 6).

In eukaryotes, during the course of development, cells of many tissues differentiate in response to the presence of gradients of substances that act as morphogens (52, 64). These developmental messenger molecules emanate from a restricted part of the tissue and spread away from their source to form a concentration gradient (52, 64). Since the pattern and the appearance of aerial fruiting body formation of the wild-type *B. subtilis* natto strain resembled the behavior and cell-coordination of a differentiating tissue, we were intrigued to know whether the pattern of *luxS* expression, and hence AI-2 production, was temporal and spatially regulated during development (25). Therefore, we analyzed the expression of *luxS* in *B. subtilis* natto during fruiting body morphogenesis. As observed in Fig. 7B, *luxS* was clearly expressed at the base of the fruiting body, supporting the view that AI-2 production was spatially restricted (probably by action of Spo0A and SinR; see Discussion and Fig. 7A and C) to a specific region of the developing colony from where it would be able to emanate. This result suggested the formation of an AI-2 gradient (bacterial morphogen) (25) that would diffuse or migrate through the incip-




FIG. 7. Spore development and AI-2-dependent cell-to-cell communication during fruiting body morphogenesis of wild-type *B. subtilis* cells. (A) Expression of the late sporulation gene *sspB* (strain RG4368) indicating that spore development takes place at the top of the developing colony in an AI-2-independent process (see the legend to Fig. 3C and the text for details). Spore morphogenesis is indicated by development of the blue color. (B) Expression of *luxS* (strain RG4366) from the bottom of the developing aerial colony, suggesting the formation of an ascending gradient of AI-2 molecules. Scale bar, 1 mm. (C) Expression of *luxS* in the absence of SinR (strain RG4381) and Spo0A (strain RG4380) after 16 h of incubation in LB at 37°C.

ient aerial colony (bacterial tissue) to coordinate appropriate gene expression during fruiting body morphogenesis.

DISCUSSION

luxS of *B. subtilis* shares a high degree of homology with *luxS* orthologs found in different bacteria, including *luxS* genes from *Escherichia coli*, *S. enterica* serovar Typhimurium, and *V. harveyi*, in which LuxS-dependent quorum sensing has been extensively studied (3, 23, 27, 45). Furthermore, the overexpression of the *B. subtilis luxS* gene in *E. coli* and the use of the produced *B. subtilis* LuxS enzyme made it possible to prove unambiguously that LuxS is a ribosylhomocysteinase that catalyzes the conversion of SRH to homocysteine and DPD, the precursor of all known active forms of AI-2 (23, 43, 45, 47, 60). However, the presence of an AI-2-dependent activity in *B. subtilis* and the impact of LuxS on the regulation of quorum sensing in this model bacterium have not yet been reported. Precisely, the main contributions of this work are (i) that it showed for the first time that *luxS* of *B. subtilis* was an active gene responsible for the production of active AI-2 able to be recognized by the AI-2-dependent signaling system of *V. harveyi* (Fig. 1) and (ii) the numerous evidence that demonstrated the key role of *luxS* in the cell density-dependent multicellular behavior of *B. subtilis*.

Even though AI-2-dependent quorum sensing has been analyzed in many bacteria, very little is known about the regulation of *luxS* expression (5, 55). Here, we demonstrated that *luxS* transcription was under AI-2-dependent quorum-sensing regulation (Fig. 2B). The dramatic overexpression of *luxS* in an AI-2-deficient background was not due to the accumulation of the toxic metabolite SAH of the AMC (Fig. 2A), because, as it was shown, the growth of the *luxS*-negative mutant strain (completely deficient in AI-2-dependent activity; see Fig. 1C) was the same as the growth of the wild-type strain (Fig. 2B). While SAH and SAM accumulated to a higher level in *luxS*-deficient cells than the levels of the AMC metabolites in wild-type cells (Fig. 2C), they were far lower than the levels of SAH reported to be toxic in bacteria (55, 60). These results (Fig. 2B and C) indicated that the levels of accumulated metabolites from the AMC in the *luxS*-deficient strain were not toxic and suggested that the concentration of the intermediates of the AMC would be cell regulated (i.e., at transcriptional level of the genes involved in the AMC and/or stability of the AMC metabolites) in order to keep these compounds at low, non-toxic concentrations.

The lack of a detrimental effect on growth resulting from the absence of LuxS activity suggested a role for this enzyme in cell-cell signaling and not in AMC-related metabolism to decrease (jointly with the activity of Pfs) the concentration of SAH. We obtained further evidence for a quorum-sensing or growth-phase regulation of *luxS* from the experiments in which we measured the effects on *luxS* expression after the addition of cell-free supernatants (CFCM) derived from AI-2-positive and AI-2-negative cultures (Fig. 2D). The observed results strongly argued for an AI-2-dependent negative inhibitory feedback loop controlling *luxS* transcription (Fig. 2A, possibility 2). In order to test this hypothesis, we predicted that an overaccumulation or overproduction of AI-2 should decrease *luxS* expression. To prove this, we overproduced AI-2 by ex-

pressing the *B. subtilis luxS* gene in the multicopy plasmid pHT315. As shown in Fig. 2E, and as was predicted from the original hypothesis, *luxS* expression was severely decreased in the strain that harbored the *luxS*-multicopy plasmid (without any detrimental effect on growth) confirming that AI-2 regulates *luxS* expression in *B. subtilis* (Fig. 2A, possibility 2).

So far, the best-characterized and most complex quorum-sensing-dependent phenomenon operating in *B. subtilis* is the process of spore development (2, 21). Therefore, we were intrigued to know whether an interrelationship would exist between sporulation and AI-2-dependent quorum sensing. As shown in Fig. 3A, LuxS activity did not affect the expression of the master regulatory protein Spo0A, and as expected from this result, spore efficiency in the *luxS*-negative mutant strain was also not affected (data not shown). In addition, LuxS did not affect the expression (Fig. 3B and C) of *abrB* and *sinR*, two other key developmental genes previously shown to be required for complete biofilm development and pluricellular behaviors in *B. subtilis* (12, 22).

In contrast, we found that Spo0A and SinR (Fig. 3E and F), but not AbrB (Fig. 3D), have a notorious negative effect on *luxS* expression. These results indicated that the regulation of *luxS* expression in *B. subtilis* was complex and intricate, being under the control of an autoregulatory AI-2-dependent feedback loop (Fig. 2) and under the control of the master regulatory proteins Spo0A and SinR (Fig. 3). Therefore, we wondered to what extent LuxS would affect the pluricellular behavior of wild isolates of *B. subtilis*. We decided to study the role of LuxS on pluricellularity by using *B. subtilis* natto, a probiotic and health-promoting strain for human consumption in Asian countries, as a model (24). We observed that AI-2-dependent activity was required for the formation of mature, robust, and highly differentiated biofilms (Fig. 4). Moreover, a significant defect of the LuxS-dependent ability of *B. subtilis* natto cells to swarm on solid medium (Fig. 5) was also noted. These results on biofilm formation and swarming motility, plus the previously shown inhibitory feedback loop on *luxS* (Fig. 2B), strongly argue in favor of a role of AI-2 as a signaling molecule in *B. subtilis*.

An equally impressive result was obtained when we analyzed the impact of LuxS-dependent activity on the ability of *B. subtilis* to make highly structured and complex communities on solid surfaces (bacterial colonies or colony biofilms). Effectively, wild-type cells of *B. subtilis* natto were able to develop into very complex and sophisticated giant aerial colonies (Fig. 6) in which spore development was temporal and spatially confined to the top of the giant colony (Fig. 7A). Importantly, we demonstrated that AI-2 activity was essential for this developmental process. Cells deficient in AI-2 production needed more time for organization to try to make the aerial colony (Fig. 6). However, this fruiting body was weaker and more undifferentiated than the fruiting bodies made by wild-type cells at all times during development. Finally, after longer incubation, the fruiting body made by the AI-2-deficient cells fell down and collapsed similar to a Babel Tower (Fig. 6). Since AI-2 expression was restricted to a specific region of the developing colony (Fig. 7B), we propose that the LuxS-produced signal (AI-2) behaves as a bacterial morphogen (25, 52, 64) that would be able to migrate (Fig. 7B) along the incipient, but well-organized, bacterial community in order to regulate an

appropriate morphogenesis and synchronization of cell activities during development (Fig. 6 and 7) (52, 64). Importantly, the master regulatory proteins Spo0A and SinR restricted the production of AI-2 to the bottom of the fruiting body (Fig. 7B). In fact, we showed that *luxS* expression was repressed by SinR (Fig. 3E) and Spo0A (Fig. 3F), and we also showed that Spo0A (sporulation) was expressed only at the top of the developing colony (Fig. 7A). Therefore, it was plausible to envision that Spo0A (and SinR) would restrict the production of the morphogen to the bottom of the developmentally committed colony from where AI-2 migrates to orchestrate cell-to-cell communication. Effectively, this prediction was confirmed by the observation of a less spatial restriction of LuxS activity in colonies deficient in *spo0A* and *sinR* expression (Fig. 7C).

Up to today, three distinct DPD-derived signals have been identified: *S*-THMF-borate (in *V. harveyi*) and *R*-THMF/*R*-THMF-P (in *S. enterica* serovar Typhimurium) (55, 57). *S*-THMF-borate and *R*-THMF chemical species may exist in equilibrium and rapidly interconvert (55, 57). Other DPD derivatives might also exist and be biologically active (see Fig. 2A and reference 57). Since *Vibrio*, *Salmonella*, and *Bacillus* organisms are inhabitants of completely different niches (sea, gut, and soil, respectively) exciting tasks for the future include elucidation of the chemical structure of the AI-2 molecule made by the soil bacterium *B. subtilis* and understanding of the AI-2-dependent signaling pathway of cell-to-cell communication that operates in this model bacterium.

ACKNOWLEDGMENTS

We are thankful for the insights of Roberto Kolter (whom we were delighted to meet in Iguazú Falls, Argentina, in 2004) and to Klaus Winzer for helpful and encouraging advice and Bonnie Bassler for the *V. harveyi* BB170 strain.

This work was supported by national grants to R.G. from FONCyT, CONICET, and Fundación Antorchas. E.L., A.R., and A.A. are doctoral fellows from CONICET, and R.G. is a career researcher from CONICET and a former Pew Latin American Scholar (Pew, Philadelphia, Pa.) and Fulbright Scholar (Washington, D.C.).

REFERENCES

- Antelmann, H., S. Engelmann, R. Schmid, A. Sorokin, A. Lapidus, and M. Hecker. 1997. Expression of stress- and starvation-induced *dps/pexB* homologous gene is controlled by the alternative sigma factor σ^B in *Bacillus subtilis*. *J. Bacteriol.* **179**:7251–7256.
- Arabolaza, A., A. Nakamura, M. E. Pedrido, L. Martelotto, L. Orsaria, and R. Grau. 2003. Characterization of a novel inhibitory feedback of the anti-sigma factor SpoIIAA on activation of Spo0A transcription factor during development in *Bacillus subtilis*. *Mol. Microbiol.* **47**:1251–1263.
- Auger, S., E. Krin, S. Aymerich, and M. Gohar. 2006. Autoinducer 2 affects biofilm formation by *Bacillus cereus*. *Appl. Environ. Microbiol.* **72**:937–941.
- Bacon Schneider, K., T. M. Palmer, and A. D. Grossman. 2002. Characterization of *comQ* and *comX*, two genes required for production of ComX pheromone in *Bacillus subtilis*. *J. Bacteriol.* **184**:410–419.
- Barák, I., E. Ricca, and S. Cutting. 2005. From fundamental studies of sporulation to applied spore research. *Mol. Microbiol.* **55**:330–338.
- Bassler, B. L., M. Wright, R. E. Showalter, and M. R. Silverman. 1993. Intercellular signalling in *Vibrio harveyi*: sequence and function of genes regulating expression of luminescence. *Mol. Microbiol.* **9**:773–786.
- Bassler, B. L. 2002. Small talk: cell-cell communication in bacteria. *Cell* **109**:421–424.
- Beeston, A. L., and M. G. Surette. 2002. *pfs*-dependent regulation of autoinducer 2 production in *Salmonella enterica* serovar typhimurium. *J. Bacteriol.* **184**:3450–3456.
- Branda, S., J. E. Gonzalez-Pastor, E. Dervyn, D. Ehrlich, R. Losick, and R. Kolter. 2004. Genes involved in formation of structured multicellular communities by *Bacillus subtilis*. *J. Bacteriol.* **186**:3970–3978.
- Branda, S., J. E. Gonzalez-Pastor, S. Ben-Yehuda, R. Losick, and R. Kolter. 2001. Fruiting body formation by *Bacillus subtilis*. *Proc. Natl. Acad. Sci. USA* **98**:11621–11626.
- Chen, X., S. Schauder, N. Potier, A. van Dorselaer, I. Pelczar, et al. 2002. Structural identification of a bacterial quorum-sensing signal containing boron. *Nature* **415**:545–549.
- Chu, F., D. Kearns, S. Branda, R. Kolter, and R. Losick. 2006. Targets of the master regulator of biofilm formation in *Bacillus subtilis*. *Mol. Microbiol.* **59**:1216–1228.
- Daniels, R., J. Vanderleyden, and J. Michiels. 2004. Quorum sensing and swarming migration in bacteria. *FEMS Microbiol. Rev.* **28**:261–289.
- Day, W. A., Jr., and A. T. Maurelli. 2001. *Shigella flexneri* LuxS quorum-sensing system modulates *virB* expression but is not essential for virulence. *Infect. Immun.* **69**:15–23.
- Federle, M. J., and B. L. Bassler. 2003. Interspecies communication in bacteria. *J. Clin. Investig.* **112**:1291–1299.
- Forsyth, M., and T. Cover. 2000. Intercellular communication in *Helicobacter pylori*: *luxS* is essential for the production of an extracellular signaling molecule. *Infect. Immun.* **68**:3193–3199.
- Fuqua, C., and P. E. Greenberg. 2002. Listening in on bacteria: acyl-homoserine lactone signalling. *Nat. Rev. Mol. Cell Biol.* **3**:685–695.
- Gottig, N., M. E. Pedrido, M. Méndez, E. Lombardía, A. Rovetto, V. Philippe, L. Orsaria, and R. Grau. 2005. The *Bacillus subtilis* SinR and RapA developmental regulators are responsible for the inhibition of spore development by alcohol. *J. Bacteriol.* **187**:2662–2672.
- Greenberg, E. P., J. W. Hastings, and S. Ulitzur. 1979. Induction of luciferase synthesis in *Bienecke harveyi* by other marine bacteria. *Arch. Microbiol.* **120**:87–91.
- Greenberg, E. P. 2003. Bacterial communication and group behaviour. *J. Clin. Investig.* **112**:1288–1290.
- Grossman, A. D. 1995. Genetic networks controlling the initiation of sporulation and the development of genetic competence in *Bacillus subtilis*. *Annu. Rev. Genet.* **29**:477–508.
- Hamon, M. A., and B. A. Lazazzera. 2001. The sporulation transcription factor Spo0A is required for biofilm development in *Bacillus subtilis*. *Mol. Microbiol.* **42**:1199–1209.
- Hilgers, M. T., and M. L. Ludwig. 2001. Crystal structure of the quorum-sensing protein LuxS reveals a catalytic metal site. *Proc. Natl. Acad. Sci. USA* **98**:11169–11174.
- Hosoi, T., and K. Kiuchi. 2004. Production and probiotic effects of natto, chapter 12. In E. Ricca, A. O. Henriques, and S. Cutting (ed.), *Bacterial spore formers: probiotics and emerging applications*. Horizon Bioscience, Norwich, United Kingdom.
- Jelback, L., and L. Søgaard-Andersen. 2002. Pattern formation by a cell surface-associated morphogen in *Myxococcus xanthus*. *Proc. Natl. Acad. Sci. USA* **99**:2032–2037.
- Jian, M., R. Grau, and M. Perego. 2000. Differential processing of propeptide inhibitors of Rap phosphatases in *Bacillus subtilis*. *J. Bacteriol.* **182**:303–310.
- Jones, M. B., and M. J. Blaser. 2003. Detection of *luxS*-signaling molecule in *Bacillus anthracis*. *Infect. Immun.* **71**:3914–3919.
- Joyce, E. A., et al. 2004. LuxS is required for persistent pneumococcal carriage and expression of virulence and biosynthesis genes. *Infect. Immun.* **72**:2964–2975.
- Kearns, D., F. Chu, R. Rudner, and R. Losick. 2004. Genes governing swarming in *Bacillus subtilis* and evidence for a phase variation mechanism controlling surface motility. *Mol. Microbiol.* **52**:357–369.
- Kearns, D., and R. Losick. 2003. Swarming motility in undomesticated *Bacillus subtilis*. *Mol. Microbiol.* **49**:581–590.
- Kim, S., et al. 2003. Regulation of *Vibrio vulnificus* virulence by the LuxS quorum-sensing system. *Mol. Microbiol.* **48**:1647–1664.
- Lazazzera, B. A., J. A. Solomon, and A. D. Grossman. 1997. An exported peptide functions intracellularly to contribute to cell density signaling in *Bacillus subtilis*. *Cell* **89**:917–925.
- Manefeld, M., and S. L. Turner. 2002. Quorum sensing in context: out of molecular biology and into microbial ecology. *Microbiology* **148**:3762–3764.
- Marouni, M. J., and S. Sela. 2003. The *luxS* gene of *Streptococcus pyogenes* regulates expression of genes that affect internalization by epithelial cells. *Infect. Immun.* **71**:5633–5639.
- McNab, R., and R. J. Lamont. 2003. Microbial dinner-party conversations: the role of LuxS in interspecies communication. *J. Med. Microbiol.* **52**:541–545.
- Méndez, M., L. Orsaria, V. Philippe, M. E. Pedrido, and R. Grau. 2004. Novel roles of the master transcription factors Spo0A and σ^B for survival and sporulation of *Bacillus subtilis* at low growth temperature. *J. Bacteriol.* **186**:989–1000.
- Merritt, J., F. Qi, S. Goodman, M. Anderson, and W. Shi. 2003. Mutation in *luxS* affects biofilm formation in *Streptococcus mutants*. *Infect. Immun.* **71**:1972–1979.
- Merritt, J., J. Kreth, W. Shi, and F. Qi. 2005. LuxS control bacteriocin production in *Streptococcus mutants* through a novel regulatory component. *Mol. Microbiol.* **57**:960–969.
- Miller, S. T., K. B. Xavier, S. R. Campagna, M. E. Taga, M. F. Semmelhack, et al. 2004. A novel form of the bacterial quorum sensing signal AI-2 recognized by *Salmonella typhimurium* receptor LsrB. *Mol. Cell* **15**:677–687.

40. Okamoto, S., A. Lezhava, T. Hosaka, Y. Okamoto-Hosoya, and K. Ochi. 2003. Enhanced expression of S-adenosylmethionine synthetase causes overproduction of actinorhodin in *Streptomyces coelicolor* A3(2). *J. Bacteriol.* **185**:601–609.
41. Pappas, K., C. Weingart, and S. Winans. 2004. Chemical communication in proteobacteria: biochemical and structural studies of signal synthases and receptors required for intercellular signaling. *Mol. Microbiol.* **53**:756–769.
42. Parsek, M. R., and E. P. Greenberg. 2005. Sociomicrobiology: the connections between quorum sensing and biofilms. *Trends Microbiol.* **13**:27–33.
43. Pei, D., and J. Zhu. 2004. Mechanism of action of S-ribosylhomocysteinase (LuxS). *Curr. Opin. Chem. Biol.* **8**:492–497.
44. Perego, M. 1997. A peptide export-import control circuit modulating bacterial development regulates protein phosphatases of the phosphorelay. *Proc. Natl. Acad. Sci. USA* **94**:8612–8617.
45. Ruzheinikov, S. N., S. K. Das, S. E. Sedelnikova, et al. 2001. The 1.2 Å structure of a novel quorum-sensing protein, *Bacillus subtilis* LuxS. *J. Mol. Biol.* **313**:111–122.
46. Schauder, S., and B. L. Bassler. 2001. The language of bacteria. *Genes Dev.* **15**:1468–1480.
47. Schauder, S., K. Shokat, M. Surette, and B. L. Bassler. 2001. The LuxS family of bacterial autoinducers: biosynthesis of a novel quorum-sensing signal molecule. *Mol. Microbiol.* **41**:463–476.
48. Senesi, S. 2004. *Bacillus* spores as probiotic products for human use, chapter 11. In E. Ricca, A. O. Henriques, and S. Cutting (ed.), *Bacterial spore formers: probiotics and emerging applications*. Horizon Bioscience, Norwich, United Kingdom.
49. Shapiro, J. A. 1998. Thinking about bacterial populations as multicellular organisms. *Annu. Rev. Microbiol.* **52**:81–104.
50. Sperandio, V., A. Torres, B. Jarvis, J. Nataro, and J. Kapers. 2003. Bacterial-host communication: the language of hormones. *Proc. Natl. Acad. Sci. USA* **100**:8951–8956.
51. Sun, J., R. Daniel, I. Wagner-Dobler, and A.-P. Zeng. 2004. Is autoinducer-2 a universal signal for interspecies communication: a comparative genomic and phylogenetic analysis of the synthesis and signal transduction pathways. *BMC Evol. Biol.* **4**:36.
52. Tabata, T., and Y. Takei. 2004. Morphogens, their identification and regulation. *Development* **131**:703–712.
53. Taga, M. E., and B. L. Bassler. 2003. Chemical communication in bacteria. *Proc. Natl. Acad. Sci. USA* **100**:14549–14554.
54. Taga, M. E., S. T. Miller, and B. L. Bassler. 2003. Lsr-mediated transport and processing of AI-2 in *Salmonella typhimurium*. *Mol. Microbiol.* **50**:1411–1427.
55. Vendeville, A., K. Winzer, K. Heurlier, C. M. Tang, and K. R. Hardie. 2005. Making “sense” of metabolism: autoinducer-2, LuxS and pathogenic bacteria. *Nat. Rev. Microbiol.* **3**:383–396.
56. Wang, L., R. Grau, M. Perego, and J. A. Hoch. 1997. A novel histidine kinase inhibitor regulating development in *Bacillus subtilis*. *Genes Dev.* **11**:2569–2579.
57. Waters, C. M., and B. L. Bassler. 2005. Quorum sensing: cell to cell communication in bacteria. *Annu. Rev. Cell Dev. Biol.* **21**:319–346.
58. Wen, Z. T., and R. A. Burne. 2004. LuxS-mediated signaling in *Streptococcus mutans* is involved in regulation of acid and oxidative stress tolerance and biofilm formation. *J. Bacteriol.* **186**:2682–2691.
59. Winans, S. C. 2002. Bacterial Esperanto. *Nat. Struct. Biol.* **9**:83–84.
60. Winzer, K., K. R. Hardie, N. Burgess, N. Doherty, D. Kirke, M. T. Holden, R. Linforth, et al. 2002. LuxS: its role in central metabolism and the in vivo synthesis of 4-hydroxy-5-methyl-3(2H)-furanone. *Microbiology* **148**:909–922.
61. Winzer, K., K. R. Hardy, and P. Williams. 2002. Bacterial cell-to-cell communication: sorry, can't talk now—gone to lunch! *Curr. Opin. Microbiol.* **5**:216–222.
62. Xavier, K. B., and B. L. Bassler. 2003. LuxS quorum sensing: more than just a numbers game. *Curr. Opin. Microbiol.* **6**:191–217.
63. Xavier, K. B., and B. L. Bassler. 2005. Interference with AI-2-mediated bacterial cell-cell communication. *Nature* **437**:750–753.
64. Zhu, A. J., and M. P. Scott. 2004. Incredible journey: how do developmental signals travel through tissue? *Genes Dev.* **18**:2985–2997.