

The Most Widespread Symbiosis on Earth

Andreas Brachmann, Martin Parniske

The interaction between plant roots and Glomeromycota fungi [1]—arbuscular mycorrhiza (AM)—is probably the most important symbiotic association in nature. More than 80% of all higher land plant species are able to benefit from the fungus' ability to extract nutrients, mostly phosphorus, from the soil. In exchange, it has been estimated that worldwide up to 20% of all photosynthetically fixed carbon might be delivered to the fungal partner. This makes it the most widespread symbiosis on earth [2–4]. Yet, despite its ecological importance, astonishingly little is known about the molecular mechanisms involved in this contact. In this issue of *PLoS Biology*, Besserer et al. [5] study how plant roots are recognized by their fungal partner and demonstrate that these events generalize across species.

Fossil evidence shows AM symbiosis to be at least as old as the earliest land plants [6,7]. It has been suggested that colonization of the land surface was dependent on the fungal symbiont's ability to forage into the soil for inorganic nutrients and water. This interaction apparently proved to be so successful that even after development of functional plant root structures, it was still retained in most plant families. However, a surprising promiscuity is observed in this symbiosis since, at least under laboratory conditions, many plant–fungus combinations lead to symbiotic structures. This lack of specificity is rather puzzling, considering the very long period of coevolution and the complete lack of sexual reproduction within the fungal partner, and points to a common mechanism of recognition.

Two important discoveries demonstrated that the early interaction between plant root and fungus involves mutual release of diffusible signaling molecules. Plant roots release “branching factors” (BFs) that induce morphological and cytological responses in the approaching fungal hyphae [8]. Among these, hyphal branching is the most prominent morphological change that can be attributed to fungus–plant recognition. In turn, fungal hyphae produce “myc factors” that lead to the transcriptional induction of symbiosis-related genes in the host root [9]. These pre-symbiotic recognition events are pre-requisite for formation of fungal appressoria structures on the root surface, invasion and colonization of fungal hyphae inside the root cortex, and, finally, the formation of highly branched, tree-like fungal structures (arbuscules) inside plant cells. The surface increase associated with arbuscule formation is believed to aid nutrient exchange between the partners.

Recently, Akiyama et al. [10] provided a major breakthrough in our understanding of the very early recognition events in this process. They identified strigolactones as the BFs released by the AM plant *Lotus japonicus* that trigger hyphal branching in the AM fungus

Gigaspora margarita. Strigolactones belong to the sesquiterpene lactones, which are believed to have a wide distribution in the plant kingdom [11]. Curiously, the same class of molecules was described exactly 40 years ago as a germination stimulant for seeds of the parasitic weeds *Striga* (witchweed) and *Orobranche* that attack plant roots and deprive them of water and nutrients [12]. Strigolactones occur with different substitutions, yet very little specificity was observed when different structures of natural and synthetic origin were tested for their activity on *Gi. margarita*. The widespread occurrence of strigolactones and this low structural specificity may underlie the promiscuity of AM fungi.

Still, open questions remained. The Glomeromycota are considered the fifth fungal phylum [1] and their common ancestor dates back 600 million years [7], yet all of these fungi exist in symbiosis with phototrophic organisms. Members of this phylum are more diverged from each other than the evolutionary younger basidiomycetes or ascomycetes, which have conquered very different niches during their evolution. Is responsiveness to strigolactones a general phenomenon within this phylum despite its old age? Besserer et al. [5] could show responses to strigolactones and synthetic derivatives, but not other sesquiterpene lactones, in three representatives of phylogenetically diverged groups of AM fungi: *Gi. rosea*, *Glomus intraradices*, and *Gl. clavoideum*, thus providing important confirmation of a general recognition mechanism. This widespread strigolactone perception system suggests that already the earliest land plants used this class of molecules to communicate with their symbiotic partners.

Interestingly, the induced morphological changes seem to be dependent on the fungal species. Whereas in both *Gi. margarita* and *Gi. rosea* spore germination occurs by itself and strigolactones produce an increase in hyphal branching, the main strigolactone effect on *Gl. intraradices* and *Gl. clavoideum* is an elevated germination rate.

Moreover, Besserer et al. [5] were able to demonstrate that application of strigolactone to these fungi led to not only morphological changes but also a rapid increase in mitochondrial density and respiration, both hallmarks of pre-symbiotic reprogramming of the fungus [8,13]. Perception of the plant signal seems to activate the breakdown of storage lipids, which enables the fungus to forage and ramify in order to enhance the chance of an encounter with plant roots. The

Citation: Brachmann A, Parniske M (2006) The most widespread symbiosis on earth. *PLoS Biol* 4(7): e239. DOI: 10.1371/journal.pbio.0040239

DOI: 10.1371/journal.pbio.0040239

Copyright: © 2006 Brachmann and Parniske. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abbreviations: AM, arbuscular mycorrhiza; BF, branching factor

Andreas Brachmann and Martin Parniske are in Genetics, Biology Department, Ludwig-Maximilians-University of Munich, Germany. E-mail: a.brachmann@lrz.uni-muenchen.de; parniske@lmu.de

Primers provide a concise introduction into an important aspect of biology highlighted by a current *PLoS Biology* research article.

high instability of strigolactones fits well with this role as a short-range signaling factor.

The finding that plant secondary compounds can modulate fungal development invites speculations about possible additional roles for strigolactones during symbiosis. For example, it is completely unclear what signal triggers the profuse hyphal branching observed during arbuscule development. Strigolactones are obvious candidates to induce such developmental changes in the fungus. What is still missing is a quantitative assessment of the importance of strigolactones for symbiosis. An important tool would be plants unable to produce strigolactones. Such genetic strategies have been very successful in eliminating a major role for flavonoids during AM development [11].

The ability to activate the pre-symbiotic stage of the fungus in a synchronized fashion will certainly help to unravel associated molecular events. Production and secretion of fungal symbiotic signaling factor(s) appears to be dependent on perception of the plant host [9] and may be induced after application of strigolactones. Thus, molecular identification of the BF and demonstration of biological activity of synthetic derivatives opens up interesting possibilities for future research on the AM fungus. ■

Acknowledgments

Funding. The authors received no specific funding for this article.

Competing interests. The authors have declared that no competing interests exist.

References

1. Schüßler A, Schwarzott D, Walker C (2001) A new fungal phylum, the Glomeromycota: Phylogeny and evolution. *Mycol Res* 105: 1413–1421.
2. Gianinazzi-Pearson V (1996) Plant responses to arbuscular mycorrhizal fungi: Getting to the roots of the symbiosis. *Plant Cell* 8: 1871–1883.
3. Bundrett M (2002) Coevolution of roots and mycorrhizas of land plants. *New Phytol* 154: 275–304.
4. Harrison MJ (2005) Signaling in the arbuscular mycorrhizal symbiosis. *Ann Rev Microbiol* 59: 19–42.
5. Besserer A, Puech-Pages V, Kiefer P, Gomez-Roldan V, Jauneau A, et al. (2006) Strigolactones stimulate arbuscular mycorrhizal fungi by activating mitochondria. *PLoS Biol* 4: e226. DOI: 10.1371/journal.pbio.0040226
6. Remy W, Taylor TN, Hass H, Kerp H (1994) Four hundred-million-year-old vesicular arbuscular mycorrhizae. *Proc Natl Acad Sci U S A* 91: 11841–11843.
7. Redecker D, Kodner R, Graham LE (2000) Glomalean fungi from the Ordovician. *Science* 289: 1920–1921.
8. Buée M, Rosignol M, Jauneau A, Ranjeva R, Bécard G (2000) The pre-symbiotic growth of arbuscular mycorrhizal fungi is induced by a branching factor partially purified from plant root exudates. *Mol Plant Microbe Interact* 13: 693–698.
9. Kosuta S, Chabaud M, Lougnon G, Gough C, Dénarié J, et al. (2003) A diffusible factor from arbuscular mycorrhizal fungi induces symbiosis-specific MtENOD expression in roots of *Medicago trunculata*. *Plant Physiol* 131: 952–962.
10. Akiyama K, Matsuzaki K, Hayashi H (2005) Plant sesquiterpenes induce hyphal branching in arbuscular mycorrhizal fungi. *Nature* 435: 824–827.
11. Akiyama K, Hayashi H (2006) Strigolactones: Chemical signals for fungal symbionts and parasitic weeds in plant roots. *Ann Bot (Lond)*. 97: 925–931.
12. Cook CE, Whichard LP, Turner B, Wall ME, Egley GH (1966) Germination of witchweed (*Striga lutea* Lour.): Isolation and properties of a potent stimulant. *Science* 154: 1189–1190.
13. Tamasloukht MB, Séjalon-Delmas N, Kluever A, Jauneau A, Roux C, et al. (2003) Root factors induce mitochondrial-related gene expression and fungal respiration during the developmental switch from asymbiosis to presymbiosis in the arbuscular mycorrhizal fungus *Gigaspora rosea*. *Plant Physiol* 131: 1468–1478.