

Research

Open Access

Isolation and characterization of a new simian rotavirus, YK-I

Larry E Westerman*^{1,2}, Baoming Jiang^{1,2}, Harold M McClure²,
Lauren J Snipes-Magaldi¹, Dixie D Griffin¹, Gary Shin³, Jon R Gentsch¹ and
Roger I Glass^{1,2}

Address: ¹Viral Gastroenteritis Team, Respiratory and Enteric Viruses Branch, Centers for Disease Control and Prevention, Atlanta, Georgia, USA, ²Yerkes National Primate Research Center, Emory University, Atlanta, Georgia, USA and ³Department of Ecology and Evolutionary Biology, University of California at Los Angeles, Los Angeles, California, USA

Email: Larry E Westerman* - larry@cidrz.org; Baoming Jiang - bjiang@cdc.gov; Harold M McClure - kays@rmy.emory.edu; Lauren J Snipes-Magaldi - lmagaldi@comcast.net; Dixie D Griffin - dixiegriffin@hotmail.com; Gary Shin - gtshin@ucla.edu; Jon R Gentsch - jgentsch@cdc.gov; Roger I Glass - rglass@cdc.gov

* Corresponding author

Published: 31 May 2006

Received: 30 June 2005

Virology Journal 2006, 3:40 doi:10.1186/1743-422X-3-40

Accepted: 31 May 2006

This article is available from: <http://www.virologyj.com/content/3/1/40>

© 2006 Westerman et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: To effectively analyze the requirements for protection to rotavirus infection, a reliable animal model that reasonably mimics infection and disease in humans is needed. A requirement for an effective animal model is the availability of appropriate rotavirus stocks for challenge.

Results: A new simian rotavirus, designated YK-I, was isolated from a 2-year-old immunodeficient pigtailed macaque with chronic diarrhea. YK-I was distinguishable by electropherotype from the other simian rotavirus strains, SA11 and RRV. One variant of YK-I, clone 311, which was isolated after adaptation and plaque purification in cell cultures, displayed an unusual RNA electropherotype with an abnormally migrating gene II segment. Sequence analysis demonstrated a genetic rearrangement that involved a partial duplication of the gene II ORF encoding NSP5. YK-I was identified as a Group A rotavirus belonging to subgroup I. To further characterize the YK-I strain, the genes encoding VP4, VP7, and NSP4 were sequenced. Analysis of VP4 and VP7 gene fragments suggests that this strain is a G3P[3] rotavirus and is closely related to the simian rotavirus strain RRV. Serotype analysis also identified YK-I as a G3 rotavirus. The NSP4 genotype of YK-I is C, the same genotype as RRV.

Conclusion: This newly isolated rotavirus, YK-I, is being used to establish a nonhuman primate model for studying the infectivity, immunity, and pathogenesis of rotavirus and for evaluating candidate rotavirus vaccines.

Background

Rotaviruses have a wide host range and can be recovered from many animal species [1]. The ability to isolate and maintain rotaviruses and to use them in animal model systems has contributed to studies of the mechanisms of

pathogenesis and immunity and to the development of vaccines. Several rotavirus isolates and animal model systems have been successfully developed, including various murine rotavirus strains in infant and adult mice [2-4],

C11 and Ala strains in rabbits [5], and human rotaviruses with piglets [6].

Two simian rotavirus strains, SA11 and RRV, have been well characterized and are currently the most widely used reference strains in laboratories throughout the world [7-9]. The sequences of all 11 genomic segments of SA11 are available. In limited studies, rotavirus infection and disease have been induced in nonhuman primates inoculated with SA11 [10-13]. Also, some human rotavirus vaccines are based on the RRV strain or reassortants of RRV with human strains [14]. The use of simian strains in human vaccines was based on a Jennerian approach prompted by studies indicating that animal and human rotaviruses share a common group antigen and that experimental animals immunized with human strains of rotavirus had a significantly lower risk of disease and infectivity when subsequently challenged with animal rotaviruses [15].

Although the two well-characterized simian rotavirus strains are readily available for use as a challenge virus, they have not been used consistently in nonhuman primate models of rotavirus infection because of their numerous passages in cell culture, which is a common method for viral attenuation. We wanted to isolate and characterize a new simian virus that is with low passage number in cell culture to be used as a challenge virus in nonhuman primates. The isolate strain, designated YK-1 and its variant clone 311, is being used to establish a non-human primate model for studying the infectivity, immunity, and pathogenesis of rotavirus and for evaluating candidate rotavirus vaccines [16,17].

Results

Rotavirus isolation and characterization

High titers of rotavirus antigen measured by an immunoassay were consistently detected in the stools of an immunodeficient pigtailed macaque, PFM-1, which was infected naturally with rotavirus and developed severe, chronic diarrhea. Rotavirus-like virus particles were detected by electron microscopy in stool extracts from this macaque (data not shown). Polyacrylamide gel electrophoresis of the viral RNA segments extracted from a stool specimen from PFM-1 revealed an electrophoretic pattern consistent with other rotaviruses, except for a lower intensity of segment 11 and an additional segment migrating slightly slower than segments 7, 8, and 9 (data not shown). This RNA electropherotype suggested that the stool extract was a mixture of subpopulations of rotavirus, as reported for other human rotavirus isolates with genome rearrangements [18,19]. The rotaviruses from PFM-1's stool extract were adapted to grow in MA-104 and plaque purified, which revealed two distinct viruses, named YK-1 and clone 311, that were distinguishable by

plaque size and electropherotype (Figure 1a and 1b) [20,21]. YK-1 produced smaller plaques and had an electropherotype typical of group A rotaviruses and that was very similar to but distinct from RRV. Variant 311 produced larger plaques and had an RNA electropherotype identical to that for YK-1, except it had an additional segment that was migrating slightly slower than segments 7, 8, and 9 and did not have a typical migrating segment 11. Both YK-1 and 311 showed a cytopathic effect typical of rotavirus grown in MA-104 cells and readily grew to titers over 10^8 ffu per ml.

Variant 311 has a rearranged segment 11

The nucleotide sequence of segment 11 from YK-1 was first determined as a reference. It consisted of 667 nt with a 594-bp ORF flanked by 5' and 3' UTRs of 21 and 52 nt, respectively. Segment 11 of YK-1 had 99% similarity to segment 11 of RRV. In the variant 311, segment 11 migrated slower than that of YK-1 and RRV, as determined by Northern blot analysis with a probe specific for segment 11 (Figure 1b). Sequence analysis of segment 11 from variant 311 identified a rearrangement consisting of a partial duplication of segment 11 from YK-1. The rearrangement occurred at nt 626 in the 3' UTR with the duplication of the ORF starting at nt 201 and included the entire 3' UTR (Figure 1c). The sequence of variant 311's segment 11 has 100% identity to YK-1's segment 11 both in the ORF and in the partial duplication.

YK-1 group, subgroup, and serotype analyses

The VP6 protein of rotavirus confers group specificity that is divided into seven groups (A to G). The commercial immunoassay Rotaclone utilizes a monoclonal antibody directed against the group A VP6 antigen and identified YK-1 as a group A rotavirus [22]. Group A rotavirus strains have been separated into four subgroups, and YK-1 is designated subgroup 1 as determined by reactivity with a subgroup 1 MAb 255/60 but not with subgroup 2 MAb 631/9.

To predict the G and P serotype specificities of the YK-1 strain, the sequences of the genes encoding both VP7 and VP4 were determined and compared with those of representatives of established G and P serotypes. The deduced VP7 amino acid sequence of YK-1 was closely related to other simian G3 rotavirus strains: 89% amino acid identity with RRV and 88% amino acid identity with SA11. A phylogenetic tree was constructed that included known VP7 amino acid sequences of G3 and other common G serotypes (Figure 2). The YK-1 strain clustered with other strains of G3 serotype, including RRV, SA11, AU1, and YO. YK-1 was also identified as a G3 serotype by reactivity with MAbs YO-1E2 (G3) and G3-159 (G3) in an immunoassay utilizing MAbs reactive to VP7-specific protein. The predicted VP4 amino acid sequence of YK-1 closely




Figure 1
 Comparison between the YK-1 strain and its variant 311 by (a) plaques size, (b) Northern blot analysis for rotavirus RNA segment 11, and (c) schematic diagram of the sequences of gene segment 11. Arrows indicate segment 11 of RRV and YK-1, or rearranged segment 11 of variant 311.

resembled that of GRV, a newly identified P[3]G3 caprine strain, and that of RRV P[3]G3 (Figure 3) [23].

NSP4 sequence analysis

The NSP4 gene segment of YK-1 was sequenced and its deduced gene product was compared with those of other known rotavirus strains. The structure of the NSP4 gene of YK-1 was similar to those of other rotavirus strains that were sequenced previously. The gene consists of a 528-bp ORF that encodes a protein with a predicted size of 175 amino acids with two conserved potential N-linked glycosylation sites. The deduced amino acid sequence of YK-1's NSP4 was 98% similar to that of the simian RRV strain. At least four genetic NSP4 groups are known, and the NSP4 of YK-1 can be classified as Group C by comparison of the amino acid sequence (aa 131–148) of the variable portion in the VP4 binding domain of various groups of NSP4 (Figure 4) [24,25].

Discussion

A rotavirus infection model using nonhuman primates offers a highly relevant system to investigate the mechanisms of disease and immunity to rotavirus and to determine vaccine effectiveness [16,17,26]. Since nonhuman primates are the animals most closely related to humans, this model may be the best predictor of infection and immunity in humans. In order to perform such studies, it is necessary to have a rotavirus strain that will consistently infect nonhuman primates after oral challenge. We describe rotavirus isolates that were obtained from a naturally infected pigtailed macaque housed in a major primate research center. This monkey was immunosuppressed and had severe chronic diarrhea possibly due to the rotavirus infection. Two isolates were obtained from a stool of this monkey and one designated YK-1 had an electropherotype typical of most group A rotaviruses and the other, designated 311, was identical to YK-1 except for a rearrangement in gene segment 11 that encodes for the NSP5 protein.

The YK-1 and its variant clone 311 were fully adapted to grow in cell culture, and both strains could produce plaques on MA-104 cells, although the plaques from the 311 variant were larger than those from the YK-1 strain. The significance of the difference in plaques size is not known. Both YK-1 and 311 were identified as a group A rotavirus, subgroup 1, genotype P[3] and serotype G3. It is of interest that these characterizations of YK-1's G and P types are very similar to another simian rotavirus strain, RRV. The nucleotide sequence of the NSP4 gene from YK-1 was also determined because of the discovery of the NSP4 gene product as a viral enterotoxin and its implication in the virulence of rotavirus [27]. YK-1 was determined to have a group C NSP4 gene, which again was similar to that of the RRV strain.




Figure 2
Phylogenetic tree based on amino acid sequences of the VP7-encoding genes for YK-1 and other established rotavirus strains.

Group A rotaviruses with atypical RNA profiles due to genomic rearrangements have been repeatedly detected in stools of chronically infected immunodeficient children [18,28]. These types of rearrangements have also been detected in rotavirus isolates from apparently immunocompetent calves and rabbits [29-31]. We have also isolated an YK-1 variant, 311, with a rearrangement in gene segment 11. With these rotaviruses, the rearrangement results from a partial duplication of the gene with a normal 5' UTR followed by a normal ORF and a duplication starting at various positions after the stop codon and extending to the 3' end and leading to a long 3' UTR. Thus, the rearranged gene expresses a normal protein product. Although the function of this rearrangement is unknown, it has been proposed to play a part in the evolution of rotaviruses and to contribute to their diversity [32]. It has also been suggested that rearranged segments containing a partial duplication might be more efficient templates for

double stranded RNA synthesis than are their wild-type counterparts and thus may be preferentially selected during viral replication [31].

Conclusion

Development of a more suitable animal model of rotavirus infection requires the identification of an appropriate challenge strain. The ideal challenge virus should be isolated from the same species as that employed in the model system because, in some systems, heterologous rotaviruses tend to undergo abortive replication. We have isolated a new rotavirus strain, designated YK-1, from fecal specimens of a 2-year-old pigtailed macaque with severe chronic diarrhea. The YK-1 strain had been used to develop a nonhuman primate model to enhance our understanding of the mechanisms of immunity to rotavirus infection [16,17]. This report describes the characterization of this new strain and a variant of this strain and




Figure 3
Phylogenetic tree based on amino acid sequences of the VP4-encoding genes for YK-1 and other established rotavirus strains.

compares the properties of this strain to those of the other simian rotavirus strains, SA11 and RRV.

Materials and methods

Rotavirus isolation

The YK-1 strain of simian rotavirus was isolated from the diarrheal stool of a 2-year-old pigtailed macaque (*Macaca nemestrina*) housed at the Yerkes National Primate Research Center, Emory University (Atlanta, GA). This

Strain	Group	Amino acid sequence																	
YK-1	C	H	D	M	L	I	I	K	P	V	D	R	I	D	M	S	Q	E	F
RRV	C	-	-	-	-	-	-	-	-	-	-	K	-	-	-	-	-	-	-
SA11	A	Y	-	K	-	T	V	Q	T	T	G	E	-	-	-	T	K	-	I
Wa	B	H	D	N	-	-	T	R	-	-	-	V	-	-	-	-	K	-	-

Figure 4
Comparison of the NSP4 deduced amino acid sequences at the variable portion in the VP4- binding domain (aa 131–148) from representative groups of rotavirus strains.

immunodeficient macaque, PFM-1, had chronic diarrhea associated with high titers of fecal rotavirus antigen detected by Rotaclone immunoassay (Meridian Diagnostics, Cincinnati, OH). The virus was isolated by previously described methods with modifications [20,21]. An extract from an antigen positive stool was prepared as a 20% (wt/vol) suspension in phosphate buffered saline (PBS, pH 7.4) and centrifuged twice at 8500 g for 10 min for clarification. The supernatant was extracted with 1,1,2-trichlorotrifluoroethane (Sigma, St. Louis, MO), and centrifuged at 4000 × g for 5 minutes. The extract was treated with trypsin (15 µg/ml) for 45 min at 37°C and inoculated onto a confluent monolayer of MA104 cells (African green monkey kidney cells) for 1 h. After being washed, the monolayer was maintained in serum-free minimal essential medium (MEM) (Gibco, Grand Island, NY) supplemented with 2 µg/ml trypsin and 50 µg/ml neomycin for 3 days. A viral lysate, obtained by freeze-thawing three times and clarification at 8500 g for 30 min, was inoculated into MA104 cells and plaque purified three times. Two distinct plaques distinguished on size were obtained and further passed in MA104 cells.

Plaque assay

Virus stocks were activated with 15 µg/ml trypsin in MEM for 45 min at 37°C. The activated virus was 10-fold serially diluted in MEM and 500 µl/well was inoculated onto 6-well tissue culture plates (Corning, Corning, NY) with a confluent monolayer of MA104. After a 1 h incubation at 37°C, the inoculum was aspirated and 4 ml of a 3.5% agarose (Seakem, Biowhittaker, Rockland, ME) in MEM was overlaid on the monolayer. The agar was allowed to solidify at room temperature (RT), after which the plates were incubated at 37°C. Plaques were visualized by adding 1 ml of MEM with 2% neutral red and 0.3% agarose 6 h prior to reading.

Purification of virus RNA

Rotavirus RNA was extracted from stools and infected cell cultures by a modification of a previously described method [33]. In brief, a 10% stool extract or 30% cell-culture suspension was prepared with Tris-buffered saline supplemented with 1% sodium dodecyl sulfate (SDS), vortexed, and incubated at RT for 10 min. Equal volumes of virus and 1,1,2-trichlorotrifluoroethane were mixed for 1 minute, and centrifuged for at 8000 g for 10 min. The supernatant was added to 2 volumes of 6 M guanidine isothiocyanate and incubated at 56°C for 10 min. Silica beads were added to each sample, vortexed, and incubated at RT for 10 min. The beads were washed once with a 2:1 solution of 6 M guanidine isothiocyanate with 50 mM Tris-HCl (pH 7.5) and then three times with 70% ethanol. After the final wash, the beads were air dried, incubated with H₂O for 10 minutes at 65°C, and centrifuged

for 2 min at 10,000 g. The extract was saved and stored at -70°C until use.

Electropherotyping

Rotavirus double-stranded genomic RNA extracted from fecal samples and cell-culture lysates were analyzed by SDS-polyacrylamide gel electrophoresis as described previously [34].

Northern blot for segment 11

The procedures employed for Northern hybridization and chemiluminescent detection of bound digoxigenin-labeled probe using a commercial reagent (ECL, Amersham, Piccataway, NJ) have been described [35]. Two digoxigenin-labeled probes, ggctttaagcgtacagtgt and ggtcacaacgggagtgaggctcc, were used to identify genomic segment 11 of rotavirus.

Subgroup and serotype analyses

Subgroup and VP7 serotyping were determined by use of a panel of monoclonal antibodies: 225/60 (Subgroup I), 631/9 (Subgroup II), KU-4 (G1), 5E8 (G1), S2-SG10 (G2), IC10 (G2), YO-1E2 (G3), G3-159 (G3), and ST-2G7 (G4) [36-38]. In brief, Immulon II plates (Nalgel Nunc, Rochester, NY) were coated overnight with serum from a rabbit hyperimmunized with purified RRV rotavirus particles for positive wells and normal rabbit sera for negative wells. After the plate was washed with wash buffer (PBS plus 0.1% Tween 20), cell-culture lysates were added to duplicate positive and negative wells. The plates were incubated for 2 h at RT and washed. Specific monoclonal antibodies were added to wells, and the plates were incubated for 1 h at RT and then washed. Biotinylated goat anti-mouse IgG (Southern Biotechnology, Birmingham, AL) was added, incubated 30 min at RT and followed by washing and the addition of streptavidin-horseradish peroxidase (Southern Biotechnology). Wells were developed by adding tetramethylbenzidine (Sigma) and stopped after 10 min with 1 N HCl. A sample was considered positive if the OD value of the positive coated well was >2 times and 0.100 greater than the negative coated well.

PCR amplification and sequence analysis of VP4, VP7, NSP4, and NSP5

The PCR products of the genes coding for VP4, VP7, and NSP4 proteins were amplified with previously described primers, Con2/Con3 for VP4, Beg9/End9 for VP7 and 10Beg16/10End722 for NSP4 [33,39,40]. Full-length PCR product for the YK-1 gene coding the NSP5 protein was amplified with primers derived from the 5' and 3' ends of the NSP5 nucleotide sequence of the SA11 strain. The nucleotide sequence of each gene was determined from gel-purified PCR products as previously described [41]. Phylogenetic relatedness of the VP4 and VP7 genes of YK-1 was examined by comparing amino acid sequences

between reference rotavirus strains by using the Wisconsin Genetics Computer Group computer program [42].

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

LEW characterized and maintained the YK-1 virus and drafted the manuscript, BJ helped draft the manuscript, HMM provided samples for virus isolation, LJSJ isolated YK-1, GS and DDG sequenced YK-1, JRG provided phylogenetic analysis of YK-1, and RIG helped draft and critically review the manuscript.

Acknowledgements

The authors thank the staff at Yerkes National Primate Research Center for their assistance with the monkeys, Harry Greenberg, Shozo Urasawa and Koki Taniguchi for providing monoclonal antibodies, and we thank Claudia Chesley for editorial assistance. Supported by CRADA with Aventis Pasteur, Lyon, France and in part by Yerkes Base Grant #RR00165.

References

1. Saif LJ, Rosen BI, Parwani AV: **Animal Rotaviruses**. In *Viral Infections of the Gastrointestinal Tract* Edited by: Kapikian AZ. New York, NY: Marcel Dekker; 1994:279-367.
2. Burns JW, Krishnaney AA, Vo PO, Rouse RV, Anderson LJ, Greenberg HB: **Analyses of homologous rotavirus infection in the mouse model**. *Virology* 1995, **207**:143-153.
3. McNeal MM, Belli J, Basu M, Choi AH, Ward RL: **Discovery of a new strain of murine rotavirus that is consistently shed in large quantities after oral inoculation of adult mice**. *Virology* 2004, **320**:1-11.
4. Ward RL, McNeal MM, Sheridan JF: **Development of an adult mouse model for studies on protection against rotavirus**. *J Virol* 1990, **64**:5070-5075.
5. Conner ME, Estes MK, Graham DY: **Rabbit model of rotavirus infection**. *J Virol* 1988, **62**:1625-1633.
6. Saif LJ, Ward LA, Yuan L, Rosen BI, To TL: **The gnotobiotic piglet as a model for studies of disease pathogenesis and immunity to human rotaviruses**. *Arch Virol Suppl* 1996, **12**:153-161.
7. Estes MK: **Rotaviruses and their replication**. In *Fields Virology* 4th edition. Edited by: Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA, Roizman B, Straus SE. Philadelphia, PA: Lippincott, Williams and Wilkins; 2001:1747-1785.
8. Malherbe HH, Strickland-Cholmley M: **Simian virus SA11 and the related O agent**. *Arch Gesamte Virusforsch* 1967, **22**:235-245.
9. Lopez S, Arias CF: **Simian rotavirus SA11 strains**. *J Virol* 1992, **66**:1832.
10. Kalter SS, Heberling RL, Rodriguez AR, Lester TL: **Infection of baboons ("Papio cynocephalus") with rotavirus (SA11)**. *Dev Biol Stand* 1983, **53**:257-261.
11. Petschow BV, Litov RE, Young LJ, McGraw TP: **Response of colostrum-deprived cynomolgus monkeys to intragastric challenge exposure with simian rotavirus strain SA11**. *Am J Vet Res* 1992, **53**:674-678.
12. Leong YK, Awang A: **Experimental group A rotaviral infection in cynomolgus monkeys raised on formula diet**. *Microbiol Immunol* 1990, **34**:153-162.
13. Soike KF, Gary GW, Gibson S: **Susceptibility of nonhuman primate species to infection by simian rotavirus SA-11**. *Am J Vet Res* 1980, **41**:1098-1103.
14. Parashar UD, Bresee JS, Gentsch JR, Glass RI: **Rotavirus**. *Emerg Infect Dis* 1998, **4**:561-570.
15. Wyatt RG, Mebus CA, Yolken RH, Kalica AR, James HD Jr, Kapikian AZ, Chanock RM: **Rotaviral immunity in gnotobiotic calves: heterologous resistance to human virus induced by bovine virus**. *Science* 1979, **203**:548-550.
16. Westerman LE, McClure HM, Jiang B, Almond JW, Glass RI: **Serum IgG mediates mucosal immunity against rotavirus infection**. *PNAS* 2005, **102**:7268-7273.
17. Westerman LE, Xu J, Jiang B, McClure HM, Glass RI: **Experimental Infection of Pigtailed Macaques with a Simian Rotavirus, YK-1**. *J Med Virol* 2005, **75**:316-325.
18. Hundley F, McIntyre M, Clark B, Beards G, Wood D, Chrystie I, Desselberger U: **Heterogeneity of genome rearrangements in rotaviruses isolated from a chronically infected immunodeficient child**. *J Virol* 1987, **61**:3365-3372.
19. Gault E, Schnepf N, Poncet D, Servant A, Teran S, Garbarg-Chenon A: **A human rotavirus with rearranged genes 7 and 11 encodes a modified NSP3 protein and suggests an additional mechanism for gene rearrangement**. *J Virol* 2001, **75**:7305-14.
20. Sato K, Inaba Y, Shinozaki T, Fujii R, Matumoto M: **Isolation of human rotavirus in cell cultures: brief report**. *Arch Virol* 1981, **69**:155-160.
21. Ward RL, Knowlton DR, Pierce MJ: **Efficiency of human rotavirus propagation in cell culture**. *J Clin Microbiol* 1984, **19**:748-753.
22. Cukor G, Perron DM, Hudson R, Blacklow NR: **Detection of rotavirus in human stools by using monoclonal antibody**. *J Clin Microbiol* 1984, **19**:888-892.
23. Lee JB, Youn SJ, Nakagomi T, Park SY, Kim TJ, Song CS, Jang HK, Kim BS, Nakagomi O: **Isolation, serologic and molecular characterization of the first G3 caprine rotavirus**. *Arch Virol* 2003, **148**:643-657.
24. Kapikian AZ, Hoshino Y, Chanock RM: **Rotaviruses**. In *Fields Virology* 4th edition. Edited by: Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA, Roizman B, Straus SE. Philadelphia, PA: Lippincott, Williams and Wilkins; 2001:1787-1833.
25. Kirkwood CD, Palombo EA: **Genetic characterization of the rotavirus nonstructural protein, NSP4**. *Virology* 1997, **236**:258-265.
26. McNeal MM, Sestak K, Choi AH, Basu M, Cole MJ, Ave PP, Bohm RP, Ward RL: **Development of a rotavirus-shedding model in rhesus macaques, using a homologous wild-type rotavirus of a new P genotype**. *J Virol* 2005, **79**:944-954.
27. Ball JM, Tain P, Zeng CQ, Morris AP, Estes MK: **Age-dependent diarrhea induced by a rotaviral nonstructural glycoprotein**. *Science* 1996, **272**:101-104.
28. Eiden J, Losonsky GA, Johnson J, Yolken RH: **Rotavirus RNA variation during chronic infection of immunocompromised children**. *Pediatr Infect Dis* 1985, **4**:632-637.
29. Pocock DH: **Isolation and characterization of two group A rotaviruses with unusual genome profiles**. *J Gen Virol* 1987, **68**:653-660.
30. Thouless ME, DiGiacomo RF, Neuman DS: **Isolation of two lapine rotaviruses: characterization of their subgroup, serotype and RNA electrophoretotypes**. *Arch Virol* 1986, **89**:161-170.
31. Pedley S, Hundley F, Chrystie I, McCrae MA, Desselberger U: **The genomes of rotaviruses isolated from chronically infected immunodeficient children**. *J Gen Virol* 1984, **65**:1141-1150.
32. Desselberger U: **Genome rearrangements of rotaviruses**. *Arch Virol Suppl* 1996, **12**:37-51.
33. Gentsch JR, Glass RI, Woods P, Gouvea V, Gorziglia M, Flores J, Das BK, Bhan MK: **Identification of group A rotavirus gene 4 types by polymerase chain reaction**. *J Clin Microbiol* 1992, **30**:1365-1373.
34. Pereira HG, Azeredo RS, Leite JP, Candeias JA, Ruez ML, Linhares AC, Gabbay YB, Trabulsi JR: **Electrophoretic study of the genome of human rotaviruses from Rio de Janeiro, Sao Paulo and Belem, Brazil**. *J of Hygiene* 1983, **90**:117-125.
35. Palombo EA, Bishop RF: **Genetic and antigenic characterization of a serotype G6 human rotavirus isolated in Melbourne, Australia**. *J Med Virol* 1995, **47**:348-354.
36. Woods PA, Gentsch J, Gouvea V, Mata L, Santosham M, Bai ZS, Urasawa S, Glass RI: **Distribution of serotypes of human rotavirus in different populations**. *J Clin Microbiol* 1992, **30**:781-785.
37. Taniguchi K, Urasawa T, Morita Y, Greenberg HB, Urasawa S: **Direct serotyping of human rotavirus in stools by an enzyme-linked immunosorbent assay using serotype 1-, 2-, 3-, and 4-specific monoclonal antibodies to VP7**. *J Infect Dis* 1987, **155**:1159-1166.
38. Coulson BS, Fowler KJ, White JR, Cotton RG: **Non-neutralizing monoclonal antibodies to a trypsin-sensitive site on the major glycoprotein of rotavirus which discriminate between virus serotypes**. *Arch Virol* 1987, **93**:199-211.

39. Cunliffe NA, Woods PA, Leite JP, Das BK, Ramachandran M, Bhan MK, Hart CA, Glass RI, Gentsch JR: **Sequence analysis of NSP4 gene of human rotavirus allows classification into two main genetic groups.** *J Med Virol* 1997, **53**:41-50.
40. Gouvea V, Glass RI, Woods P, Taniguchi K, Clark HF, Forrester B, Fang ZY: **Polymerase chain reaction amplification and typing of rotavirus nucleic acid from stool specimens.** *J Clin Microbiol* 1990, **28**:276-282.
41. Laird AR, Gentsch JR, Nakagomi T, Nakagomi O, Glass RI: **Characterization of serotype G9 rotavirus strains isolated in the United States and India from 1993 to 2001.** *J Clin Microbiol* 2003, **41**:3100-3111.
42. Devereux J, Haeberli P, Smithies O: **A comprehensive set of sequence analysis programs for the VAX.** *Nucleic Acids Res* 1984, **12**:387-395.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

