

Research article

Open Access

Cytokine polymorphisms influence treatment outcomes in SLE patients treated with antimalarial drugsPatricia López¹, Jesús Gómez², Lourdes Mozo², Carmen Gutiérrez^{1,2} and Ana Suárez¹¹Department of Functional Biology, Area of Immunology, University of Oviedo. Spain²Department of Immunology, Hospital Universitario Central de Asturias, Oviedo. SpainCorresponding author: Ana Suárez, anasua@uniovi.es

Received: 2 Nov 2005 Revisions requested: 6 Dec 2005 Revisions received: 20 Dec 2005 Accepted: 11 Jan 2006 Published: 13 Feb 2006

Arthritis Research & Therapy 2006, **8**:R42 (doi:10.1186/ar1897)This article is online at: <http://arthritis-research.com/content/8/2/R42>© 2006 López *et al.*; licensee BioMed Central Ltd.This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**Abstract**

Antimalarial agents have been widely used as disease-modifying antirheumatic drugs in the treatment of systemic lupus erythematosus (SLE) and other rheumatological diseases, although their mechanism of action has not yet been fully defined. It is known, however, that effective response to treatment is variable among patients. Thus, the identification of genetic predictors of treatment response would provide valuable information for therapeutic intervention. The aim of the present study was to analyze the effect of antimalarial treatment on tumor necrosis factor (TNF) α serum levels and evaluate the possible influence of TNF α and IL-10 functional genetic polymorphisms on the response to antimalarial drugs. To this end, TNF α serum levels were quantified in 171 SLE patients and 215 healthy controls by ELISA techniques and polymorphisms at positions -1,082 and -308 of the IL-10 and TNF α gene promoters were determined by PCR amplification followed by hybridization with fluorescent-labeled allele-specific probes in 192 SLE patients and 343 matched controls. Data were related to clinical features and treatment at the time of sampling and during the course of the disease. Results showed a significantly higher amount of serum TNF α in the entire SLE population

compared with controls. However, TNF α serum levels correlated negatively with the use of antimalarial treatment during at least three months before sampling. Patients under single or combined treatment with these drugs had TNF α serum levels similar to healthy controls, whereas untreated patients and those under corticosteroid or immunosuppressive therapies had increased amounts of this cytokine. This suggests, however, that antimalarial-mediated inhibition of TNF α was only significant in patients who were genetically high TNF α or low IL-10 producers. In addition, evaluation of SLE patients administered antimalarial drugs for three or more years who did not require any other specific SLE treatment indicates that patients with the combined genotype low IL-10/high TNF α are the best responders to antimalarial therapy, developing mild disease with a good course under this treatment. In conclusion, we proposed that an antimalarial-mediated downregulation of TNF α levels in SLE patients is influenced by polymorphisms at IL-10 and TNF α promoters. Our results may thus find important clinical application through the identification of patients who are the most likely to benefit from antimalarial therapy.

Introduction

Systemic lupus erythematosus (SLE) is a disorder of immune regulation resulting in chronic inflammation that affects many organs. Treatment of lupus disease must be determined individually, since different patients may have diverse and multiple symptoms with variable severity. Mild disease requires no or little therapy, usually nonsteroidal anti-inflammatory medica-

tions (NSAIDs). Articular and skin symptoms are frequently treated with antimalarial drugs, especially hydroxychloroquine, alone or with low dose corticosteroids when required, whereas severe lupus must be treated with corticosteroids and/or immunosuppressive drugs. However, it is assumed that responses to specific agents may be variable among SLE patients. Consequently, the identification of genetic predictors

CI = confidence interval; ELISA = enzyme-linked immunosorbent assay; IL = interleukin; LPS = lipopolysaccharide; NSAIDs = nonsteroidal anti-inflammatory drugs; OR = odds ratio; PCR = polymerase chain reaction; SLE = systemic lupus erythematosus; SNP = single nucleotide polymorphism; TNF = tumor necrosis factor.

Table 1**Systemic lupus erythematosus patient characteristics and disease parameters**

Total SLE patients	192
Sex (female/male)	178/14
Age (mean \pm sd)	47.24 \pm 14.63
Age at diagnosis (mean \pm sd)	33.88 \pm 14.53
Disease duration (mean \pm sd)	13.16 \pm 7.83
Clinical manifestations, n (%)	
Malar rash	102 (53.1)
Discoid lesions	38 (19.8)
Subacute cutaneous lesions	33 (17.2)
Photosensitivity	99 (51.6)
Oral ulcers	63 (32.8)
Arthritis	139 (72.4)
Serositis	48 (25.0)
Renal disorder	56 (29.2)
Neurological disorder	15 (7.8)
Hematological disorder	111 (57.8)
Treatment, n (%)	
None/NSAIDS	23 (12.0)
Antimalarials	114 (59.4)
Corticosteroids	126 (65.6)
Immunosuppressive drugs ^a	41 (21.4)

^aMethotrexate, azathioprine, cyclophosphamide, cyclosporine A or mycophenolate mophetil. NSAIDS, nonsteroidal anti-inflammatory drugs; SD, standard deviation; SLE, systemic lupus erythematosus.

of treatment response would provide valuable clinical information, since these can be determined at the time of diagnosis, when therapeutic intervention has the potential to offer the greatest benefits.

Antimalarial drugs (hydroxychloroquine, chloroquine and quinacrine) have been used as disease-modifying antirheumatic agents in the treatment of several autoimmune diseases, usually associated with increased secretion of pro-inflammatory cytokines. However, despite an extensive clinical history of use in rheumatoid arthritis and SLE [1], their mechanisms of anti-inflammatory action have not yet been fully defined. Chloroquine is thought to concentrate in acidic subcellular compartments, such as endolysosomes, where it inhibits acidic proteases [2]. As lysosomal enzymes are involved in antigen processing and presentation, an antirheumatic effect might be mediated by a downregulation of the immune response

against autoantigens. However, the effects of chloroquine and related drugs may extend beyond this. Interestingly, antimalarials have been shown to inhibit the release of the pro-inflammatory cytokines IL-1, IL-6 and tumor necrosis factor (TNF) α by monocytes activated with lipopolysaccharide (LPS) or CpG oligonucleotides [3-6] and it has been reported that chloroquine interferes with LPS-induced expression of the gene encoding TNF α in human blood monocytes by a nonlysosomal mechanism [7]. However, it is not yet known whether treatment of humans with antimalarial drugs is capable of reducing pro-inflammatory cytokines *in vivo*.

Considering the central role that IL-10 and TNF α cytokines play in the pathogenesis of SLE, it is possible that different cytokine production may not only affect the natural course of the disease, but also the response to therapy. Genetic polymorphisms at the promoter of the genes encoding IL-10 and TNF α have been associated with different constitutive and induced cytokine production. The genetic variant at position -308 (G/A) of the gene encoding TNF α was found to have functional effects on gene transcriptional activity, carriers of the uncommon TNF2 allele (-308A*) being considered as genetically high TNF α producers [8-10]. Similarly, IL-10 basal and induced production presented interindividual variations that were genetically regulated by three single nucleotide polymorphisms (SNPs) at positions -1,082(G/A), -819(C/T) and -592(C/A) of the IL-10 promoter. In Caucasian populations, only three haplotypes have been found (GCC, ACC and ATA), the individuals GCC/GCC being considered as genetically high IL-10 producers [10-12]. Several studies have analyzed the association of IL-10 or TNF α genetic variants with susceptibility to and outcome of SLE and other autoimmune diseases, showing variable results in most cases. However, it is known that the actions of cytokines may be profoundly conditioned by the presence of other cytokines, this being particularly true in the case of IL-10 and TNF α , which are mutually regulated and have complex and predominantly opposing roles in systemic inflammatory responses. In a previous study, we found that carriers of the combined genotype high TNF α /low IL-10 have the highest risk factor for developing SLE and producing antibodies to the SSa antigen [10], suggesting that interactions between IL-10 and TNF α cytokine genes may influence susceptibility to SLE, its phenotype and possibly the clinical response to disease modifying antirheumatic drugs. The aim of the present study was to detect the effect of SLE treatments on TNF α serum levels and to evaluate the possible influence of IL-10 and TNF α functional genetic polymorphisms on the response to antimalarial drugs.

Table 2**Tumor necrosis factor- α serum levels in healthy controls and systemic lupus erythematosus patients**

	<i>n</i>	TNF α (pg/ml) ^a	<i>p</i>
Healthy controls	215	19.66 (53.93)	0.020
SLE patients	171	33.57 (132.55)	
Patient treatment:			
None/NSAIDS	21	60.78 (158.15)	0.136
Antimalarials	36	16.64 (165.57)	
Antimalarials and corticosteroids	54	24.95 (88.49)	
Corticosteroids	26	60.01 (145.67)	
Immunosuppressive drugs ^b alone or with corticosteroids	22	105.34 (195.29)	
Antimalarials, corticosteroids and immunosuppressive drugs ^b	12	16.89 (50.70)	

^aTNF α values are median (interquartile range). ^bMethotrexate, azathioprine, cyclophosphamide, cyclosporine A or mycophenolate mophetil. Differences were evaluated by Mann-Whitney or Kruskal-Wallis nonparametric testing. NSAIDS, nonsteroidal anti-inflammatory drugs; SLE, systemic lupus erythematosus; TNF, tumor necrosis factor.

Materials and methods


Patients

Approval for this study was obtained from the Regional Ethics Committee for Clinical Investigation. Patients included in the study ($n = 192$) were from the Asturian Register of SLE [13]; all of them were Caucasian in origin and fulfilled the American College of Rheumatology (ACR) criteria for SLE [14]. At the time of serum sampling for TNF α quantification, patients were asked precise questions regarding the treatment received during the past three months. All untreated or NSAID treated patients presented inactive SLE. In addition, for genotype associations, information on clinical manifestations (age at diagnosis, disease duration, malar rash, discoid or subacute cutaneous lesions, photosensitivity, oral ulcers, arthritis, serositis, renal, neurological or hematological disorder) and treatments followed during the course of the disease was obtained after a detailed review of clinical histories. Those patients receiving antimalarial agents for three or more years without requiring any other specific treatment were defined as good responders to antimalarial therapy. The demographic and clinical characteristics of the patients are shown in Table 1. Matched healthy controls ($n = 343$) were obtained from the Asturian Blood Transfusion Center. Consent was obtained from all individuals prior to participation in the study.

TNF α quantification

Serum samples for TNF α quantification were collected from 171 SLE patients and 215 healthy controls. TNF α concentration was determined by an in-house ELISA test, as follows. Microtiter wells were coated overnight with affinity purified anti-human TNF α monoclonal antibody (R&D Systems, Abingdon, UK) and blocked with 1% casein in Tris Buffered Saline (TBS) for two hours at 37°C. Samples and TNF α standards (R&D) were diluted in blocking solution and incubated for 18

hours at 4°C. After washing with TBS/Tween 20 (0.05%), wells were incubated for two hours with biotinylated anti-human TNF α monoclonal antibody (R&D), washed, incubated for one hour with streptavidin-alkaline phosphatase conjugate and revealed using *p*-nitrophenyl phosphate as substrate. Absorbance was determined at a wavelength of 405 nm. Quantities of serum TNF α were calculated according to the standard curves. The assay has a detection limit of 7.5 pg/ml, a within-run imprecision (coefficient of variation) of <7%, and a between-run coefficient of variation of <10%.

Figure 1

Antimalarial treatment downregulates tumor necrosis factor (TNF) α serum levels in systemic lupus erythematosus (SLE) patients. TNF α serum levels were quantified by ELISA techniques in 171 SLE patients and 215 healthy controls. All patients were classified as users or non-users of antimalarial drugs during the last three months before sampling. Differences were evaluated by the Mann-Whitney *U* test.

Table 3**TNF α levels in high and low TNF α producing patient users and nonusers of antimalarial treatment**

TNF α genotype	Healthy controls		Patients without antimalarial drugs		Patients with antimalarial drugs		<i>p</i>
	<i>n</i>	TNF α (pg/ml) ^a	<i>n</i>	TNF α (pg/ml) ^a	<i>n</i>	TNF α (pg/ml) ^a	
High (AA/AG)	53	20.06 (125.63)	24	83.55 (171.64)	48	16.88 (54.89)	0.001
Low (GG)	162	18.83 (40.67)	45	58.56 (168.94)	54	26.00 (154.67)	0.516

^aValues are median (interquartile range). Differences between users and nonusers of antimalarial therapy were evaluated by the Mann-Whitney *U* test. TNF, tumor necrosis factor.

Promoter polymorphism genotyping

DNA was obtained from the peripheral blood cells of 192 SLE patients and 343 local Caucasian unrelated healthy blood donors by standard procedures. SNPs at positions -1,082 on the gene encoding IL-10 and -308 on the gene encoding TNF α were determined by analyzing the Tm of the probe/target duplex after PCR amplification and hybridization with fluorescent-labeled probes matched with one sequence variant (LighCycler, Roche Diagnostics, Mannheim, Germany), as was previously reported [10]. The primers used were: 5'-ATC CAA GAC AAC ACT ACT AAG GC and 5'-ATG GGG TGG AAG AAG TTG AA for -1,082 IL-10 and 5'-CCT GCA TCC TGT CTG GAA GTT A and 5'-CTG CAC CTT CTG TCT CGG TTT for -308 TNF α . The hybridization probes (designed by TIB MOLBIOL, Berlin, Germany) were: GGA TAG GAG GTC CCT TAC TTT CCT CTT ACC-F and LC Red 640-CCC TAC TTC CCC CTC CCA AA for -1,082 IL-10 and AAC CCC GTC CCC ATG CCC C-F and LC Red 640-CCA AAC CTA TTG CCT CCA TTT CTT TTG GGG AC for -308 TNF α .

Statistical analysis

As serum TNF α levels were not distributed normally, nonparametric testing was used throughout (Mann-Whitney *U* test or Kruskal-Wallis test). Correlations between TNF α concentration and clinical parameters were performed using Spearman's rank correlation test. TNF α values were described by median and interquartile range. Univariate and multivariate analyses were performed by unconditional logistic regression to define the impact of specific single or combined functional genotypes on the response to antimalarial treatment, calculating odds ratios (ORs) and 95% confidence intervals (95% CI). Single-locus regression models were run to estimate the effects of IL-10 and TNF α cytokine polymorphisms separately, comparing the high producer genotypes with the most common low producers. A combined two-loci model was developed, including both cytokine polymorphisms to estimate individual effects of each combined genotype, using the common low/low producer genotype as referent. Covariates for the multivariate analyses included sex, age, disease duration and the clinical parameters: age at diagnosis, malar rash, discoid or subacute cutaneous lesions, photosensitivity, oral ulcers, arthritis, serositis, renal, neurological or hematological

disorder. The SPSS 12.0 statistical software package (SPSS Inc., Chicago, IL) was used for all calculations.

Results**Antimalarial treatment associates with low TNF α serum levels in SLE patients**

Quantification of TNF α levels in the serum of 171 SLE patients and 215 healthy controls (Table 2) showed a significantly higher amount of this cytokine in the entire patient population compared with controls ($p = 0.020$, Mann-Whitney *U* test). Spearman's rank correlation test did not show any significant relationship between treatment with corticosteroids or immunosuppressive drugs or the clinical features age at diagnosis, malar rash, discoid or subacute cutaneous lesions, photosensitivity, oral ulcers, arthritis, serositis, or renal, neurological or hematological disorder and TNF α serum levels. However, a highly significant negative correlation was detected between the use of antimalarial drugs during at least three months before sampling and the concentration of serum TNF α ($\rho = -0.296$, $p = 0.008$, Spearman's test). In fact, when patients were stratified according to treatment (Table 2), no differences were detected between controls and patients under antimalarial treatment, either alone or combined with corticosteroids, whereas increased levels were observed in untreated patients and those with corticosteroid or other immunosuppressive therapies. Kruskal-Wallis test analysis did not show significant differences among treatments, probably due to the reduced number of patients in each group after stratification. Thus, when patients were classified as users or nonusers of antimalarial drugs (Figure 1), we found that patients without antimalarial treatment had significantly higher levels of serum TNF α (median value, 61.45) than both healthy controls (19.66, $p = 0.00034$) and SLE patients receiving this drug (20.60, $p = 0.008$). These results suggest that antimalarial treated SLE patients do not have the increased TNF α production usually found in lupus patients, showing serum levels similar to healthy controls.

Antimalarial-mediated inhibition of TNF α is associated with TNF α and IL-10 promoter genotypes

It has been previously shown that the TNF α genotype at the -308 position (A/G) regulates basal TNF α mRNA levels, although no significant differences were detected in healthy

Table 4**TNF α levels in high and low IL-10 producing patient users and nonusers of antimalarial treatment**

IL-10 genotype	Healthy controls		Patients without antimalarial drugs		Patients with antimalarial drugs		<i>p</i>
	<i>n</i>	TNF α (pg/ml) ^a	<i>n</i>	TNF α (pg/ml) ^a	<i>n</i>	TNF α (pg/ml) ^a	
Low (AA/AG)	179	18.55 (49.27)	56	80.88 (201.48)	79	20.94 (84.97)	0.005
High (GG)	36	21.74 (58.42)	12	51.91 (51.75)	23	20.03 (134.44)	0.797

^aValues are median (interquartile range). Differences between users and nonusers of antimalarial therapy were evaluated by the Mann-Whitney *U* test. TNF, tumor necrosis factor.

controls at the protein level [10]. To ascertain the possible influence of this functional SNP on the association between antimalarial treatment and decreased TNF α levels, patients and controls were genotyped and classified as genetically high (-308AA or AG) or low (-308GG) TNF α producers and TNF α serum levels were evaluated in patients who used antimalarial drugs and those who did not (Table 3). Highly significant differences between the two groups were detected among high TNF α producer patients ($p = 0.001$), whereas no differences between users and nonusers were observed among genetically low TNF α producers.

It is known that the two cytokines TNF α and IL-10 are mutually regulated and that, similar to TNF α , IL-10 levels are genetically determined. We thus wished to evaluate the possible role of functional IL-10 genotypes on the suggested antimalarial-mediated TNF α downregulation. All individuals were accordingly classified as high (GG) and low (AA/AG) genetic IL-10 producers by determination of the allele present at the -1,082 position. Table 4 indicates that IL-10 genotype is able to influence TNF α serum levels in SLE patients, as significant differences between users and nonusers of antimalarial treatment were detected among low IL-10 producers ($p = 0.005$). No significant variations were observed among high IL-10 producers. In conjunction, these results suggest a relationship between antimalarial treatment and low TNF α serum levels in genetically high TNF α and low IL-10 producing SLE patients. This association was probably due to the high TNF α levels of this group. In fact, when patients were classified in the four possible combined IL-10/TNF α genotypes, we found that TNF α serum levels in SLE patients without antimalarial treatment were influenced by both cytokine polymorphisms (Figure 2). Low IL-10/high TNF α patients presented significantly higher levels than high IL-10/low TNF α producers (93.19 versus 22.19, $p = 0.018$) whereas patients with balanced cytokine production (low/low and high/high) presented intermediate values (73.61 and 59.97, respectively).

Combined IL-10 and TNF α genotype influences response to antimalarial treatment in SLE patients

Finally, in order to examine the role of IL-10 and TNF α genetic polymorphisms as predictors of response to treatment, we selected those patients who had been users of antimalarial

agents for more than three years without the need for any other specific SLE therapy, thus indicating a successful response to treatment. Among the 192 patients previously genotyped for SNPs at both cytokine genes, we found that 40 patients (20.83%) were good responders to antimalarial therapy whereas another 74 (38.54%) were also users of this treatment but required the combination with corticosteroids or immunosuppressive drugs. Table 5 shows an overrepresentation of the high TNF α genotype in the good responder group. Therefore, using logistic regression modeling, we evaluated the influence of single and combined IL-10 and TNF α functional genotypes on the response to antimalarial therapy. Table 6 shows a significant association between carriage of the high TNF α producer genotype and good response to antimalarial drugs (OR 2.25, 95%CI 1.11–4.58, $p = 0.024$), whereas the IL-10 genotype did not show any significant association. However, when combined genotypes were analyzed, only the low IL-10/high TNF α genotype was significantly associated (OR 3.13, 95%CI 1.41–6.92, $p = 0.005$). Analysis of clinical features indicated that the group of 40 patients who were long-term users of antimalarial drugs without requiring any other specific treatment was characterized by lower frequency of serositis and nephritis when compared with the remainder of the patients ($p = 0.014$ and 0.003 , respectively). No significant differences were detected with respect to patient age (46.25 ± 12.95 years versus 47.50 ± 15.07) or disease duration (12.68 ± 7.75 years versus 13.28 ± 7.86). Moreover, previous associations were sustained in the multivariate analysis after adjusting for sex, age, disease duration and clinical parameters. Therefore, these results indicate that, in addition to the TNF2 allele, carriage of the low IL-10 producer genotype is required to become a very good responder patient to antimalarial treatment.

Discussion

In this study we show that SLE patients receiving single or combined treatment with antimalarial drugs have TNF α serum levels similar to healthy controls, whereas untreated patients and those receiving corticosteroid or immunosuppressive therapies presented increased amounts of this cytokine. These results suggest a very valuable effect of antimalarial treatment by means of the downregulation of *in vivo* TNF α levels. Although these pharmacological agents have been widely

Table 6**Association between IL-10 and TNF α genotypes and response to antimalarial treatment**


Genotype	Healthy controls n (%)	SLE patients		Univariate analysis		Multivariate analysis ^a	
		Good responders to antimalarials n (%)	Others n (%)	OR (95% CI)	<i>p</i>	OR (95% CI)	<i>p</i>
-1,082 IL-10							
Low (AA/AG)	292 (85.1)	34 (85.0)	119 (78.8)	Referent		Referent	
High (GG)	51 (14.9)	6 (15.0)	32 (21.2)	0.66 (0.25–1.70)	0.386	0.59 (0.20–1.78)	0.354
-308 TNFα							
Low (GG)	265 (77.3)	17 (42.5)	95 (62.5)	Referent		Referent	
High (AA/GA)	78 (22.7)	23 (57.5)	57 (37.5)	2.25 (1.11–4.58)	0.024	2.67 (1.20–5.97)	0.016
Combined IL-10/TNFα							
Low/Low	224 (65.3)	12 (30.0)	75 (49.7)	Referent		Referent	
Low/High	68 (19.8)	22 (55.0)	44 (29.1)	3.13 (1.41–6.92)	0.005	4.20 (1.66–10.63)	0.002
High/Low	41 (12.0)	5 (12.5)	19 (12.6)	1.64 (0.52–5.24)	0.400	1.77 (0.47–6.63)	0.397
High/High	10 (2.9)	1 (2.5)	13 (8.6)	0.48 (0.06–4.02)	0.499	0.43 (0.04–4.21)	0.472
Trend test				0.021		0.011	

Association was calculated by unconditional logistic regression modeling using good response to antimalarial treatment as the dependent variable. ^aAdjusted for sex, age, disease duration and clinical parameters: malar rash, discoid lesions, subacute cutaneous lesions, photosensitivity, oral ulcers, arthritis, serositis, renal disorder, neurological disorder, hematological disorder and age at systemic lupus erythematosus (SLE) diagnosis (continuously). CI, confidence interval; OR, odds ratio; TNF, tumor necrosis factor.

used as disease-modifying antirheumatic drugs mainly in the treatment of SLE and rheumatoid arthritis [1], their mechanisms of anti-inflammatory actions have not yet been completely understood. Several *in vitro* experiments have demonstrated that quinacrine and related drugs decreased the release of pro-inflammatory cytokines induced by LPS in macrophages [4-7,15]. In mouse models, it has been reported that chloroquine may protect mice from lethal challenge by CpG oligonucleotides and LPS and may decrease serum TNF α and IL-6 in rats injected with sublethal doses of both stimuli [16]. To the best of our knowledge, however, this is the first report demonstrating an *in vivo* association between the use of antimalarial therapy and low levels of serum TNF α , suggesting that the disease-modifying antirheumatic effect of these drugs may be mediated, at least in part, by a strong downregulatory effect on TNF α production.

However, association between antimalarial treatment and TNF α serum levels seems to be influenced by polymorphisms of the genes encoding TNF α and IL-10, indicating that this advantageous connection may only be completely valuable for patients with a specific genotype. Constitutive and induced TNF α and IL-10 production have important interindividual variations that are genetically regulated by SNP at their promoters [8-12]. Our data indicate that antimalarial therapy plays a role

in the TNF α production of patients who are genetically high TNF α producers. These patients probably have the highest TNF α transcription rates [9] and consequently the highest serum levels. Although the mechanisms of *in vitro* antimalarial-mediated TNF α inhibition [4-7] are not yet entirely known, it has been reported that chloroquine inhibited LPS-induced TNF α transcription [6,7], interfering with mitogen-activated protein kinase signaling [17]. Thus, our results suggest that antimalarial agents require a high rate of TNF α transcription to achieve the maximal inhibitory effect. On the other hand, though the relevance of IL-10 was already known in lupus disease, the influence of genetic polymorphisms at the IL-10 promoter on treatment outcome after the use of antimalarials was surprising. Results indicated an involvement of antimalarial treatment in the amount of serum TNF α in SLE patients with a low IL-10 genotype. The limited TNF α downregulatory effect observed in genetically high IL-10 producing patients might be explained by the regulatory feedback mechanism that controls the production of both cytokines, which would lead to a decrease in the TNF α transcription rate in patients producing elevated amounts of IL-10. Supporting this, it has been reported that high IL-10 levels were associated with less effective clearance of *Plasmodium falciparum* parasites in patients receiving antimalarial therapy [18].

Figure 2

Combined IL-10/tumor necrosis factor (TNF) α genotype influences TNF α serum levels in SLE patients without antimalarial treatment. After determination of the allele present at positions -1,082 and -308 of the IL-10 and TNF α gene promoters, respectively, patients were classified as genetically high (GG) or low (AA/AG) IL-10 producers and high (AA/AG) or low (GG) TNF α producers. Box plots represent TNF α serum levels in SLE patients not receiving antimalarial therapy classified according to the four possible IL-10/TNF α combined genotypes. Differences were evaluated by the Mann-Whitney *U* test.

Our data support the idea that the actions of cytokines are profoundly conditioned by the presence of other cytokines, particularly in the case of IL-10 and TNF α , which have opposing roles in systemic inflammatory responses. Thus, on the basis of our previous results, we evaluated the role of the interaction between IL-10 and TNF α genotypes in regulating the response to antimalarial treatment in SLE patients. A strong association was found between carriage of the combined genotype low IL-10/high TNF α and the use of antimalarials for more than three years without the need for any other specific SLE treatment (good responder patients), although the single analysis of the IL-10 genotype did not show significant results. Moreover, the relationship between this combined genotype and treatment outcome was higher than that obtained after

single analysis of the gene encoding TNF α , since high IL-10/high TNF α producers were not overrepresented among good responder patients. Taken in conjunction, these results thus indicate that determination of TNF α and IL-10 alleles at the onset of the disease may help identify more suitable candidates for antimalarial treatment and could be used as a genetic predictor of clinical outcome. We would expect SLE patients who are carriers of the pro-inflammatory genotype low IL-10/high TNF α to develop a mild disease presenting a good course under antimalarial therapy. Most of these patients probably developed SLE due to the effect of environmental factors added to their genetically determined high TNF α levels, which could not be modulated by the low production of IL-10. TNF α is a pro-inflammatory cytokine that has been found at elevated levels in the serum of patients suffering SLE and other autoimmune diseases [19,20]; it has also been suggested that TNF α genotype influences their susceptibility [10,21,22] and, possibly, their clinical response to treatment. In these patients, the elevated TNF α levels may be involved in diverse pathological mechanisms and, therefore, a clinical benefit is to be expected under a treatment that diminishes TNF α production.

Under the assumption that the elevated TNF α levels found in patients with various chronic inflammatory diseases are deleterious, several anti-TNF α therapies are now available to block the action of TNF α . Actually, TNF α -blockage with antibodies has been of unquestionable clinical benefit to many patients with rheumatoid arthritis, Crohn's disease, psoriasis and, more recently, ankylosing spondylitis [20,23-26]. However, it has been reported that response to this treatment is also influenced by genetic polymorphisms at Fc γ RIII [27], HLA-DRB1 [28], lymphotoxin- α [29] and TNF α alone [30] or combined with IL-10 [31] or lymphotoxin- α [28]. In fact, TNF α antagonists seem to be more effective in genetically low TNF α producer patients [30] or with the combined high IL-10/low TNF α genotype [31]. Several authors have also proposed the use of

Table 5**IL-10 and TNF α genotypes of systemic lupus erythematosus patients stratified by treatment**

Treatment	TNF α genotype		IL-10 genotype	
	High (AA/AG)	Low (GG)	High (GG)	Low (AA/AG)
None/NSAIDS	6 (26.1)	17 (73.9)	4 (17.4)	19 (82.6)
Antimalarials	23 (57.5)	17 (42.5)	6 (15.0)	34 (85.0)
Corticosteroids	8 (28.6)	20 (71.4)	6 (22.2)	21 (77.8)
Antimalarials and corticosteroids	25 (41.7)	35 (58.3)	16 (26.7)	44 (73.3)
Immunosuppressive drugs ^a alone or with corticosteroids	13 (48.1)	14 (51.9)	3 (11.1)	24 (88.9)
Antimalarials, corticosteroids and immunosuppressive drugs ^a	5 (35.7)	9 (64.3)	3 (21.4)	11 (78.6)

Values are n (%). ^aMethotrexate, azathioprine, cyclophosphamide, cyclosporine A or mycophenolate mophetil. NSAIDS, nonsteroidal anti-inflammatory drugs; TNF, tumor necrosis factor.

TNF α lowering agents in the treatment of lupus disease [20,32,33]. Supporting this, an open label study of infliximab in six patients with SLE indicated that TNF α blockage might have a therapeutically beneficial effect, although autoantibodies to double-stranded DNA and cardiolipin were increased [34]. This trend toward an augmented autoantibody production frequently accompanying this treatment was not reported after the use of antimalarials. Given the therapeutic relevance of these results, new studies need to be designed to evaluate the use of these drugs as TNF α downregulators in the treatment of genetically low IL-10/high TNF α producer patients with SLE and other inflammatory diseases who, in addition, are poor responders to TNF α blockage with antibodies.

The limitations of the work were the lack of a prospective longitudinal study analyzing TNF α levels before and after treatment with antimalarials and other SLE therapies and the absence of standardized validated measures of SLE activity.

Conclusion

Our results demonstrate a relationship between antimalarial treatment and low TNF α serum levels in SLE patients that were influenced by polymorphisms at the IL-10 and TNF α gene promoters. Therefore, our findings may have an important clinical application through the identification of patients who are the most likely to benefit from antimalarial therapy.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PL performed genetic and immunological assays, data collection and statistical analyses and participated in interpretation of data. JG carried out a detailed review of clinical histories and participated in data collection. LM performed patient selection and participated in data collection. CG participated in the study design, interpretation of data and helped to draft the manuscript. AS conceived the study, participated in its design and coordination, performed genetic assays and statistical analyses and drafted the manuscript.

Acknowledgements

The authors wish to thank ALAS (Asociación Lúpicas de Asturias) for its continuous encouragement. Supported by Grant SV-04-FMM-01 from the Fundación Médica Mutua Madrileña.

References

- Rynes RI: **Antimalarial drugs in the treatment of rheumatological diseases.** *Br J Rheumatol* 1997, **36**:799-805.
- MacIntyre AC, Cutler DJ: **Role of lysosomes in hepatic accumulation of chloroquine.** *J Pharm Sci* 1988, **77**:196-199.
- Macfarlane DE, Manzel L: **Antagonism of immunostimulatory CpG-oligodeoxynucleotides by quinacrine, chloroquine, and structurally related compounds.** *J Immunol* 1998, **160**:1122-1131.
- Karres I, Kremer JP, Dielt I, Steckholzer U, Jochum M, Ertel W: **Chloroquine inhibits proinflammatory cytokine release into human whole blood.** *Am J Physiol* 1998, **274**:R1058-R1064.
- Seitz M, Valbracht J, Quach J, Lotz M: **Gold sodium thiomalate and chloroquine inhibit cytokine production in monocytic THP-1 cells through distinct transcriptional and posttranslational mechanisms.** *J Clin Immunol* 2003, **23**:477-484.
- Zhu X, Ertel W, Ayala A, Morrison MH, Perrin MM, Chaudry IH: **Chloroquine inhibits macrophage tumour necrosis factor-alpha mRNA transcription.** *Immunology* 1993, **80**:122-126.
- Weber SM, Levitz SM: **Chloroquine interferes with lipopolysaccharide-induced TNF-alpha gene expression by a nonlyso-somotropic mechanism.** *J Immunol* 2000, **165**:1534-1540.
- Kroeger KM, Steer JH, Joyce DA, Abraham LJ: **Effects of stimulus and cell type on the expression of the -308 tumour necrosis factor promoter polymorphism.** *Cytokine* 2000, **12**:110-119.
- Wilson AG, Symons JA, McDowell TL, McDevitt HO, Duff GW: **Effects of a polymorphism in the human tumor necrosis factor alpha promoter on transcriptional activation.** *Proc Natl Acad Sci USA* 1997, **94**:3195-3199.
- Suarez A, Lopez P, Mozo L, Gutierrez C: **Differential effect of IL10 and TNF[alpha] genotypes on determining susceptibility to discoid and systemic lupus erythematosus.** *Ann Rheum Dis* 2005, **64**:1605-1610.
- Suarez A, Castro P, Alonso R, Mozo L, Gutierrez C: **Interindividual variations in constitutive interleukin-10 messenger RNA and protein levels and their association with genetic polymorphisms.** *Transplantation* 2003, **75**:711-717.
- Turner DM, Williams DM, Sankaran D, Lazarus M, Sinnott PJ, Hutchinson IV: **An investigation of polymorphism in the interleukin-10 gene promoter.** *Eur J Immunogenet* 1997, **24**:1-8.
- Lopez P, Mozo L, Gutierrez C, Suarez A: **Epidemiology of systemic lupus erythematosus in a northern Spanish population: gender and age influence on immunological features.** *Lupus* 2003, **12**:860-865.
- Tan EM, Cohen AS, Fries JF, Masi AT, McShane DJ, Rothfield NF, Schaller JG, Talal N, Winchester RJ: **The 1982 revised criteria for the classification of systemic lupus erythematosus.** *Arthritis Rheum* 1982, **25**:1271-1277.
- Salmeron G, Lipsky PE: **Immunosuppressive potential of antimalarials.** *Am J Med* 1983, **75**:19-24.
- Hong Z, Jiang Z, Liangxi W, Guofu D, Ping L, Yongling L, Wendong P, Minghai W: **Chloroquine protects mice from challenge with CpG ODN and LPS by decreasing proinflammatory cytokine release.** *Int Immunopharmacol* 2004, **4**:223-234.
- Weber SM, Chen JM, Levitz SM: **Inhibition of mitogen-activated protein kinase signaling by chloroquine.** *J Immunol* 2002, **168**:5303-5309.
- Hugosson E, Montgomery SM, Premji Z, Troye-Blomberg M, Bjorkman A: **Higher IL-10 levels are associated with less effective clearance of Plasmodium falciparum parasites.** *Parasite Immunol* 2004, **26**:111-117.
- Aringer M, Feierl E, Steiner G, Stummvoll GH, Hofler E, Steiner CW, Radda I, Smole JS, Graninger WB: **Increased bioactive TNF in human systemic lupus erythematosus: associations with cell death.** *Lupus* 2002, **11**:102-108.
- Atzeni F, Sarzi-Puttini P, Doria A, Iaccarino L, Capsoni F: **Potential off-label use of infliximab in autoimmune and non-autoimmune diseases: a review.** *Autoimmun Rev* 2005, **4**:144-152.
- Rood MJ, van Krugten MV, Zanelli E, van der Linden MW, Keijsers V, Schreuder GM, Verduyn W, Westendorp RG, de Vries RR, Breedveld FC, Verweij CL, Huizinga TW: **TNF-308A and HLA-DR3 alleles contribute independently to susceptibility to systemic lupus erythematosus.** *Arthritis Rheum* 2000, **43**:129-134.
- Sullivan KE, Wooten C, Schmeckpeper BJ, Goldman D, Petri MA: **A promoter polymorphism of tumor necrosis factor alpha associated with systemic lupus erythematosus in African-Americans.** *Arthritis Rheum* 1997, **40**:2207-2211.
- Davis JC, van der Heijde DM, Braun J, Dougados M, Cush J, Clegg D, Inman RD, Kivitz A, Zhou L, Solinger A, Tsuji W: **Sustained durability and tolerability of etanercept in ankylosing spondylitis for 96 weeks.** *Ann Rheum Dis* 2005, **64**:1557-1562.
- Maini R, St Clair EW, Breedveld F, Furst D, Kalden J, Weisman M, Smolen J, Emery P, Harriman G, Feldmann M, Lipsky P: **Infliximab (chimeric anti-tumour necrosis factor alpha monoclonal antibody) versus placebo in rheumatoid arthritis patients receiv-**

- ing concomitant methotrexate: a randomised phase III trial. **ATTRACT Study Group.** *Lancet* 1999, **354**:1932-1939.
25. Reimold AM: **New indications for treatment of chronic inflammation by TNF-alpha blockade.** *Am J Med Sci* 2003, **325**:75-92.
 26. Suryaprasad AG, Prindiville T: **The biology of TNF blockade.** *Autoimmun Rev* 2003, **2**:346-357.
 27. Tutuncu Z, Kavanaugh A, Zvaifler N, Corr M, Deutsch R, Boyle D: **Fcgamma receptor type IIIA polymorphisms influence treatment outcomes in patients with inflammatory arthritis treated with tumor necrosis factor alpha-blocking agents.** *Arthritis Rheum* 2005, **52**:2693-2696.
 28. Criswell LA, Lum RF, Turner KN, Woehl B, Zhu Y, Wang J, Tiwari HK, Edberg JC, Kimberly RP, Moreland LW, Seldin MF, Bridges SL Jr: **The influence of genetic variation in the HLA-DRB1 and LTA-TNF regions on the response to treatment of early rheumatoid arthritis with methotrexate or etanercept.** *Arthritis Rheum* 2004, **50**:2750-2756.
 29. Taylor KD, Plevy SE, Yang H, Landers CJ, Barry MJ, Rotter JI, Targan SR: **ANCA pattern and LTA haplotype relationship to clinical responses to anti-TNF antibody treatment in Crohn's disease.** *Gastroenterology* 2001, **120**:1347-1355.
 30. Mugnier B, Balandraud N, Darque A, Roudier C, Roudier J, Reviron D: **Polymorphism at position -308 of the tumor necrosis factor alpha gene influences outcome of infliximab therapy in rheumatoid arthritis.** *Arthritis Rheum* 2003, **48**:1849-1852.
 31. Padyukov L, Lampa J, Heimbürger M, Ernestam S, Cederholm T, Lundkvist I, Andersson P, Hermansson Y, Harju A, Klareskog L, Bratt J: **Genetic markers for the efficacy of tumour necrosis factor blocking therapy in rheumatoid arthritis.** *Ann Rheum Dis* 2003, **62**:526-529.
 32. Anolik JH, Aringer M: **New treatments for SLE: cell-depleting and anti-cytokine therapies.** *Best Pract Res Clin Rheumatol* 2005, **19**:859-878.
 33. Smolen JS, Steiner G, Aringer M: **Anti-cytokine therapy in systemic lupus erythematosus.** *Lupus* 2005, **14**:189-191.
 34. Aringer M, Graninger WB, Steiner G, Smolen JS: **Safety and efficacy of tumor necrosis factor alpha blockade in systemic lupus erythematosus: an open-label study.** *Arthritis Rheum* 2004, **50**:3161-3169.