

Oxidative Stress in Toxicology: Established Mammalian and Emerging Piscine Model Systems

Sue A. Kelly,^{1,2} Christine M. Havrilla,^{2,3} Todd C. Brady,^{2,4} Kimberly Harris Abramo,^{2,3} and Edward D. Levin²

¹Nicholas School of the Environment, ²Integrated Toxicology Program, ³Department of Chemistry, and ⁴Department of Pathology, Duke University, Durham, NC 27708 USA

Interest in the toxicological aspects of oxidative stress has grown in recent years, and research has become increasingly focused on the mechanistic aspects of oxidative damage and cellular responses in biological systems. Toxic consequences of oxidative stress at the subcellular level include lipid peroxidation and oxidative damage to DNA and proteins. These effects are often used as end points in the study of oxidative stress. Typically, mammalian species have been used as models to study oxidative stress and to elucidate the mechanisms underlying cellular damage and response, largely because of the interest in human health issues surrounding oxidative stress. However, it is becoming apparent that oxidative stress also affects aquatic organisms exposed to environmental pollutants. Research in fish has demonstrated that mammalian and piscine systems exhibit similar toxicological and adaptive responses to oxidative stress. This suggests that piscine models, in addition to traditional mammalian models, may be useful for further understanding the mechanisms underlying the oxidative stress response. *Key words:* antioxidants, fish, lipid peroxidation, mammals, oxidative DNA damage, oxidative stress, reactive oxygen species. *Environ Health Perspect* 106:375–384 (1998). [Online 9 June 1998] <http://ehpnet1.niehs.nih.gov/docs/1998/106p375-384kelly/abstract.html>





The discovery of superoxide dismutase by McCord and Fridovich (1) provided the first evidence implicating radicals and oxidative stress as important sources of biological injury. Oxidative stress occurs when there is an imbalance in the generation and removal of radical species within an organism. The majority of these radicals involve oxygen and are referred to as reactive oxygen species (ROS) (2). Because of the potential for ROS to damage tissues and cellular components such as membranes (3,4), DNA (5–7), and proteins (8–10), oxidative stress has become a topic of significant interest in both mammalian and aquatic toxicology.

The majority of animal studies focusing on oxidative stress have employed mammalian cell or organ models to analyze the oxidative capacity of a particular toxicant or to evaluate the efficacy of a compound, such as an antioxidant, to protect from toxicant-mediated oxidative damage. Inflammation, ischemic reperfusion injury, aging, and carcinogenesis have all been linked to oxidative stress in mammalian systems (10–13). However, while the breadth of knowledge in mammalian systems is great, a considerable amount remains to be understood regarding cellular response mechanisms, disease etiology, and repair processes within cells. Notably, there is a growing literature on the effects of oxidative stress in aquatic organisms, and it is

becoming apparent that the general mechanisms of oxidative toxicity in both mammalian and piscine systems are similar, with comparable lesions and responses serving as markers of oxidative stress (14). Consequently, evidence suggests that fish have the potential to serve as suitable model organisms for studying the mechanistic aspects of oxidative toxicity, such as chemical carcinogenesis and damage repair, that are also relevant to biomedical toxicology.

Sources of Reactive Oxygen Species

Several basic cellular processes lead to the production of ROS within a cell. Cellular respiration involves the reduction of molecular oxygen (O₂) to water in the electron transport chain. This reduction occurs through four one-electron reductions resulting in the formation of reactive, partially reduced intermediates such as the superoxide anion radical (·O₂⁻), hydrogen peroxide (H₂O₂), and the hydroxyl radical (HO·) that may act as prooxidants:



One to 5% of these ROS may escape the electron transport chain and damage cellular components (15).

Several oxidizing enzymes produce ROS, comprising a second source of ROS within cells. Diamine oxidase, tryptophan dioxygenase, xanthine oxidase, and cytochrome P450 reductase can generate ·O₂⁻, while enzymes such as guanyl cyclase and glucose oxidase generate H₂O₂ (16–18). Similarly, nitric oxide synthase produces ·O₂⁻ under low arginine conditions (19). Other enzymatic processes produce extracellular ·O₂⁻ and include the activity of the leukotriene generator lipoxygenase (20) and the prostaglandin generator cyclooxygenase (21). The enzymatic production of ROS is of additional interest in toxicology because some of these enzymes, such as cytochrome P450, are involved in the metabolism of xenobiotics (14,22).

Some radicals serve useful purposes. For instance, during the respiratory burst of leukocytes in immune-mediated host defense, monocytes, neutrophils, and macrophages produce ·O₂⁻, nitric oxide (·NO), and peroxynitrite (ONOO·) (23). This response is triggered by stimulation of enzymes such as NADPH-oxidase (24), and these ROS are used as cytotoxic agents against pathogenic organisms (23). In addition to its use in the immune response, ·NO can act as a neurotransmitter and muscle relaxant (25), while levels of other radicals play a role in the control of the transcription factors nuclear factor κB (NF-κB) and activator protein-1 (AP-1) (26). Additionally, superoxide has been proposed to terminate lipid peroxidation and may be mitogenic as well (27). The mitogenic function of ·O₂⁻ is supported by a recent finding suggesting that ·O₂⁻ may be a mitogenic signal in *ras*-mediated cellular transformation (28).

Address correspondence to S.A. Kelly, Nicholas School of the Environment - LSRC Rm A141, Research Drive, Duke University, Durham, NC 27708 USA.

The authors wish to thank Irwin Fridovich and Richard Di Giulio for technical review of the manuscript.

Received 12 November 1997; accepted 16 March 1998.



Xenobiotics comprise a third source of ROS because some xenobiotics can enhance the production of oxyradicals within cells. This may occur through processes such as inhibition of mitochondrial electron transport and subsequent accumulation of reduced intermediates (29), inactivation of antioxidant enzymes such as catalase (30,31), and depletion of radical scavengers (32–35). Of particular interest in toxicology are xenobiotics that are capable of redox cycling (Fig. 1). Quinones, some dyes, bipyridyl herbicides, some transition metals, and aromatic nitro compounds comprise classes of compounds known to redox cycle. Redox cycling involves the univalent reduction of the xenobiotic to a radical intermediate by enzymes such as xanthine oxidase and NADPH-cytochrome P450 reductase. This radical intermediate then transfers an electron to O_2 , producing $\cdot O_2^-$ and regenerating the parent compound. Thus, a single molecule of parent compound can generate many oxyradicals. Furthermore, this process occurs at the expense of cellular reducing equivalents, such as NADPH, which can have consequences for other metabolic processes (36–38).

Significance of Reactive Oxygen Species



Once produced, ROS may damage cellular components and tissues. Superoxide and

$HO\cdot$ are two of the most studied ROS in this respect. Superoxide oxidizes catecholamines, tocopherols, ascorbate, and thiols (39,40). Superoxide can also inactivate enzymes such as catalase (31) and peroxidases (30). Furthermore, $\cdot O_2^-$ inactivates [4Fe-4S]-containing dehydratases (e.g., aconitase, fumarase) and releases Fe^{2+} (41).

Much of the toxicity of $\cdot O_2^-$ may be explained by its reaction with other ROS, such as H_2O_2 , to form more reactive ROS. Of particular biological importance is the production and toxicity of $HO\cdot$ (Eq. 5 and 6). One of the most potent oxidants known, $HO\cdot$ reacts indiscriminately with cellular components such as lipids, DNA, and proteins. Because of its high reactivity, $HO\cdot$ has an *in vivo* lifetime of only a few nanoseconds (42). $HO\cdot$ may be produced by the reaction of $\cdot O_2^-$ and H_2O_2 in the Haber-Weiss reaction (Eq. 6) (43).



Because this reaction is kinetically slow (16), transition metals (e.g., iron and copper) often serve as catalysts. The iron-catalyzed Haber-Weiss reaction is depicted in Equations 7 and 8. Equation 8 is also known as the Fenton reaction (44).



The polyanionic nature of cell membranes and DNA provides a structure for the adherence of metal cations. Thus, $HO\cdot$ produced in the Fenton reaction would be expected to occur adjacent to these critical biological targets, resulting in subsequent oxidation of lipids and DNA (45–47). The mechanistic aspects of oxidative damage to lipids and DNA are discussed below.

Lipid peroxidation. The important role of lipids in cellular components emphasizes the significance of understanding of the mechanisms and consequences of lipid peroxidation in biological systems. Polyunsaturated fatty acids (PUFAs) serve as excellent substrates for lipid peroxidation because of the presence of active bis-allylic methylene groups. The carbon–hydrogen bonds on these activated methylene units have lower bond dissociation energies, making these hydrogen atoms more easily abstracted in radical reactions (48). The susceptibility of a particular PUFA toward peroxidation increases with an increase in the number of unsaturated sites in the lipid chain (49).

Autoxidation of lipids in biological systems proceeds via a chain reaction consisting of three phases: initiation, propagation, and termination. Initiation of lipid peroxidation *in vivo* has not been extensively studied and is under much debate (50,51). This phase of lipid peroxidation may proceed by the reaction of an activated oxygen species such as singlet oxygen (1O_2), $\cdot O_2^-$, or $HO\cdot$ with a lipid substrate or by the breakdown of pre-existing lipid hydroperoxides by transition metals. In the former case, peroxidation occurs by abstraction of a hydrogen atom from a methylene carbon in the lipid substrate (LH) to generate a highly reactive carbon-centered lipid radical ($L\cdot$).

In the propagation phase of lipid peroxidation, molecular oxygen adds rapidly to




Figure 1. Redox cycling, which includes radical formation, scavenging, and effects. Abbreviations: GSH, glutathione; GSSG, glutathione disulfide. Adapted from Kappus (163).




Figure 2. Overview of lipid peroxidation. Abbreviations: NRP, nonradical product; LOOH, lipid hydroperoxide; α -TOH, α -tocopherol; α - $TO\cdot$, α -TOH radical; LH, lipid substrate; $LOO\cdot$, lipid peroxyl radical. Adapted from Waldeck and Stocker (164).

L[•] at a diffusion controlled rate to produce the lipid peroxy radical (LOO[•]) (4). The peroxy radical can abstract a hydrogen atom from a number of *in vivo* sources, such as DNA and proteins, to form the primary oxidation product, a lipid hydroperoxide (LOOH). Alternatively, antioxidants such as α -tocopherol (α -TOH) can act as excellent hydrogen atom donors (52), generating LOOH and the relatively inert α -tocopherol phenoxyl radical (α -TO[•]). In the absence of antioxidants or other inhibitors, LOO[•] can abstract a hydrogen from another lipid molecule (LH), producing another highly reactive carbon centered radical (L[•]), which then propagates the radical chain (Fig. 2).

Transition metals are of particular interest to lipid peroxidation. In transition metal-catalyzed lipid peroxidation, HO[•] is thought to be the primary initiating radical species (51). However, both ferrous (Fe²⁺) and ferric (Fe³⁺) iron, in addition to increasing the production of initiating hydroxyl radicals, can catalyze the propagation of the lipid peroxidation chain by decomposing LOOH oxidation products (53,54). The resulting alkoxy (LO[•]) and peroxy (LOO[•]) radicals are able to initiate new radical chains by interacting with additional lipid molecules. Other redox active metals such as copper ions may catalyze this reaction, but the availability of these ions under physiological conditions is yet unclear (23).

Termination of lipid peroxidation occurs via the coupling of any two radicals to form nonradical products. Nonradical products are stable and unable to propagate lipid peroxidation chains. Several extensive reviews of the chemistry of free radical oxidation of monoene and polyene unsaturated lipids have been discussed by Porter (55–57).

Repair of lipid peroxidation. Lipid peroxidation products modify the physical characteristics of biological membranes (58). For instance, incorporation of LOOH changes the physical structure of the membrane by decreasing the fluidity and increasing the permeability (59). Thus, the removal of the lipid peroxidation products from the membrane is necessary to repair the membrane damage (58) and is accomplished by two separate enzymatic systems: the sequential action of phospholipase A₂ with glutathione peroxidase (60) and phospholipid hydroperoxide glutathione peroxidase (61).

Phospholipases are activated by lipid peroxidation (62). Phospholipase A₂ (PLA₂) shows a substrate specificity for peroxidized phospholipids in membranes, catalyzing the hydrolysis of the phospholipid hydroperoxides to the hydroperoxy fatty acids (63–65).

Once released, the fatty acid hydroperoxides may undergo a reaction with glutathione peroxidase (GPx) to form stable, reduced hydroxy products (58,66). This terminates lipid peroxidation. A second enzymatic system eliminates phospholipid hydroperoxides from lipid membranes through the direct reaction of phospholipid hydroperoxide glutathione peroxidase with the esterified phospholipid hydroperoxides. This *in situ* reduction of the phospholipid hydroperoxides to phospholipid hydroxides while still incorporated in the membrane structure also results in the halt of membrane lipid peroxidation (67).

Detection of lipid peroxidation. Because direct analysis of endogenous primary lipid peroxidation products is complicated, the extent of lipid peroxidation is typically assessed by measuring levels of secondary oxidation products. The primary lipid oxidation products, LOOH, are unstable and decompose to form secondary products such as aldehydes and ketones through a multitude of reaction pathways (68). The resulting diverse array of breakdown products, coupled with the small *in vivo* concentration of these products, presents a challenge for accurate quantification of lipid peroxidation. The most used assay for lipid peroxidation is the thiobarbituric acid reactive substances (TBARS) test. Aldehydic compounds are reactive and highly cytotoxic (69–71), and the TBARS test relies on the production of a colored adduct from the reaction of lipid peroxidation products and thiobarbituric acid (72). This test is most frequently used as a measure of malondialdehyde (MDA), a secondary lipid oxidation product. However, because thiobarbituric acid reacts with a number of other oxidation products including 4-hydroxynoneal, other unsaturated aldehydes (73), and endoperoxides from enzymatic routes (74), this test is nonspecific (75). Therefore, the TBARS test can only give a crude measure of lipid peroxidation. Consequently, as an assay for MDA, the TBARS test generally gives higher values for TBARS reactivity compared to more rigorous tests for MDA (76,77).

Another crude measure of lipid peroxidation is the determination of diene conjugation. Generation of lipid peroxides typically involves the formation of conjugated dienes from the PUFA substrate. These conjugated hydroperoxide products absorb UV light in the region of 230–235 nm. However, many other endogenous materials also absorb UV light in the region of interest, creating a high background absorbance. Thus, this analytical technique is relatively insensitive to small changes in product formation (78). While these crude methods of measuring lipid peroxidation may be less

sensitive, they are often adequate for the question at hand and are more easily performed than more rigorous alternatives.

Oxyradical-generated DNA damage. DNA is another key cellular component that is particularly susceptible to oxidative damage. The primary oxidant responsible for DNA damage is HO[•], as neither H₂O₂ nor peroxy radicals react directly with DNA (79). The heterogeneity of DNA allows for many susceptible sites for HO[•] attack, including the nitrogenous bases and the sugar–phosphate backbone. The rate of HO[•] reaction with bases is approximately five times greater than that with the backbone (80). Lesion production for both portions of the DNA structure have been thoroughly reviewed (6,7).

Generally speaking, HO[•] attack on DNA bases leads to three classes of damage: hydroxylation, ring opening, and fragmentation. The resulting lesions are usually products of the secondary reactions that occur after the initial radical attack. Although over 100 different DNA lesions may result from HO[•] attack on DNA bases (81), the lesions most commonly used as biomarkers are 8-hydroxy-2'-deoxyguanosine (8-OHdG) and 2,6-diamino-4-hydroxy-5-formamidopyrimidine (FapyGua). Both 8-OHdG and FapyGua result from HO[•] attack on deoxyguanosine, and the differentiating step between them is a secondary reaction. The potential steps of formation for both biomarkers are illustrated in Figure 3. These lesions may be directly detected in the DNA or assayed in urine. Specific analytical methods that detect biomarkers of oxidative damage to DNA bases have been reviewed (82).

In addition to producing direct damage to bases, a transient radical species generated within the DNA base may covalently bind with other macromolecules within the cell, forming intermolecular complexes. Perhaps the most important example of this phenomena is the protein–DNA crosslink. While the formation of such crosslinks has been reviewed (83), little is known about the significance of the different classes and structures (84).

Hydroxyl radicals may also attack the sugar–phosphate backbone of DNA, causing a different variety of lesions, including base free sites or apurinic sites (AP sites) where the base has been removed by oxidant-mediated reactions. An indicator of such damage is the presence of free bases in urine. Another indicator for radical attack on the DNA backbone is the fragmentation of deoxyribose. Single strand breaks occur via hydrogen abstraction at the C-4 position, leading to oxidation of the sugar moiety. This may be coupled with a second sugar oxidation on

the complimentary strand, causing a double strand break. Strand breaks in the DNA molecule may prove mutagenic or even lethal for the cell. (79).

Repair of oxidative DNA damage.


There are a variety of cellular defense mechanisms that are designed to protect the genome from radical attack or repair oxidative damage. Some of these defenses include compartmentalization of the sensitive target molecules (81) and shielding of nonreplicating DNA by histones and polyamines (6). If protection of the genetic material is not successful, cellular regulatory mechanisms such as apoptotic induction and inhibition of cell cycle progression may prevent transfer of genetic damage to daughter cells (81). Alternatively, DNA repair processes may

correct the damage. Excision repair is performed by enzymes such as DNA glycosylases and AP endonucleases and occurs before replication, while postreplication repair provides a method for repairing lesions during or after the replication process. Both of these processes are believed to be constitutive and inducible (85). Unlike the repair of damaged bases, many of the processes that are involved in repair of strand breaks are not fully understood (81).

Antioxidant Defenses in Eukaryotes


To prevent damage to cellular components, there are numerous enzymatic antioxidant defenses designed to scavenge ROS in the cell. Superoxide dismutases (SODs) are

enzymes that scavenge $\cdot\text{O}_2^-$ by a rapid dismutation reaction ($k > 10^9 \text{ M}^{-1}\text{sec}^{-1}$) (1):



There are three SOD isoforms in eukaryotes: manganese SOD (Mn-SOD), copper/zinc SOD (Cu/Zn-SOD), and extracellular SOD (EC-SOD) (41). Mn-SOD, which contains a manganese prosthetic group, resides in the mitochondria, perhaps because of the need to protect mitochondrial proteins, membranes, and DNA from $\cdot\text{O}_2^-$ generated as a result of the respiratory chain. The Cu/Zn-SOD, containing copper and zinc prosthetic groups, is a cytosolic SOD. EC-SOD, also containing copper and zinc prosthetic groups, is secreted and binds to the plasma membrane and heparin-containing elements of the extracellular matrix (86). All forms of SODs are thought to dismutate $\cdot\text{O}_2^-$ via a ping-pong mechanism whereby the transition metal prosthetic group is reduced by $\cdot\text{O}_2^-$, forming O_2^- . The metal in the prosthetic group is then immediately re-oxidized by another $\cdot\text{O}_2^-$ molecule, resulting in the production of H_2O_2 (17).

As mentioned previously, the production of H_2O_2 within the cell may lead to the production of $\text{HO}\cdot$ and subsequent cellular damage via the metal-catalyzed Haber-Weiss reaction. Thus, it is important to remove H_2O_2 . Catalase, a manganese or heme-containing enzyme, functions to rapidly dismutate H_2O_2 to water and oxygen ($k > 10^7 \text{ M}^{-1} \text{sec}^{-1}$) (87):



Catalase is mainly found in peroxisomes, perhaps because of the large number of H_2O_2 -producing oxidases found in these organelles, while lower levels are also found in mitochondria and the cytosol. Several reports suggest that catalase can metabolize alkyl peroxides, such as methyl hydroperoxide and ethyl hydroperoxide in addition to H_2O_2 (88), but these events are thought to be an insignificant component of the activity of the enzyme (89).

A third enzymatic antioxidant, GPx, catalyzes the reduction of organic peroxides (ROOH), which are potential radical forming species within the cell. Glutathione (GSH), a nonenzymatic antioxidant, provides the reducing equivalents needed by GPx to carry out the reduction of organic peroxides to the corresponding alcohols and water:






Figure 3. Mechanisms of formation of the common DNA lesions 2,6-diamino-4-hydroxy-5-formamidopyrimidine (FapyGua) and 8-hydroxy-2'-deoxyguanosine (8-OHdG). Adapted from Cadet (165).

There are several cytosolic, selenium-containing GPxs as well as an extracellular form. These enzymes also have high affinity for H_2O_2 and may be important for maintenance of low intracellular levels of H_2O_2 in the cytosol (88), where catalase levels are low. An enzyme important to the function of GPx is glutathione reductase (GR), which reduces oxidized glutathione (GSSG, glutathione disulfide) in the presence of NADPH in order to regenerate levels of GSH (90,91):



Levels of antioxidant enzymes, including SODs, catalases, GPx, and GR, are low in the extracellular space (50,92). Similarly, plasma levels of SODs and catalases are particularly low when compared to intracellular levels (92). Thus, a diverse array of small molecule, nonenzymatic antioxidants complete an efficient network of intra- and extracellular defenses necessary to modulate the levels of ROS. Nonenzymatic antioxidants may be classified in terms of their location and mode of action. Some small molecule antioxidants can act by scavenging oxidants or chelating transition metal ions (93).

Lipid soluble membrane-bound antioxidants are efficient in preventing oxidative damage to biomolecules. Vitamin E is an important lipid soluble antioxidant present in cells, as it is the major chain terminating antioxidant in biological membranes (94) and scavenges a wide array of ROS including 1O_2 , HO^\cdot , O_2^\cdot , peroxy, and alkoxyl radicals. Vitamin E is composed of a number of derivatives of tocopherols and tocotrienols. The major isomer in humans is α -TOH, which also possesses the greatest antioxidant activity of any vitamin E derivative. In homogeneous solutions, α -TOH is a strong inhibitor of polyunsaturated lipid peroxidation (52) and *in vivo*, most of the cellular vitamin E is concentrated in the lipid membranes (95,96). The primary antioxidant activity of tocopherols is to stop chain propagation of peroxy radicals ($k = 6 \times 10^3 - 3 \times 10^6 M^{-1} sec^{-1}$) (97). Tocopherols can typically scavenge two ROS per tocopherol molecule (98).

Other lipid soluble antioxidants contribute to the protection of lipid bilayers from oxidative stress. Ubiquinol Q_{10} , which plays an important role in mitochondrial electron transport (99), acts as a potent antioxidant in lipoproteins (92) and other lipid membranes (100). Ubiquinols are redox active quinone derivatives with a hydrophobic isoprenyl tail. They can rapidly react with oxygen, peroxy, and alkoxyl radicals, preventing initiation of biomolecule damage. Ubiquinol can also

directly interact with α -TO $^\cdot$, regenerating active α -TOH (101). Carotenoids are also lipid soluble antioxidants that are present in lipid membranes. Two of the main carotenoids, α - and β -carotene, serve as provitamin A analogues and are effective antioxidants as 1O_2 quenchers and peroxy radical scavengers (102).

Ascorbic acid (vitamin C), a water soluble antioxidant, exists primarily as ascorbate at physiological pH. Ascorbate is a powerful reducing agent capable of rapidly scavenging a number of ROS including O_2^\cdot ($k = 2.7 \times 10^5 M^{-1} sec^{-1}$) (103,104). In addition, ascorbate reacts with other cellular prooxidants such as 1O_2 , hypochlorous acid (HOCl), and thiol radicals (RS^\cdot) (103,105). Ascorbate is also able to reduce the vitamin E derived α -TO $^\cdot$. Through this mechanism, ascorbate in the aqueous phase is able to regenerate membrane-bound α -TOH, prolonging the lifetime of this important antioxidant in the lipid phase and effecting removal of the radical from the lipid to the aqueous phase. In its action as an antioxidant, ascorbate is rapidly oxidized to dehydroascorbate, which shows little antioxidant activity. Dehydroascorbate may be reduced back to ascorbate via a GSH-dependent reductase (106).

In addition to acting as a reducing agent, GSH also provides antioxidant protection in the aqueous phase of cellular systems. GSH is synthesized in several different cell types and is present intracellularly in millimolar concentrations, making it the most abundant thiol with levels up to 10 mM (90). The cysteine thiol moiety of GSH imparts the antioxidant activity of this molecule. Like ascorbate, GSH can directly reduce a number reactive oxygen species including 1O_2 , HO^\cdot , and O_2^\cdot , and GSH is oxidized to GSSG in this process. Relatively high ratios of GSH/GSSG are maintained intracellularly through the action of GR in an NADPH-dependent reaction (90,91). As mentioned previously, GSH also acts as a substrate or cosubstrate in many essential enzymatic reactions, such as with the antioxidant enzyme GPx. Thus, the depletion of GSH during oxidative stress could have a significant impact on the antioxidant poise within a cell.

It should be noted that antioxidants also have the potential to act as prooxidants under certain conditions. For example, ascorbate, in the presence of a high concentration of ferric iron (Fe^{3+}), is a potent potentiator of lipid peroxidation. This behavior results from the reduction of Fe^{3+} to Fe^{2+} by the strong reducing agent ascorbate (107), enabling redox cycling of the iron. Other antioxidants may behave similarly; Okada et al. (108) demonstrated the physiologic relevance of such reactions in the proximal tubule of ddY mice exposed

to Fe^{3+} -nitrilotriacetate. The release of cysteine during the hydrolysis of GSH in the tubule, followed by the reduction of Fe^{3+} by this cysteine, appears to lead to lipid peroxidation in the proximal tubule.

Piscine Systems as Models for Oxidative Stress Studies

Typically, mammalian species have been used as models to study oxidative stress and to elucidate the mechanisms behind cellular damage and response, largely because of the interest in human health issues surrounding oxidative stress. However, the aquatic environment provides a sink for many environmental contaminants that have the potential to cause oxidative stress in aquatic organisms. Some common pollutants in this category include polyaromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs), quinones, aromatic nitro compounds, aromatic hydroxylamines, bipyridyls, and transition metals (14). These compounds may cause adverse effects in aquatic organisms (e.g., lipid and DNA damage). Thus, a better understanding of oxidative stress in aquatic organisms is essential to understanding the impacts of these pollutants on aquatic systems.

However, with the growing interest in oxidative toxicity, the potential benefits of using fish as model organisms to study oxidative stress reach beyond applications to aquatic ecotoxicology. The similarities in the types of oxidative damage and defenses between mammalian and piscine systems suggest that fish can provide model systems for further understanding how ROS damage cellular components, how cells respond to and repair this damage, and how oxidative damage can lead to disease. This type of information could lead to the development of more sensitive markers for exposure and toxicity, and these markers may prove invaluable for evaluating subtle effects of oxidative stress, such as genotoxicity (35), in both mammalian and aquatic systems. Additionally, fish may provide an alternative system to rodents for evaluating the link between exposure and effect for diseases related to oxidative toxicity. For example, in some populations of aquatic organisms, the link between neoplasia and environmental pollution is more obvious than that in mammalian studies (109). Combined with other benefits, such as the potential for decreased cost, the ability to directly observe embryonic development in some fish species, and the advent of transgenic fish, piscine systems have the potential to provide suitable and beneficial models for studying oxidative stress in vertebrate systems and may help to bridge the gap between ecological and human health issues in oxidative toxicology.

Lipid peroxidation and DNA damage as indicators of oxidative stress. In both mammalian and piscine systems, oxidative stress causes toxic and adaptive responses within a cell (14). Toxic responses include damage to cellular components such as lipid peroxidation and DNA damage, as described earlier. In mammalian systems, transition metals are a classic source of radical generation and subsequent cellular damage. Lipid peroxidation has been used as a measure of this xenobiotic-induced oxidative stress. For example, iron and/or copper have been shown to induce lipid peroxidation in rat hepatocytes (110), rat testes (111), macrophages (112), and rat liver mitochondria (113).

Lipid peroxidation has also been used as an indicator of oxidative damage in fish. In a study of Atlantic croaker (*Micropogonias undulatus*), cadmium exposure resulted in an increase in MDA levels in homogenates of hepatic and ovarian tissues and in microsomes. Similarly, Aroclor 1254 caused an increase in MDA in croaker liver *in vivo* and *in vitro*, as well as in ovarian homogenates (32). Parihar and Dubey (114) found that freshwater Indian catfish (*Heteropneustes fossilis*) exposed to a temperature increase (from 25 to 37°C) exhibited significant increases in MDA levels in gill and air sac tissues. A different study in channel catfish (*Ictalurus punctatus*) evaluated lipid peroxidation following exposure to sediments contaminated with aromatic hydrocarbons. At all time points evaluated, the concentration of MDA in hepatic tissue was significantly elevated in fish exposed to contaminated sediments as compared to fish exposed to noncontaminated sediments (115).

Oxidative damage to DNA is another consequence of oxidative stress that has been used as a marker of this phenomenon. There are numerous recent reports in the literature indicating that iron exposure leads to DNA damage via oxidative mechanisms in mammalian systems. Examples of such studies include DNA-protein cross-links in a murine cell line (116), as well as 8-OHdG in the rat testis (111) and human pulmonary epithelia after exposure to asbestos (117). The iron-associating properties of asbestos are also related to double-strand DNA breaks in pulmonary epithelia (118), and double-strand breaks caused by iron complexed with certain chelators, such as 8-hydroxyquinolone, have been demonstrated in human lung (119). Similarly, copper causes elevated levels of 8-OHdG in rat liver and kidney (120), DNA-protein cross-linking in human lymphocytes (121), and increased levels of 8-OHdG in murine keratinocytes exposed to copper and benzoyl peroxide (122).

As with lipid peroxidation, oxidative damage to DNA has received increasing attention in aquatic organisms in recent years. For example, juvenile English sole (*Parophrys vetulus*) treated with nitrofurantoin, a known redox cycling xenobiotic, demonstrated an increase in hepatic 8-OHdG (123). Another DNA lesion linked to oxidative stress and recently found in fish is FapyGua. Malins et al. (124) found the FapyGua lesion in the DNA of neoplastic hepatic tissues of feral fish (English sole) but not in the nonneoplastic tissues. A similar study of feral English sole also found that fish from sites contaminated with PAHs and PCBs showed a high degree of hepatic DNA damage in the form of 8-OH adducts of guanine and adenine as well as FapyGua (125). These studies on lipid peroxidation and oxidative DNA damage demonstrate that mammals and fish exhibit similar lesions and that the markers used to detect toxic consequences of oxidative stress in mammalian systems are also useful in piscine models.

Cellular defenses. Fish exhibit many of the same defenses against oxidative stress as do mammals, suggesting that fish and mammalian systems have similar cellular responses to oxidative stress. These defenses include both low molecular weight free radical scavengers such as GSH, α -TOH, and ascorbic acid, as well as enzymatic defenses such as SOD, catalase, and GPx (35,126). Many studies in the aquatic literature provide baseline information for comparative purposes and have been reviewed by Winston and Di Giulio (35). It is clear from these studies that there is a significant degree of variation in the basal activities of antioxidant defenses between fish species, and much of this work has focused on enzymatic defenses. In comparison to other vertebrate systems, fish appear to exhibit lower basal activities of SOD and catalase, but have increased activities of GPx (127-131) and this should be taken into consideration when designing studies with fish.

Several studies in mammals and fish have demonstrated the importance of antioxidant defenses in protecting cells and organisms from oxidative damage and toxicity. For instance, tocopherol diminished lipid peroxidation in rat hepatocytes treated with iron and copper (110) and in rat lung and liver treated with iron (132), while both tocopherol and β -carotene were effective at reducing hepatic lipid peroxidation in rats fed high iron diets (133). In aquatic systems, studies have centered more on the effects of depleting antioxidant defenses rather than the effects of supplementation as seen in mammalian studies. For instance, a study of blue gill sunfish BF-2 fibroblasts

showed that depletion of GSH by agents such as buthionine sulfoximine, maleic acid, or 1-chloro-2,4-dinitrobenzene increased the toxicity of H_2O_2 and nitrofurantoin to fibroblasts. Furthermore, inhibition of SOD with diethylthiocarbamate also enhanced the toxicity of H_2O_2 and paraquat (134). In freshwater Indian catfish, a decline in levels of ascorbic acid following temperature stress was correlated with an increase in lipid peroxidation in gill and air sac membranes, suggesting that ascorbic acid plays a protective role in preventing lipid peroxidation (114), perhaps through regeneration of tocopherol. Williams et al. (135) examined the importance of vitamin E in protecting rainbow trout (*Oncorhynchus mykiss*) from lipid peroxidation. Feeding trout a vitamin E-deficient diet resulted in an 18-fold decrease in liver microsomal levels of vitamin E and a significant increase in susceptibility to Fe^{3+} -ascorbate dependent lipid peroxidation (measured as MDA) in liver microsomes. Additionally, vitamin E depletion also enhanced the amount of lipid hydroperoxides generated *in vivo* as seen in extracts of microsomes and plasma (135).

In addition to providing protection in both mammalian and piscine systems, some of the low molecular weight ROS scavengers and enzymatic antioxidant defenses are responsive to challenge with compounds that cause oxidative stress (i.e., they are induced), representing an adaptive consequence of exposure. In mammals, numerous studies have demonstrated that antioxidant enzymes are up regulated, particularly in response to toxicant-induced inflammation. The most studied of these enzymes is Mn-SOD, the activity and transcription of which is increased following exposure to toxins such as asbestos (136) and lipopolysaccharide (137), as well as inflammatory cytokines such as TNF- α (136). Similarly, phagocytosis of particulate matter leads to increased SOD activity in macrophages (138). Cu/Zn-SOD, EC-SOD, and glutathione-related enzymes are less commonly up regulated following toxic exposures, but lipopolysaccharide has been shown to induce transcription of both GPx and Cu/Zn-SOD (139), while toxicants that induce the inflammatory cytokine IFN- γ may up regulate transcription and secretion of EC-SOD (140,141).

Similar responses have been reported in aquatic species, and these may serve as biomarkers of exposure for certain compounds in aquatic organisms (14). For instance, Pedrajas et al. (142) found that sea bream (*Sparus aurata*) treated with copper chloride demonstrated increased SOD activity. Otto and Moon (143) found that PCBs altered

antioxidant defenses in trout by demonstrating that trout treated with 3,3',4,4'-tetrachlorobiphenyl showed increased GPx, GR, SOD, and catalase activities as well as increases in GSH and GSSG.

Notably, sometimes reductions are seen in the levels of antioxidants with or without an increase in others. For example, in a study of channel catfish exposed to bleached kraft mill effluent, the fish exhibited increased hepatic catalase activity and decreased hepatic GSH (33). Similarly, Thomas and Wofford (32) showed that Atlantic croaker exposed to cadmium exhibited a decrease in GPx activity and ascorbic acid content in hepatic and ovarian tissues while croaker in the same study exposed to Aroclor 1254 showed an increase in GPx activity in hepatic and ovarian tissues without any change in ascorbic acid content of either tissue. While not necessarily an adaptive response, a decrease in levels of antioxidant defenses is biologically significant and can also serve as a marker of oxidative stress. Combined with the presence of antioxidant inductions, it may be possible to develop a response profile indicative of some xenobiotics.

Field studies. In addition to laboratory studies, field studies are common for evaluating the impacts of oxidative stress in natural populations. Measurements of lipid and DNA damage combined with changes in the levels of antioxidant defenses can serve as a means of detecting exposure to xenobiotics that induce oxidative stress. However, the choice of end points in field studies of fish may be complicated by a history of exposure to xenobiotics causing oxidative stress. For instance, grey mullet (*Mugil* sp.) collected from an estuary polluted with metals, PAHs, PCBs, and pesticides demonstrated evidence of oxidative stress as indicated by oxidized glutathione redox status. However, these fish did not show elevated MDA levels while showing elevated activities of antioxidant enzymes (GPx, SOD, catalase, GR). It is possible that an adaptive response had occurred and that repair of lipid peroxidation may have taken place (144).

Because the response of the organism may be xenobiotic and tissue specific, it is also important to examine several end points. Otto and Moon (145) compared brown bullhead (*Ameiurus nebulosus*) that were collected from a system polluted with PCBs to bullhead collected from a relatively nonpolluted aquatic system. Fish from the polluted site had a 22-fold increase in PCB concentrations in white muscle compared to fish from the nonpolluted site. Cytosolic SOD activity was increased in the kidney of fish from the polluted site. Conversely, catalase activity in the kidney,

GPx activity in the red and white muscle, and total glutathione in the liver, kidney, and white muscle were decreased relative to fish from the nonpolluted site. Studies such as these advocate the use of a group of end points when evaluating oxidative stress in feral organisms.

Oxidative stress and disease. The link between oxidative stress and disease is also of interest to oxidative stress toxicology. Cancer is one disease for which fish may be especially well suited to serve as model organisms, and the appeal of using fish as models for studying the molecular basis of carcinogenesis is growing. The benefits of using rainbow trout in addition to rodent models when evaluating environmental carcinogenesis have been discussed by Bailey et. al. (146,147). Some of these benefits include low rearing costs that enable quantitative studies with as many as 10,000 fish as well as well-described tumor pathology. *Xiphophorus* fish models have been used to study the roles of genetics and UV radiation in melanoma tumorigenesis (148–151). Studies of oncogenes and tumor suppressor genes have also been conducted in fish. For instance, two *ras* genes are expressed in rainbow trout liver, and both contain a high degree of homology to the human *ras* genes. The predicted amino acid sequence of trout *ras-1* suggests a highly conserved protein and suggests that the function of the *ras* p21 protein is the same in both higher and lower vertebrates (152). Similarly, the tumor suppressor gene *p53* has been cloned in Japanese medaka (*Oryzias latipes*) (153) and rainbow trout (154) and shown to be strongly conserved in functional domains critical to tumor suppressor function in other vertebrates, including humans. Furthermore, Winn et. al. (155) have created transgenic medaka and mummichog (*Fundulus heteroclitus*) that contain multiple copies of the bacteriophage Phi X174am3cs70 that will enable comparisons of mutations in the same target gene sequence across species following xenobiotic exposure.

Studies that pertain specifically to the role of oxidative stress in cancer have also been conducted in fish. For instance, comparative studies between fish species have been done that evaluate changes in antioxidant defenses that may be linked to disease etiology. Channel catfish, which seldom express pollutant-mediated neoplasms, and brown bullhead, which often exhibit neoplasms in contaminated waters, differ greatly in their glutathione-dependent defense systems. Channel catfish have higher levels of hepatic GSH and γ -glutamylcysteine synthase activity, while bullhead have more hepatic GSSG and increased GPx activity (34). In a comparative study with menadione (a quinone) as the toxicant, channel

catfish exhibited a more rapid induction of total glutathione than did bullhead, likely due to the increased hepatic γ -glutamylcysteine synthase activity. Furthermore, when challenged with an organic peroxide (*tert*-butyl hydroperoxide), hepatic GSH was depleted in bullhead but not in channel catfish. This difference in GSH depletion was attributed to the differences in glutathione peroxidase and glutathione reductase activities between species (34). A different study of the same two fish species exposed to β -naphthoflavone showed that catfish exhibited significantly higher hepatic catalase and GR activities as well as total glutathione content, whereas bullhead exhibited higher hepatic GPx activity. Catfish showed higher levels of menadione-mediated cytosolic $\cdot\text{O}_2^-$ and H_2O_2 production, whereas bullhead demonstrated higher rates of $\cdot\text{O}_2^-$ and H_2O_2 production in microsomes. Liver homogenate from bullhead was also less effective at inhibiting Fe^{3+} -ascorbate-mediated production of 8-OHdG (156). The results from these two studies suggest that differences in antioxidant poise may result in differential abilities to defend against certain types of oxidative stress and may account for the difference in susceptibility to cancer between these two species (34,156).

Other studies have linked oxidative DNA damage to neoplasms in fish. As mentioned previously, the FapyGua lesion has been found in the DNA of neoplastic hepatic lesions in English sole (124), and hepatic DNA damage, in the form of both 8-OH adducts of guanine and adenine as well as FapyGua, has been reported in a population of English sole contaminated with PAHs and PCBs that also show a high incidence of liver cancer (125). The comparative studies of antioxidant poise in catfish, the studies on DNA damage in English sole, the demonstration of conserved oncogenes and tumor suppressor genes between mammals and fish, and advances in the use of fish as models for carcinogenicity suggest that markers of oxidative stress may be predictive of cancer risk in some instances and that fish may be useful in further elucidating the mechanisms of carcinogenesis.

Conclusions

Oxidative stress comprises an important aspect of toxicology and, consequently, has received increasing attention in recent years. During this time, research has moved from observing the effects of oxidative stress at the level of the organism to elucidating the mechanisms behind the responses and damages seen at the cellular and biochemical levels. Much of the research on oxidative stress has taken place in mammalian models as the

interest in oxidative stress has centered primarily around human health issues. However, increasing evidence suggests that oxidative stress is also of ecological significance, particularly in the aquatic environment which provides a sink for many pollutants capable of causing oxidative stress. Comparing the data available for both mammalian and piscine systems demonstrates that the toxic and adaptive mechanisms of oxidative stress are similar across species. This suggests that fish may serve as good biomonitoring tools, and, in addition to mammals, may provide useful models for further research into understanding oxidative stress.

There are several advantages to using piscine systems to study oxidative stress, such as the reduction in the number of mammals used in research, the potential for a reduction in the cost of animal maintenance, and the ability to increase the power of the experiment by increasing the number of organisms per study. Additionally, the developmental biology of some fish species, such as zebra fish (*Brachydanio rerio*) and mummichog, have been well studied (157–159). These species provide excellent models for studying the developmental aspects of oxidative stress toxicology, as the growing embryo can be observed throughout development. Moreover, most fish species are oviparous, enabling the evaluation of development in isolation as fertilization and development occur externally. With the recent advent of transgenic fish (160–162) comes the potential to perform knock-out experiments and evaluate how oxidative stress may influence gene expression in piscine models. Furthermore, the use of piscine models to study oxidative stress allows for the evaluation of environmental issues from both a human health and ecological perspective.


Many research opportunities still exist in oxidative stress toxicology and present the occasion to study oxidative stress in piscine systems. For example, little is known specifically about the repair processes that are utilized by cells to repair damage induced by oxidative stress. In addition to aiding in the understanding of how cells cope with oxidative damage, these processes may involve inducible enzymes or other responses that could serve as biomarkers of exposure or effect. Similarly, in light of evidence that oxidative stress can damage DNA and gametes, the impact of oxidative stress on processes such as reproduction and development warrants further study. Finally, there is still a need for research that definitively links oxidative stress and disease. Performing these studies in fish will increase our knowledge of oxidative stress mechanisms while expanding the ability to generalize the phenomenon of oxidative stress across species.

REFERENCES AND NOTES

- McCord JM, Fridovich I. Superoxide dismutase. An enzymatic function for erythrocyte (hemocypre). *J Biol Chem* 244:6049–6055 (1969).
- Seis H, ed. *Oxidative Stress: Oxidants and Antioxidants*. London:Academic Press, 1991.
- Kellogg EW, Fridovich I. Liposome oxidation and erythrocyte lysis by enzymatically generated superoxide and hydrogen peroxide. *J Biol Chem* 252:6721–6728 (1977).
- Howard JA. *Advances in Free Radical Chemistry*. New York:Academic Press, 1972.
- Imlay JA, Linn S. DNA damage and oxygen radical toxicity. *Science* 240:1302–1309 (1988).
- von Sonntag C. *The Chemical Basis of Radiation Biology*. New York:Taylor & Francis, 1987.
- Steenken S. Purine bases, nucleosides, and nucleotides: aqueous solution redox chemistry and transformation reactions of their radical cations and e- and OH adducts. *Chem Rev* 89:503–520 (1989).
- Park EM, Thomas JA. S-Thiolation of creatine kinase and glycogen phosphorylase b initiated by partially reduced oxygen species. *Biochim Biophys Acta* 964:151–160 (1988).
- Kono Y, Fridovich I. Superoxide radical inhibits catalase. *J Biol Chem* 257:5751–5754 (1982).
- Stadtman ER, Berlett BS. Reactive oxygen-mediated protein oxidation in aging and disease. *Chem Res Toxicol* 10:485–494 (1997).
- Kappus H. Overview of enzyme systems involved in bio-reduction of drugs and in redox cycling. *Biochem Pharmacol* 35:1–6 (1986).
- Cohen GM, d'Arcy Doherty M. Free radical mediated cell toxicity by redox cycling chemicals. *Br J Cancer* 55:46–52 (1987).
- Berlett BS, Stadtman ER. Protein oxidation in aging, disease, and oxidative stress. *J Biol Chem* 272:20313–20316 (1997).
- Di Giulio RT, Washburn PC, Wenning RJ, Winston GW, Jewell CS. Biochemical responses in aquatic animals: a review of determinants of oxidative stress. *Environ Toxicol Chem* 8:1103–1123 (1989).
- Punchard NA, Kelly FJ, eds. *Free Radicals: A Practical Approach*. Oxford:IRL Press, 1996.
- Halliwell B. Superoxide-dependent formation of hydroxyl radicals in the presence of iron chelates: is it a mechanism for hydroxyl radical production in biochemical systems? *FEBS Lett* 92:321–326 (1978).
- Fridovich I. Superoxide dismutases. *Adv Enzymol Relat Areas Mol Biol* 41:35–97 (1974).
- Fridovich I. The biology of oxygen radicals. *Science* 201:875–880 (1978).
- Xia Y, Dawson VL, Dawson TM, Snyder SH, Zweier JL. Nitric oxide synthase generates superoxide and nitric oxide in arginine-depleted cells leading to peroxynitrite-mediated cellular injury. *Proc Natl Acad Sci USA* 93:6770–6774 (1996).
- Baud L, Hagege J, Sraer J, Rondeau E, Perez J, Ardaillou R. Reactive oxygen production by cultured rat glomerular mesangial cells during phagocytosis is associated with stimulation of lipoxygenase activity. *J Exp Med* 158:1836–1852 (1983).
- Leroyer V, Werner L, Shaughnessy S, Goddard GJ, Orr FW. Chemiluminescence and oxygen radical generation by Walker carcinosarcoma cells following chemotactic stimulation. *Cancer Res* 47:4771–4775 (1987).
- Washburn PC, Di Giulio RT. Nitrofurantoin-stimulated superoxide production by channel catfish (*Ictalurus punctatus*) hepatic microsomal and soluble fractions. *Toxicol Appl Pharmacol* 95:363–377 (1988).
- Halliwell B, Gutteridge JMC. Role of free radicals and catalytic metal ions in human disease: an overview. In: *Oxygen Radicals in Biological Systems, Part B: Oxygen Radicals and Antioxidants*, Vol 186 (Packer L, Glazer A, eds). San Diego, CA:Academic Press, 1990:1–85.
- Roos D, Bolscher B, DeBoer M. Generation of reactive oxygen species by phagocytes. In: *Mononuclear Phagocytes* (van Furth R, ed). Dordrecht, Netherlands:Kluwer Academic Publishers, 1992:243–253.
- Bruhwyler J, Chleide E, Liegeois JF, Carreer F. Nitric oxide: a new messenger in the brain. *Neurosci Biobehav Rev* 17:373–384 (1993).
- Remacle J, Raes M, Toussaint O, Renard P, Rao G. Low levels of reactive oxygen species as modulators of cell function. *Mutat Res* 316:103–122 (1995).
- McCord JM. Superoxide radical: controversies, contradictions, and paradoxes. *Proc Soc Exp Biol Med* 209:112–117 (1995).
- Irani K, Xia Y, Zweier L, Sollot SJ, Der CJ, Fearon ER, Sundaresan M, Finkel T, Goldschmidt-Clearmont PJ. Mitogenic signaling mediated by oxidants in Ras-transformed fibroblasts. *Science* 275:1649–1652 (1997).
- Nohl H, Hegner D. Identification of free hydroxyl radicals in respiring heart mitochondria by spin trapping with nitron DPMO. *FEBS Lett* 123:241–244 (1981).
- Odajima T, Yamazaki I. Myeloperoxidase of the leukocyte of normal blood. III. The reaction of ferric myeloperoxidase with superoxide anion. *Biochim Biophys Acta* 284:355–359 (1972).
- Kono Y, Fridovich I. Inhibition and reactivation of Mn-catalase: implications for valence changes at the active site manganese. *J Biol Chem* 258:13646–13648 (1983).
- Thomas P, Wofford HW. Effects of cadmium and Aroclor 1254 on lipid peroxidation, glutathione peroxidase activity, and selected antioxidants in Atlantic croaker tissues. *Aquat Toxicol* 27:159–178 (1993).
- Mather-Mihaich E, Di Giulio RT. Oxidant, mixed-function oxidase, and peroxisomal responses in channel catfish exposed to bleached kraft mill effluent. *Arch Environ Contam Toxicol* 20:391–397 (1991).
- Hasspieler BM, Behar JV, Di Giulio RT. Glutathione-dependent defense in channel catfish (*Ictalurus punctatus*) and brown bullhead (*Ameriurus nebulosus*). *Ecotoxicol Environ Saf* 28:82–90 (1994).
- Winston GW, Di Giulio RT. Prooxidant and antioxidant mechanisms in aquatic organisms. *Aquat Toxicol* 19:137–161 (1991).
- Bachur NR, Gordon SL, Gee MV. A general mechanism for microsomal activation of quinone anticancer agents to free radicals. *Cancer Res* 38:1745–1750 (1978).
- Kappus H, Sies H. Toxic drug effects associated with oxygen metabolism, redox cycling and lipid peroxidation. *Experientia* 37:1233–1241 (1981).
- Thor H, Smith MT, Hartzell P, Bellomo G, Jewell SA, Orrenius S. The metabolism of menadione (2-methyl-1,4-naphthoquinone) by isolated hepatocytes. *J Biol Chem* 257:12419–12425 (1982).
- Fridovich I. Superoxide dismutases: regularities and irregularities. *Harvey Lect* 79:51–75 (1983).
- Mashino T, Fridovich I. Superoxide radical initiates the autoxidation of dihydroxyacetone. *Arch Biochem Biophys* 254:547–554 (1987).
- Fridovich I. Superoxide radical and superoxide dismutases. *Annu Rev Biochem* 64:97–112 (1995).
- Pryor WA. Oxy-radicals and related species: their formation, lifetimes and reactions. *Annu Rev Physiol* 48:657–667 (1986).
- Haber F, Weiss J. The catalytic decomposition of hydrogen peroxide by iron salts. *Proc R Soc Lond* 147:332–351 (1934).
- Walling C. Fenton's reagent revisited. *Accounts Chem Res* 8:125–131 (1975).
- Mello Filho AC, Meneghini R. *In vivo* formation of single-strand breaks in DNA by hydrogen peroxide is mediated by the Haber-Weiss reaction. *Biochim Biophys Acta* 781:56–63 (1984).
- Korbashi P, Kohen R, Katchendler J, Chevon M. Iron mediates paraquat toxicity in *Escherichia coli*. *J Biol Chem* 261:12472–12476 (1986).
- Fridovich I. Superoxide dismutases. *J Biol Chem* 264:7761–7764 (1989).
- Davies AG, Griller D, Ingold KU, Lindsay DA, Walton JC. An electron spin resonance study of pentadienyl and related radicals: homolytic fission of cyclobut-2-enylmethyl radicals. *J Chem Soc Perkin Trans II*:633–641 (1981).
- Nagaoka S, Okauchi Y, Urano S, Nagashima U, Mukai K. Kinetic and *ab initio* study of the prooxidant effect of vitamin E: hydrogen abstraction from fatty acid esters and egg yolk lecithin. *J Am Chem Soc* 112:8921–8924 (1990).
- Halliwell B, Gutteridge JMC. The antioxidants of human extracellular fluids. *Arch Biochem Biophys* 280:1–8 (1990).

51. Samokyszyn VM, Miller DM, Reif DW, Aust SD. Iron-catalyzed lipid peroxidation. In: *Membrane Lipid Oxidation*, Vol 1 (Vigo-Pelfrey C, ed). Boca Raton, FL: CRC Press, 1990;101-127.
52. Burton GW, Ingold KU. Vitamin E: application of the principles of physical organic chemistry to the exploration of its structure and function. *Acc Chem Res* 19:194-201 (1986).
53. Waters WA. Kinetics and mechanism of metal catalyzed autooxidation. *J Am Oil Chem Soc* 48:427-433 (1976).
54. O'Brien PJ. Intracellular mechanisms for the decomposition of a lipid peroxide. I. Decomposition of a lipid hydroperoxide by metal ions, heme compounds, and nucleophiles. *Can J Biochem* 47:485-492 (1969).
55. Porter NA. Mechanisms for the autooxidation of polyunsaturated lipids. *Acc Chem Res* 19:262-268 (1986).
56. Porter NA. Autooxidation of polyunsaturated fatty acids: initiation, propagation, and product distribution (Basic chemistry). In: *Membrane Lipid Oxidation*, Vol 1 (Vigo-Pelfrey C, ed). Boca Raton, FL: CRC Press, 1990;33-62.
57. Porter NA, Caldwell SE, Mills KA. Mechanisms of free radical oxidation of unsaturated lipids. *Lipids* 30:277-290 (1995).
58. Ursini F, Maiorino M, Sevanian A. Membrane hydroperoxides. In: *Oxidative Stress: Oxidants and Antioxidants* (Sies H, ed). London: Harcourt Brace Jovanovich, 1991;319-336.
59. Kagan VE. Molecular mechanisms of biomembrane damage caused by LPD. In: *Lipid Peroxidation in Biomembranes* (Kagan VE, ed). Boca Raton, FL: CRC Press, 1988;55-117.
60. Tappel AL. Measurement of and protection from *in vivo* lipid peroxidation. In: *Free Radicals in Biology*, Vol 4 (Pryor WA, ed). New York: Academic Press, 1980;1-47.
61. Ursini F, Maiorino M, Gregolin C. The selenoenzyme phospholipid hydroperoxide glutathione peroxidase. *Biochim Biophys Acta* 839:62-70 (1985).
62. Sevanian A, Stein RA, Mead JF. Metabolism of epoxidized phosphatidylcholine by phospholipase A₂ and epoxide hydrolase. *Lipids* 16:781-789 (1981).
63. Van Kuijk FJGM, Sevanian A, Handelman GJ, Dratz EA. A new role for phospholipase A₂: protection of membranes from lipid peroxidation damage. *Trends Biochem Sci* 12:31-34 (1987).
64. Sevanian A, Kim E. Phospholipase A₂ dependent release of fatty acids from peroxidized membranes. *J Free Radical Biol Med* 1:263-271 (1985).
65. Salgo MG, Corongiu FP, Sevanian A. Enhanced interfacial catalysis and hydrolytic specificity of phospholipase A₂ toward peroxidized phosphatidylcholine vesicles. *Arch Biochem Biophys* 304:123-132 (1993).
66. Sevanian A, Muakkassah-Kelly SF, Montestruque S. The influence of phospholipase A₂ and glutathione peroxidase on the elimination of membrane lipid peroxides. *Arch Biochem Biophys* 223:441-452 (1983).
67. Ursini F, Maiorino M, Valente M, Ferri L, Gregolin C. Purification from pig liver of a protein which protects liposomes and biomembranes from peroxidative degradation and exhibits glutathione peroxidase activity on phosphatidylcholine hydroperoxides. *Biochim Biophys Acta* 710:197-211 (1982).
68. Esterbauer H, Zollner H, Schaur RJ. Aldehydes formed by lipid peroxidation: mechanisms of formation, occurrence, and determination. In: *Membrane Lipid Oxidation*, Vol 1 (Vigo-Pelfrey C, ed). Boca Raton, FL: CRC Press, 1990;239-268.
69. Comporti M. Biology of disease: lipid peroxidation and cellular damage in toxic liver injury. *Lab Invest* 53:599-623 (1985).
70. Esterbauer H. Aldehydic products of lipid peroxidation. In: *Free Radicals, Lipid Peroxidation and Cancer* (McBrien DCH, Slater TF, eds). London: Academic Press, 1982;101.
71. Esterbauer H. Lipid peroxidation products: formation, chemical properties and biological activities. In: *Free Radicals in Liver Injury* (Poli G, Cheeseman KH, Dianzani MU, Slater TF, eds). Oxford: IRL Press, 1985;29.
72. Kohn H, Liversedge M. On a new aerobic metabolite whose production by brain is inhibited by apomorphine, emetine, ergotamine, epinephrine, and menadione. *J Pharmacol Exp Ther* 82:292-300 (1944).
73. Kosugi H, Kato T, Kikugawa K. Formation of yellow, orange, and red pigments in the reaction of alk-2-enals with 2-thiobarbituric acid. *Anal Biochem* 165:456-464 (1987).
74. Shimizu T, Kondo K, Hayaishi O. Role of prostaglandin endoperoxides in the serum thiobarbituric acid reaction. *Arch Biochem Biophys* 206:271-276 (1981).
75. Janero DR. Malondialdehyde and thiobarbituric acid-reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury. *Free Radic Biol Med* 9:515-540 (1990).
76. Marshall PJ, Warso MA, Lands WEM. Selective microdetermination of lipid hydroperoxides. *Anal Biochem* 145:192-199 (1985).
77. Largilliere C, Melancon SB. Free malondialdehyde determination in human plasma by high-performance liquid chromatography. *Anal Biochem* 170:123-126 (1988).
78. Gutteridge JMC. Aspects to consider when detecting and measuring lipid peroxidation. *Free Radical Res Commun* 1:173-184 (1986).
79. Friedberg EC, Walker GC, Siede W. *DNA Repair and Mutagenesis*. Washington, DC: ASM Press, 1995.
80. Buxton GV, Greenstock CL, Helman WP, Ross AB. Critical review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals. *Phys Chem Ref Data* 17:513-517 (1988).
81. Demple B, Levin J. Repair systems for radical-damaged DNA. In: *Oxidative Stress: Oxidants and Antioxidants* (Sies H, ed). New York: Academic Press, 1991;119-154.
82. Favier AE. How to demonstrate the occurrence of an oxidative stress in humans? In: *Analysis of Free Radicals in Biological Systems* (Favier AE, Cadet J, Kalyanaram B, Fontecave M, Pierre J-L, eds). Basel, Switzerland: Birkhauser Verlag, 1995;99-117.
83. Oleinick NL, Chiu S, Friedman LR, Xue L, Ramakrishnan N. DNA-protein cross-links: new insights into their formation and repair in irradiated mammalian cells. In: *Mechanisms of DNA Damage and Repair* (Sicim MG, Grossman L, Upton AC, eds). New York: Plenum Press, 1986;181-192.
84. Dizdaroglu M. Oxidative damage to DNA in mammalian chromatin. *Mutat Res* 275:331-342 (1992).
85. Brusick D. *Principles of Genetic Toxicology*. New York: Plenum Press, 1987.
86. Karlsson K, Marklund SL. Binding of human extracellular-superoxide dismutase C to cultured cell lines and to blood cells. *Lab Invest* 60:659-666 (1989).
87. Krinsky NI. Mechanism of action of biological antioxidants. *Proc Soc Exp Biol Med* 200:248-254 (1992).
88. Quinlan T, Spivak S, Mossman BT. Regulation of antioxidant enzymes in lung after oxidant injury. *Environ Health Perspect* 102(suppl 2):79-87 (1994).
89. Kinnula VL, Crapo JD, Ravio KO. Generation and disposal of reactive oxygen metabolites in the lung. *Lab Invest* 73:3-19 (1995).
90. Anderson ME. Glutathione. In: *Free Radicals: A Practical Approach* (Punchar NA, Kelly FJ, eds). Oxford: IRL Press, 1996;213-226.
91. Akerboom TPM, Sies H. Assay of glutathione, glutathione disulfide, and glutathione mixed disulfides in biological samples. In: *Detoxication and Drug Metabolism: Conjugation and Related Systems*, Vol 77 (Jakoby W, ed). New York: Academic Press, 1981;373-382.
92. Stocker R, Frei B. Endogenous antioxidant defenses in human blood plasma. In: *Oxidative Stress: Oxidants and Antioxidants* (Sies H, ed). London: Harcourt Brace Jovanovich, 1991;213-244.
93. Frei B, Stocker R, Ames BN. Small molecule antioxidant defenses in human extracellular fluids. In: *Molecular Biology of Free Radical Scavenging Systems*, Vol 5 (Scandalios JG, ed). Plainview, NY: Cold Spring Harbor Laboratory Press, 1992;23-46.
94. Burton GW, Joyce A, Ingold KU. Is vitamin E the only lipid soluble, chain-breaking antioxidant in human blood plasma and erythrocyte membranes? *Arch Biochem Biophys* 221:281-290 (1983).
95. Kagan VE, Bakalova RA, Serbinova EA, Stoychev ES. Fluorescence measurements of incorporation and hydrolysis of tocopherol esters in biomembranes. In: *Oxygen Radicals in Biological Systems*, Part B: Oxygen Radicals and Antioxidants, Vol 186 (Packer L, Glazer A, eds). San Diego, CA: Academic Press, 1990;355-367.
96. Bjorneboe A, Bjorneboe GE, Drevon CA. Absorption, transport and distribution of vitamin E. *J Nutr* 120:233-242 (1990).
97. Niki E, Matsuo M. Rates and products of reactions of vitamin E with oxygen radicals. In: *Vitamin E in Health and Disease* (Packer L, Fuchs J, eds). New York: Dekker, 1993;121-130.
98. Horswill EC, Howard JA, Ingold KU. Alpha-tocopherol. *Can J Chem* 44:985-991 (1966).
99. Mitchell P. Protonmotive redox mechanism of the cytochrome *b-c1* complex in the respiratory chain: protonmotive ubiquinone cycle. *FEBS Lett* 56:1-6 (1975).
100. Sustray PC, Jayaraman J, Ramasarma T. Distribution of coenzyme Q in rat liver cell fractions. *Nature* 189:577-580 (1961).
101. Landi L, Cabrini L, Fiorentini D, Stefanelli C, Pedulli GF. The antioxidant activity of ubiquinol-3 in homogenous solution and in liposomes. *Chem Phys Lipids* 61:121-130 (1992).
102. Burton GW, Ingold KU. β -Carotene: an unusual type of lipid antioxidant. *Science* 224:569-573 (1984).
103. Halliwell B. How to characterize a biological antioxidant. *Free Radical Res Commun* 9:1-32 (1990).
104. Nishikimi M. Oxidation of ascorbic acid with superoxide anion generated by the xanthine-xanthine oxidase system. *Biochem Biophys Res Commun* 63:463-468 (1975).
105. Anaber M, Neta P. A compilation of specific biomolecular rate constants for the reactions of hydrated electrons, hydrogen atoms, and hydroxyl radicals with inorganic and organic compounds in aqueous solution. *Int J Appl Radiat Isot* 18:495-523 (1967).
106. Sharma MK, Buetner GR. Interaction of vitamin C and vitamin E during free radical stress in plasma: an ESR study. *Free Radical Biol Med* 14:649-653 (1993).
107. Scarpa M, Stevanato R, Viglino P, Rigo A. Superoxide as an active intermediate in the autooxidation of ascorbate by molecular oxygen. *J Biol Chem* 258:6695-6697 (1983).
108. Okada S, Minamiyama Y, Hamazaki S, Toyokuni S, Sotomatsu A. Glutathione cycle dependency of ferric nitrilotriacetate induced lipid peroxidation in mouse proximal renal tubules. *Arch Biochem Biophys* 301:138-142 (1993).
109. Malins DC, McCain BB, Lindahl JT, Meyers MS, Krahn MM, Brown DW, Chan S-L, Roubal WT. Neoplastic and other diseases in fish in relation to toxic chemicals: an overview. *Aquat Toxicol* 11:43-67 (1988).
110. Furono K, Suetsuga T, Sugihara N. Effects of metal ions on lipid peroxidation in cultured rat hepatocytes loaded with alpha-linolenic acid. *J Toxicol Environ Health* 48:121-129 (1996).
111. Lucesoli F, Fraga CG. Oxidative damage to lipids and DNA concurrent with decrease of antioxidants in rat testes after acute iron intoxication. *Arch Biochem Biophys* 316:567-571 (1995).
112. Fuhrman B, Oiknine J, Aviram M. Iron induces lipid peroxidation in cultured macrophages, increases their ability to oxidatively modify LDL, and affects their secretory properties. *Atherosclerosis* 111:65-78 (1994).
113. Reinheckel T, Wiswedel I, Noack H, Augustin W. Electrophoretic evidence for the impairment of complexes of the respiratory chain during iron/ascorbate induced peroxidation in isolated rat liver mitochondria. *Biochim Biophys Acta* 1239:45-50 (1995).
114. Parihar MS, Dubey AK. Lipid peroxidation and ascorbic acid status in respiratory organs of male and female freshwater catfish *Heteropneustes fossilis* exposed to temperature increase. *Comp Biochem Physiol C* 112:309-313 (1995).
115. Di Giulio RT, Habig C, Gallagher EP. Effects of Black Rock Harbor sediments on indices of biotransformation, oxidative stress, and DNA integrity in channel catfish. *Aquat Toxicol* 26:1-22 (1993).
116. Altman SA, Zastawny TH, Randers-Eichhorn L, Cacciuto MA, Akman SA, Dizdaroglu M, Rao G. Formation of DNA-protein cross-links in cultured mammalian cells upon treatment with iron ions. *Free Radical Biol Med* 19:897-902 (1995).

117. Chao CC, Aust AE. Photochemical reduction of ferric iron by chelators results in DNA strand breaks. *Arch Biochem Biophys* 300:544–550 (1993).
118. Kamp DW, Israbian VA, Preussen SE, Zhang CX, Weitzman SA. Asbestos causes DNA strand breaks in cultured pulmonary epithelial cells: role of iron-catalyzed free radicals. *Am J Physiol* 268:L471–480 (1995).
119. Leanderson P, Tagesson C. Iron bound to the lipophilic iron chelator, 8-hydroxyquinoline, causes DNA strand breakage in cultured lung cells. *Carcinogenesis* 17:545–550 (1996).
120. Toyokuni S, Sagripanti JL. Increased 8-hydroxydeoxyguanosine in kidney and liver of rats continuously exposed to copper. *Toxicol Appl Pharmacol* 126:91–97 (1994).
121. Costa M, Zhitkovich A, Gargas M, Paustenbach D, Finley B, Kuykendall J, Billings R, Carlson TJ, Wetterhahn K, Xu J, et al. Interlaboratory validation of a new assay for DNA–protein crosslinks. *Mutat Res* 369:13–21 (1996).
122. King JK, Egner PA, Kensler TW. Generation of DNA base modification following treatment of cultured murine keratinocytes with benzoyl peroxide. *Carcinogenesis* 17:317–320 (1996).
123. Nishimoto M, Roubal WT, Stein JE, Varanasi U. Oxidative DNA damage in tissues of English sole (*Parophrys vetulus*) exposed to nitrofurantoin. *Chem Biol Interact* 80:317–326 (1991).
124. Malins DC, Ostrander GK, Haimanot R, Williams P. A novel DNA lesion in neoplastic livers of feral fish: 2,6-diamino-4-hydroxy-5-formamidopyrimidine. *Carcinogenesis* 11:1045–1047 (1990).
125. Malins DC, Gunselman SJ. Fourier-transform infrared spectroscopy and gas chromatography-mass spectrometry reveal a remarkable degree of structural damage in the DNA of wild fish exposed to toxic chemicals. *Proc Natl Acad Sci USA* 91:13038–13041 (1994).
126. Winston GW. Oxidants and antioxidants in aquatic animals. *Comp Biochem Physiol C* 100:173–176 (1991).
127. Matkovic B, Novak R, Hanh HD, Szabo L, Varga SI, Zalesna G. A comparative study of some more important experimental animal peroxide metabolism enzymes. *Comp Biochem Physiol B* 56:31–34 (1977).
128. Mazeaud F, Maral J, Michelson AM. Distribution of superoxide dismutase and glutathione peroxidase in the carp: erythrocyte manganese SOD. *Biochem Biophys Res Commun* 86:1161–1168 (1979).
129. Wdzięcsak J, Zalesna G, Bartkowiak A, Witas H, Leydo W. Comparative studies on superoxide dismutase, catalase and peroxidase levels in erythrocytes of different fish species. *Comp Biochem Physiol B* 68:357–358 (1981).
130. Tappel ME, Chaudiere J, Tappel AL. Glutathione peroxidase activities of animal tissues. *Comp Biochem Physiol B* 73:945–949 (1982).
131. Wdzięcsak J, Zalesna G, Wujec E, Peres G. Comparative studies on superoxide dismutase, catalase and peroxidase levels in erythrocytes and livers of different freshwater and marine fish species. *Comp Biochem Physiol B* 73:361–365 (1982).
132. Schweich MD, Lison D, Lauwerys R. The role of vitamin E in the susceptibility of rat lung and liver microsomes to iron-stimulated peroxidation. *Environ Res* 70:62–69 (1995).
133. Whittaker P, Wamer WG, Chanderbhan RF, Dunkel VC. Effects of alpha-tocopherol and beta-carotene on hepatic lipid peroxidation and blood lipids in rats with dietary iron overload. *Nutr Cancer* 25:119–128 (1996).
134. Babich H, Palace MR, Stern A. Oxidative stress in fish cells: *in vitro* studies. *Arch Environ Contam Toxicol* 24:173–178 (1993).
135. Williams DE, Carpenter HM, Buhler DR, Kelly JD, Dutchuk M. Alterations in lipid peroxidation, antioxidant enzymes, and carcinogen metabolism in liver microsomes of vitamin E-deficient trout and rat. *Toxicol Appl Pharmacol* 116:78–84 (1992).
136. Pietarinen-Runtti P, Raivio KO, Linnainmaa K, Ekman A, Saksela M, Kinnula VL. Differential effects of tumor necrosis factor and asbestos fibers on manganese superoxide dismutase induction and oxidant-induced cytotoxicity in human mesothelial cells. *Cell Biol Toxicol* 12:167–175 (1996).
137. Mitchell J, Jiang H, Berry L, Meyrick B. Effect of antioxidants on lipopolysaccharide-stimulated induction of manganese superoxide dismutase mRNA in bovine pulmonary artery endothelial cells. *J Cell Physiol* 169:333–340 (1996).
138. Kantengwa S, Polla BS. Phagocytosis of *Staphylococcus aureus* induces a selective stress response in human monocytes-macrophages (Mφ): modulation by Mφ differentiation and by iron. *Infect Immun* 61:1281–1287 (1993).
139. Spolarics Z. Endotoxin stimulates gene expression of ROS-eliminating pathways in rat hepatic endothelial and Kupffer cells. *Am J Physiol* 270:G660–G666 (1996).
140. Marklund SL. Regulation by cytokines of extracellular superoxide dismutase and other superoxide dismutase isoenzymes in fibroblasts. *J Biol Chem* 267:6696–6701 (1992).
141. Brady TC, Chang LY, Day BJ, Crapo JD. Extracellular superoxide dismutase is upregulated with inducible nitric oxide synthase following NF-κB activation. *Am J Physiol* 273:L1002–L1006 (1997).
142. Pedrajas JR, Peinado J, Lopez-Barea J. Oxidative stress in fish exposed to model xenobiotics: oxidatively modified forms of Cu,Zn-superoxide dismutase as potential biomarkers. *Chem Biol Interact* 98:267–282 (1995).
143. Otto DME, Moon TW. 3,3',4,4'-Tetrachlorobiphenyl effects on antioxidant enzymes and glutathione status in different tissues of rainbow trout. *Pharmacol Toxicol* 77:281–287 (1995).
144. Rodriguez-Ariza A, Peinado J, Pueyo C, Lopez-Barea J. Biochemical indicators of oxidative stress in fish from polluted littoral areas. *Can J Fish Aquat Sci* 50:2568–2573 (1993).
145. Otto DME, Moon TW. Phase I and II enzymes and antioxidant responses in different tissues of brown bullheads from relatively polluted and non-polluted systems. *Arch Environ Contam Toxicol* 31:141–147 (1996).
146. Bailey GS, Williams DE, Hendricks JD. Fish models for environmental carcinogenesis: the rainbow trout. *Environ Health Perspect* 104(suppl 1):5–21 (1996).
147. Bailey G, Hendricks J, Dashwood R. Anticarcinogenesis in fish. *Mutat Res* 267:243–250 (1992).
148. Dooley TP. Recent advances in cutaneous melanoma oncogenesis research. *Oncol Res* 6:1–9 (1994).
149. Maueler W, Schartl A, Schartl M. Different expression patterns of oncogenes and proto-oncogenes in hereditary and carcinogen-induced tumors of *Xiphophorus*. *Int J Cancer* 55:288–296 (1993).
150. Setlow RB, Woodhead AD, Grist E. Animal model for ultraviolet radiation-induced melanoma: platyfish–swordtail hybrid. *Proc Natl Acad Sci USA* 86:8922–8926 (1989).
151. Setlow RB, Grist E, Thompson K, Woodhead AD. Wavelengths effective in induction of malignant melanoma. *Proc Natl Acad Sci USA* 90:6666–6670 (1993).
152. Mangold K, Chang Y-J, Mathews C, Marien K, Hendricks J, Bailey G. Expression of *ras* genes in rainbow trout liver. *Mol Carcinog* 4:97–102 (1991).
153. Krause MK, Rhodes LD, Van Beneden RJ. Cloning of the *p53* tumor suppressor gene from the Japanese medaka (*Oryzias latipes*) and evaluation of mutational hotspots in MNNG-exposed fish. *Gene* 189:101–106 (1997).
154. Caron de Fromental C, Pakdel F, Chapus A, Baney C, May P, Soussi T. Rainbow trout *p53*: cDNA cloning and biochemical characterization. *Gene* 112:241–245 (1992).
155. Winn RN, Van Beneden RJ, Burkhart JGA. Transfer, methylation and spontaneous mutation frequency of Phi X174am3cs70 sequences in medaka (*Oryzias latipes*) and mummichog (*Fundulus heteroclitus*): implications for gene transfer and environmental mutagenesis in aquatic species. *Mar Environ Res* 40:247–265 (1995).
156. Hasspieler BM, Behar JV, Carlson DB, Watson DE, Di Giulio RT. Susceptibility of channel catfish (*Ictalurus punctatus*) and brown bullhead (*Ameiurus nebulosus*) to oxidative stress: a comparative study. *Aquat Toxicol* 28:53–64 (1994).
157. Easter SS Jr. An Oregonian school of zebra fish. *Trends Neurosci* 9:246–248 (1986).
158. Trinkaus JP. *Fundulus*. In: *Methods in Developmental Biology* (Wilt FH, Wessels NK, eds). New York:Thomas Y. Cromwell Company, 1967:113–122.
159. Armstrong PB, Child JS. Stages in the normal development of *Fundulus heteroclitus*. *Biol Bull* 128:143–169 (1965).
160. Iyengar A, Muller F, Maclean N. Regulation and expression of transgenes in fish—a review. *Transgenic Res* 5:147–166 (1996).
161. Kuo CH, Uetsuki T, Kim CH, Tanaka H, Li BS, Taira E, Higuchi H, Okamoto H, Yoshikawa K, Miki N. Determination of a neocin cis-acting element required for neuron specific expression using zebra fish. *Biochem Biophys Res Commun* 211:438–446 (1995).
162. Kavumpurath S, Andersen O, Kisen G, Alestrom P. Gene transfer methods and luciferase gene expression in zebra fish, *Brachydanio rerio* (Hamilton). *Isr J Aquacult* 45:154–163 (1993).
163. Kappus H. Lipid peroxidation: mechanisms, analysis, enzymology and biological relevance. In: *Oxidative Stress* (Sies H, ed). London:Academic Press, 1985:273–310.
164. Waldeck AR, Stocker R. Radical-initiated lipid peroxidation in low density lipoproteins: insights obtained from kinetic modeling. *Chem Res Toxicol* 9:954–964 (1996).
165. Cadet J, Berger M, Morin B, Raoul S, Wagner JR. Oxidative damage to DNA. In: *Analysis of Free Radicals in Biological Systems* (Favier AE, Cadet J, Kalyanaram B, Fontecave M, Pierre J-L, eds). Boston, MA:Birkhauser Verlag, 1995:51–64.



The EHIS puts even more environmental health information right at your fingertips!

EHIS articles contain convenient links to PubMed—the National Library of Medicine's free on-line search service of more than 9 million citations! Search MEDLINE and Pre-MEDLINE (including links to other online journals and databases) for information directly related to each EHIS article's topic!

Subscribe to the EHIS today at: <http://ehis.niehs.nih.gov/>