

## References

- Day, R. E., and White, R. H. R. (1977). Peritoneal dialysis in children. Review of 8 years' experience. *Archives of Disease in Childhood*, **52**, 56-61.
- Leigh, D. A. (1969). Peritoneal infections in patients on long-term peritoneal dialysis before and after human cadaveric renal transplantation. *Journal of Clinical Pathology*, **22**, 539-544.

V. P. CHOUDHRY  
Department of Paediatrics,  
All-India Institute of Medical Sciences,  
New Delhi 110016, India.

Dr R. H. R. White comments:

It is difficult to compare our findings with those of Dr Choudhry in the absence of more detailed information regarding (1) the age range of the patients dialysed, which Dr Day and I found influenced the infection rate, and (2) the causes of renal failure, which would have some bearing on the duration of dialysis. However, we would not disagree with his view that removal of the cannula after a short period of continuous dialysis might reduce the incidence of infection; indeed we suggested that this might be the case but felt that, since reinsertion of the cannula is an unpleasant experience for the child, such perfection might not always be practicable.

In our hands early detection and treatment of infection had a successful outcome except in one child with rapidly progressive glomerulonephritis, whose *Candida* peritonitis contributed to death at a time when we had no facilities for haemodialysis and transplantation in children.

R. H. R. WHITE  
Nephrology Department,  
The Children's Hospital,  
Birmingham B16 8ET.

## Assessment of total body fat in infancy from skinfold thickness measurements

Sir,

We have read the paper by Dauncey *et al.* (*Archives*, 1977, **52**, 223) and wish to make a few comments. In their paper these authors have related subcutaneous fat layer to skinfold thickness. They refer to the study by Hammond (1955) which examined the relation between uncompressed fat thickness  $T$  measured by x-ray and skinfold thickness  $S$  measured by calipers. Hammond finds that  $T = 0.95S - 0.0074S^2$ . For the range values encountered in the newborn, i.e.  $S \approx 5$  mm, Hammond's results (Tables IV and V) show  $T = S - 0.3$  mm. If we regard  $\theta$  as representing the true uncompressed subcutaneous fat thickness, i.e.  $T$  less the thickness of the dermis, then  $\theta = T - 1$ ; using Dauncey's value for thickness of the dermal layer (1 mm). Hence  $\theta = S - 1.3$  mm.

Dauncey *et al.* graph  $S$  against  $T$  using Hammond's data (see curve B, Figure), and by allowing 1 mm for the dermis derive curve C which relates subcutaneous fat to compressed skinfold thickness. Then, by assuming that

subcutaneous fat is 2 mm less than the skinfold thickness they show a line A,  $\theta = S - 2$ , and claim that, since line A lies fairly close to curve C, there is some justification for the empirical relation they have used to derive uncompressed fat thickness. In fact, as shown in our Figure, the line  $A_1$  with the equation  $\theta = S - 1.3$  is a better fit to the data which produce curve C since most of Hammond's observations occur with  $4.5 < S < 7$ ; that is within the region of contact of the curve and the straight line. We therefore believe that Dauncey *et al.* are in error in using  $\theta = S - 2$ . As shown in our first paragraph  $\theta$  is, in fact,  $S - 1.3$ .




Figure Thickness of subcutaneous fat  $\theta$  related to skinfold thickness  $S$  measured by calipers. Line A, equation  $\theta = S - 2$  (Dauncey *et al.*, 1977); curve B from Hammond's data; curve C, allowing 1 mm for the thickness of the dermis; line  $A_1$  equation  $\theta = S - 1.3$ .

Hammond's work shows that the Harpenden caliper, exerting as it does a constant force of 10 g, compresses a fold of skin to almost half its uncompressed value. Just as the calipers compress the fat layer by a factor of almost two, so they also compress the dermis. Knowing this, one would empirically suggest that the true uncompressed subcutaneous fat is 1 mm less (two thicknesses of compressed dermis) than the reading given on the calipers while compressing the fold of skin.

We, therefore, recommend that in using the body fat formula derived by Dauncey *et al.* one should use  $\theta = S - 1.3$ .

## Reference

- Hammond, W. H. (1955). Measurement and interpretation of subcutaneous fat, with norms for children and young adult males. *British Journal of Preventive and Social Medicine*, **9**, 201-211.

B. RICHARDS and S. W. DE SOUZA  
Department of Computation,  
University of Manchester Institute of  
Science and Technology, and  
Department of Child Health,  
University of Manchester.

Drs Dauncey and Gairdner comment:

We are unable to agree with Richards and De Souza's reasoning for several reasons. Their choice of the figure