

Characterization of Poly- γ -Glutamate Hydrolase Encoded by a Bacteriophage Genome: Possible Role in Phage Infection of *Bacillus subtilis* Encapsulated with Poly- γ -Glutamate

Keitarou Kimura and Yoshifumi Itoh*

Division of Applied Microbiology, National Food Research Institute, Tsukuba 305-8642, Japan

Received 20 September 2002/Accepted 7 February 2003

Some *Bacillus subtilis* strains, including natto (fermented soybeans) starter strains, produce a capsular polypeptide of glutamate with a γ -linkage, called poly- γ -glutamate (γ -PGA). We identified and purified a monomeric 25-kDa degradation enzyme for γ -PGA (designated γ -PGA hydrolase, PghP) from bacteriophage Φ NIT1 in *B. subtilis* host cells. The monomeric PghP internally hydrolyzed γ -PGA to oligopeptides, which were then specifically converted to tri-, tetra-, and penta- γ -glutamates. Monoiodoacetate and EDTA both inhibited the PghP activity, but Zn^{2+} or Mn^{2+} ions fully restored the enzyme activity inhibited by the chelator, suggesting that a cysteine residue(s) and these metal ions participate in the catalytic mechanism of the enzyme. The corresponding *pghP* gene was cloned and sequenced from the phage genome. The deduced PghP sequence (208 amino acids) with a calculated M_r of 22,939 was not significantly similar to any known enzyme. Thus, PghP is a novel γ -glutamyl hydrolase. Whereas phage Φ NIT1 proliferated in *B. subtilis* cells encapsulated with γ -PGA, phage BS5 lacking PghP did not survive well on such cells. Moreover, all nine phages that contaminated natto during fermentation produced PghP, supporting the notion that PghP is important in the infection of natto starters that produce γ -PGA. Analogous to polysaccharide capsules, γ -PGA appears to serve as a physical barrier to phage absorption. Phages break down the γ -PGA barrier via PghP so that phage progenies can easily establish infection in encapsulated cells.

Bacterial capsules that are exposed to the outermost cell surface play important roles in cell attachment for colonization, biofilm formation, and protecting cells from serum and phagocytosis (11, 12, 20, 22). Since capsules coat phage receptors on the cell surface, they also function as physical barriers against bacteriophages. The protective function of capsules against phage infection has been demonstrated with the capsular polysaccharides (CPSs) of several types of bacteria (5, 6, 9, 23, 37). However, CPS barriers are not always effective for all types of phages. Some *Escherichia coli* phages possess an enzyme that can degrade CPSs (4, 16, 18, 20, 24). For instance, *E. coli* phage 29 has an endo-*N*-acetylneuraminidase in its spike that hydrolyzes repeating poly- α -2,8-linked sialosyl units of K1 capsules (7, 16). Some phages of *Klebsiella* and *Streptococcus* strains also have CPS degradation activities associated with their particles (10, 25, 27).

Some strains of *B. subtilis*, *B. licheniformis*, and *B. anthracis* produce a unique capsule polymer of γ -linked glutamate, poly- γ -glutamate (γ -PGA) (30). The γ -PGA of *B. subtilis* contains almost equimolar amounts of L- and D-glutamate and the ratio of the D-isomer increases to about 80% when cells are cultured in the presence of $MnCl_2$ (21). The *capABC* genes encoding the γ -PGA synthetic system were originally identified in *B. anthracis* (19). Thereafter, their homologues were found in many *Bacillus* strains including *B. subtilis* (2, 3, 34). *B. subtilis* CapABC proteins are structurally and functionally equivalent to the *B. anthracis* counterparts (2, 3, 34), but the *B. subtilis*

genes are referred to as *pgsABC* or *ywsC-ywtAB* (2, 3, 34). The *ywsC*, *ywtA*, and *ywtB* genes have been renamed *capB*, *capC*, and *capA*, respectively (35). To avoid confusion with *pgsA* that has been assigned to a gene for phosphatidylglycerophosphate synthase (15), we also refer to the capsule synthetic genes of *B. subtilis* NAFM5 as *capABC* (DDJB accession number AB039950). Unlike *B. anthracis*, which forms γ -PGA in response to carbon dioxide (30), *B. subtilis* exclusively produces the capsule during the stationary growth phase through regulation by the ComQXPA quorum-sensing machinery (32). Because γ -PGA is very viscous, this capsule polypeptide, like CPSs, could perform a barrier function against phagocytosis and phage infection. Indeed, Makino et al. (19) have demonstrated a protective function of *B. anthracis* γ -PGA against phagocytosis by leukocytes. No evidence yet supports the notion of a hypothetical barrier function against phages. Rather, incidents contradictory to a barrier function are often experienced in natto factories. Although *B. subtilis* starters produce voluminous amounts of γ -PGA during fermentation, natto products are often contaminated with phages and then tend to rapidly lose γ -PGA viscosity when mixed for serving. An early study (14) identified and partially purified a γ -PGA depolymerase in a *B. natto* (= *B. subtilis*) culture infected with phage NP-1 cl, but did not investigate the role of the enzyme in phage infection and the corresponding gene. These findings indicated that phages produce a γ -PGA-degradation enzyme and that this enzyme may contribute to the infection of encapsulated host cells by eliminating the capsule.

To further characterize the γ -PGA degradation enzyme, we isolated phage Φ NIT1 from a natto product, purified its γ -PGA hydrolase (PghP) to homogeneity, and identified the enzyme gene in the phage genome. We found that the enzyme

* Corresponding author. Mailing address: Division of Applied Microbiology, National Food Research Institute, Kannondai 2-1-12, Tsukuba, Ibaraki 305-8642, Japan. Phone: 81-298-38-8075. Fax: 81-298-38-7996. E-mail: yosifumi@nfri.affrc.go.jp.

is a novel γ -glutamyl hydrolase that randomly hydrolyzes γ -PGA into oligo- γ -glutamates and then specifically into tri-, tetra-, and penta- γ -glutamate. We showed that Φ NIT1 has an apparent advantage over a phage that lacks PghP in terms of proliferation on encapsulated host cells. These findings support the notion that γ -PGA protects cells from phage adsorption and that PghP degrades the capsular barrier to allow phage progenies to infect encapsulated host cells.

MATERIALS AND METHODS

Strains, plasmids, phages and media. *B. subtilis* NAFM5 (Rif^r) is a spontaneous rifampin-resistant mutant of the natto starter strain Miyagino (Miura Natto starter) cured of cryptic plasmids pLS20 and pTA1015 as described (21). Bacteriophages Φ NIT1, FSG, KKP, MOP, ONPC, ONPB, P-1, SUP, SS2P, and THP were isolated from independent natto products with suspected phage contamination. *B. subtilis* typing phages (BS5, CS₁, F, Phi105, Phi29, Phi3T, PS10, PS50, S-a, and SPP1) and their host strains were obtained from Ackermann et al. (1). *E. coli* and *B. subtilis* strains were cultured in Luria-Bertani (LB) medium (26). *B. subtilis* NAFM5 was cultured on GSP for the production of γ -PGA as described (21). *E. coli* was transformed and selected on LB agar containing an appropriate antibiotic as described previously (32).

Assay for PghP. γ -PGA (5×10^6 Da) was extracted and purified from *B. subtilis* NAFM5 culture incubated on GSP with or without 0.1 mM MnCl₂ (21). The D-glutamate contents of the polypeptides from MnCl₂-free and 0.1 mM MnCl₂ media were 56 and 80%, respectively, as determined according to Nagai et al. (21). A standard mixture for the PghP assay (1 ml; 1 mg of γ -PGA with 56% of D-glutamate, 10 mM sodium phosphate [pH 7.5], 150 mM NaCl, enzyme) was incubated at 37°C for the indicated periods. The viscosity of reaction mixtures was measured using a falling ball viscometer (Haake) equipped with a syringe (0.5 ml) containing a stainless steel ball (3 mm in diameter) and is expressed as falling velocity (centimeters per second) of the ball. Degradation products of γ -PGA were detected by agarose gel electrophoresis. Portions (10 μ l) of reaction mixtures sampled after the indicated incubation periods were resolved by electrophoresis at 6 V/cm for 30 min on 1.0% agarose gels using TAE (40 mM Tris-hydroxylaminomethane, 1 mM EDTA, 0.14% [vol/vol] acetic acid) running buffer. Resolved degradation products on the gels were visualized by staining with methylene blue (0.23% [wt/vol] methylene blue, 23% [vol/vol] ethanol, 0.008% [wt/vol] KOH) for 10 min, followed by destaining with water.

Purification of PghP. *B. subtilis* NAFM5 cultured overnight in 20 ml of LB medium was inoculated into 1 liter of fresh LB medium along with phage Φ NIT1 at a multiplicity of infection of 5×10^{-4} . The culture was vigorously shaken at 37°C for 5 h. Cells and cell debris were removed by centrifugation, and then the enzyme was precipitated from the supernatant by adding ammonium sulfate (40 to 80% saturation), dissolved in 35 ml of buffer A (25 mM sodium phosphate buffer, pH 6.8, and 10 mM NaCl) and dialyzed against buffer A. After centrifugation at $100,000 \times g$ for 1 h, dialyzed proteins were loaded on a DEAE-Sepharose column equilibrated with buffer A. Active fractions eluted from the column using a gradient of NaCl (0 to 0.4 M) in buffer A were pooled and dialyzed against 10 mM sodium phosphate buffer, pH 6.8 (buffer B). Proteins were precipitated with ammonium sulfate (70% saturation), dialyzed against the same buffer, and fractionated by DEAE-Sepharose column chromatography as described above. Active fractions were combined, dialyzed against buffer B containing 1 M ammonium sulfate, and then loaded onto a Butyl-Toyopearl column (Tosoh). Active proteins were eluted using a reverse gradient (1 to 0 M) of ammonium sulfate in buffer B. Fractions containing the enzyme were pooled, dialyzed in buffer B, and fractionated by chromatography through a MonoQ column (Amersham Biosciences). The enzyme was eluted from the column using a gradient of 0 to 0.4 M NaCl in buffer B, concentrated with a Centrprep-10 (Millipore), and finally purified by Superose 12 (Amersham Biosciences) gel filtration column chromatography using buffer C (10 mM sodium phosphate, pH 6.9, 150 mM NaCl).

Analysis of reaction products. γ -PGA was incubated with purified PghP (25 ng) in the standard reaction mixture (1.5 ml) at 37°C. A portion (0.27 ml) was withdrawn from the reaction mixture after various incubation periods, boiled for 15 min to terminate the reaction, and fractionated through Superdex peptide HR10/30 (Amersham Biosciences) using buffer C. Degradation products were detected by absorbance at 210 nm. The molecular mass of reaction products was estimated from their elution volumes relative to those of glutamate ($M_r = 147$), di- γ -glutamate ($M_r = 276$; Bachem), tetra- γ -glutamate ($M_r = 535$; Hokkaido System Science), and gastrin ($M_r = 2,126$; Bachem). A mass spectrometer (Apex

II 70e; Bruker Daltonics) also determined the molecular masses of reaction products. Salts in reaction samples were removed by filtration through TSK-GEL ALPHA-2500 (Tosoh) before mass spectrometry. Amounts of L-glutamate were measured using NAD-L-glutamate dehydrogenase as described previously (36).

Cloning of pghP. We initially identified restriction fragments of Φ NIT1 DNA carrying *pghP* appropriate for cloning, by Southern blotting. Φ NIT1 DNA was digested with various restriction enzymes and blotted onto a Hybond N+ membrane (Amersham Biosciences) after resolution by agarose gel electrophoresis (26). Southern blotting using ³²P-end-labeled oligonucleotides, with 5'-TA(C/T)CCGAA(C/T)ATTGA(A/G)GC-3' as the probe, identified a 250-bp *SphI* fragment and a 3.2-kb *EcoRV* fragment that carry the relevant region of *pghP*: the nucleotide probes correspond to the 6th to 11th residues (YPNIEA) of the determined amino terminal sequence for PghP. The *SphI* fragment was extracted from an agarose gel and cloned into plasmid pKF19 (Takara Shuzo) at the *SphI* site. *E. coli* DH5 α (Bethesda Research Laboratories) transformants harboring the 250-bp *pghP* fragment on the plasmid were screened by colony hybridization using ³²P-end-labeled oligonucleotides as described previously (26). A 3.2-kb *EcoRV* fragment carrying *pghP* was also extracted from an agarose gel and cloned into plasmid pKF19 at the *HincII* site and then transformed into *E. coli* DH5 α . Transformants harboring *pghP* on the plasmid were selected by colony hybridization using the ³²P-labeled 250-bp *SphI* fragment as the probe. The *pghP* sequence was determined using plasmid pNAG201 harboring *pghP* in a positive clone.

Phage titration. Either phage Φ NIT1 (*pghP*⁺) or BS5 (*pghP*⁻) (10^4 PFU/ml) was inoculated into log-phase (optical density at 600 nm = 0.3) or stationary-phase cultures (24 h incubation) of *B. subtilis* NAFM5 shaken in 20 ml of GSP medium at 37°C. The incubation was continued for the indicated periods at 37°C with shaking. Aliquots (0.5 ml each) were withdrawn from the cultures and mixed with a few drops of chloroform. After removing cells and cell debris by centrifugation and subsequent filtration through a membrane filter (pore size, 0.45 μ m), phage samples were diluted as required with LB medium. Portions (0.1 ml) of diluted phage were added to 3 ml of soft LB agar containing 0.8% (wt/vol) agarose and 2×10^7 cells of strain NAFM5. Soft agar was quickly mixed and overlaid on bottom LB layers (1.5% [wt/vol] agarose) in plates. After incubation at 37°C overnight, the number of plaques that developed was counted. Strain NAFM5 does not produce γ -PGA in LB medium.

Nucleotide and amino acid sequencing. Nucleotides were sequenced using a Dye terminator cycle sequencing kit (Applied Biosystems) and an ABI 310A DNA sequencer (Applied Biosystems). The *pghP* sequence was deposited in DDJB/EMBL/GenBank databases under accession number AB091475. Purified PghP (10 pmol) was resolved by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) (17) and blotted onto a polyvinylidene difluoride membrane (Millipore). The amino terminus was then determined using a HP G10000A protein sequencer (Hewlett-Packard).

RESULTS

Identification of a γ -PGA-degradation enzyme in a culture infected with Φ NIT1. Bacteriophage Φ NIT1 was isolated from a natto product that contained small amounts of γ -PGA, an indication of phage contamination. To examine whether this phage produces a γ -PGA degradation enzyme, we cultured *B. subtilis* NAFM5, a derivative of a natto starter, along with phage Φ NIT1 at a multiplicity of infection of 5×10^{-4} in LB medium at 37°C for 5 h. Incubation with the culture supernatant caused γ -PGA to rapidly lose viscosity (Fig. 1A) and undergo fragmentation as confirmed by agarose gel electrophoresis (Fig. 1B). *B. subtilis* strains produce an unidentified γ -PGA degradation enzyme during the late stationary growth phase (31). However, neither viscometry nor agarose gel electrophoresis detected γ -PGA degradation activity in the culture supernatant or in extracts of host cells. In addition, γ -PGA degradation activity was undetectable by the phage particles. These results indicated that the γ -PGA degradation enzyme is specifically synthesized in host cells harboring the phage and that the enzyme is not associated with the phage particle. We designated the enzyme PghP, for γ -PGA hydrolase of phage.




FIG. 1. γ -PGA degradation activity in a *B. subtilis* NAFM5 culture infected with Φ NIT1 phage measured by viscometry (A) and by agarose gel electrophoresis (B). Reactions proceeded as described in Materials and Methods, and portions of mixtures were withdrawn after the indicated incubation periods. (A) Viscosity was measured using a falling-ball viscometer and is expressed as relative viscosity to the mixture at zero time. (B) Samples (10 μ l) were resolved by 1.0% agarose gel electrophoresis, and degradation products were visualized by staining with methylene blue. Positions of intact γ -PGA (5×10^6 Da) and of 10^4 Da and 10^3 Da γ -PGA (fractionated from partial hydrolysates of γ -PGA by PghP through gel filtration and high-performance liquid chromatography) (21) are indicated at left.

Purification and characterization of PghP. We purified PghP from a Φ NIT1-infected NAFM5 culture through five chromatographic steps as described in Materials and Methods. The purified enzyme migrated as a single band of 25-kDa protein on SDS-PAGE (Fig. 2). This value was the same as the M_r (25,000) determined for native PghP by Superose 12 gel filtration chromatography, indicating that PghP is a monomer of 25 kDa. The first 25 amino acids of the amino-terminal sequence of PghP was AQTDTYPNIEALENAETVGVA YNIE. As described below, the deduced nucleotides corresponding to residues 6 to 11 (YPNIEA) of the amino terminus were used as probes in Southern blotting and in *phgP* cloning. EDTA (1 mM) powerfully inhibited the enzyme, and an excess of $MnCl_2$ and $ZnCl_2$, but not $MgCl_2$ and $CaCl_2$, fully restored the activity inhibited by the chelator (Fig. 3). Monoiodoacetate also inhibited the enzyme (Fig. 3).

Analysis of degradation products. Glutamate was undetectable even after prolonged incubation or after incubation with more of the purified enzyme, implying that this enzyme does not degrade γ -PGA into the monomer. We accordingly determined the sizes of the smallest degradation products by gel filtration chromatography. Like crude enzyme (Fig. 1A and B), purified PghP rapidly degraded γ -PGA of 5×10^6 Da into smaller fragments. The substrate was degraded into fragments




FIG. 2. SDS-PAGE of purified PghP. PghP (1 μ g, lane P) from Superose 12 column chromatography was analyzed together with molecular mass markers (1 μ g each, lane M) by SDS-10% PAGE (17). Molecular markers: phosphorylase *b* (97 kDa), bovine serum albumin (67 kDa), ovalbumin (45 kDa), carbonic anhydrase (30 kDa), and catalase (21 kDa).




FIG. 3. Effects of monoiodoacetate, EDTA, and divalent cations on PghP activity. Reactions proceeded at 37°C for 60 min in PghP assay mixture (see Materials and Methods) containing 50 mM Tris-HCl (pH 7.5) instead of 50 mM sodium phosphate (pH 7.5), and reaction products were analyzed using agarose gel electrophoresis. Lane 1, no enzyme (control). Lanes 2 through 8 contained 45 ng of purified PghP/ml plus the following: lane 2, no further addition; lane 3, 1 mM EDTA; lane 4, 1 mM EDTA + 5 mM CaCl₂; lane 5, 1 mM EDTA + 5 mM MgCl₂; lane 6, 1 mM EDTA + 5 mM MnCl₂; lane 7, 1 mM EDTA + 5 mM ZnCl₂; lane 8, 1 mM monoiodoacetate. Reaction mixtures for lanes 4 to 7 were incubated for 5 min before adding chlorides.

of diverse sizes within 10 min (Fig. 4 [10 min]). The average molecular mass of the products in the first peak was approximately 10⁵ Da as determined by comparison with the elution volume relative to those of pullulans in gel filtration chromatography (21). The smallest peptide was a trimer, and relatively large amounts of pentamers and heptamers were generated (Fig. 4 [10 min]). Degradation intermediates larger than octamers were further degraded to accumulate trimers, tetramers, pentamers, and heptamers within a further 20 min of incubation (Fig. 4 [30 min]). By 60 min, only trimers, tetramers, and pentamers were detected (Fig. 4 [60 min]). These oligo- γ -glutamates were not further cleaved by prolonged incubation or by more of the enzyme. Hydrolysis of heptamers did not yield monomers or dimers, and the amounts of trimers and tetramers significantly increased relative to that of pentamers (Fig. 4 [30 and 60 min]), suggesting that heptamers are specifically converted into trimers and tetramers. Mass spectroscopic analyses confirmed the presence of these oligomers in the 60-min reaction mixture. We propose a hydrolytic mechanism for PghP on oligo- γ -glutamates in the Discussion.

Cloning and structure of *pghP*. Southern blotting using oligonucleotides deduced from the determined amino terminal sequence of PghP showed that the 5'-region of *pghP* is located on a 250-bp *SphI* segment of a 3.2-kb *EcoRV* fragment of Φ NIT1 DNA (Fig. 5). We initially cloned the *SphI* segment and sequenced the nucleotides to confirm that the segment contains the amino terminal region of *pghP* that is identical to the determined sequence, except for the first methionine, which appears to be eliminated posttranslationally (Fig. 5). We subsequently cloned the 3.2-kb *EcoRV* fragment and determined the entire sequence of *pghP* (DDJB/EMBL/GenBank accession number AB091475) located between 1.6 and 2.3 kb (Fig. 5). The predicted M_r of PghP (203 amino acids) was




FIG. 4. Analysis of γ -PGA degradation products generated by PghP over reaction time. Reactions proceeded under standard assay conditions (see Materials and Methods). Samples (0.2 ml) after various incubation periods were applied onto Superdex peptide columns, and reaction products were monitored by absorbance at 210 nm. Molecular sizes of oligomers were estimated from elution volumes relative to those of glutamate ($M_r = 147$), di- γ -glutamate ($M_r = 276$), tetra- γ -glutamate ($M_r = 535$), and gastrin ($M_r = 2,126$). Abbreviations: Vo, void volume; G₃, tri- γ -glutamate; G₄, tetra- γ -glutamate; G₅, penta- γ -glutamate; G₇, hepta- γ -glutamate.

22,939, which was in agreement with the M_r (25,000) of purified PghP. A BLAST search using the deduced PghP sequence did not reveal any similar proteins.

Infection of encapsulated cells with phage Φ NIT1. For the above experiments we incubated phage Φ NIT1 in log-phase cultures in which strain NAFM5 does not produce capsule (32). Thus, whether phage Φ NIT1 could infect encapsulated cells remained unclear. To investigate the role of PghP in the infection of encapsulated host cells, we compared the growth of phage Φ NIT1 with that of a phage lacking PghP. To achieve this, we required a phage that can infect strain NAFM5 but which does not produce PghP. We tested the ability of 10 *B. subtilis* typing phages (1) to infect strain NAFM5 and to pro-

access to their cell surface receptors. We showed that like CSPs (5, 6, 9, 23, 27, 37), the γ -PGA capsule of *B. subtilis* also functions as a physical barrier against phages (Fig. 6B). In addition, analogous to phages that can infect CPS-encapsulated cells by dismantling the capsule with a degradation enzyme (4, 7, 10, 16, 24, 25, 27, 37), phage Φ NIT1 achieves absorption by eliminating γ -PGA on host cells with PghP (Fig. 6A and B). In contrast to CPS degradation enzymes that are associated with phage particles (4, 10, 16, 18, 25, 27), PghP is produced in the absence of such association. The different locations of degradation enzymes may correlate with distinct mechanisms of capsule synthesis by host cells. In contrast to *E. coli* that forms capsule independently of growth phase (8), *B. subtilis* strains form γ -PGA under the high-cell-density conditions of stationary phase (32). Phage Φ NIT1, which does not have PghP as its component, may not easily access receptors on encapsulated cells. However, an infective phage produces the enzyme in host cells that will be released outside host cells during lysis, which would eliminate γ -PGA on neighboring cells and thereby allow its progenies to easily access cell surface receptors. PghP would efficiently degrade capsules when the cell density is high, particularly in colonies or layers on solid surfaces such as that of fermenting soybeans, where diffusion of the enzyme is limited.

Phages often cause serious damage in natto factories. Problems associated with phage contamination are not only poor fermentation but also the rapid decrease in viscosity of natto products (an important factor in natto quality). Contamination with a few phages does not cause an apparent failure of fermentation but can cause a significant decrease in slime viscosity when mixed for serving. Phage Φ NIT1 produces about 1 mg of PghP per liter under appropriate culture conditions, and 1 μ g of the enzyme completely hydrolyzes 1.5 mg of γ -PGA within 1 min to the final products. Our present results showed that PghP, which is produced at high levels and has powerful degradation activity, is responsible for the rapid decay of γ -PGA slime in natto that is contaminated with phage.

Other reported enzymes that cleave the γ -glutamyl linkage include γ -PGA depolymerase of *B. anthracis* (33), γ -PGA hydrolase of *Myrothecium* sp. (29), carboxypeptidase G (EC 3.4.17.11) of *Pseudomonas* sp. (28), animal γ -glutamyl hydrolase (EC 3.4.19.9) (13, 38), and animal glutamate carboxypeptidase II (EC 3.4.17.21) (13). Among these, carboxypeptidase G, γ -glutamyl hydrolase, and glutamate carboxypeptidase II have been well characterized and their corresponding genes have been cloned. γ -Glutamyl hydrolase is a thiol enzyme that externally and internally hydrolyzes the γ -glutamyl tail of folyl- γ -PGAs (38). Glutamate carboxypeptidase II is an endopeptidase of folyl- γ -PGA that requires Zn^{2+} , but not cysteine, for activity (13). *Pseudomonas* carboxypeptidase G (EC 3.4.17.11) is also a zinc metalloenzyme that releases carboxyl-terminal glutamate residues from γ -glutamyl peptides and folyl- γ -PGAs (28). Besides the carboxypeptidases, many peptidyl and non-peptidyl C-N hydrolases use Zn^{2+} or Mn^{2+} to activate the hydroxyl group of water that makes a nucleophilic attack on the carbon atom of a C-N bond. PghP requires both cysteine and Zn^{2+} or Mn^{2+} for activity (Fig. 3), implying that they participate in the reaction mechanism. PghP has two cysteine residues at positions 91 and 156. However, these regions do not show any local homology with the region of γ -glutamyl hydro-

lase containing the essential cysteine residue (Cys110) (38). The cysteine residue(s) of PghP might be involved either in formation of a thioester enzyme-substrate intermediate or in binding to metal ions. This requires elucidation.

PghP internally and randomly degrades γ -PGA and large oligo- γ -glutamates (Fig. 1B and Fig. 4 [10 min]). Within 30 min of incubation, trimers, tetramers, pentamers, and heptamers become dominant (Fig. 4 [30 min]). PghP therefore appears inert to trimers, tetramers, and pentamers, thus accumulating these peptides as the final product (Fig. 4 [60 min]). High levels of heptamers relative to larger oligopeptides suggest that PghP hydrolyzes heptamers more slowly than it does the other larger oligopeptides. The absence of monomers and dimers in the final product indicates the specific conversion of hexamers to two trimers. Heptamers and octamers may also be specifically cleaved into trimers and tetramers and into trimers and pentamers or into two tetramers. Thus, the cleavage of oligo- γ -glutamates by PghP is not random. The specific hydrolysis of oligo- γ -glutamates suggests that this enzyme binds to oligo- γ -glutamates at six γ -glutamyl residues and therefore may have low binding-affinity for oligopeptides that are smaller than pentamers. Assuming that hydrolysis occurs at the center of the binding site, this substrate-binding mode is quite consistent with the product profiles of PghP (Fig. 4 [60 min]). Alternatively, PghP might recognize the γ -linkages of either D- or L-glutamate in γ -PGA. However, this is less likely because PghP generated tri-, tetra-, and penta- γ -glutamates from γ -PGA consisting of 80% D-glutamate in the same amounts as those from the polypeptide consisting of 56% D-glutamate (Fig. 4 [60 min]).

The γ -PGA degradation enzyme of phage NP-1 cl produces dimers and trimers as the final product (14). Perhaps the substrate-binding site of this enzyme corresponds to a tetrapeptide. It may therefore be able to bind pentamers and tetramers and to cleave them into dimers and trimers and into two dimers, respectively. Our preliminary Southern blot experiments showed that Φ NIT1 *pghP* hybridizes with the counterparts of phages KKP, MOP, ONPC, ONPB, SUP, SS2P, THP, and SP50, but not with those of phages CS₁, FSG, P-1, S-a, and SP10 (data not shown). Phage PghPs appear to be heterogeneous with respect to primary structure as well as mode of oligopeptide cleavage. *B. subtilis* typing phages are not related to each other or to natto phages in terms of morphology, restriction profiles of genomic DNA, and host range (1). Nevertheless, the *pghP* gene also occurs in some typing phages. In conclusion, PghP is widely distributed in *B. subtilis* phages and significantly contributes to the infection of *B. subtilis* strains that produce γ -PGA.

ACKNOWLEDGMENTS

We thank K. Nagashima for providing phage Φ NIT1 and H.-W. Ackerman for the gift of typing phages and their host strains.

This study was supported in part by a grant-in-aid from the Ministry of Agriculture, Forestry and Fisheries.

REFERENCES

- Ackermann, H.-W., R. R. Azizbekyan, R. L. Bernier, H. de Barje, S. Saindoux, J.-R. Valéro, and M.-X. Yu. 1995. Phage typing of *Bacillus subtilis* and *B. thuringensis*. Res. Microbiol. 146:643-657.
- Ashiuchi, M., C. Nawa, T. Kamei, J.-J. Song, S.-P. Hong, M.-H. Sung, K. Soda, T. Yagi, and H. Misono. 2001. Physical and biochemical characterization of poly- γ -glutamate synthase complex of *Bacillus subtilis*. Eur. J. Biochem. 268:5321-5328.

3. Ashiuchi, M., K. Soda, and H. Misono. 1999. A poly- γ -glutamate synthetic complex of *Bacillus subtilis* IFO3336: gene cloning and biochemical analysis of poly- γ -glutamate produced by *Escherichia coli* clone cells. *Biochem. Biophys. Res. Commun.* **263**:6–12.
4. Bayer, M. E., H. Thurow, and M. H. Bayer. 1979. Penetration of the polysaccharide capsule of *Escherichia coli* (Bi161/42) by bacteriophage K29. *Virology* **94**:95–118.
5. Benedi, V. J., B. Ciurana, and J. M. Tomas. Isolation and characterization of *Klebsiella pneumoniae* unencapsulated mutants. *J. Clin. Microbiol.* **27**:82–87.
6. Bernheimer, H. P., and J. G. Tiraby. 1976. Inhibition of phage infection by pneumococcus capsule. *Virology* **73**:308–309.
7. Bessler, W., F. Fehmel, E. Freund-Molbert, H. Knufemann, and S. Stirm. 1975. *Escherichia coli* capsule bacteriophages. IV. Free capsule depolymerase 29. *J. Virol.* **15**:976–984.
8. Bliss, J. M., and R. P. Silver. 1996. Coating the surface: a model for expression of capsular polysialic acid in *Escherichia coli* K1. *Mol. Microbiol.* **21**:221–231.
9. Burt, S., S. Meldrum, D. R. Woods, and D. T. Jones. 1978. Colonial variation, capsule formation, and bacteriophage resistance in *Bacteroides thetaiotaomicron*. *Appl. Environ. Microbiol.* **35**:439–443.
10. Cescutti, P., and S. Paoletti. 1994. On the specificity of a bacteriophage-borne endoglycanase for the native capsular polysaccharide produced by *Klebsiella pneumoniae* SK1 and its derived polymers. *Biochem. Biophys. Res. Commun.* **198**:1128–1134.
11. Cross, A. S., K. S. Kim, D. C. Wright, J. C. Sadoff, and P. Gemski. 1986. Role of lipopolysaccharide and capsule in the serum resistance of bacteremic strains of *Escherichia coli*. *J. Infect. Dis.* **154**:497–503.
12. Czuprynski, C. J., and A. K. Sample. 1990. Interactions of *Haemophilus Actinobacillus-Pasteurella* bacteria with phagocytic cells. *Can. J. Vet. Res.* **54**(Suppl.):S34–S40.
13. Galivan, J., T. J. Ryan, K. Chave, M. Rhee, R. Yao, and D. Yin. 2000. Glutamyl hydrolase: pharmacological role and enzymatic characterization. *Pharmacol. Ther.* **85**:207–215.
14. Hongo, M., and A. Yoshimoto. 1970. Bacteriophage of *Bacillus natto*. part III. Action of phage-induced γ -polyglutamic acid depolymerase on γ -polyglutamic acid and the enzymatic hydrolyzates. *Agric. Biol. Chem.* **34**:1055–1063.
15. Kunst, F., et al. 1997. The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*. *Nature* **390**:249–256.
16. Kwiatkowski, B., B. Boschek, H. Thiele, and S. Stirm. 1982. Endo-*N*-acetylneuraminidase associated with bacteriophage particles. *J. Virol.* **43**:697–704.
17. Laemmlli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**:680–685.
18. Long, G. S., J. M. Bryant, P. W. Taylor, and J. P. Luzio. 1995. Complete nucleotide sequence of the gene encoding bacteriophage E endosialidase: implications for K1E endosialidase structure and function. *Biochem. J.* **309**:543–550.
19. Makino, S., I. Uchida, N. Terakado, C. Sasakawa, and M. Yoshikawa. 1989. Molecular characterization and protein analysis of the *cap* region, which is essential for encapsulation in *Bacillus anthracis*. *J. Bacteriol.* **171**:722–730.
20. McCallum, K. L., D. H. Laakso, and C. Whitfield. 1989. Use of a bacteriophage-encoded glycanase enzyme in the generation of lipopolysaccharide O side chain deficient mutants of *Escherichia coli* O9:K30 and *Klebsiella* O1:K20: role of O and K antigens in resistance to complement-mediated serum killing. *Can. J. Microbiol.* **35**:994–999.
21. Nagai, T., K. Koguchi, and Y. Itoh. 1997. Chemical analysis of poly- γ -glutamic acid produced by plasmid-free *Bacillus subtilis* (*natto*): Evidence that plasmids are not involved in poly- γ -glutamic acid production. *J. Gen. Appl. Microbiol.* **43**:139–143.
22. Ohman, D. E., K. Kalai, C. J. McPherson, C. A. DeVries, S. Ma, D. J. Wozniak, and M. J. Franklin. 1996. Regulation of the alginate (*algD*) operon in *Pseudomonas aeruginosa*, p. 472–483. In T. Nakazawa, K. Furukawa, D. Haas, and S. Silver (ed.) *Molecular biology of pseudomonads*. ASM Press, Washington, D.C.
23. Ohshima, Y., F. Schumacher-Perdreau, G. Peters, and G. Pulverer. 1988. The role of capsule as a barrier to bacteriophage adsorption in an encapsulated *Staphylococcus simulans* strain. *Med. Microbiol. Immunol.* **177**:229–233.
24. Pelkonen, S., J. Aalto, and J. Finne. 1992. Differential activities of bacteriophage depolymerase on bacterial polysaccharide: binding is essential but degradation is inhibitory in phage infection of K1-defective *Escherichia coli*. *J. Bacteriol.* **174**:7757–7761.
25. Rieger-Hug, D., and S. Stirm. 1981. Comparative study of host capsule depolymerases associated with *Klebsiella* bacteriophages. *Virology* **113**:363–378.
26. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
27. Saxelin, M. L., E. L. Nurmiaho, M. P. Korhola, and V. Sundman. 1979. Partial characterization of a new C3-type capsule-dissolving phage of *Streptococcus cremoris*. *Can. J. Microbiol.* **25**:1182–1187.
28. Sherwood, R., R. G. Melton, S. M. Alwan, and P. Hughes. 1985. Purification and properties of carboxypeptidase G₂ from *Pseudomonas aeruginosa* sp. strains RS-16. **148**:447–453.
29. Tanaka, T., O. Hirata, T. Futamura, K. Uotani, A. Satoh, M. Taniguchi, and S. Oi. 1993. Purification and characterization of poly(γ -glutamic acid) hydrolase from a filamentous fungus, *Myrothecium* sp. TM-422. *Biosci. Biotech. Biochem.* **57**:2148–2153.
30. Thorne, C. B. 1993. *Bacillus anthracis*. P. 113–124. In A. L. Sonenshein, J. A. Hock, and R. Losick (ed.), *Bacillus subtilis* and other Gram-positive bacteria, American Society for Microbiology, Washington, D.C.
31. Throne, C. B., C. G. Gómez, H. E. Noyes, and R. D. Housewright. 1954. Production of glutamyl polypeptide by *Bacillus subtilis*. *J. Bacteriol.* **68**:307–315.
32. Tran, L.-S. P., T. Nagai, and Y. Itoh. 2000. Divergent structure of the *comQXPA* quorum-sensing components: molecular basis of strain-specific communication mechanism in *Bacillus subtilis*. *Mol. Microbiol.* **37**:1159–1171.
33. Uchida, I., S. Makino, C. Sasakawa, M. Yoshikawa, C. Sugimoto, and N. Terakado. 1993. Identification of a novel gene, *dep*, associated with depolymerization of the capsule polymer in *Bacillus anthracis*. *Mol. Microbiol.* **9**:487–496.
34. Urushibata, Y., S. Tokuyama, and Y. Tahara. 2002. Characterization of the *Bacillus subtilis ywsC* gene, involved in γ -polyglutamic acid production. *J. Bacteriol.* **184**:337–343.
35. Urushibata, Y., S. Tokuyama, and Y. Tahara. 2002. Difference in transcription levels of *cap* genes for γ -polyglutamic acid production between *Bacillus subtilis* IFO16449 and Marburg 168. *J. Biosci. Bioeng.* **93**:252–254.
36. Valero, E., and F. Garcia-Carmona. 1998. A continuous spectrophotometric method based on enzymatic cycling for determining L-glutamate. *Anal. Biochem.* **259**:265–271.
37. Wilkinson, B. J., and K. M. Holmes. 1979. *Staphylococcus aureus* cell surface: capsule as a barrier to bacteriophage adsorption. *Infect. Immun.* **23**:549–552.
38. Yao, R., E. Schneider, T. J. Ryan, and J. Galivan. 1996. Human γ -glutamyl hydrolase: cloning and characterization of the enzyme expressed *in vitro*. *Proc. Natl. Acad. Sci. USA* **93**:10134–10138.