

Gene transfer from a parasitic flowering plant to a fern

Charles C. Davis^{1,*}, William R. Anderson² and Kenneth J. Wurdack³

¹Department of Organismic and Evolutionary Biology, Harvard University Herbaria, 22 Divinity Avenue, Cambridge, MA 02138, USA

²University of Michigan Herbarium, 3600 Varsity Drive, Ann Arbor, MI 48108-2287, USA

³Department of Botany and Laboratories of Analytical Biology, Smithsonian Institution, 4210 Silver Hill Road, Suitland, MD 20746, USA

The rattlesnake fern (*Botrychium virginianum* (L.) Sw.) is obligately mycotrophic and widely distributed across the northern hemisphere. Three mitochondrial gene regions place this species with other ferns in Ophioglossaceae, while two regions place it as a member of the largely parasitic angiosperm order Santalales (sandalwoods and mistletoes). These discordant phylogenetic placements suggest that part of the genome in *B. virginianum* was acquired by horizontal gene transfer (HGT), perhaps from root-parasitic Loranthaceae. These transgenes are restricted to *B. virginianum* and occur across the range of the species. Molecular and life-history traits indicate that the transfer preceded the global expansion of *B. virginianum*, and that the latter may have happened very rapidly. This is the first report of HGT from an angiosperm to a fern, through either direct parasitism or the mediation of interconnecting fungal symbionts.

Keywords: biogeography; ferns; horizontal gene transfer; Ophioglossaceae; parasitic plants; phylogeography

1. INTRODUCTION

Horizontal gene transfer (HGT) appears to be common in plant mitochondrial (mt) genomes (Bergthorsson *et al.* 2003), but until very recently explanations of how such transfers occur have been speculative (Bergthorsson *et al.* 2003; Won & Renner 2003). Two studies show that parasitism of one plant by another is potentially an important mode of HGT in angiosperms. The first documented HGT from a host to its parasite (Davis & Wurdack 2004), and the second from a parasite to its host (Mower *et al.* 2004). This paper reports the first case of HGT from an angiosperm parasite to a putative fern host, and offers the first data on the geographic distribution of transgenes in a recipient species.

2. MATERIAL AND METHODS

(a) Taxon sampling

Our analyses included four mt gene regions (*atp1*, *atp6*, *matR* and *nad1B-C*) spanning all major vascular plant clades (i.e. ferns, cycads, gymnosperms and angiosperms). The *matR* dataset included nearly all orders of flowering plants (APG 2003) and *nad1B-C* was similarly sampled for angiosperms and spanned most major eudicot clades. Each of these gene trees was rooted with *Huperzia* (*atp1*, *matR* and *nad1B-C*), *Lycopodium* (*atp1*), and *Marchantia* (*atp6*) following the large-scale phylogenetic studies by Karol *et al.* (2001) and Pryer *et al.* (2001). All primers and protocols used to generate these data, including those for RT-PCR and cDNA analyses described in §3 below, can be found in

the supplementary online version accompanying this article.

Focused sampling for these four gene regions included 24 species (in 21 genera) representing all five families of Santalales, including representatives of the earliest diverging members of the order (Nickrent & Malécot 2001; APG 2003; Malécot *et al.* 2004). Our sampling of Loranthaceae spans the basal node of the family: the monospecific Australian root-parasitic *Nuytsia* is sister to the rest of the family and is included in our analyses (Vidal-Russell & Nickrent 2005). Along with the more derived aerial-parasitic loranths, root-parasitic *Gaiadendron* and *Nuytsia* are also included. We also sampled 12 fern species representing all three genera of Ophioglossaceae and species from all subgenera of *Botrychium* (Hauk *et al.* 2003), including the previously unplaced Afroasian subgenus *Japanobotrychium* (Hauk *et al.* 2003).

Population sampling of *Botrychium virginianum* (L.) Sw. for transgenic regions, *matR* and *nad1B-C*, included 34 accessions spanning its worldwide distribution, including North America (Canada, Mexico, United States), Central America (Costa Rica, Panama), South America (Bolivia, Brazil, Peru), the Greater Antilles (Dominican Republic), Europe (Austria, Germany, Switzerland) and Asia (China, Japan). For complete sampling information see table 1 in the Electronic Appendix.

We also included a fifth dataset, plastid *rbcL*, that included most major fern clades plus horsetails, and was rooted with *Huperzia*, *Lycopodium*, gymnosperms and cycads (Pryer *et al.* 2001). The purpose of including *rbcL* was to infer the biogeography of *Botrychium*, and to determine the approximate maximum age estimate for this HGT event (see §2*d*). The *rbcL* data are especially

* Author for correspondence (cdavis@oeb.harvard.edu).

amenable to biogeographic analysis because: (i) the plastid genome does not appear to be susceptible to HGT across species boundaries and (ii) *rbcL* is sufficiently well sampled in ferns to provide a solid foundation for inferring the phylogeny and biogeography of Ophioglossaceae—the *rbcL* data set includes most major fern lineages, all genera of Ophioglossaceae, and nearly all species of *Botrychium* throughout their worldwide range (Hauk *et al.* 2003).

(b) Phylogenetic analysis

Maximum likelihood (ML) optimization was implemented for all independent (*atp1*, *atp6*, *matR* and *nad1B-C*) and combined (*matR* plus *nad1B-C*) datasets in PHYML *ver.* 2.4.4 (Guindon & Gascuel 2003) under the general time reversible model (GTR), or a submodel of the GTR model, as determined by ModelTest 3.6 (Posada & Crandall 1998). ML support values were estimated from 100 bootstrap replicates in PHYML, and parsimony support was estimated from 1000 bootstrap replicates in PAUP*4b10 (Swofford 2003). Parsimony bootstrap replicates were implemented with heuristic searches using 10 random taxon additions per replicate, tree-bisection–reconnection, MULPARS, and holding 10 trees at each taxon addition. To assess the alternate topological placement of *B. virginianum* in single data set analyses we employed the Kishino–Hasegawa (Kishino & Hasegawa 1989), Shimodaira–Hasegawa (Kadowaki *et al.* 1996) and parametric bootstrap tests (Huelsenbeck *et al.* 1996) using ML.

(c) Statistical phylogeography

We calculated Tajima's *D* and mismatch distribution with ARLEQUIN (Schneider *et al.* 2000) for all samples of *B. virginianum*. The demographic expansion model of Rogers and Harpending (Harpending 1994; Rogers 1995) was used to analyse the pairwise mismatch distributions. Twenty-five polymorphic sites were obtained for combined *matR* and *nad1B-C*. There were 19 and 6 polymorphic sites representing 11 and 7 haplotypes across individuals for transgenic *matR* and *nad1B-C*, respectively.

(d) Biogeographic Analyses

To infer the location of disjunctions of Ophioglossaceae clades, ancestral areas were reconstructed on the ML *rbcL* topology with dispersal–vicariance analysis (DIVA) as implemented in DIVA *ver.* 1.1 (Ronquist 1997). Our data matrix used to assess ancestral areas was constructed by scoring each species for its presence in seven major continental areas of endemism: Africa, Asia, Australia, Central America, Europe, North America and South America. Distributional data were obtained from Clausen (1938) and Wagner & Wagner (1993) (see supplementary online materials for trees and scorings). To corroborate results obtained using DIVA we similarly analysed these data using parsimony as implemented with the default 'trace' optimization of MacClade.

To estimate divergence times, branch lengths and an associated likelihood score were calculated for the respective models of sequence evolution in the absence of a molecular clock for plastid *rbcL* data following Davis *et al.* (2002). Finding that a clock was rejected ($p < 0.05$) we used penalized likelihood (Sanderson 2002) to estimate divergence times, focusing specifically on the split between *B. virginianum* and its closest *Botrychium*

relatives. Since this transfer event is restricted to *B. virginianum* (see §3), the divergence time for this node provides a maximum age estimate for this event. The transfer can be younger than this node, but not older. To estimate standard errors associated with divergence times, we used the parametric bootstrapping strategy outlined in Davis *et al.* (2002).

We applied 12 minimum fossil age constraints recently reviewed by Schneider *et al.* (2004) to the *rbcL* data to estimate absolute divergence times (see Electronic Appendix). In addition, we also included the minimum age constraint of 57.8 Myr (time-scale *sensu* Berggren *et al.* 1995) for stem group *Botrychium* based on reliable Paleocene *Botrychium* fossils from western North America (Rothwell & Stockey 1989). We also separately enforced two maximum age constraints for the root of our tree, which corresponds to the euphyllophyte clade (Pryer *et al.* 2001)—380 and 450 Myr. The first age constraint corresponds to the earliest known euphyllophyte fossils (Schneider *et al.* 2004), and the second corresponds to the approximate maximum age for all land plants based on fossil and molecular divergence time estimates (Sanderson 2003). The cross validation procedure for these data yielded an optimal smoothing value of 1000.

3. RESULTS AND DISCUSSION

While reconstructing the phylogenies of all four mt gene regions we uncovered two strongly discordant placements for the rattlesnake fern, *B. virginianum*. The genes *atp1*, *atp6*, and one copy of *matR* placed *B. virginianum* with its closest relatives as a member of the fern family Ophioglossaceae (figure 1*b–d*). In contrast, *nad1B-C* and a second copy of *matR* placed it as a member of the parasitic angiosperm order Santalales (figure 1*a,b*), which includes the sandalwoods and mistletoes. We sought angiosperm *matR* and *nad1B-C* in other Ophioglossaceae (i.e., representatives of all three genera of Ophioglossaceae plus all subgenera of *Botrychium sensu* Hauk *et al.* 2003, including the closest relatives of *B. virginianum*) using a battery of angiosperm specific primers for these gene regions. In all sampled Ophioglossaceae, angiosperm-like copies were detected only in *B. virginianum*. In addition, universal *matR* primers designed to amplify both native and transgene copies of *matR* uncovered only the native copy in other Ophioglossaceae.

This anomalous phylogenetic placement is robust, according to the Kishino–Hasegawa, Shimodaira–Hasegawa and parametric bootstrap tests: transgenic *matR* and *nad1B-C* favoured ($p < 0.05$ for all tests) the placement of *B. virginianum* with Santalales rather than with Ophioglossaceae; similarly, native *matR*, *atp1* and *atp6* favoured ($p < 0.05$ for all tests) the placement of *B. virginianum* with Ophioglossaceae rather than with Santalales. These results cannot be attributed to contamination, which can be ruled out for these reasons: (i) DNA extractions of *B. virginianum* were done independently in the laboratories of Davis and Wurdack, prior to any extraction of Santalales; (ii) all samples of *B. virginianum* possessed the same transgenes and (iii) these transgenes are different from any Santalales we sampled—if contamination had occurred we would expect sequences from *B. virginianum* to match those of Santalales extracted in our labs, but they do not. The most reasonable explanation for our results is




Figure 1. Phylogenetic evidence for the discordant placements of *Botrychium virginianum*. Central tree depicts our current understanding of major land plant relationships. Each pair of trees in the corners is extracted from the larger ML analyses of (a) *nad1B-C* (from 100 total taxa), (b) *matR* (from 208 total taxa), (c) *atp1* (from 145 total taxa), and (d) *atp6* (from 74 total taxa). Santalales are depicted in red, Ophioglossaceae in blue. Transgenic copies of *matR* and *nad1B-C* are shown in Santalales in blue. Asterisks indicate the placement of *B. virginianum* in Ophioglossaceae based on vertically inherited copies of these genes; we did not recover a native copy of *nad1B-C*. See Electronic Appendix for detailed tree topologies and statistics.

that part of the genome in *B. virginianum* was acquired from Santalales via HGT. While other studies have reported HGT between angiosperms and gymnosperms (Won & Renner 2003) and between angiosperms and mosses (Berghthorsson *et al.* 2004), this is the first report of gene transfer between angiosperms and ferns.

The two transgenic regions found in *B. virginianum*, *nad1B-C* and *matR*, both reside within the *nad1* gene (Dombrowska & Qiu 2004). The similar phylogenetic placement of *B. virginianum* using each gene region indicates that transgenes in this species were most likely transferred together in a single event. The native *matR* gene that places *B. virginianum* in Ophioglossaceae appears to be functional, while the copy of *matR* that nests within Santalales is not. RT-PCR products and cDNA sequences were only recovered for native *matR*, and were absent for the Santalalean copy. The pseudogenetic nature of the latter is further confirmed by the presence of several internal termination codons and by the loss of reading frame in this sequence. The exonic region of *nad1B-C* was similarly undetected in RT-PCR/cDNA analysis suggesting that the coding portion of this transgene region is also non-functional.

Botrychium virginianum is a terrestrial fern common in temperate forests throughout the northern hemisphere and extends south through America in moist upland habitats to Bolivia and Brazil (Clausen 1938; Hauk *et al.* 2003). The *nad1B-C* and *matR* transgenic regions are restricted to *B. virginianum* within Ophioglossaceae and

were present in all 34 individuals we sampled across the range of *B. virginianum*. Therefore, HGT probably occurred after *B. virginianum* diverged from its closest relatives and before it expanded into its present global distribution. Ancestral area reconstructions (using DIVA and MACCLADE) and molecular divergence time estimates based on plastid *rbcL* data indicate that this divergence most likely occurred in Asia, and that the transfer event is no older than the Eocene (34.9 ± 3.5 Myr, or 44.2 ± 6 Myr; older maximum age constraints (e.g. 500 Myr) for the root node in our analysis push this age estimate slightly older, but the age of interest is still within the Eocene). Alternatively, the transfer event may have occurred in a single population of an already widespread *B. virginianum*, followed by spread of the transgenic regions throughout the species via selective gene sweep. Since neither of these transgene regions appears to be functional in *B. virginianum*, they are unlikely to confer a selective advantage, which favours the hypothesis that the transfer preceded the expansion of *B. virginianum*.

Two lines of evidence suggest that *B. virginianum* may have achieved its global distribution very rapidly, perhaps in thousands of years rather than millions. A rapid expansion would fit with the life history of this species. Like many ferns, including Ophioglossaceae, *B. virginianum* is almost certainly easily dispersed over long distances by its small spores (Peck *et al.* 1990; Barrington 1993). Additionally, the ability of single spores to establish new colonies through its inbreeding bisexual gametophytes

(Soltis & Soltis 1986) should also speed the spread of this species. Rapid expansion is similarly supported at the molecular level by pairwise mismatch distributions (Rogers & Harpending 1992) and Tajima's *D* (Tajima 1989) for all sampled individuals of these transgenes (analysed independently and in combination). When we pooled all samples and used mismatch distributions to infer population expansion within *B. virginianum* none of the datasets were able to reject a unimodal distribution (i.e. expanding population model), providing evidence for a rapid population expansion in *B. virginianum* (*matR* ($p=0.10$), *nad1B-C* ($p=0.59$) and combined *matR* plus *nad1B-C* ($p=0.94$)). Similarly, significantly negative values of Tajima's *D* provide evidence of rapidly expanding populations, and were suggested for *matR* ($D=-1.62$, $p=0.03$), and combined *matR* plus *nad1B-C* ($D=-1.82$, $p=0.02$), and were marginally insignificant for *nad1B-C* ($D=-1.35$, $p=0.07$).

Phylogenetic analyses of the *nad1B-C* and *matR* transgenes (analysed independently and in combination) place *B. virginianum* as sister to the hemiparasitic family Loranthaceae within the order Santalales (figure 2). While most Loranthaceae are aerial stem parasites, three monospecific genera (*Atkinsonia*, *Gaiadendron* and *Nuytsia*) are root parasites (Kuijt 1969), and both morphological (Feuer & Kuijt 1980) and molecular (Nickrent 2001; Nickrent 2002) evidence indicates that root parasitism is ancestral in the family. *B. virginianum* is a terrestrial fern that spends part of its life cycle as a subterranean gametophyte and juvenile sporophyte (Johnson-Groh *et al.* 2002), and the rhizome is hardly, if at all, emergent at maturity (Gifford & Foster 1989). While it follows that a root parasite is the most likely donor of the transgenes found in *B. virginianum*, none of the three root-parasitic lorch species are presently found in Asia. Given the evidence cited above that *B. virginianum* originated in Asia, and the fact that our analyses (figure 2) show that the transgenes are not sister to any extant genus of Loranthaceae, it seems most likely that the transgenic donor was a root-parasitic Asian lorch that is now extinct. Nevertheless, the Neotropical *Gaiadendron punctatum* (Ruiz & Pavón) G. Don may serve as a model for how this gene transfer could have occurred. *Gaiadendron* represents one of the earliest-diverging extant members of Loranthaceae (Kuijt 1963; Feuer & Kuijt 1980; Nickrent 2002), and is the only root-parasitic lorch sympatric with *B. virginianum* (TROPICOS 2005). While we do not know any report of a parasitic relationship between Loranthaceae and *B. virginianum*, root parasitism is very cryptic and would probably go unnoticed if such a relationship existed, and *Gaiadendron* is known to parasitize ferns in the mountains of Costa Rica, where *B. virginianum* occurs (Kuijt 1963; TROPICOS 2005).

Another possibility is that there was never a direct parasitic connection between *B. virginianum* and terrestrial Loranthaceae. Instead, HGT may have been indirect via a shared fungal symbiont. The subterranean gametophyte of *B. virginianum* lacks chlorophyll and must be infected by an endophytic fungus in order to grow (Gifford & Foster 1989). The intracellular fungus forms a mycorrhiza-like association with the fern gametophyte and transfers to it carbohydrates from nearby mycorrhizal photosynthetic plants, a relationship dubbed epiparasitism or mycoheterotrophy (Schmid & Oberwinkler 1994;




Figure 2. Phylogeny of Santalales based on combined *nad1B-C* and *matR*. Likelihood and parsimony bootstrap values, respectively, are given for those clades supported at greater than 50%. Asterisks indicate less than 50% bootstrap support. See Electronic Appendix for detailed tree topologies and statistics.

Smith & Read 1997). At maturity, the roots of the sporophyte of *B. virginianum* lack root hairs and depend on their fungal symbiont for water and minerals (Kovács *et al.* 2003). Mycorrhizae have been reported in Santalales (Landis *et al.* 2002), but have not been demonstrated in root-parasitic Loranthaceae, although all of the latter lack root hairs like many mycorrhizal plants (Kuijt 1963, 1969). If there was a fungal bridge between a terrestrial lorch and *B. virginianum*, the HGT postulated here may have been mediated by the fungus.

If fungi are functioning as a conduit for gene transfer, that may help to explain some of the many reported HGTs for which no mechanism has been obvious (Bergthorsson *et al.* 2003; Won & Renner 2003). Mycorrhizal fungi are notoriously non-specific in their host selection and connect many distantly related plants in the same community (Smith & Read 1997). This 'wood-wide web' (Simard *et al.* 1997) may be facilitating rapid and widespread exchange of DNA across phylogenetic distances spanning all green plants. If so, such a finding would affect our thinking about the long-term evolution of the terrestrial biota, and may also have commercial implications. Fungus-mediated HGT could make it very difficult to restrict transgenes to their genetically modified organisms—if this is happening, the transgenic would be out of the bottle.

We thank B. Carstens, J. Hall, W. Hauk, J. Johnson, M. Latvis, K. Lewis, L. Knowles, D. Mindell, R. Moran, J. Palmer and P. Tucker. C.D. was supported by NSF AtoL EF 04-31242 and by the Michigan Society of Fellows. The curators of the following herbaria kindly permitted us to remove leaf material from specimens: A, GH, M, MICH, NY, US, Z. This paper is dedicated to W. H. Wagner and S. Wonder.

REFERENCES

- APG 2003 An update of the angiosperm phylogeny group classification for the orders and families of flowering plants: APG II. *Bot. J. Linn. Soc.* **141**, 399–436. (doi:10.1046/j.1095-8339.2003.t01-1-00158.x.)
 Barrington, D. S. 1993 Ecological and historical factors in fern biogeography. *J. Biogeogr.* **20**, 275–280.

- Berggren, W. A., Kent, D. V., Swisher II, C. C. & Aubry, M. P. 1995 A revised Cenozoic geochronology and chronostratigraphy. In *Geochronology, time scales and global stratigraphic correlation* (ed. W. A. Berggren, D. V. Kent, M. P. Aubry & J. Hardenbol), pp. 129–212. Tulsa: SEPM (Society for Sedimentary Geology).
- Berghthorsson, U., Adams, K. L., Thomason, B. & Palmer, J. D. 2003 Widespread horizontal transfer of mitochondrial genes in flowering plants. *Nature* **424**, 197–201. (doi:10.1038/nature01743.)
- Berghthorsson, U., Richardson, A. O., Young, G. J., Goertzen, L. R. & Palmer, J. D. 2004 Massive horizontal transfer of mitochondrial genes from diverse land plant donors to the basal angiosperm *Amborella*. *Proc. Natl Acad. Sci. USA* **101**, 17 747–17 752. (doi:10.1073/pnas.0408336102.)
- Clausen, R. T. 1938 A monograph of the Ophioglossaceae. *Mem. Torrey Bot. Club* **19**, 1–177.
- Davis, C. C. & Wurdack, K. J. 2004 Host-to-parasite gene transfer in flowering plants: phylogenetic evidence from Malpighiales. *Science* **305**, 676–678. (doi:10.1126/science.1100671.)
- Davis, C. C., Bell, C. D., Mathews, S. & Donoghue, M. J. 2002 Laurasian migration explains Gondwanan disjunctions: evidence from Malpighiaceae. *Proc. Natl Acad. Sci. USA* **99**, 6833–6837. (doi:10.1073/pnas.102175899.)
- Dombrowska, O. & Qiu, Y. L. 2004 Distribution of introns in the mitochondrial gene *nad1* in land plants: phylogenetic and molecular evolutionary implications. *Mol. Phylogenet. Evol.* **32**, 246–263. (doi:10.1016/j.ympev.2003.12.013.)
- Feuer, S. M. & Kuijt, J. 1980 Fine structure of mistletoe pollen. III. Large-flowered neotropical Loranthaceae and their Australian relatives. *Am. J. Bot.* **67**, 34–50.
- Gifford, E. M. & Foster, A. S. 1989 *Morphology and evolution of vascular plants*. New York: W. H. Freeman and Co.
- Guindon, S. & Gascuel, O. 2003 A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.* **52**, 696–704. (doi:10.1080/10635150390235520.)
- Harpending, H. C. 1994 Signature of ancient population growth in a low-resolution mitochondrial DNA mismatch distribution. *Hum. Biol.* **66**, 591–600.
- Hauk, W. D., Parks, C. R. & Chase, M. W. 2003 Phylogenetic studies of Ophioglossaceae: evidence from *rbcl* and *trnL-F* plastid DNA sequences and morphology. *Mol. Phylogenet. Evol.* **28**, 131–151. (doi:10.1016/S1055-7903(03)00032-0.)
- Huelsenbeck, J. P., Hillis, D. M. & Jones, R. 1996 Parametric bootstrapping in molecular phylogenetics: applications and performance. In *Molecular zoology: advances, strategies, and protocols* (ed. J. D. Ferris & S. R. Palumbi), pp. 19–45. New York: Wiley-Liss.
- Johnson-Groh, C., Riedel, C., Schoessler, L. & Skogen, K. 2002 Belowground distribution and abundance of *Botrychium* gametophytes and juvenile sporophytes. *Am. Fern J.* **92**, 80–92.
- Kadowaki, K.-I., Kubo, N., Ozawa, K. & Hirai, A. 1996 Targeting presequence acquisition after mitochondrial gene transfer to the nucleus occurs by duplication of existing targeting signals. *EMBO J.* **15**, 6652–6661.
- Karol, K. G., McCourt, R. M., Cimino, M. T. & Delwiche, C. F. 2001 The closest living relatives of land plants. *Science* **294**, 2351–2353. (doi:10.1126/science.1065156.)
- Kishino, H. & Hasegawa, M. 1989 Evaluation of the maximum likelihood estimates of the evolutionary tree topologies from sequence data, and the branching order in Hominoidea. *J. Mol. Evol.* **29**, 170–179.
- Kovács, G. M., Kottke, I. & Oberwinkler, F. 2003 Light and electron microscopic study on the mycorrhizae of sporophytes of *Botrychium virginianum*—arbuscular structures resembling fossil forms. *Plant Biol.* **5**, 574–580. (doi:10.1055/s-2003-44786.)
- Kuijt, J. 1963 On the ecology and parasitism of the Costa Rican tree mistletoe, *Gaiadendron punctatum* (Ruiz and Pavón) G. Don. *Canad. J. Bot.* **41**, 927–938.
- Kuijt, J. 1969 *The biology of parasitic flowering plants*. Berkeley: University of California Press.
- Landis, F., Gargas, A. & Givnish, T. 2002 *The plant tree, roots and clades: Mycorrhizae and plant phylogeny*. Annual Meeting of the Botanical Society of America 2002, Madison, Wisconsin p. 174. See <http://www.botany2002.org/section13/abstracts/4.shtml>.
- Malécot, V., Nickrent, D. L., Baas, P., Oever, L. v. d. & Lobreau-Callen, D. 2004 A morphological cladistic analysis of Olacaceae. *Syst. Bot.* **29**, 569–586. (doi:10.1600/0363644041744301.)
- Mower, J. P., Stefanovic, S., Young, G. J. & Palmer, J. D. 2004 Gene transfer from parasitic to host plants. *Nature* **432**, 165–166. (doi:10.1038/432165b.)
- Nickrent, D. L. 2001 Santalales (Mistletoe); treatment A3714. In *Encyclopedia of life sciences*. New York: Macmillan Publishers Ltd.
- Nickrent, D. L. 2002 Mistletoe phylogenetics: current relationships gained from analysis of DNA sequences. In *Proc. 48th Annual Western Int. Forest Disease Work Conf.* (ed. P. Angwin), pp. 48–57. Redding: USDA Forest Service.
- Nickrent, D. L. & Malécot, V. 2001 A molecular phylogeny of Santalales. In *Proc. 7th Int. Parasitic Weed Symp.* (ed. A. Fer, P. Thalouarn, D. M. Joel, L. J. Musselman, C. Parker & J. A. C. Verkleij), pp. 69–74. Nantes, France: Faculté des Sciences, Université de Nantes.
- Peck, J. H., Peck, C. J. & Farrar, D. R. 1990 Influence of life history attributes on formation of local and distant fern populations. *Am. Fern J.* **80**, 126–142.
- Posada, D. & Crandall, K. A. 1998 Modeltest: testing the model of DNA substitution. *Bioinformatics* **14**, 817–818. (doi:10.1093/bioinformatics/14.9.817.)
- Pryer, K. M., Schneider, H., Smith, A. R., Cranfill, R., Wolf, P. G., Hunt, J. S. & Sipes, S. D. 2001 Horsetails and ferns are a monophyletic group and the closest living relatives to seed plants. *Nature* **409**, 618–622. (doi:10.1038/35054555.)
- Rogers, A. 1995 Genetic evidence for a Pleistocene population explosion. *Evolution* **49**, 608–615.
- Rogers, A. R. & Harpending, H. 1992 Population growth makes waves in the distribution of pairwise genetic differences. *Mol. Biol. Evol.* **9**, 552–569.
- Ronquist, F. 1997 Dispersal–vicariance analysis: a new approach to the quantification of historical biogeography. *Syst. Biol.* **46**, 195–203.
- Rothwell, G. W. & Stockey, R. A. 1989 Fossil Ophioglossales in the Paleocene of Western North America. *Am. J. Bot.* **76**, 637–644.
- Sanderson, M. J. 2002 Estimating absolute rates of molecular evolution and divergence times: a penalized likelihood approach. *Mol. Biol. Evol.* **19**, 101–109.
- Sanderson, M. J. 2003 Molecular data from 27 proteins do not support a Precambrian origin of land plants. *Am. J. Bot.* **90**, 954–956.
- Schmid, E. & Oberwinkler, F. 1994 Light and electron microscopy of the host–fungus interaction in the achlorophyllous gametophyte of *Botrychium lunaria*. *Canad. J. Bot.* **72**, 182–188.
- Schneider, S., Roessli, D. & Excoffier, L. 2000 Arlequin v2.0: Documentation and program. <http://anthro.unige.ch/arlequin>.
- Schneider, H., Schuettpelz, E., Pryer, K. M., Cranfill, R., Magallón, S. & Lupia, R. 2004 Ferns diversified in the shadow of angiosperms. *Nature* **428**, 553–557. (doi:10.1038/nature02361.)

- Simard, S. W., Perry, D. A., Jones, M. D., Myrold, D. D., Durall, D. M. & Molina, R. 1997 Net transfer of carbon between ectomycorrhizal tree species in the field. *Nature* **388**, 579–582. (doi:10.1038/41557.)
- Smith, S. E. & Read, D. J. 1997 *Mycorrhizal symbiosis*. London: Academic Press.
- Soltis, D. E. & Soltis, P. S. 1986 Electrophoretic evidence for inbreeding in the fern *Botrychium virginianum* (Ophioglossaceae). *Am. J. Bot.* **73**, 588–592.
- Swofford, D. L. 2003 *PAUP*: phylogenetic analysis using parsimony (*and other methods)*, v. 4.0.610. Sunderland, MA: Sinauer Associates.
- Tajima, F. 1989 The effect of change in population size on DNA polymorphism. *Genetics* **123**, 597–601.
- TROPICOS. 2005. Database of the Missouri Botanical Garden; <http://mobot.mobot.org/W3T/Search/vast.html>.
- Vidal-Russell, R. & Nickrent, D. 2005 *A molecular phylogeny of the mistletoe family Loranthaceae*. Annual Meeting of the Botanical Society of America 2005 Austin, Texas p. 101. (See <http://www.2005.botanyconference.org>)
- Wagner, W. H. & Wagner, F. S. 1993 Ophioglossaceae Agardh. In *Pteridophytes and gymnosperms* (ed. F.O.N.A.E. Committee), vol. 2, pp. 102–105. New York: Oxford University Press.
- Won, H. & Renner, S. S. 2003 Horizontal gene transfer from flowering plants to Gnetum. *Proc. Natl Acad. Sci. USA* **100**, 10 824–10 829. (doi:10.1073/pnas.1833775100.)

The supplementary Electronic Appendix is available at <http://dx.doi.org/10.1098/rspb.2005.3226> or via <http://www.journals.royalsoc.ac.uk>.