

The distributions of the duplicate oestrogen receptors ER- β a and ER- β b in the forebrain of the Atlantic croaker (*Micropogonias undulatus*): evidence for subfunctionalization after gene duplication

M. B. Hawkins^{1,*}, J. Godwin², D. Crews³ and P. Thomas¹

¹Marine Science Institute, University of Texas at Austin, Port Aransas, TX 78373, USA

²Department of Zoology, North Carolina State University, PO Box 7617, Raleigh, NC 27695, USA

³Patterson Labs, Department of Zoology, University of Texas at Austin, Austin, TX 78712, USA

Teleost fishes have three distinct oestrogen receptor (ER) subtypes: ER- α , ER- β a (or ER- γ) and ER- β b. ER- β a and ER- β b arose from a duplication of an ancestral ER- β gene early in the teleost lineage. Here, we describe the distribution of the three ER mRNAs in the hypothalamus and cerebellum of the Atlantic croaker to address two issues: the specific functions of multiple ERs in the neuroendocrine system and the evolution and fate of duplicated genes. ER- α was detected in nuclei of the preoptic area (POA) and hypothalamus previously shown to possess ER- α s in teleosts. AcER- β b, but not ER- β a, labelling was detected in the magnocellular neurons of the POA, nucleus posterior tuberis, the nucleus recessus posterior and cerebellum. By contrast, acER- β a, but not ER- β b, was detected in the dorsal anterior parvocellular POA and suprachiasmatic nucleus. Both ER- β s were found in posterior parvocellular and ventral anterior POA nuclei, the ventral hypothalamus, and periventricular dorsal hypothalamus. The differences we observed in ER subtype mRNA distribution within well-characterized brain nuclei suggest that ER- β a and ER- β b have distinct functions in the neuroendocrine control of reproduction and behaviour, and provide evidence that the teleost ER- β paralogues have partitioned functions of the ancestral ER- β gene they shared with tetrapods.

Keywords: oestrogen receptor; gene duplication; teleost fishes; neuroendocrine regulation; hypothalamus; brain

1. INTRODUCTION

Steroid hormone receptors are members of a large family of ligand-activated nuclear transcription factors that are critical to the reproduction, differentiation and development of vertebrates. All steroid receptors, including androgen, progesterone, glucocorticoid and mineralcorticoid receptors, are derived from an ancient oestrogen receptor (ER) that duplicated multiple times during early vertebrate evolution (Thornton 2001).

The primary physiological ligand for ERs is the sex steroid oestradiol-17 β (oestradiol, E₂). Oestradiol, synthesized in the gonads or locally in the brain by aromatization of testosterone, plays an important role in the neuroendocrine control of reproduction and reproductive behaviour in vertebrates (Knobil & Neill 1994). Oestradiol acts in the pituitary and the brain to influence the secretion of gonadotropin-releasing hormone (GnRH) and other neuropeptides that control reproduction and behaviour (Gore 2002; Shupnik 2002). In addition to nuclear ERs, there is evidence for membrane localization of ERs and novel ERs in

brain tissues (Blaustein 2004; Thomas *et al.* 2004; Toran-Allerand 2004).

Despite intense focus on oestrogen actions, the presence of more than one ER subtype escaped notice for decades. The discovery of a second ER subtype, (ER- β Kuiper *et al.* 1996), in addition to the initially described ER- α , has expanded the potential mechanisms by which oestrogens can elicit their effects. Teleost fishes have two types of ER- β : ER- β a (formerly ER- γ ; see Hawkins *et al.* 2000; Hawkins & Thomas 2004) and ER- β b (Menuet *et al.* 2002) as well as an ER- α . The two ER- β subtypes arose from the duplication of an ancestral ER- β gene early in the teleost lineage after the split of tetrapods and fishes (Hawkins *et al.* 2000).

The fate of genes after a duplication event is of fundamental importance in evolutionary biology because it is believed that this process is the predominant mechanism through which novel proteins evolve and new species arise (Ohno 1970; Amores *et al.* 1998). After a duplication event, genes are subject to mutations that result in three possible outcomes: gene silencing of one member of the pair (Mighell *et al.* 2000), the acquisition of a novel function by one copy (neofunctionalization; Ohno 1970) or the partitioning of the original functions between the two copies (subfunctionalization; Force *et al.* 1999). The two ER- β subtypes of teleosts

* Author for correspondence (beth_hawkins@ncsu.edu).

† Present address: Department of Zoology, North Carolina State University, PO Box 7617, Raleigh, NC 27695, USA.

provide an excellent system to investigate the fate of duplicated genes in vertebrate genomes, because ERs are such critical and extensively studied proteins (Postlethwait *et al.* 2004). Our sequence and phylogenetic analyses showed that after the duplication event, *ER-βa* underwent rapid change followed by strict conservation of certain amino acids (Hawkins *et al.* 2000). This suggested the acquisition of novel functions, or neofunctionalization, after the duplication event. In addition, *ER-βa* has unusual binding affinities for the endogenous steroid oestriol and other oestrogenic compounds, further suggesting novel functions for this teleost-specific subtype (Hawkins & Thomas 2004).

In this study, we describe the distribution of *ER-α*, *ER-βa* and *ER-βb* mRNA in the hypothalamus and cerebellum of the Atlantic croaker. This is an important first step toward characterizing the role of ER subtypes in the neuroendocrine regulation of reproduction in this well-established model species (Khan & Thomas 1999, 2001; Khan *et al.* 1999). In addition, comparing the distribution of *ER-βa* and *ER-βb* in teleosts to that of *ER-β* in tetrapods could provide insights into the functional evolution of duplicated genes.

In mammals and birds, the distribution of *ER-β* partially overlaps that of *ER-α* in the brain, but some nuclei express *ER-β* exclusively. These include the paraventricular nuclei of the preoptic area (POA), the suprachiasmatic nucleus, the tuberal hypothalamic nuclei and the cerebellum (Shughrue *et al.* 1997; Bernard *et al.* 1999; Foidart *et al.* 1999; Hileman *et al.* 1999; Mitra *et al.* 2003; Mercantheiler *et al.* 2004). By contrast, *ER-α* mRNA is found exclusively in the ventromedial hypothalamus and the subfornical organ (Shughrue *et al.* 1997).

Little information is currently available on the distribution of the three ER subtypes in the teleost brain. *ER-α* protein and mRNA distribution have been mapped in the forebrain of rainbow trout (*Oncorhynchus mykiss*; Kah *et al.* 1997; Menuet *et al.* 2003), but the neuroanatomical distribution of a possible *ER-βa* or *ER-βb* has not been investigated in this species. Our preliminary studies showed that *ER-α*, *ER-βa* and *ER-βb* mRNA expression patterns differ in the suprachiasmatic nucleus of the POA of Atlantic croaker *Micropogonias undulatus* (Hawkins *et al.* 2000). Similarly, all three subtypes show partially overlapping distributions in the anterior preoptic area, the ventral hypothalamus and the posterior tuberculum of the zebrafish *Danio rerio* (Menuet *et al.* 2002). These differences in distribution suggest that the three ER subtypes have distinct roles in the reproductive physiology and behaviour of fishes. It also raises the possibility that the functions of *ER-βa* and *ER-βb* have diverged from their ancestral *ER-β* gene. A more detailed description of the distribution of the three subtypes will begin to address their specific functions in the neuroendocrine system as well as the more general issue of the evolution and fate of duplicated genes.

2. MATERIAL AND METHODS

(a) *Animal and tissue collection*

Adult female Atlantic croakers were collected from the waters surrounding Port Aransas, TX, USA, and maintained

in tanks at the University of Texas Marine Science Institute under natural cycles of light and salinity. Six fishes with regressing gonads were injected intraperitoneally with E_2 at a concentration (1 mg kg^{-1}) that results in high physiological levels of E_2 and upregulation of ER in fishes (Hawkins *et al.* 2000). Fishes were deeply anaesthetized 48 h later in a seawater : phenoxy ethanol bath (1 : 2000) and quickly killed by severing the spinal cord. Brains were removed and frozen in dry ice-chilled isopentane within 2 min. Brains were stored at -70°C until they were sectioned.

(b) *Creation of sequence-specific ER probes in Atlantic croaker*

The three ER constructs used as probes were designed and prepared in the same manner as described previously (Hawkins *et al.* 2000). The Atlantic croaker *ER-α*, *ER-βa* and *ER-βb* probes encompassed amino acids 177–293, 266–406 and 443–552, respectively. [^{35}S]CTP-labelled antisense riboprobes (New England Nuclear) were transcribed from each ER cDNA subclone.

(c) *In situ hybridization*

Frozen whole brains were embedded in optimal cutting temperature (OCT) compound (Tissutek) and cryosectioned at $20 \mu\text{m}$ (Frigocut, Reichert-Jung). Consecutive sections were transferred to a series of six poly-L-lysine-treated slides to localize the ER subtypes relative to each other in the same brain area. *In situ* hybridizations were conducted as described previously (Young *et al.* 1994; Hawkins *et al.* 2000).

After development, slides were visualized under dark field illumination to assess labelling in specific areas. In regions of low specific labelling or high background (cerebellum, NRL, PMm), positive labelling was confirmed via analysis with the grain counting software BRAIN as described previously (Young *et al.* 1994). Areas with silver grain density greater than $3\times$ background were considered positively labelled. Four different animals were assessed for each area with the exception of the suprachiasmatic nucleus ($n=3$).

(d) *Nomenclature*

The terminology of Braford & Northcutt is followed in the text (Braford & Northcutt 1983), except in some cases where the equivalent terms of Peter & Gill are more specific (Peter *et al.* 1975). The terminology and abbreviations of both research groups are given in table 1.

3. RESULTS

(a) *The anterior hypothalamus/preoptic area*

The distribution of *ER-α*, *ER-βa* and *ER-βb* mRNA labelling in the croaker brain is summarized and compared with findings for other teleost species in table 1. Hybridization signals for the three forms of ER exhibited some overlap in the POA but also showed several areas with marked differences. Most rostrally in the anterior preopticus parvocellularis (PPa), *ER-α* showed strong labelling in both the ventral and dorsal POA regions, whereas *ER-βb* showed more restricted labelling and *ER-βa* demonstrated weaker labelling (figure 1*b–d*). Interestingly, *ER-βb* labelling was restricted to the ventral portion of the PP nucleus (PPa_v), while some *ER-βa* labelling was also observed dorsal to the *ER-βb* labelling (PPa_d; figure 1*c,d*). Proceeding caudally, all the ER probes

Table 1. The distribution of Atlantic croaker (ac)ER- α , acER- β a and acER- β b mRNA in the brain compared with the distribution of ERs and E2-concentrating cells in other teleosts. (Terminology of Peter & Gill (Peter *et al.* 1975) is given in italicized parentheses. The oyster toadfish (OTF; Fine *et al.* 1990), paradise fish (PF; Davis *et al.* 1977), goldfish (GF; Kim *et al.* 1979) and platyfish (PLF; Kim *et al.* 1979) tubular autoradiography data of E₂-concentrating cells were taken in part from Fine *et al.* (1990). Autoradiography data do not differentiate ER subtypes. The zebrafish (zf) ER mRNA data are from Meneuet *et al.* (2002). The existence of few cells is labelled (.). Superscript v indicates ventral labelling; superscript d indicates dorsal labelling; superscript 1 indicates rainbow trout (RT) ER- α mRNA (Kah *et al.* 1997); superscript 2 is (RT) ER- α antibody (Navas *et al.* 1995; Linard *et al.* 1996); superscript a means anatomical designation is not given in the original reference, but oestrogen-concentrating cells are shown in appropriate areas (Fine *et al.* 1990); superscript b means that no distinction was made between parvo and magnocellular (Fine *et al.* 1990); superscript c means designated n. saccus vasculosus in the original reference and a question mark identifies an area not investigated.)

	ACERs			ZfERs			RT	GF	OTF	PF	PLF
	α	β a	β b	α	β a	β b					
preoptic area (POA)											
n. preopticus parvocellularis anterior (PPa)	+ ^{v,d}	+ ^{(v)d}	+ ^v	+ ^v	+ ^{v,(d)}	+ ^{v,d}	+ ^{1,2}	+	+	+	+ ^a
(= n. <i>preopticus periventricularis</i> , NPP)											
n. preopticus magnocellularis pars parvocellularis (PMp) (= n. <i>preopticus parvocellularis</i> , NPOp)	+	+	?	?	?	?	(+) ¹	+ ^b	(+)	+	+
n. preopticus magnocellularis pars magnocellularis (PMm)	+	-	+	?	?	?	(+) ^{1,2}	+ ^b	(+)	(+)	+
(= n. <i>preopticus magnocellularis</i> , NPOm)											
n. preopticus parvocellularis posterioris (PPp)	+	-	-	?	?	?	+ ¹	-	(+)	-	-
(= n. <i>anterior hypothalami periventricularis</i> , NAPv)											
n. suprachiasmaticus (SCN)	+	+	-	?	?	?	- ²	?	-	?	?
ventral hypothalamus (HV):											
(n. <i>lateralis tubercis pars anterior</i> , NLTa)	+	+	+	?	?	?	?	(+)	(+)	(+)	-
(n. <i>lateralis tubercis pars posterior</i> , NLTp)	+	+	+	?	?	?	+ ^{1,2}	+	+	+	+
(n. <i>lateralis tubercis pars inferior</i> , NLTi)	+	+	+	?	?	?	+ ²	+	+	+	+
(n. <i>lateralis tubercis pars lateralis</i> , NLTL)	+	+	+	?	?	?	+ ^{1,2}	?	?	?	?
n. anterior tubercis (TA)	+	+	+	?	?	?	+ ²	-	(+)	-	-
n. lateralis hypothalami (LH)	+	+	+	+	+	-	?	-	+	-	-
dorsal hypothalamus (Hd):											
(n. <i>posterioris periventricularis</i> , NPPv)	+	+	+	?	?	?	(+) ^{1,2}	+	+	(+)	+ ^a
(n. <i>recessus lateralis</i> , NRL)	+	-	(+)	?	?	?	+ ¹	-	+	+	+
caudal hypothalamus (Hc):											
(n. <i>recessus posterior</i> , NRP)	+	-	-	?	?	?	(+) ^{1,2}	-	-	-	-
posterior tuberculum											
n. tubercis posterior (TP)	+	-	+	+	-	+	+ ^{2c}	- ^c	+ ^c	- ^c	+ ^c
valvular cerebelli (VC)	-	-	+	?	?	?	?	?	?	?	?
corpus cerebellum (CC)	-	-	+	?	?	?	?	?	?	?	?




Figure 1. ER subtype mRNA expression patterns differ in the n. preopticus parvocellularis anterior (PPa) of the Atlantic croaker. (a) Arrows indicate the location of the dorsal PPa_d and ventral PPa_v in relation to the third ventricle (3V) in the most rostral portion of the preoptic area. (b) acER- α labelling is seen in both the PPa_d and PPa_v. (c) acER- β b labelling is detected only in the PPa_v. (d) acER- β a labelling is detected in the PPa_d, and there is some labelling in the PPa_v.

labelled the parvocellular portion of the magnocellular POA (PMp; table 1). More caudal and dorsal in the magnocellular nucleus (PMm), many large neurosecretory neurons were consistently and strongly labelled with the ER- β b probe, while ER- β a only very weakly labelled (less than $3\times$ background) a few of these cells in some sections (figure 2c, d). In contrast to the patterns for ER- β b, the ER- α probe labelling over large neurons was more diffuse and was not discernibly stronger than that in the immediately surrounding tissue (figure 2b). The PPa is the most caudal nucleus of the POA and is arranged in laminae 2–5 cells thick. The PPa was labelled with ER- α




Figure 2. ER subtype mRNA expression patterns differ in the n. preopticus magnocellularis pars magnocellularis (PMm) of the Atlantic croaker. (a) Arrows indicate the location of large neurosecretory neurons along the third ventricle (3V). (b) acER- α labelling in the PMm is diffuse over and around large neurons. (c) acER- β b labelling is strong over magnocellular perikarya. (d) The acER- β a labelling is less than $3\times$ background.

probe but not with the ER- β a or ER- β b probe. ER- α and ER- β a labelling were seen in the suprachiasmatic nucleus as reported previously (Hawkins *et al.* 2000).

(b) *The hypothalamus and posterior tuberculum*

Most nuclei of the ventral and dorsal hypothalamus showed labelling for all three ERs (table 1). There were two notable exceptions: we found no ER- β a or ER- β b labelling in the NRP or the NRL. In the posterior tuberculum, the posterior tuberal nucleus (NPT) located dorsal to the ventricle was strongly labelled with the ER- β b probe (figure 3c). However, this nucleus was negative for ER- β a (figure 3d).




Figure 3. ER subtype mRNA expression patterns differ in the n. tuberis posterior (TP) of the posterior tuberculum of the Atlantic croaker. (a) Arrows indicate the location of the TP dorsal to the caudal portion of the third ventricle (3V) and the n. lateralis tuberis (NLT). (b) The acER- α probe diffusely labels the TP. (c) Strong acER- β b labelling is detected in the TP. (d) acER- β a labelling is not detected in the TP.

(c) The cerebellum

In the cerebellum, strong ER- β b labelling was observed in a bead-like arrangement in what appears to be the Purkinje cell layer (figure 4). Similar labelling was also observed in the valvular cerebelli. Neither the ER- α nor ER- β a probes labelled these regions.

4. DISCUSSION

Major differences were seen in the distribution of ER- β a and ER- β b subtypes in the parvo- and magnocellular preoptic areas, the suprachiasmatic nucleus, the posterior




Figure 4. Distribution of acER mRNA in the corpus cerebellum (CC) of the Atlantic croaker. (a) Arrows indicate the location of the Purkinje cell layer. (b) acER- α labelling is less than $3\times$ background. 'VC' is the valvular cerebelli. (c) The acER- β b riboprobe is specifically labelling Purkinje cells of the cerebellum. (d) acER- β a labelling is less than $3\times$ background.

tuberis and the cerebellum of the Atlantic croaker brain. The PPa of the POA showed strong labelling with ER- β b in the ventral portion but no dorsal labelling, whereas ER- β a showed moderate labelling in both regions. There were also high levels of ER- β b expression in the PMm portion of the POA, but little or no ER- β a expression. In addition, ER- β b was expressed in the NPT and the cerebellum, while ER- β a was not detected in these areas. The only nucleus found to express ER- β a exclusively was the suprachiasmatic nucleus of the preoptic area. However, ER- β a distribution overlapped with that of ER- α and ER- β b in many areas (table 1).

The present study found that magnocellular neurons of the PM portion of the POA (PMm) express high amounts of ER- β b mRNA but little or no ER- β a or ER- α mRNA. This agrees with findings in mammals and birds, where

the magnocellular region of the paraventricular nucleus (PVN) expresses ER- β , but not ER- α (Shughrue *et al.* 1997; Bernard *et al.* 1999; Foidart *et al.* 1999; Hileman *et al.* 1999; Mitra *et al.* 2003; Mercanteller *et al.* 2004). ER- β is coexpressed in magnocellular neurons of the mammalian PVN that also produce and release arginine vasopressin (AVP) or oxytocin (OT) (Hrabovsky *et al.* 2004). These neurohypophysial hormones are involved in modulating reproductive behaviours in vertebrates (Moore 1992). Oestrogen treatment alters AVP and OT mRNA levels in the PVN of mice but has no effect in ER- β knockout mice (Nomura *et al.* 2002; Patisaul *et al.* 2003). It is likely that some of the cells expressing ER- β in croaker are also AVT or OT neurons.

Some neurons of the rat POA that express ER- β produce GnRH, a neuropeptidergic hormone which is central to the control of vertebrate reproductive function and behaviour and whose release and production is modulated by E₂ (Herbison & Pape 2001; Hrabovszky *et al.* 2001; Gore 2002). ER- α is not expressed in GnRH neurons, so until the recent discovery of ER- β in mammalian GnRH neurons (Herbison & Pape 2001), it was thought that in mammals and fishes, all of the oestrogenic effects on GnRH were indirect via interneurons that are regulated by E₂ and whose neurons possess ERs (Flugge *et al.* 1986; Linard *et al.* 1996; Herbison 1998; Senthilkumaran *et al.* 2001). However, the recent demonstrations that GnRH neurons express ER- β in mammals suggest that E₂ could have direct, as well as indirect, effects on GnRH regulation. The presence of ER- β a and ER- β b in the POA of croaker reopens the possibility that GnRH neurons in fishes may express an ER- β . This finding would suggest that ER expression in GnRH neurons is an ancient feature of vertebrate neuroendocrinology.

ER- α and ER- β b, but not ER- β a mRNA was detected in the NTP of the posterior tuberculum. These findings agree with those for zebrafish (Menuet *et al.* 2002). The TP as reported here appears to be at least in part the nucleus saccus vasculosus (SV) of Braford & Northcutt (1983). This nucleus is more appropriately named the TP because it does not innervate the SV organ as originally thought, but instead innervates the more rostral nucleus tuberis posterior of the posterior tuberculum. Oestradiol binding was detected here in oyster toadfish and platyfish (Kim *et al.* 1979; Fine *et al.* 1990). ER- α protein was also found in the nucleus SV of rainbow trout (Kah *et al.* 1997). This nucleus contains dopaminergic neurons and may be analogous to the nigrostriatal dopaminergic system of tetrapods (Rink & Wullimann 2001). In mammals, these dopaminergic neurons are responsive to oestrogen, and E₂ protects them from degeneration induced by the neurotoxins 6-hydroxydopamine and 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP; Arvin *et al.* 2000; Murray *et al.* 2003).

High levels of ER- β b mRNA were detected in the Purkinje cells of the cerebellum, while no ER- α or ER- β a labelling was observed. ER- β b was also detected in the goldfish and bird cerebellum by RT/PCR (Bernard *et al.* 1999; Foidart *et al.* 1999; Ma *et al.* 2000), and ER- β , but not ER- α , mRNA is present in Purkinje cells of the rat and mouse cerebellum (Price & Handa 2000; Mitra *et al.* 2003). In addition, ER- β may be involved in the E₂-induced

changes in transcription of GABA_A receptors in rats (Barami *et al.* 1993). The presence of an ER- β in the cerebellum of species in three different vertebrate classes suggests an ancient function for this subtype in motor learning and muscle coordination. GnRH and 5-HT fibres were also detected in the cerebellum of the Atlantic croaker, but their function here is unknown (Khan & Thomas 1993).

The distribution of E₂-concentrating cells in the brain is extremely conserved across species (Pfaff *et al.* 1988). In mammals, this conservation basically extends to ER- α and ER- β distribution (Shughrue *et al.* 1997; Hileman *et al.* 1999; Mitra *et al.* 2003; Mercanteller *et al.* 2004). In addition to our results presented here for croaker, there is one other study investigating the three ER subtypes' distribution in the brain of zebrafish (Menuet *et al.* 2002). Our results now provide a framework for comparing ER subtype distribution across fish species as well as across other vertebrate taxa. For example, in the PPa, we found ER- α and ER- β a in both the dorsal and ventral regions of the nucleus, while ER- β b was restricted to the ventral portion (table 1). In zebrafish, there is also a distinct, yet overlapping distribution of the three subtypes in the PPa. However, in this species ER- β 1 (ER- β b) and ER- β 2 (ER- β a) are found in both regions, whereas ER- α is restricted to the ventral region (Menuet *et al.* 2002). In the Hv we found all three subtypes in the nuclei investigated. However, zebrafish do not express ER- β b in the Hv. In the NPT, croaker and zebrafish have similar distributions of ERs. They show a more diffuse labelling with ER- α and very restricted labelling with ER- β b, while this region is negative for ER- β a in both species. Thus, while both species differentially express ER subtypes in many brain regions, the specific ER subtypes expressed in these areas are not conserved. A comparison incorporating more species is needed to assess the general conservation of ER subtype distributions among teleosts.

Since the ER subtypes were often found together in the same brain nuclei, it is possible that they colocalize to the same cells. If the ER subtypes are coexpressed, they could have either the same functions in these areas, or they could have different roles in the E₂-controlled regulation of physiology and behaviour. There is evidence that there are regulatory differences between ER- α and ER- β in mammals and fishes (Paech *et al.* 1997; Menuet *et al.* 2002; Schultz *et al.* 2002). Evidence from rats and humans indicate that ERs may form heterodimers, which may result in distinct interactions with DNA and transcriptional cofactors, and cause alternative gene expression patterns (Pace *et al.* 1997; Pettersson *et al.* 1997; Ogawa *et al.* 1998). The three croaker ERs have different affinities for E₂ and oestriol, which could allow for different levels of receptor activation in a given cell (Hawkins & Thomas 2004). Our recent findings have shown that the three ERs have distinct binding profiles for phytoestrogens and xenoestrogens (Hawkins & Thomas 2004, unpublished observations). Given the differential distribution of these receptor subtypes within the CNS, environmental exposure to oestrogenic substances may have profoundly different effects on neuroendocrine function.

ER- β a has the most restricted distribution of the three subtypes, suggesting that it has very specific functions within the brain. ER- β a (as well as ER- β b) appears to have arisen in a duplication of the ancestral ER- β early in the teleost

lineage (Hawkins *et al.* 2000). Structural comparisons and phylogenetic analyses among teleost ER- β s indicate that ER- β a quickly diverged from ER- β b after the duplication, suggesting the acquisition of novel functions by ER- β a. The exclusive presence of ER- β a in a region lacking ER- β s in other species would lend further support to the hypothesis that neofunctionalization occurred after the duplication event. Alternatively, ER- β a and ER- β b may have partitioned the ancestral ER- β functions within tissues (subfunctionalization). In croaker, the only brain areas investigated that express ER- β a and not ER- β b are the SCN (table 1; Hawkins *et al.* 2000) and the dorsal PPa. Conversely, ER- β b, but not ER- β a, was found in the ventral PPa, the magnocellular neurons of the POA and the Purkinje cells of the cerebellum. In rats and mice, both the SCN and the ventral and dorsal portions of the anterior preoptic area (PPa) express ER- β (Shughrue *et al.* 1997; Mitra *et al.* 2003; Merchenthaler *et al.* 2004). Other regions that express ER- β in mammals and birds include the magnocellular region of the PVN (Shughrue *et al.* 1997; Foidart *et al.* 1999; Hileman *et al.* 1999; Mitra *et al.* 2003; Mercanteller *et al.* 2004) and the cerebellum (Shughrue *et al.* 1997; Bernard *et al.* 1999; Foidart *et al.* 1999; Mitra *et al.* 2003). Thus ER- β a and ER- β b are found in different regions that express ER- β in tetrapods, suggesting that each ER- β subtype has lost ancestral ER- β functions that their duplicated counterpart (paralogue) has retained. Therefore, our data indicate that after the gene duplication event, subfunctionalization may have occurred because the distributions of ER- β a and ER- β b in croaker appear to reflect a partitioning of the functions of the ancestral ER- β gene. It is important to note that homologies for brain nuclei both within the fishes and across vertebrates are somewhat uncertain, so these regions may not be comparable. In addition, the oestrogen treatments given to upregulate ER levels in this study may have differential effects on ER subtype expression patterns (Menuet *et al.* 2004). Alternatively, it is also possible that ER- β b and ER- β a are expressed in different subsets of neurons within brain nuclei and have acquired novel neuron-specific functions that have resulted in neofunctionalization.

We conclude that the striking differences in the distributions of the three subtypes within brain nuclei of the POA and hypothalamus in the Atlantic croaker indicate that ER- α , ER- β a and ER- β b have distinct functions in the neuroendocrine control of reproduction and behaviour. The differential expression of the ER subtypes in other brain regions, such as the cerebellum and the posterior tuberculum, indicate that ER multiplicity is important for additional neural pathways. The combined neuroanatomical distribution of ER- β a and ER- β b follows the distribution pattern of mammalian ER- β . This conserved distribution pattern reflects the origin of ER- β a and ER- β b as a duplication of ER- β early in the teleost lineage and suggests that ancestral ER- β functions were subfunctionalized between teleost ER- β a and ER- β b after the duplication event. However, our results indicate that the largely conserved distribution patterns seen between tetrapod ER- α and ER- β may not hold true for euteleost ER- α , ER- β a and ER- β b. Additional studies of ER subtype distributions in fishes are needed to resolve the differential patterns of the three ER subtypes in

teleosts and ultimately interpret their roles in the evolution of vertebrate neural function.

We thank Russell Borski, Brenda Brizuela and Craig Sullivan for laboratory support. M.B.H. was supported by The University of Texas Marine Science Institute Harry Page Marine Science Fellowship and the Houston Livestock Show and Rodeo Natural Sciences Fellowship. This work was supported by Texas Sea grant R/ES-92, EPA STAR grant R827399, NIEHS grant ESO4214 and ESO7672 (to P.T.); NIH grant MH58271 (to JG); and NIH grants MH41770 and MH57874 (to D.C.).

REFERENCES

- Amores, A. *et al.* 1998 Zebrafish hox clusters and vertebrate genome evolution. *Science* **282**, 1711–1714.
- Arvin, M., Fedorkova, L., Disshon, K. A., Dluzen, D. E. & Leipheimer, R. E. 2000 Estrogen modulates responses of striatal dopamine neurons to MPP⁺: evaluations using *in vitro* and *in vivo* techniques. *Brain Res.* **872**, 160–171.
- Barami, S., Amiri, Z., Weizman, A., Fares, F. & Gavish, M. 1993 Long-term testosterone or diethylstilbestrol treatment affects gamma-aminobutyric-acid and central-type benzodiazepine receptors but not peripheral-type benzodiazepine receptors in the female rat-brain. *Neuroendocrinology* **57**, 1114–1118.
- Bernard, D. J., Bentley, G. E., Balthazart, J., Turek, F. W. & Ball, G. F. 1999 Androgen receptor, estrogen receptor alpha, and estrogen receptor beta show distinct patterns of expression in forebrain song control nuclei of European starlings. *Endocrinology* **140**, 4633–4643.
- Blaustein, J. D. 2004 Minireview: neuronal steroid hormone receptors: they're not just for hormones anymore. *Endocrinology* **145**, 1075–1081.
- Braford, M. R. J. & Northcutt, R. G. 1983 Organization of the diencephalon and pretectum of the ray-finned fishes. In *Fish neurobiology*, Vol. 2 (ed. R.E. Davis & R.G. Northcutt), pp. 117–163. Ann Arbor, MI: University of Michigan Press.
- Davis, R. E., Morrell, J. I. & Pfaff, D. W. 1977 Autoradiographic localization of sex steroid-concentrating cells in brain of teleost *Macropodus opercularis* (Osteichthyes: Belontiidae). *Gen. Comp. Endocrinol.* **33**, 496–505.
- Fine, M. L., Keefer, D. A. & Russelmergenthal, H. 1990 Autoradiographic localization of estrogen-concentrating cells in the brain and pituitary of the oyster toadfish. *Brain Res.* **536**, 207–219.
- Flugge, G., Oertel, W. H. & Wuttke, W. 1986 Evidence for estrogen-receptive gabaergic neurons in the preoptic-anterior hypothalamic area of the rat-brain. *Neuroendocrinology* **43**, 1–5.
- Foidart, A., Lakaye, B., Grisar, T., Ball, G. F. & Balthazart, J. 1999 Estrogen receptor- β in quail: cloning, tissue expression and neuroanatomical distribution. *J. Neurobiol.* **40**, 327–342.
- Force, A., Lynch, M., Pickett, F. B., Amores, A., Yan, Y. L. & Postlethwait, J. 1999 Preservation of duplicate genes by complementary, degenerative mutations. *Genetics* **151**, 1531–1545.
- Gore, A. C. 2002 *GnRH, the master molecule of reproduction*. Boston: Kluwer Academic Publishers.
- Hawkins, M. B. & Thomas, P. 2004 The unusual binding properties of the third distinct teleost estrogen receptor subtype ER β a are accompanied by highly conserved amino acid changes in the ligand binding domain. *Endocrinology* **145**, 2968–2977.

- Hawkins, M. B., Thornton, J. W., Crews, D., Skipper, J. K., Dotte, A. & Thomas, P. 2000 Identification of a third distinct estrogen receptor and reclassification of estrogen receptors in teleosts. *Proc. Natl Acad. Sci. USA* **97**, 10751–10756.
- Herbison, A. E. 1998 Multimodal influence of estrogen upon gonadotropin-releasing hormone neurons. *Endocr. Rev.* **19**, 302–330.
- Herbison, A. E. & Pape, J. R. 2001 New evidence for estrogen receptors in gonadotropin-releasing hormone neurons. *Front. Neuroendocrinol.* **22**, 292–308.
- Hileman, S. M., Handa, R. J. & Jackson, G. L. 1999 Distribution of estrogen receptor-beta messenger ribonucleic acid in the male sheep hypothalamus. *Biol. Reprod.* **60**, 1279–1284.
- Hrabovszky, E., Steinhauser, A., Barabas, K., Shughrue, P. J., Petersen, S. L., Merchenthaler, I. & Liposits, Z. 2001 Estrogen receptor-beta immunoreactivity in luteinizing hormone-releasing hormone neurons of the rat brain. *Endocrinology* **142**, 3261–3264.
- Hrabovszky, E., Kalló, I., Steinhauser, A., Merchenthaler, I., Coen, C. W., Petersen, S. L. & Liposits, Z. 2004 Estrogen receptor-beta in oxytocin and vasopressin neurons of the rat and human hypothalamus: immunocytochemical and *in situ* hybridization studies. *J. Comp. Neurol.* **473**, 315–333.
- Kah, O. *et al.* 1997 Estrogen receptors in the brain-pituitary complex and the neuroendocrine regulation of gonadotropin release in rainbow trout. *Fish Physiol. Biochem.* **17**, 53–62.
- Khan, I. A. & Thomas, P. 1993 Immunocytochemical localization of serotonin and gonadotropin-releasing-hormone in the brain and pituitary gland of the Atlantic croaker *Micropogonias undulatus*. *Gen. Comp. Endocrinol.* **91**, 167–180.
- Khan, I. A. & Thomas, P. 1999 GABA exerts stimulatory and inhibitory influences on gonadotropin II secretion in the Atlantic croaker (*Micropogonias undulatus*). *Neuroendocrinology* **69**, 261–268.
- Khan, I. A. & Thomas, P. 2001 Disruption of neuroendocrine control of luteinizing hormone secretion by Aroclor 1254 involves inhibition of hypothalamic tryptophan hydroxylase activity. *Biol. Reprod.* **64**, 955–964.
- Khan, I. A., Hawkins, M. B. & Thomas, P. 1999 Gonadal stage-dependent effects of gonadal steroids on gonadotropin II secretion in the Atlantic croaker (*Micropogonias undulatus*). *Biol. Reprod.* **61**, 834–841.
- Kim, Y. S., Stumpf, W. E. & Sar, M. 1979 Topographical distribution of estrogen target-cells in the forebrain of platyfish, *Xiphophorus maculatus*, studied by autoradiography. *Brain Res.* **170**, 43–59.
- Knobil, E. & Neill, J. D. 1994 *The physiology of reproduction*, New York: Raven Press.
- Kuiper, G., Enmark, E., PeltoHuikko, M., Nilsson, S. & Gustafsson, J. A. 1996 Cloning of a novel estrogen receptor expressed in rat prostate and ovary. *Proc. Natl Acad. Sci. USA* **93**, 5925–5930.
- Laflamme, N., Nappi, R. E., Drolet, G., Labrie, C. & Rivest, S. 1998 Expression and neuropeptidergic characterization of estrogen receptors (ER alpha and ER beta) throughout the rat brain: anatomical evidence of distinct roles of each subtype. *J. Neurobiol.* **36**, 357–378.
- Linard, B., Anglade, I., Corio, M., Navas, J. M., Pakdel, F., Saligaut, C. & Kah, O. 1996 Estrogen receptors are expressed in a subset of tyrosine hydroxylase-positive neurons of the anterior preoptic region in the rainbow trout. *Neuroendocrinology* **63**, 156–165.
- Ma, C. H., Dong, K. W. & Yu, K. L. 2000 cDNA cloning and expression of a novel estrogen receptor beta-subtype in goldfish (*Carassius auratus*). *Biochim. Biophys. Acta-Gene Struct. Expr.* **1490**, 145–152.
- Menuet, A., Pellegrini, E., Anglade, I., Blaise, O., Laudet, V., Kah, O. & Pakdel, F. 2002 Molecular characterization of three estrogen receptor forms in zebrafish: binding characteristics, transactivation properties, and tissue distributions. *Biol. Reprod.* **66**, 1881–1892.
- Menuet, A., Anglade, I., Le Guevel, R., Pellegrini, E., Pakdel, F. & Kah, O. 2003 Distribution of aromatase mRNA and protein in the brain and pituitary of female rainbow trout: comparison with estrogen receptor alpha. *J. Comp. Neurol.* **462**, 180–193.
- Menuet, A., Le Page, Y., Torres, O., Kern, L., Kah, O. & Pakdel, F. 2004 Analysis of the estrogen regulation of the zebrafish estrogen receptor (ER) reveals distinct effects of ER alpha, ER beta 1 and ER beta 2. *J. Mol. Endocrinol.* **32**, 975–986.
- Merchenthaler, I., Lane, M. V., Numan, S. & Dellovade, T. L. 2004 Distribution of estrogen receptor α and β in the mouse central nervous system: *in vivo* autoradiographic and immunocytochemical analyses. *J. Comp. Neurol.* **473**, 270–291. (doi:10.1002/cne.20128.)
- Mighell, A. J., Smith, N. R., Robinson, P. A. & Markham, A. F. 2000 Vertebrate pseudogenes. *FEBS Lett.* **468**, 109–114.
- Mitra, S. W., Hoskin, E., Yudkovitz, J., Pear, L., Wilkinson, H. A., Hayashi, S., Pfaff, D. W., Ogawa, S., Rohrer, S. P., Schaeffer, J. M., McEwen, B. S. & Alves, S. E. 2003 Immunolocalization of estrogen receptor beta in the mouse brain: comparison with estrogen receptor alpha. *Endocrinology* **144**, 2055–2067.
- Moore, F. L. 1992 Evolutionary precedents for behavioral actions of oxytocin and vasopressin. *Ann. NY Acad. Sci.* **652**, 156–165.
- Murray, H. E., Pillai, A. V., McArthur, S. R., Razvi, N., Datla, K. P., Dexter, D. T. & Gillies, G. E. 2003 Dose- and sex-dependent effects of the neurotoxin 6-hydroxydopamine on the nigrostriatal dopaminergic pathway of adult rats: differential actions of estrogen in males and females. *Neuroscience* **116**, 213–222.
- Navas, J. M., Anglade, I., Bailhache, T., Pakdel, F., Breton, B., Jego, P. & Kah, O. 1995 Do gonadotrophin-releasing hormone neurons express estrogen receptors in the rainbow trout? A double immunohistochemical study. *J. Comp. Neurol.* **363**, 461–474.
- Nomura, M., McKenna, E., Korach, K. S., Pfaff, D. W. & Ogawa, S. 2002 Estrogen receptor-beta regulates transcript levels for oxytocin and arginine vasopressin in the hypothalamic paraventricular nucleus of male mice. *Mol. Brain Res.* **109**, 84–94.
- Ogawa, S., Inoue, S., Watanabe, T., Hiroi, H., Orimo, A., Hosoi, T., Ouchi, Y. & Muramatsu, M. 1998 The complete primary structure of human estrogen receptor beta (hER beta) and its heterodimerization with ER alpha *in vivo* and *in vitro*. *Biochem. Biophys. Res. Commun.* **243**, 122–126.
- Ohno, S. 1970 *Evolution by gene duplication*. Berlin: Springer-Verlag.
- Pace, P., Taylor, J., Suntharalingam, S., Coombes, R. C. & Ali, S. 1997 Human estrogen receptor beta binds DNA in a manner similar to and dimerizes with estrogen receptor alpha. *J. Biol. Chem.* **272**, 25 832–25 838.
- Paech, K., Webb, P., Kuiper, G., Nilsson, S., Gustafsson, J. A., Kushner, P. J. & Scanlan, T. S. 1997 Differential ligand activation of estrogen receptors ER alpha and ER beta at AP1 sites. *Science* **277**, 1508–1510.

- Patisaul, H. B., Scordalakes, E. M., Young, L. J. & Rissman, E. F. 2003 Oxytocin, but not oxytocin receptor, is regulated by oestrogen receptor beta in the female mouse hypothalamus. *J. Neuroendocrinol.* **15**, 787–793.
- Peter, R. E., Macey, M. J. & Gill, V. E. 1975 Stereotaxic atlas and technique for forebrain nuclei of killifish, *Fundulus heteroclitus*. *J. Comp. Neurol.* **159**, 103–127.
- Pettersson, K., Grandien, K., Kuiper, G. & Gustafsson, J. A. 1997 Mouse estrogen receptor beta forms estrogen response element-binding heterodimers with estrogen receptor alpha. *Mol. Endocrinol.* **11**, 1486–1496.
- Pfaff, D. W. & Schwartz-Giblin, S. 1988 Cellular mechanisms of female reproductive behaviors. In *The physiology of reproduction*, vol. 2 (ed. E. Knobil & J. Neil), pp. 1487–1568. New York: Raven Press.
- Postlethwait, J., Amores, A., Cresko, W., Singer, A. & Yan, Y. L. 2004 Subfunction partitioning, the teleost radiation and the annotation of the human genome. *Trends Genet.* **20**, 481–490.
- Price, R. H. & Handa, R. J. 2000 Expression of estrogen receptor-beta protein and mRNA in the cerebellum of the rat. *Neurosci. Lett.* **288**, 115–118.
- Rink, E. & Wullimann, M. F. 2001 The teleostean (zebrafish) dopaminergic system ascending to the subpallium (striatum) is located in the basal diencephalon (posterior tuberculum). *Brain Res.* **889**, 316–330.
- Schultz, J. R., Loven, M. A., Melvin, V. M. S., Edwards, D. P. & Nardulli, A. M. 2002 Differential modulation of DNA conformation by estrogen receptors alpha and beta. *J. Biol. Chem.* **277**, 8702–8707.
- Senthilkumaran, B., Okuzawa, K., Gen, K. & Kagawa, H. 2001 Effects of serotonin, GABA and neuropeptide Y on seabream gonadatropin releasing hormone release *in vitro* from preoptic-anterior hypothalamus and pituitary of red seabream, *Pagrus major*. *J. Neuroendocrinol.* **13**, 395–400.
- Shughrue, P. J., Lane, M. V. & Merchenthaler, I. 1997 Comparative distribution of estrogen receptor-alpha and -beta mRNA in the rat central nervous system. *J. Comp. Neurol.* **388**, 507–525.
- Shupnik, M. A. 2002 Oestrogen receptors, receptor variants and oestrogen actions in the hypothalamic-pituitary axis. *J. Neuroendocrinol.* **14**, 85–94.
- Thomas, P., Pang, Y., Filardo, E. J. & Dong, J. 2004 Identity of an estrogen membrane receptor coupled to a G-protein in human breast cancer cells. *Endocrinology* **146**, 624–632.
- Thornton, J. W. 2001 Evolution of vertebrate steroid receptors from an ancestral estrogen receptor by ligand exploitation and serial genome expansions. *Proc. Natl Acad. Sci. USA* **98**, 5671–5676.
- Toran-Allerand, C. D. 2004 Minireview: a plethora of estrogen receptors in the brain: where will it end? *Endocrinology* **145**, 1069–1074.
- Young, L. J., Lopreato, G. F., Horan, K. & Crews, D. 1994 Cloning and *in-situ* hybridization analysis of estrogen-receptor, progesterone-receptor, and androgen receptor expression in the brain of whiptail lizards (*Cnemidophorus uniparens* and *C. inornatus*). *J. Comp. Neurol.* **347**, 288–300.

As this paper exceeds the maximum length normally permitted, the authors have agreed to contribute to production costs.