

*Review*

# Stress-induced variation in evolution: from behavioural plasticity to genetic assimilation

Alexander V. Badyaev\*

*Department of Ecology and Evolutionary Biology, University of Arizona,  
Tucson, AZ 85721-0088, USA*

Extreme environments are closely associated with phenotypic evolution, yet the mechanisms behind this relationship are poorly understood. Several themes and approaches in recent studies significantly further our understanding of the importance that stress-induced variation plays in evolution. First, stressful environments modify (and often reduce) the integration of neuroendocrinological, morphological and behavioural regulatory systems. Second, such reduced integration and subsequent accommodation of stress-induced variation by developmental systems enables organismal ‘memory’ of a stressful event as well as phenotypic and genetic assimilation of the response to a stressor. Third, in complex functional systems, a stress-induced increase in phenotypic and genetic variance is often directional, channelled by existing ontogenetic pathways. This accounts for similarity among individuals in stress-induced changes and thus significantly facilitates the rate of adaptive evolution. Fourth, accumulation of phenotypically neutral genetic variation might be a common property of locally adapted and complex organismal systems, and extreme environments facilitate the phenotypic expression of this variance. Finally, stress-induced effects and stress-resistance strategies often persist for several generations through maternal, ecological and cultural inheritance. These transgenerational effects, along with both the complexity of developmental systems and stressor recurrence, might facilitate genetic assimilation of stress-induced effects. Accumulation of phenotypically neutral genetic variance by developmental systems and phenotypic accommodation of stress-induced effects, together with the inheritance of stress-induced modifications, ensure the evolutionary persistence of stress–response strategies and provide a link between individual adaptability and evolutionary adaptation.

**Keywords:** adaptation; developmental plasticity; genetic and phenotypic variation; stress

## 1. INTRODUCTION

Environments outside the range normally experienced by a population, and the associated changes in organisms’ morphological, physiological or behavioural homeostasis, accompany most evolutionary changes (Bijlsma & Loeschcke 1997; Hoffmann & Parsons 1997; Hoffmann & Hercus 2000). Depending on the intensity, predictability and recurrence of stress, responses might range from stress tolerance and avoidance at the organismal level to the rapid appearance of novel traits or extinction at a population level. Yet, moderate stress is essential for normal growth and differentiation of many organismal systems (Huether 1996; Clark & Fucito 1998; Muller 2003). For example, a large part of skeletal development is directed by exposure to tension and mechanical overloads in excess of those normally experienced in the organism’s functioning (Hall 1986; Carter 1987; Herring 1993). Stress plays an important role in facilitating local adaptation by enabling better adjustments, synchronization and functioning of many organismal systems (Simons & Johnston 1997; Emlen *et al.* 2003; Rutherford 2003; Wingfield 2003; Robertson 2004). Anyone who has experienced the invigorating effects of diving into icy-cold water after a sauna, the health benefits of rigorous exercise, as well as

analgesic and attention-sharpening effects accompanying stressful encounters (McEwen & Sapolsky 1995; Shors & Servatious 1997) will testify to these effects of stress. On the other hand, response to an acute and unfamiliar stressor precludes normal organismal functions (Sibly & Calow 1989), and the high cost of stress tolerance or lack of evolved stress response strategies leads to evolutionary stasis (Parsons 1994).

Extreme environments not only disrupt normal development and induce large phenotypic changes in novel directions, they also simultaneously exert strong phenotypic selection that favours changes in these directions (Waddington 1941; Schmalhausen 1949; Bradshaw & Hardwick 1989; Jablonka *et al.* 1995). Not surprisingly, evolutionary diversification, the appearance of phenotypic novelties and mass extinction are all closely associated with extreme environmental changes (e.g. Howarth 1993; Guex 2001; reviewed in West-Eberhard 2003). Although it is widely recognized (especially in physiological and embryological studies) that stress plays an important role in directing and organizing the adaptive adjustment of an organism to ever-changing environments, very little is known about the evolution of mechanisms that enable the organismal accommodation of stress-induced variation.

The lack of a developmental perspective in evolutionary studies of stress has left us with several unresolved

\* (abadyaev@email.arizona.edu).

Table 1. Empirical patterns of the stress-induced phenotypic and genetic variation.

general phenomenon	specific pattern (organisms)	references
stress-induced generation of novel genetic variation	exposure to stress induces directional and locally adaptive mutations (green alga, bacteria, yeast, <i>Daphnia</i> )	Ruvinsky <i>et al.</i> (1983), Cairns <i>et al.</i> (1988), Hall (1990), Steele & Jinks-Robertson (1992), Foster (2000), Goho & Bell (2000), Sniegowski <i>et al.</i> (2000), Bjedov <i>et al.</i> (2003) and Wright (2004)
	increase in evolutionary rate of a gene (cyanobacteria; human)	Prody <i>et al.</i> (1989) and Dvornyk <i>et al.</i> (2002)
	increase in frequency of sexual recombination ( <i>Volvox</i> )	Nedelcu & Michod (2003)
	increase in mutation and/or recombination rates (many species)	Belyaev & Borodin (1982), Parsons (1988), Selker (1990) and Imasheva (1999)
	increase in stress-induced transposition (plants, <i>Drosophila</i> )	McClintock (1984), Bownes (1990), Rather <i>et al.</i> (1992) and Wessler (1996)
stress challenge of general homeostasis releases hidden variation	appearance of primitive, ancestor-like forms	Guex (2001)
	phenotypic responses to stress mimic the expression of mutation	Goldschmidt (1940), Waddington (1941, 1953), Milkman (1965), Ho <i>et al.</i> (1983), Chow & Chan (1999) and Schlichting & Smith (2002)
	phenotypically neutral genetic variance in ancestral forms of domesticated organisms becomes adaptive in the hybrid backgrounds, including domesticated forms (soybeans, maize, sunflowers) and other organisms	Roth <i>et al.</i> (1989), Lauter & Doebley (2002), Rieseberg <i>et al.</i> (2003), Innan & Kim (2004), Pelabon <i>et al.</i> (2004), Piffanell <i>et al.</i> (2004) and Seehausen (2004)
	environment dependency and context dependency in expression of genetic variation	Kondrashov & Houle (1994), Threadgill <i>et al.</i> (1995), Leips & MacKay (2000) and Keller <i>et al.</i> (2002)
stress challenge of specific buffering mechanisms releases hidden variation	complex and redundant developmental systems enable accumulation of mutational variance	Szafraniec <i>et al.</i> (2001)
	stress-induced changes in regulation of chaperone proteins releases normally unexpressed genetic variation	Rutherford & Lindquist (1998), Queitsch <i>et al.</i> (2002) and Ruden <i>et al.</i> (2003)
	release of cryptic genetic variation by artificial selection ( <i>Drosophila</i> )	Gibson & Hogness (1996) and Dworkin <i>et al.</i> (2003)
	epigenetic regulation of genes uncovers normally unexpressed phenotypic variation	Rutherford (2003), Sollars <i>et al.</i> (2003) and True <i>et al.</i> (2004)

questions. First, how can organisms prepare for novel and extreme environmental change? Organismal ability to mount an appropriate reaction to a stressor requires recognition and evaluation of the extreme environment. How can this ability evolve in relation to stressors that are short and rare in relation to a species' generation time? Second, numerous studies have documented an increase in phenotypic and genotypic variance under stress, and it is suggested that this variance is a source of novel adaptations under changed environments. Yet, for stress-induced modifications to have evolutionary importance, they have to be inherited and must persist in a sufficient number of individuals within a population. This requires an organism to survive stress and reproduce; that is, to accommodate stress-induced variation without reduction in the organism's functionality. How is such accommodation accomplished? Moreover, could existing organismal systems channel accumulation of stress-induced variance in some directions, but not others and, thus, direct evolutionary change in response to stress? The perspective outlined here, with a particular focus on the effect of stress during development in animals, suggests that these questions are resolved by considering: (i) the organization of developmental systems that enable accommodation

and channelling of stress-induced variation without compromising organismal functionality; (ii) the significance of phenotypic and genetic assimilation of neurological, physiological, morphological and behavioural responses to stressors; and (iii) multiple inheritance systems that transfer the wide array of developmental resources and conditions between the generations enabling long-term persistence and evolution of stress-induced adaptations.

## 2. SOURCES OF STRESS-INDUCED VARIATION

A stress-induced increase of phenotypic and genetic variation in a population has three main sources. First, directional selection imposed by a stressor can result in faster rates of mutation and recombination. Second, stress challenges to homeostatic mechanisms can release and amplify phenotypic variation. Third, stressful environments can facilitate developmental expression of novel genetic variation that had accumulated, but was phenotypically neutral under a normal range of environments. These sources of variation can be adaptive under stressful conditions when they facilitate the development of novel adaptations to changed environments.

**(a) 'Generated' variance**

Organismal reaction to a stressor is often associated with generation of variation in a directional and locally adaptive manner (Jablonka & Lamb 1995; West-Eberhard 2003; see table 1). In some cases, such directionality is attributed to the channelling effects of complex developmental networks (e.g. Walker 1979; Roth & Wake 1985). In other cases, it is associated directly with a stressful environment (e.g. Wills 1983). An extreme environment often increases genetic variation because of the elevation in mutation and recombination rates (Hoffmann & Parsons 1997; Hoffmann & Merilä 1999; Imasheva 1999; Wright 2004). When such mutations are directional (or 'focused'; Caropale 1999) in relation to a stressor—that is, when the stressful environment causes a mutation and favours phenotypic change associated with this mutation—such an increase in mutation rate might result in greater similarity among individuals within a population in response to a stressor facilitating evolutionary adaptation to novel environments (Shapiro 1992; Jablonka & Lamb 1995; Foster 2000; Wright 2000; Brisson 2003; Rutherford 2003; Wright 2004). Table 1 shows examples of stress-induced generation of novel genetic variation. At the level of phenotype, induction of a phenotypic trait by a stressor and concurrent selection on the induced trait are common (Price *et al.* 2003; West-Eberhard 2003).

**(b) 'Hidden' variance**

Stressful environments reveal greater phenotypic and genetic variability than is seen under normal conditions, and it is commonly suggested that such hidden variation results from stress-induced challenge to organismal homeostasis (Scharloo 1991). In turn, an increase in variation and subsequent reorganization of organismal systems are thought to enable the formation of novel adaptations (Bradshaw & Hardwick 1989; Eshel & Matessi 1998; Gibson & Wagner 2000; Schlichting & Smith 2002). The idea that the extreme environments' challenge to previously canalized systems is the source of such hidden variation is corroborated by numerous empirical observations of environment- and context-dependency in expression of novel genetic variation (table 1). However, it is not clear how genetic and developmental systems can accumulate and store phenotypically neutral genetic variation while not expressing it (Wagner & Mezey 2000; Hermisson *et al.* 2003; Masel & Bergman 2003; Gibson & Dworkin 2004). Specifically, the discussion has focused on the existence of 'evolutionary capacitors' (Rutherford 2000) and 'adaptively inducible canalizers' (Meiklejohn & Hartl 2002)—specific mechanisms that buffer and accumulate developmental variation, producing 'hidden reaction norms' of a phenotype under stress. A debated question is whether these 'evolutionary capacitors' are stressor-specific regulatory systems or whether evolutionary capacity is a property of any complex and locally adapted organismal system. Rutherford & Lindquist (1998) found that mutations at the gene for the stress-induced chaperone proteins harbours abundant but normally unexpressed genetic variation, which, when selected, leads to the appearance and assimilation of novel phenotypes in the population (Queitsch *et al.* 2002; Ruden *et al.* 2003; Rutherford 2003; Sollars *et al.* 2003). However, other studies suggested that 'evolutionary capacity' is a property of most locally adapted

developmental systems that, when challenged by a novel environment, reveal large genetic variation (Kirschner & Gerhart 1998; Rutherford 2000; Bergman & Siegal 2003). Complex developmental processes and genetic networks can constrain variation in individual traits (Rice 2004), and phenotypically neutral genetic variation is commonly accumulated in such systems given sufficient time and population size (Hermisson & Wagner 2004).

**3. EVOLUTION OF RESPONSE TO STRESS**

Stress occurs when changes in the external or internal environment are interpreted by an organism as a threat to its homeostasis (e.g. Greenberg *et al.* 2001; McEwen & Wingfield 2003). Thus, the ability of an organism to mount an appropriate response to potentially stressful environmental changes requires correct recognition of environmental change and the activation of a stress response (e.g. Johnson *et al.* 1992). However, it is unclear how the ability to recognize and assess potentially stressful environments can evolve. How can organisms judge the appropriate reaction to a stressor, as is required in order to select between stressor avoidance and stress tolerance? Are the mechanisms of assessment and avoidance specific to a particular stressor?

**(a) Insights from cognitive and physiological assimilations of a stressful event: what does not kill you makes you stronger**

Generally, repeated exposure to a particular stressor favours the evolution of mechanisms that suppress an organism-wide stress reaction and, instead, activate stress-specific responses (Johnson *et al.* 1992; Barclay & Robertson 2001; Veenema *et al.* 2003). For example, in mammals and birds, stress-induced activation of the neuroendocrinological system increases its reactivity to internal and external stimuli, facilitates the processing of sensory information, and ultimately enables the formation of a behavioural or physiological strategy of dealing with a stressor. Furthermore, stress-induced activation of neuroendocrinological systems facilitates long-term retention of information about a stressful event and corresponding organismal response. Once formed, the maintenance of such 'memory' can be accomplished by periodic exposure to different stressors. For example, hormones associated with stress detection and avoidance also play a major role in modifications of neural circuits (Gold & McGaugh 1978). Once the stress-avoidance strategy is formed, exposure to even a low concentration of these hormones maintains the strategy (McGaugh *et al.* 1982; for similar examples, see Sockman *et al.* 2002, 2004).

Animal physiology studies show that the repeated experience of successfully overcoming social stresses during ontogeny is a prerequisite for the acquisition of a normal repertoire of behavioural strategies (Huether 1996; e.g. Gans 1979). An insightful example is provided by experiments that show that individuals exposed to repeatable, but consistently unfamiliar (and thus 'uncontrollable' by an animal) stressors develop 'stressful helplessness' (i.e. they lose their ability to react to any stressor; Katz *et al.* 1981; Johnson *et al.* 1992; Avitsur *et al.* 2001). In contrast, individuals that were allowed to develop a stress-avoidance strategy by exposure to a previously encountered stressor not only developed stress

tolerance to a particular stressor, but also actively sought out other mild stressors. In the absence of other stressors, their stress-avoidance abilities diminished (Katz *et al.* 1981; Johnson *et al.* 1992; Avitsur *et al.* 2001). Similarly, in insects, exposure to temperature stress facilitated not only subsequent long-term tolerance of extreme temperatures, but also performance under anoxia and various motor stresses (Karunanithi *et al.* 1999; Ramirez *et al.* 1999; Wu *et al.* 2002). The interchangeability of distinct stressors in maintaining and exerting organismal stress response, and long-term persistence of stress effects, have also been documented in the studies of amphibians, where organismal stress from food deprivation and from a water level reduction exerted similar endocrinological effects during larval metamorphosis (Denver 1999; Boorse & Denver 2004). These results suggest that, once originated, a stress-response strategy can be maintained by other environments, and adaptation to one type of stressor facilitates adaptation to other stressors.

Phenotypic assimilation of the appropriate stress response is further facilitated by a common involvement of neural and endocrine pathways of the stress response in other organismal functions (Aston-Jones *et al.* 1986; Greenberg *et al.* 2001; Robertson 2004). In such cases, even a single stressful experience during development is often enough to induce changes that, in the future, will prevent organism-wide stressful reactions and will instead activate stress-specific behavioural and physiological responses (Levine *et al.* 1967, 1989). Generally, stress-induced reorganization of existing developmental pathways and organismal function, rather than the production of novel stress-specific pathways, accounts for the ease with which individuals and populations lose and gain the ability to resist stress in laboratory settings (Chapin *et al.* 1993; Hoffmann & Merilä 1999).

#### **(b) Stress-avoidance strategies**

The ability actively to remove a stressor by either relocation or avoidance requires an evolved ability to detect or anticipate stressful changes and the 'knowledge' or 'memory' of stress avoidance strategies or adjustments (Bradshaw & Hardwick 1989; Jablonka *et al.* 1995; Denver 1999). Therefore, the evolution of stress avoidance is more probable when stressful events are predictable, prolonged and frequent in relation to generation time (Ancel Meyers & Bull 2002). Alternatively, the short-term avoidance of a frequent but mild stressor might be accommodated by behavioural or physiological plasticity of an organism (Schlichting & Smith 2002; Nicolakakis *et al.* 2003; Piersma & Drent 2003; Wingfield & Sapolsky 2003). For example, repeated challenges of an organism's immune system and challenges to skeletal tissues caused by mechanical overload during growth lower developmental errors and enable a more precise reaction to a specific pathogen (e.g. Graham *et al.* 2000; Hallgrímsson *et al.* 2004). Generally, organisms can activate stress reactions when there is discordance between environments during their development and their current external and internal environments (Meaney 2001; Bateson *et al.* 2004). On the longer time-scale, avoidance of a predictable stressor can be accomplished by changes in an organism's life history, especially by altering the timing of reproduction or duration of development. Common cases include stress-induced modification of the timing of metamorphosis

in amphibians, changes in the duration of gestation in mammals and the timing of flowering and seeding in plants (e.g. Bradshaw & Hardwick 1989; Denver 1999; Stanton *et al.* 2000; Boorse & Denver 2004).

In sum, initial behavioural accommodations of stress (e.g. hiding, relocation, lowering metabolism) may set the stage for the evolution of adaptive stress-avoidance strategies (e.g. periodic hibernation, migration, torpor). When a stressor is reliably preceded by other environmental changes, their mutual recurrence facilitates the establishment of stressor recognition, assessment and avoidance strategies, such that an evolved stress-specific strategy does not involve an activation of an organism-wide stress response. When individuals vary in their reaction to stress and when stress-induced strategies are favoured by natural selection during and after a stressful event, these strategies can become phenotypically and genetically assimilated in a population (see below; Baldwin 1896; Hinton & Nolan 1980; Oyama 2000; West-Eberhard 2003).

#### **(c) Stress buffering as a by-product of developmental complexity**

Organismal functions most closely related to fitness are thought to be the most buffered against internal and external stressors (Waddington 1941; Schmalhausen 1949; Stearns & Kawecki 1994). However, an organism's functioning in changing environments requires the ability to track and respond to these environments, and evolved buffering from stressors would also restrict an organism's ability and capacity to adapt to continuously changing environments (Wagner *et al.* 1997; Eshel & Matessi 1998; Ancel 1999; Schlichting & Smith 2002). For example, suppression of stress-induced activation of the sensory systems limits an organism's ability to acquire and retain the sensory cues and behavioural strategies necessary for stress avoidance (Huether 1996). On the one hand, a lack of phenotypic plasticity results in population extinction under stress (Gavrilets & Scheiner 1993; Ancel 1999). On the other hand, extensive phenotypic variability in organismal functions weakens the effects of directional selection imposed by stressful environments, and thus lessens the opportunity for genetic assimilation and evolution of adaptations to stress (Fear & Price 1998; Ancel 2000; Huey *et al.* 2003). Recent studies suggest that buffering is an emerging property of developmental complexity rather than an evolved stress-resistance mechanism (see above). The increasing complexity of developmental pathways and networks leads directly to environmental and genetic stability and canalization (Baatz & Wagner 1997; Rice 1998, 2004; Waxman & Peck 1998; Meiklejohn & Hartl 2002; Siegal & Bergman 2002; Ruden *et al.* 2003). Complex genetic and developmental networks can accommodate the effects of stressful perturbations without the loss of function or structure, while building up neutral genetic variation (Rutherford 2000; Bergman & Siegal 2003; Masel 2004).

#### **(d) Accommodation of stress-induced variation by changes in an organism's integration**

Organisms might accommodate stress-induced variation without the loss of function by lessening homeostasis of individual systems. For example, individual hormonal systems have a far greater potential range of performances

Table 2. Empirical patterns of the relationship between stress and homeostasis.

general phenomenon	specific pattern	references
stress increases homeostasis	stress-induced increase in resource exchange leads to a greater developmental integration among normally independent organismal components in animals raised under stress exposure to stress prevents the expression of deleterious mutations in <i>Escherichia coli</i> increased homeostatic interactions within reproductive systems counteracts the stress effects on the organism when breeding opportunities are limited	Sciulli <i>et al.</i> (1979), Schandorff (1997) and Klingenberg <i>et al.</i> (2001)  Kishony & Leibler (2003)  Wingfield & Sapolsky (2003)
stress decreases homeostasis	increase in developmental noise and variation in numerous organismal traits  destruction of the nest site leads to temporary breaking of the strict hierarchical social structure and rapid proliferation of random individual search routes that facilitate finding of a new nest site in social ant species	Parsons (1990), Hoffmann & Parsons (1997), Møller & Swaddle (1997), Polak (2003) and Badyaev (2004)  Britton <i>et al.</i> (1998) and Couzin & Franks (2003)

and can remain functional under a wider range of environments than is allowed by homeostasis under a normal range of environments (Dickinson 1988; Johnson *et al.* 1992; Greenberg *et al.* 2001; Robertson 2004; Romero 2004). Frequently documented suppression of immunocompetence under stressful conditions might facilitate novel adaptations to a stressor by realizing full capabilities of individual immune systems (Råberg *et al.* 1998). When stress is associated with damage of tissues, suppression of immunological functions enables individual organismal systems to respond to a stressor without activation of organism-wide autoimmunological response (Dickinson 1988; Råberg *et al.* 1998; Avitsur *et al.* 2001). Examples of stress-induced increases in organismal integration, mostly in response to mild stressors, and corresponding suppression of random developmental variation under stress are shown in table 2.

The importance of the timing of stress for directing the evolution of morphological traits is well documented. When stressors are mild and occur during ontogeny, individual organismal systems often accommodate stress-induced variation without the reduction in functionality (Bradshaw & Hardwick 1989; Herring 1993; Huether 1996; Schandorff 1997; Hallgrímsson 1999; Badyaev & Foresman 2004). For example, when components of foraging structures differ in patterns of ossification, morphological variation in later ossified components can be directed by stress-induced modifications of earlier ossified components (Mabee *et al.* 2000; Badyaev *et al.* in press). In addition, when stress occurs early in ontogeny, accommodation and channelling of stress-induced variation by existing organismal structures causes similar reorganization in many individuals simultaneously (Roth & Wake 1985; Chapin *et al.* 1993), thus facilitating adaptive evolution (Goldschmidt 1940; West-Eberhard 2003).

#### (e) *Channelling of stress-induced variation*

Natural selection favours organismal homeostasis that maintains some degree of developmental variation necessary for adjustment of the organism to its environments (Simons & Johnston 1997; Wagner *et al.* 1997; Eshel & Matessi 1998; Emlen *et al.* 2003; Nanjundiah 2003). Stressful conditions can increase this variation, and

differences among organismal systems in their reaction to a stressor might bias the introduction and expression of variation available for selection and, thus, ultimately, affect evolutionary change (Bonner 1965; Roth & Wake 1985; Jablonka & Lamb 1995; West-Eberhard 2003). Empirical studies show that the coordinated development of morphological traits often leads to their similarity in expression of stress-induced developmental variation (Leamy 1993; Badyaev & Foresman 2000, 2004; Klingenberg *et al.* 2001; Badyaev *et al.* in press). These examples not only confirm a strong effect of functional complexes on directing and incorporating stress-induced variation during development, but might also explain the historical persistence of complex groups of traits through stressful environments.

#### 4. INHERITANCE OF STRESS-INDUCED VARIATION

For a stress-induced modification to be preserved in a lineage, it needs to be accommodated by an organism and transmitted between generations (i.e. inherited). This presents two problems. First, how can environmentally induced effects become inherited? Second, if each organism accommodates a stressor by different adjustments, then how can this diversity enable directional evolution of a stress–response strategy?

Stress-induced phenotypic changes commonly persist across several generations. Such across-generation carry-over effects (*sensu* Jablonka *et al.* 1995) can be a result of the transfer of physical substances, inheritance and developmental incorporation of a stressor, hormonal effects that influence expression of genetic variance in subsequent generations, epigenetic inheritance of stress-induced variation and structures, as well as behavioural effects (Jablonka & Lamb 1995; Rossiter 1996; Oyama 2000; Odling-Smee *et al.* 2003; West-Eberhard 2003). For example, inheritance of dominant–subdominant relationships in groups of many social mammals is accomplished by mechanisms different from original stressful encounters that established the dominance structure (Creel *et al.* 1996; Fairbanks 1996). Similarly, maternal care often sets the stage for a lifelong reaction to stressors by offspring by modifying the expression of

genes that regulate behavioural, physiological and endocrinological responses to stressors (Mousseau & Fox 1998; Meaney 2001; Badyaev *in press*). At the individual level, stress-induced changes in neuroendocrinological systems often occur with significant delay after the exposure to stress, and persist for a long time. This observation led to the suggestion that the primary function of such delayed changes is integration of past stress-induced responses and sensitization of the organism to future occurrences of similar stressors (Huether 1996; Ramirez *et al.* 1999; Romero 2004). In turn, within- and between-generation maintenance of stress-induced changes in neurophysiological systems is accomplished by similar hormonal mechanisms (see above; McGaugh *et al.* 1982; Meaney 2001). More generally, short-term and non-genetic inheritance might be beneficial when the frequency of stress recurrence is greater than the generation time, but shorter than is necessary for the spread and fixation of adaptive mutation (i.e. the evolution of genetic adaptation; Levins 1963; Ancel Meyers & Bull 2002; Gavrilets 2004).

In sum, accommodation of stress-induced variance by an organism can be facilitated by recurrent developmental stressors and genetic assimilation replaces stress-induced developmental modification if this modification has a fitness advantage in both stressful and post-stress environments (Schmalhausen 1949; Waddington 1953). Even when the short-term organismal responses to a stressor are not genetically heritable, differences among organisms in their ability to survive stress and the recurrence of stressful environments will canalize stress-induced responses developmentally (Baldwin 1896; Schlichting & Pigliucci 1998; Ancel 1999; West-Eberhard 2003). An excellent example is a recent comprehensive review by Palmer (2004), where he shows that genetic assimilation enables the evolutionary establishment of environmentally induced novelty—directional asymmetry (for other examples of genetic assimilation of stress-induced developmental variation see Ho *et al.* 1983; Chapman *et al.* 2000; Heil *et al.* 2004).

## 5. STRESS-INDUCED EVOLUTION VERSUS STRESS-INDUCED STASIS

Only a subset of stressful environments—narrowly fluctuating and slowly changing in relation to generation time—are thought to be associated with a rapid evolutionary change, whereas extreme and rapidly changing environments might result in morphological stasis because of the costs associated with stress tolerance (Parsons 1994). Furthermore, only stressors specific to an organismal system are expected to enable phenotypic assimilation and evolutionary persistence of stress-induced adaptations because a more general stressor favours stress tolerance by increasing homeostasis, which, in turn, leads to a reduction in organismal metabolism and fitness. Thus, among the array of organismal responses to stressful environments reviewed above, only accommodation of stress-induced variation and stress avoidance are expected to lead to significant evolutionary change (Parsons 1993). On the contrary, because of its association with lower metabolism and stronger regulatory systems, stress tolerance is unlikely to be associated with greater organismal plasticity, and instead, leads to stasis under

extreme environments which is observed in ‘living fossils’ (Parsons 1993, 1994).

I thank J. Hermisson, J. Masel and G. Wagner for discussions, L. Carsten, R. Duckworth, K. Oh, H. Maughan, L. Reed, D. Seaman, R. Young and two anonymous reviewers for comments that significantly improved this manuscript. The expanded treatment of points raised in this review will be published in *Variation* (ed. B. Hallgrímsson & B. K. Hall, Academic Press). I thank the US National Science Foundation (DEB-0075388, DEB-0077804, and IBN-0218313) and University of Arizona for partly funding this work.

## REFERENCES

- Ancel, L. W. 1999 A quantitative model of the Simpson–Baldwin effect. *J. Theor. Biol.* **196**, 197–209.
- Ancel, L. W. 2000 Undermining the Baldwin expediting effect: does phenotypic plasticity accelerate evolution? *Theor. Popul. Biol.* **58**, 307–319.
- Ancel Meyers, L. & Bull, J. J. 2002 Fighting change with change: adaptive variation in an uncertain world. *Trends Ecol. Evol.* **17**, 551–557.
- Aston-Jones, G., Ennis, M., Pieribone, V. A., Nickel, W. T. & Shipley, M. T. 1986 The brain nucleus locus coeruleus: restricted afferent control of a broad efferent network. *Science* **234**, 734–737.
- Avitsur, R., Stark, J. & Sheridan, J. 2001 Social stress induces glucocorticoid resistance in subordinate animals. *Horm. Behav.* **39**, 247–257.
- Baatz, M. & Wagner, G. P. 1997 Adaptive inertia caused by hidden pleiotropic effects. *Theor. Popul. Biol.* **51**, 49–66.
- Badyaev, A. V. 2004 Integration and modularity in the evolution of sexual ornaments: an overlooked perspective. In *Phenotypic integration: the evolutionary biology of complex phenotypes* (ed. M. Pigliucci & K. Preston), pp. 50–79. Oxford University Press.
- Badyaev, A. V. *In press*. Maternal inheritance and rapid evolution of sexual size dimorphism: passive effects or active strategies? *Am. Nat.*
- Badyaev, A. V. & Foresman, K. R. 2000 Extreme environmental change and evolution: stress-induced morphological variation is strongly concordant with patterns of evolutionary divergence in shrew mandibles. *Proc. R. Soc. B* **267**, 371–377. (doi:10.1098/rspb.2000.1011)
- Badyaev, A. V. & Foresman, K. R. 2004 Evolution of morphological integration. I. Functional units channel stress-induced variation. *Am. Nat.* **163**, 868–879.
- Badyaev, A. V., Foresman, K. R. & Young, R. L. *In press*. Evolution of morphological integration. II. Developmental accommodation of stress-induced variation. *Am. Nat.*
- Baldwin, J. M. 1896 A new factor in evolution. *Am. Nat.* **30**, 441–451.
- Barclay, J. W. & Robertson, R. M. 2001 Enhancement of short-term synaptic plasticity by prior environmental stress. *J. Neurophysiol.* **85**, 1332–1335.
- Bateson, P. *et al.* 2004 Developmental plasticity and human health. *Nature* **430**, 419–421.
- Belyaev, D. K. & Borodin, P. M. 1982 The influence of stress on variation and its role in evolution. *Biol. Zentralbl.* **100**, 705–714.
- Bergman, A. & Siegal, M. L. 2003 Evolutionary capacitance as a general feature of complex gene networks. *Nature* **424**, 549–552.
- Bijlsma, R. & Loeschcke, V. (ed.) 1997 *Environmental stress, adaptation and evolution*. Basel, Switzerland: Birkhauser.
- Bjedov, I., Tenaillon, O., Gerard, B., Souza, V., Denamur, E., Radman, M., Taddei, F. & Matic, I. 2003 Stress-induced mutagenesis in bacteria. *Science* **300**, 1404–1409.
- Bonner, J. T. 1965 *Size and cycle*. Princeton University Press.

- Boorse, G. C. & Denver, R. J. 2004 Endocrine mechanisms underlying plasticity in metamorphic timing in spadefoot toads. *Integr. Comp. Biol.* **43**, 646–657.
- Bownes, M. 1990 Preferential insertion of P elements into genes expressed in the germ line of *Drosophila melanogaster*. *Mol. Gen. Genet.* **222**.
- Bradshaw, A. D. & Hardwick, K. 1989 Evolution and stress—genotypic and phenotypic components. *Biol. J. Linn. Soc.* **37**, 137–155.
- Brisson, D. 2003 The directed mutation controversy in an evolutionary context. *Crit. Rev. Microbiol.* **29**, 25–35.
- Britton, N. F., Stickland, T. R. & Franks, N. R. 1998 Analysis of ant foraging algorithms. *J. Biol. Syst.* **6**, 315–336.
- Cairns, J., Overbaugh, J. & Miller, S. 1988 The origin of mutants. *Nature* **335**, 142–145.
- Caropale, L. H. 1999 Chance favors the prepared genome. *Ann. NY Acad. Sci.* **870**, 1–21.
- Carter, D. R. 1987 Mechanical loading history and skeletal biology. *J. Biomech.* **20**, 1095–1109.
- Chapin III, F. S., Autumn, K. & Pugnnaire, F. 1993 Evolution of suites of traits in response to environmental stress. *Am. Nat.* **142**(Suppl.), S78–S92.
- Chapman, L. J., Galis, F. & Shinn, J. 2000 Phenotypic plasticity and the possible role of genetic assimilation: hypoxia-induced trade-offs in the morphological traits of an African cichlid. *Ecol. Lett.* **3**, 387–393.
- Chow, K. L. & Chan, K. W. 1999 Stress-induced phenocopy of *C. elegans* defines functional steps of sensory organ differentiation. *Dev. Growth Differ.* **41**, 629–637.
- Clark, A. G. & Fucito, C. D. 1998 Stress tolerance and metabolic response to stress in *Drosophila melanogaster*. *Heredity* **81**, 514–527.
- Couzin, I. D. & Franks, N. R. 2003 Self-organized lane formation and optimized traffic flow in army ants. *Proc. R. Soc. B* **270**, 139–146. (doi:10.1098/rspb.2002.2210)
- Creel, S., Creel, N. M. & Monfort, S. L. 1996 Social stress and dominance. *Nature* **379**, 212.
- Denver, R. J. 1999 Evolution of the corticotropin-releasing hormone signaling system and its role in stress-induced phenotypic plasticity. *Ann. NY Acad. Sci.* **897**, 46–53.
- Dickinson, W. J. 1988 On the architecture of regulatory systems: evolutionary insights and implications. *BioEssays* **8**, 204–208.
- Dvornyk, V., Vinogradova, O. & Nevo, E. 2002 Long-term microclimatic stress causes rapid adaptive radiation of kaiABC clock gene family in a cyanobacterium, *Nostoc linckia*, from “Evolution Canyons” I and II, Israel. *Proc. Natl Acad. Sci. USA* **99**, 2082–2087.
- Dworkin, I., Palsson, I., Birdsall, K. & Gibson, G. 2003 Evidence that *Egfr* contributes to cryptic genetic variation for photoreceptor determination in natural populations of *Drosophila melanogaster*. *Curr. Biol.* **13**, 1888–1893.
- Emlen, J. M., Freeman, D. C. & Graham, J. H. 2003 The adaptive basis of developmental instability: a hypothesis and its implication. In *Developmental instability: causes and consequences* (ed. M. Polak), pp. 51–61. Oxford University Press.
- Eshel, I. & Matessi, C. 1998 Canalization, genetic assimilation and preadaptation: a quantitative genetic model. *Genetics* **149**, 2119–2133.
- Fairbanks, L. A. 1996 Individual differences in maternal style: causes and consequences for mothers and offspring. *Adv. Study Behav.* **25**, 579–611.
- Fear, K. K. & Price, T. 1998 The adaptive surface in ecology. *Oikos* **82**, 440–448.
- Foster, P. L. 2000 Adaptive mutation: implications for evolution. *BioEssays* **22**, 1067–1074.
- Gans, C. 1979 Momentary excessive construction as the basis for protoadaptation. *Evolution* **33**, 227–233.
- Gavrilets, S. 2004 *Fitness landscapes and the origin of species*. Princeton University Press.
- Gavrilets, S. & Scheiner, S. 1993 The genetics of phenotypic plasticity. V. Evolution of reaction norm shape. *J. Evol. Biol.* **6**, 31–48.
- Gibson, G. & Dworkin, I. 2004 Uncovering cryptic genetic variation. *Nat. Rev.* **5**, 681–690.
- Gibson, G. & Hogness, D. S. 1996 Effect of polymorphism in the *Drosophila* regulatory gene *Ultrabithorax* on homeotic stability. *Science* **271**, 200–203.
- Gibson, G. & Wagner, G. P. 2000 Canalization in evolutionary genetics: a stabilizing theory? *BioEssays* **22**, 372–380.
- Goho, S. & Bell, G. 2000 Mild environmental stress elicits mutations affecting fitness in *Chlamydomonas*. *Proc. R. Soc. B* **267**, 123–129. (doi:10.1098/rspb.2000.0976)
- Gold, P. E. & McGaugh, J. L. 1978 Neurobiology and memory: modulators, correlates and assumptions. In *Brain and learning* (ed. T. Teilor), pp. 93–104. Stanford, CT: Greylock Publishers.
- Goldschmidt, R. B. 1940 *The material basis of evolution*. New Haven, CT: Yale University Press.
- Graham, J. H., Fletcher, D., Tigue, J. & McDonald, M. 2000 Growth and developmental stability of *Drosophila melanogaster* in low frequency magnetic fields. *Bioelectromagnetics* **21**, 465–472.
- Greenberg, N., Carr, J. A. & Summers, C. H. 2001 Ethological causes and consequences of the stress response. *Integr. Comp. Biol.* **42**, 508–516.
- Guex, J. 2001 Environmental stress and atavism in ammonoid evolution. *Ecol. Geol. Helv.* **94**, 321–328.
- Hall, B. K. 1986 The role of movement and tissue interactions in the development and growth of bone and secondary cartilage in the clavicle of the embryonic chick. *J. Embryol. Exp. Morph.* **93**, 133–152.
- Hall, B. G. 1990 Spontaneous point mutations that occur more often when advantageous than when neutral. *Genetics* **126**, 5–16.
- Hallgrímsson, B. 1999 Ontogenetic patterning of skeletal fluctuating asymmetry of rhesus macaques and human: evolutionary and developmental implications. *Int. J. Primatol.* **20**, 121–151.
- Hallgrímsson, B., Willmore, K., Dorval, C. & Cooper, D. M. L. 2004 Craniofacial variability and modularity in macaques and mice. *J. Exp. Zool.* **302B**, 207–225.
- Heil, M., Greiner, S., Meimberg, H., Kruger, R., Noyer, J.-L., Heubl, G., Linsenmair, K. E. & Boland, W. 2004 Evolutionary change from induced to constitutive expression of an indirect plant resistance. *Nature* **430**, 205–208.
- Hermisson, J. & Wagner, G. P. 2004 The population genetic theory of hidden variation and genetic robustness. *Genetics* **168**, 2271–2284.
- Hermisson, J., Hansen, T. F. & Wagner, G. P. 2003 Epistasis in polygenic traits and the evolution of genetic architecture under stabilizing selection. *Am. Nat.* **161**, 708–734.
- Herring, S. W. 1993 Epigenetic and functional influences on skull growth. In *The skull, vol. 1* (ed. J. Hanken & B. K. Hall), pp. 153–206. Chicago and London: University of Chicago Press.
- Hinton, S. J. & Nolan, G. E. 1980 How learning can guide evolution. *Complex Syst.* **1**, 495–502.
- Ho, M. W., Tucker, C., Keeley, D. & Saunders, P. T. 1983 Effects of successive generations of ether treatment on penetrance and expression of the *Bithorax* phenocopy in *Drosophila melanogaster*. *J. Exp. Zool. (Mol. Dev. Evol.)* **225**, 357–368.
- Hoffmann, A. A. & Hercus, M. J. 2000 Environmental stress as an evolutionary force. *BioScience* **50**, 217–226.

- Hoffmann, A. A. & Merilä, J. 1999 Heritable variation and evolution under favourable and unfavourable conditions. *Trends Ecol. Evol.* **14**, 96–101.
- Hoffmann, A. A. & Parsons, P. A. 1997 *Extreme environmental change and evolution*. Cambridge University Press.
- Howarth, F. G. 1993 High-stress subterranean habitats and evolutionary change in cave-inhabiting arthropods. *Am. Nat.* **142**(Suppl.), S65–S77.
- Huether, G. 1996 The central adaptation syndrome: psychosocial stress as a trigger for adaptive modifications of brain structure and brain function. *Prog. Neurobiol.* **48**, 569–612.
- Huey, R. B., Hertz, P. E. & Sinervo, B. 2003 Behavioral drive versus behavioral inertia in evolution: a null model approach. *Am. Nat.* **161**, 357–366.
- Imasheva, A. G. 1999 Environmental stress and genetic variation in animal populations. *Russ. J. Genet.* **35**, 343–351.
- Innan, H. & Kim, Y. 2004 Pattern of polymorphism after strong artificial selection in a domestication event. *Proc. Natl Acad. Sci. USA* **101**, 10 667–10 672.
- Jablonka, E. & Lamb, M. J. 1995 *Epigenetic inheritance and evolution: the Lamarckian dimension*. Oxford University Press.
- Jablonka, E., Oborny, B., Molnar, I., Kisdi, E., Hofbauer, J. & Czaran, T. 1995 The adaptive advantage of phenotypic memory in changing environments. *Phil. Trans. R. Soc. B* **350**, 133–141.
- Johnson, E. O., Kamilaris, T. C., Chrousos, G. P. & Gold, P. W. 1992 Mechanisms of stress: a dynamic overview of hormonal and behavioral homeostasis. *Neurosci. Biobehav. Rev.* **16**, 115–130.
- Karunanithi, S., Barclay, J. W., Robertson, R. M., Brown, I. R. & Atwood, H. L. 1999 Neuroprotection of *Drosophila* synapses conferred by prior heat shock. *J. Neurosci.* **19**, 4360–4369.
- Katz, R. J., Roth, K. A. & Carrol, B. J. 1981 Acute and chronic stress effects on open field activity in the rats. Implication for a model of depression. *Neurosci. Biobehav. Rev.* **5**, 247–251.
- Keller, L. F., Grant, P. R., Grant, B. R. & Petren, K. 2002 Environmental conditions affect the magnitude of inbreeding depression in survival of Darwin's finches. *Evolution* **56**, 1229–1239.
- Kirschner, M. & Gerhart, J. 1998 Evolvability. *Proc. Natl. Acad. Sci. USA* **95**, 8420–8427.
- Kishony, R. & Leibler, S. 2003 Environmental stresses can alleviate the average deleterious effect of mutations. *J. Biol.* **2**, 14–19.
- Klingenberg, C. P., Badyaev, A. V., Sowry, S. M. & Beckwith, N. J. 2001 Inferring developmental modularity from morphological integration: analysis of individual variation and asymmetry in bumblebee wings. *Am. Nat.* **157**, 11–23.
- Kondrashov, A. S. & Houle, D. 1994 Genotype–environment interactions and the estimation of the genomic mutation rate in *Drosophila melanogaster*. *Proc. R. Soc. B* **258**, 221–227.
- Lauter, N. & Doebley, J. 2002 Genetic variation for phenotypically invariant traits detected in teosine: implications for the evolution of novel forms. *Genetics* **160**, 333–342.
- Leamy, L. J. 1993 Morphological integration of fluctuating asymmetry in the mouse mandible. *Genetica* **89**, 139–153.
- Leips, J. & MacKay, T. F. C. 2000 Quantitative trait loci for life span in *Drosophila melanogaster*: interactions with genetic background and larval density. *Genetics* **160**, 333–342.
- Levine, S., Haltmeyer, G. C., Karas, G. G. & Denenberg, V. H. 1967 Physiological and behavioral effects of infantile stimulation. *Physiol. Behav.* **2**, 55–63.
- Levine, S., Coe, C. H. & Wiener, S. 1989 Psychoneuroendocrinology of stress: a psychobiological perspective. In *Psychoendocrinology* (ed. F. R. Brush & S. Levine), pp. 341–380. San Diego: Academic Press.
- Levins, R. 1963 Theory of fitness in a heterogeneous environment. II. Developmental flexibility and niche selection. *Am. Nat.* **97**, 75–90.
- Mabee, P. M., Olmstead, K. L. & Cabbage, C. C. 2000 An experimental study of intraspecific variation, developmental timing, and heterochrony in fishes. *Evolution* **54**, 2091–2106.
- McClintock, B. 1984 The significance of responses of the genome to challenge. *Science* **226**, 792–801.
- McEwen, B. S. & Sapolsky, R. M. 1995 Stress and cognitive function. *Curr. Opin. Neurobiol.* **5**, 205–216.
- McEwen, B. S. & Wingfield, J. C. 2003 The concept of allostasis in biology and biomedicine. *Horm. Behav.* **43**, 2–15.
- McGaugh, J. L., Martinez, J. L., Jensen, R. A., Hannan, T. J., Vasquez, B. J., Messing, R. B., Liang, K. C., Brewton, K. C. & Spiehl, V. 1982 Modulation of memory storage by treatments affecting peripheral catecholamines. In *Neuronal plasticity and memory formation* (ed. C. Ajmone-Marsan & H. Matthies), pp. 129–141. New York: Raven Press.
- Masel, J. 2004 Genetic assimilation can occur in the absence of selection for the assimilating phenotype, suggesting a role for the canalization heuristic. *J. Evol. Biol.* **17**, 1106–1110.
- Masel, J. & Bergman, A. 2003 The evolution of evolvability properties of the yeast prion [PSI+]. *Evolution* **57**, 1498–1512.
- Meaney, M. J. 2001 Maternal care, gene expression, and the transmission of individual differences in stress reactivity across generations. *Annu. Rev. Neurosci.* **24**, 1161–1192.
- Meiklejohn, C. D. & Hartl, D. L. 2002 A single mode of canalization. *Trends Ecol. Evol.* **17**, 468–473.
- Milkman, R. D. 1965 Genetic basis of natural variation. VI. Selection of a crossveinless strain of *Drosophila* by phenocopying at high temperatures. *Genetics* **51**, 87–96.
- Møller, A. P. & Swaddle, J. P. 1997 *Asymmetry, developmental stability, and evolution*. Oxford University Press.
- Mousseau, T. A. & Fox, C. W. 1998 The adaptive significance of maternal effects. *Trends Ecol. Evol.* **13**, 403–407.
- Muller, G. B. 2003 Embryonic motility: environmental influences and evolutionary innovation. *Evol. Dev.* **5**, 56–60.
- Nanjundiah, V. 2003 Phenotypic plasticity and evolution by genetic assimilation. In *Origination of organismal form: beyond the gene in developmental and evolutionary biology* (ed. G. B. Muller & S. Newman), pp. 245–263. Cambridge, MA: MIT Press.
- Nedelcu, A. M. & Michod, R. E. 2003 Sex as a response to oxidative stress: the effect of antioxidants on sexual induction in a facultatively sexual lineage. *Proc. R. Soc. B* **270**(Suppl. 2), S136–S139. (doi:10.1098/rsbl.2003.0062)
- Nicolakakis, N., Sol, D. & Lefebvre, L. 2003 Behavioural flexibility predicts species richness in birds, but not extinction risk. *Anim. Behav.* **65**, 445–452.
- Odling-Smee, F. J., Laland, K. N. & Feldman, M. W. 2003 *Niche construction: the neglected process in evolution*. Princeton University Press.
- Oyama, S. 2000 *The ontogeny of information: developmental systems and evolution*. Durham, NC: Duke University Press.
- Palmer, A. R. 2004 Symmetry breaking and the evolution of development. *Science* **306**, 828–833.
- Parsons, P. A. 1988 Evolutionary rates: effects of stress upon recombination. *Biol. J. Linn. Soc.* **35**, 49–68.



- Parsons, P. A. 1990 Fluctuating asymmetry: an epigenetic measure of stress. *Biol. Rev.* **65**, 131–145.
- Parsons, P. A. 1993 The importance and consequences of stress in living and fossil populations: from life-history variation to evolutionary change. *Am. Nat.* **142**(Suppl.), S5–S20.
- Parsons, P. A. 1994 Morphological stasis: an energetic and ecological perspective incorporating stress. *J. Theor. Biol.* **171**, 409–414.
- Pelabon, C., Carlson, M., Hansen, T. F., Yoccoz, N. G. & Armbruster, W. S. 2004 Consequences of inter-population crosses on developmental stability and canalization of floral traits in *Dalechampia scandens* (Euphorbiaceae). *J. Evol. Biol.* **17**, 19–32.
- Piersma, T. & Drent, J. 2003 Phenotypic flexibility and the evolution of organismal design. *Trends Ecol. Evol.* **18**, 228–233.
- Piffanell, P., Ramsay, L., Waugh, R., Benabdelmouna, A., D'Hont, A., Hollricher, K. & Jørgensen, J. H. 2004 A barley cultivation-associated polymorphism conveys resistance to powdery mildew. *Nature* **430**, 887–891.
- Polak, M. 2003 *Developmental instability: causes and consequences*. Oxford University Press.
- Price, T. D., Qvarnström, A. & Irwin, D. E. 2003 The role of phenotypic plasticity in driving genetic evolution. *Proc. R. Soc. B* **270**, 1433–1440. (doi:10.1098/rspb.2003.2372)
- Prody, C. A., Dreyfus, P., Zamir, R., Zakut, H. & Soreq, H. 1989 *De novo* amplification within a “silent” human cholinesterase gene in a family subjected to prolonged exposure to organophosphorus insecticides. *Proc. Natl Acad. Sci. USA* **86**, 690–694.
- Queitsch, C., Sangster, T. A. & Lindquist, S. 2002 Hsp90 as a capacitor of phenotypic variation. *Nature* **418**, 618–624.
- Råberg, L., Grahn, M., Hasselquist, D. & Svensson, E. 1998 On the adaptive significance of stress-induced immunosuppression. *Proc. R. Soc. B* **265**, 1637–1641. (doi:10.1098/rspb.1998.0482)
- Ramirez, J. M., Elsen, F. P. & Robertson, R. M. 1999 Long-term effects of prior heat shock on neuronal potassium currents recorded in a novel insect ganglion slice preparation. *J. Neurophysiol.* **81**, 795–802.
- Rather, V. A., Zabanov, S. A., Kolesnikova, O. V. & Vasilyeva, L. A. 1992 Induction of the mobile genetic element Dm-412 transpositions in the *Drosophila* genome by heat shock treatment. *Proc. Natl Acad. Sci. USA* **89**, 5650–5654.
- Rice, S. H. 1998 Evolution of canalization and the breaking of von Baer's laws: modeling the evolution of development with epistasis. *Evolution* **52**, 647–656.
- Rice, S. H. 2004 Developmental associations between traits: covariance and beyond. *Genetics* **166**, 513–526.
- Rieseberg, L. H. *et al.* 2003 Major ecological transitions in wild sunflowers facilitated by hybridization. *Science* **301**, 1211–1216.
- Robertson, R. M. 2004 Modulation of neural circuit operation by prior environmental stress. *Integr. Comp. Biol.* **44**, 21–27.
- Romero, L. M. 2004 Physiological stress in ecology: lessons from biomedical research. *Trends Ecol. Evol.* **19**, 249–255.
- Rossiter, M. 1996 Incidence and consequences of inherited environmental effects. *Annu. Rev. Ecol. Syst.* **27**, 451–476.
- Roth, G. & Wake, D. B. 1985 Trends in the functional morphology and sensorimotor control of feeding behavior in salamanders: an example of the role of internal dynamics in evolution. *Acta Biotheor.* **34**, 175–192.
- Roth, E. J., Frazier, B. L., Apuya, N. R. & Lark, K. G. 1989 Genetic variation in an inbred plant: variation in tissue cultures of soybean *Glycine max* L. *Genetics* **121**, 359–368.
- Ruden, D. M., Garfinkel, M. D., Sollars, V. E. & Lu, X. 2003 Waddington's widget: Hsp90 and the inheritance of acquired characters. *Semin. Cell Dev. Biol.* **14**, 301–310.
- Rutherford, S. L. 2000 From genotype to phenotype: buffering mechanisms and the storage of genetic information. *BioEssays* **22**, 1095–1105.
- Rutherford, S. L. 2003 Between genotype and phenotype: protein chaperones and evolvability. *Nature Rev.* **4**, 263–274.
- Rutherford, S. L. & Lindquist, S. 1998 Hsp90 as a capacitor for morphological evolution. *Nature* **396**, 336–343.
- Ruvinsky, A. O., Lobkov, Y. I. & Belyaev, D. K. 1983 Spontaneous and induced activation of genes affecting the phenotypic expression of glucose 6-phosphate dehydrogenase in *Daphnia pulex*. II. Glucose induced changes in the electrophoretic mobility of G6PD. *Mol. Gen. Genet.* **189**, 490–494.
- Schandorff, S. 1997 Developmental stability and skull lesions in the harbour seal (*Phoca vitulina*) in the 19th and 20th centuries. *Ann. Zool. Fennici* **34**, 151–166.
- Scharloo, W. 1991 Canalization: genetic and developmental aspects. *Annu. Rev. Ecol. Syst.* **22**, 65–93.
- Schlichting, C. D. & Pigliucci, M. 1998 *Phenotypic evolution: a reaction norm perspective*. Sunderland, MA: Sinauer Associates.
- Schlichting, C. D. & Smith, H. 2002 Phenotypic plasticity: linking mechanisms with evolutionary outcomes. *Evol. Ecol.* **16**, 189–211.
- Schmalhausen, I. I. 1949 *Factors of evolution*. Philadelphia, PA: Blakiston.
- Sciulli, P. W., Doyle, W. J., Kelley, C., Siegel, P. & Siegel, M. I. 1979 The interaction of stressors in the induction of increased levels of fluctuating asymmetry in the laboratory rat. *Am. J. Phys. Anthropol.* **50**, 279–284.
- Seehausen, O. 2004 Hybridization and adaptive radiation. *Trends Ecol. Evol.* **19**, 198–206.
- Selker, E. U. 1990 Premeiotic instability of repeated sequences in *Neurospora crassa*. *Annu. Rev. Genet.* **24**, 579–613.
- Shapiro, J. A. 1992 Natural genetic engineering in evolution. *Genetica* **86**, 99–111.
- Shors, T. J. & Servatius, R. J. 1997 The contribution of stressor intensity, duration, and context to the stress-induced facilitation of associative learning. *Neurobiol. Learn. Mem.* **67**, 92–96.
- Sibly, R. M. & Calow, P. 1989 A life-cycle theory of responses to stress. *Biol. J. Linn. Soc.* **37**, 101–116.
- Siegal, M. L. & Bergman, A. 2002 Waddington's canalization revisited: developmental stability and evolution. *Proc. Natl Acad. Sci. USA* **99**, 10 528–10 532.
- Simons, A. M. & Johnston, M. O. 1997 Developmental instability as a bet-hedging strategy. *Oikos* **80**, 401–405.
- Sniegowski, P., Gerrish, P., Johnson, T. & Shever, A. 2000 The evolution of mutation rates: separating causes from consequences. *BioEssays* **22**, 1057–1066.
- Sockman, K. W., Gentner, T. Q. & Ball, G. F. 2002 Recent experience modulates forebrain gene-expression in response to mate-choice cues in European starlings. *Proc. R. Soc. B* **269**, 2479–2485. (doi:10.1098/rspb.2002.2180)
- Sockman, K. W., Williams, T. D., Dawson, A. & Ball, G. F. 2004 Prior experience with photostimulation enhances photo-induced reproductive development in female European starlings: a possible basis for the age-related increase in avian reproductive performance. *Biol. Reprod.* **71**, 979–986.
- Sollars, V., Lu, X., Xiao, L., Wang, X., Garfinkel, M. D. & Ruden, D. M. 2003 Evidence for an epigenetic mechanism by which Hsp90 acts as a capacitor for morphological evolution. *Nature Genet.* **33**, 70–74.

- Stanton, M. L., Roy, B. A. & Thiede, D. A. 2000 Evolution in stressful environments. I. Phenotypic variability, phenotypic selection, and response to selection in five distinct environmental stresses. *Evolution* **54**, 93–111.
- Stearns, S. C. & Kawecki, T. J. 1994 Fitness sensitivity and the canalization of life-history traits. *Evolution* **48**, 1438–1450.
- Steele, D. F. & Jinks-Robertson, S. 1992 An examination of adaptive reversion in *Saccharomyces cerevisiae*. *Genetics* **132**, 9–21.
- Szafraniec, K., Borts, R. H. & Korona, R. 2001 Environmental stress and mutational load in diploid strains of the yeast *Saccharomyces cerevisiae*. *Proc. Natl Acad. Sci. USA* **98**, 1107–1112.
- Threadgill, D. W. *et al.* 1995 Targeted disruption of mouse EGF receptor: effect of genetic background on mutant phenotype. *Science* **269**, 230–234.
- True, H. L., Berlin, I. & Lindquist, S. L. 2004 Epigenetic regulation of translation reveals hidden genetic variation to produce complex traits. *Nature* **431**, 184–187.
- Veenema, A. H., Meijer, O. C., de Kloet, E. R., Koolhaas, J. M. & Bohus, B. G. 2003 Differences in basal and stress-induced HPA regulation of wild house mice selected for high and low aggression. *Horm. Behav.* **43**, 197–204.
- Waddington, C. H. 1941 Evolution of developmental systems. *Nature* **147**, 108–110.
- Waddington, C. H. 1953 Genetic assimilation of an acquired character. *Evolution* **7**, 119–127.
- Wagner, G. P. & Mezey, J. G. 2000 Modeling the evolution of genetic architecture: a continuum of alleles model with pairwise A×A epistasis. *J. Theor. Biol.* **203**, 163–175.
- Wagner, G. P., Booth, G. & Bacheri-Chaichian, H. 1997 A population genetic theory of canalization. *Evolution* **51**, 329–347.
- Walker, I. 1979 The mechanical properties of proteins determine the laws of evolutionary change. *Acta Biotheor.* **28**, 239–282.
- Waxman, D. & Peck, J. R. 1998 Pleiotropy and the preservation of perfection. *Science* **279**, 1210–1213.
- Wessler, S. R. 1996 Plant retrotransposons: turned on by stress. *Curr. Biol.* **6**, 959–961.
- West-Eberhard, M. J. 2003 *Developmental plasticity and evolution*. Oxford University Press.
- Wills, C. 1983 The possibility of stress-triggered evolution. *Lect. Notes Biomath.* **53**, 299–312.
- Wingfield, J. C. 2003 Control of behavioural strategies for capricious environments. *Anim. Behav.* **66**, 807–816.
- Wingfield, J. C. & Sapolsky, R. M. 2003 Reproduction and resistance to stress: when and how. *J. Neuroendocrinol.* **15**, 711–724.
- Wright, B. E. 2000 Minireview: a biochemical mechanism for nonrandom mutations and evolution. *J. Bacteriol.* **182**, 2993–3002.
- Wright, B. E. 2004 Stress-directed adaptive mutations and evolution. *Mol. Microbiol.* **52**, 643–650.
- Wu, B. S., Lee, J. K., Thompson, K. M., Walker, V. K., Moyes, C. D. & Robertson, R. M. 2002 Anoxia induces thermotolerance in the locust flight system. *J. Exp. Biol.* **205**, 815–827.