



## SPECIAL REPORT

# Augmented acetylcholine-induced, Rho-mediated $\text{Ca}^{2+}$ sensitization of bronchial smooth muscle contraction in antigen-induced airway hyperresponsive rats

\*<sup>1</sup>Yoshihiko Chiba, <sup>1</sup>Yuka Takada, <sup>2</sup>Shigeki Miyamoto, <sup>2</sup>Minori Mitsui-Saito, <sup>2</sup>Hideaki Karaki & <sup>1</sup>Miwa Misawa

<sup>1</sup>Department of Pharmacology, School of Pharmacy, Hoshi University, 2-4-41 Ebara, Shinagawa-ku, Tokyo 142-8501, Japan and

<sup>2</sup>Department of Veterinary Pharmacology, Graduate School of Agriculture and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

Treatment with acetylcholine (ACh) of a  $\beta$ -escin-permeabilized intrapulmonary bronchial smooth muscle of the rat induced force when the  $\text{Ca}^{2+}$  concentration was clamped at  $1 \mu\text{M}$ . The ACh-induced  $\text{Ca}^{2+}$  sensitization of myofilaments was significantly greater in antigen-induced airway hyperresponsive rats than in control rats. The ACh-induced  $\text{Ca}^{2+}$  sensitization was completely blocked by treatment with *Clostridium botulinum* C3 exoenzyme, an inactivator of Rho family of proteins. Moreover, the protein level of RhoA in the intrapulmonary bronchi was significantly increased in the airway hyperresponsive rats. Thus, increased airway smooth muscle contractility observed in asthmatics may be related to augmented agonist-induced, Rho-mediated  $\text{Ca}^{2+}$  sensitization of myofilaments.

**Keywords:** Asthma; airway hyperresponsiveness; intrapulmonary bronchial smooth muscle;  $\text{Ca}^{2+}$  sensitization; Rho;  $\beta$ -escin

**Abbreviations:** ACh, acetylcholine; AHR, airway hyperresponsiveness; DNP-Asc, 2,4-dinitrophenylated *Ascaris suum* extract; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; pCa,  $-\log [\text{Ca}^{2+}]$

**Introduction** In general, smooth muscle contraction has been thought to be induced by an increase in cytosolic  $\text{Ca}^{2+}$  via the activation of plasma membrane  $\text{Ca}^{2+}$  channels and/or  $\text{Ca}^{2+}$  release from sarcoplasmic reticulum. However, additional mechanisms have been suggested in agonist-induced smooth muscle contraction by studies which used the simultaneous measurements of force development and intracellular  $\text{Ca}^{2+}$  concentration (Sato *et al.*, 1988), and chemically permeabilized preparations (Fujita *et al.*, 1995) in various types of smooth muscles including airways (Ozaki *et al.*, 1990). It has been demonstrated that agonist stimulation increases myofilament  $\text{Ca}^{2+}$  sensitivity in  $\beta$ -escin-permeabilized smooth muscles of the rat coronary artery (Satoh *et al.*, 1994), guinea-pig vas deferens (Fujita *et al.*, 1995), canine trachea (Bremerich *et al.*, 1997), and so on. Although the detailed mechanism is not fully understood, a participation of Rho protein, a monomeric GTP binding protein, in the agonist-induced  $\text{Ca}^{2+}$  sensitization has been suggested by many investigators (e.g., Fujita *et al.*, 1995; Otto *et al.*, 1996; Gong *et al.*, 1997).

Asthmatic patients have an increased contractility of airway smooth muscle (Roberts *et al.*, 1984), which might be a major cause of airway hyperresponsiveness (AHR). Similarly, an increased responsiveness of bronchial smooth muscle has been demonstrated in a rat model of AHR induced by repeated antigen inhalation (Misawa & Chiba, 1993; Chiba & Misawa, 1995a,b). In this animal model of AHR, the bronchial smooth muscle contraction induced by receptor agonists such as acetylcholine (ACh), but not by high  $\text{K}^+$  depolarization, is markedly augmented (Misawa & Chiba, 1993; Chiba & Misawa, 1995a,b). Moreover, it has also been demonstrated that muscarinic receptor density and antagonist affinity of airway smooth muscle are normal (Chiba & Misawa, 1995a).

Thus, it is possible that the mechanisms responsible for the AHR exist, at least in part, in the downstream pathway of muscarinic receptor signaling, including ACh-mediated  $\text{Ca}^{2+}$  sensitization.

In the present study, we examined participation of ACh-induced  $\text{Ca}^{2+}$  sensitization in the augmented contraction of airway smooth muscle at the antigen-induced AHR by using  $\beta$ -escin-permeabilized muscle strips.

**Methods** Male Wistar rats (170–190 g, specific pathogen-free) were used. The induction of AHR was performed as described previously (Chiba & Misawa, 1995a,b). Briefly, the rats were sensitized with 2,4-dinitrophenylated *Ascaris suum* extract (DNP-Asc) together with *Bordetella pertussis* and were boosted 5 days later. Eight days after the first immunization, the rats were challenged by inhaling DNP-Asc for 20 min under conscious state. Then the animals were subjected to totally three times repeated antigen challenge every 48 h with the same inhalational challenge method.

Twenty-four hours after the last antigen challenge, the third branch of intrapulmonary bronchus was isolated, carefully cleaned of lung parenchyma and adhering connective tissue, and then cut into ring strips (about 200  $\mu\text{m}$  width, 500  $\mu\text{m}$  diameter). The epithelium was removed by gently rubbing with keen-edged tweezers. The ring strips were then permeabilized by a 30-min treatment with 10  $\mu\text{M}$   $\beta$ -escin (Sigma) at room temperature in relaxing solution.

Relaxing solution contained: (in mM): PIPES 20,  $\text{Mg}^{2+}$ -dimethanesulphonate 7.1,  $\text{K}^+$ -methanesulphonate 108, EGTA 2,  $\text{Na}_2\text{ATP}$  5.875, creatine phosphate 2, creatine phosphokinase 4  $\text{u ml}^{-1}$ , carbonyl cyanide p-trifluoromethoxyphenylhydrazone (FCCP) 1  $\mu\text{M}$  and E-64 (pH 6.8) 1  $\mu\text{g ml}^{-1}$  containing 10  $\mu\text{M}$  A23187. Free  $\text{Ca}^{2+}$  concentration was changed by adding an appropriate amount of  $\text{CaCl}_2$ . The apparent binding constant of EGTA for  $\text{Ca}^{2+}$  was considered to be  $10^6 \text{ M}^{-1}$

\*Author for correspondence; E-mail: [chiba@hoshi.ac.jp](mailto:chiba@hoshi.ac.jp)

(Hori *et al.*, 1993). The permeabilized muscle strip was then suspended in a 400  $\mu$ l organ bath at room temperature. The contractile force developed was measured by an isometric transducer under a resting tension of 50 mg. To determine the involvement of Rho in the ACh-induced myofilament Ca<sup>2+</sup> sensitization, the  $\beta$ -escin-permeabilized muscle strips were treated with *Clostridium botulinum* C3 exoenzyme (1  $\mu$ g ml<sup>-1</sup>; Calbiochem) in the presence of 100  $\mu$ M NAD for 20 min at room temperature.

To quantify the expression of Rho proteins, Western blot was performed in the homogenates of intrapulmonary bronchi that were dissected from lung parenchyma. Briefly, the samples (10  $\mu$ g of total protein per lane) were subjected to 12% SDS-PAGE and the proteins were then electrophoretically transferred to a nitrocellulose membrane. After blocking with 3% gelatin, the nitrocellulose membrane was incubated with primary antibody (polyclonal rabbit anti-human RhoA [amino acids 119–132] or anti-human RhoB [amino acids 119–132]; 1:2500 dilution, respectively; Santa Cruz). Then the membrane was incubated with horseradish peroxidase-conjugated goat anti-rabbit IgG (1:2500 dilution; Amersham), detected by an enhanced chemiluminescent system (Amersham) and analysed by a densitometry system. Thereafter, the primary and secondary antibodies were stripped and the membrane was reprobed by using monoclonal mouse anti-glyceraldehyde-3-phosphate dehydrogenase (GAPDH; 1:1000 dilution; Chemicon) to confirm the same amount of proteins loaded. Rat whole brain was used as a positive control for RhoA and RhoB (Olenic *et al.*, 1997).

All the data are expressed as the mean  $\pm$  s.e.mean. Statistical significance of difference was determined by Dunnett's multiple analysis.

**Results** Our previous study revealed that the sensitization procedure to antigen used in the present study had no significant effect by itself on the ACh responsiveness of the bronchial muscle and muscarinic receptors property (Chiba & Misawa, 1995a) in rats. So in the present study, the age-matched nonsensitized normal rats were used as control.

In all tissue preparations treated with 10  $\mu$ M  $\beta$ -escin (for 30 min), the application of free Ca<sup>2+</sup> (pCa = 6.5, 6.0, 5.5 and 5.0) induced a concentration-dependent reproducible contractile response, indicating successful permeabilization. In the  $\beta$ -escin-permeabilized intrapulmonary bronchial smooth muscle, no significant difference between groups was observed in the Ca<sup>2+</sup> responsiveness and the maximal contractile response induced by pCa = 5.0 (Table 1). When the Ca<sup>2+</sup> concentration

**Table 1** Summary of the Ca<sup>2+</sup>-induced contractile responses of the permeabilized bronchial smooth muscle in rats

Groups	n	$-\log EC_{50}[Ca^{2+}]$ (M)	Max. response (mg)
Control	6	6.01 $\pm$ 0.09	115.5 $\pm$ 9.4
AHR	5	5.90 $\pm$ 0.12	112.4 $\pm$ 24.8


Intrapulmonary bronchial smooth muscles were isolated from normal (Control) and the repeatedly antigen challenged rats (AHR) and were permeabilized by 30-min treatment with 10  $\mu$ M  $\beta$ -escin. The isometric force developed was measured. Data represent the mean  $\pm$  s.e.mean. EC<sub>50</sub>[Ca<sup>2+</sup>]; 50% effective concentration of Ca<sup>2+</sup> (M), Max. response; maximal contractile responses induced by 10<sup>-5</sup> M Ca<sup>2+</sup>. No significant difference in the Ca<sup>2+</sup>-induced contractile response was observed between groups.

was clamped at pCa = 6.0, the application of 100  $\mu$ M ACh in the presence of 100  $\mu$ M GTP caused a further contraction (i.e. ACh-induced Ca<sup>2+</sup> sensitization) in both the control and AHR groups (Figure 1). However, ACh-induced Ca<sup>2+</sup> sensitization was significantly greater in the AHR group (Figure 1). The ACh-induced Ca<sup>2+</sup> sensitization was completely antagonized by 10<sup>-6</sup> M atropine (data not shown).


To determine the involvement of Rho proteins in the ACh-induced Ca<sup>2+</sup> sensitization, the effect of pretreatment with C3 exoenzyme, an inactivator of Rho family of proteins (Fujita *et al.*, 1995), on the contractile responses of the  $\beta$ -escin-permeabilized muscle was estimated. In both groups, the C3 treatment had no effect on the contractile response induced by pCa = 6.0 (typical recordings being shown in Figure 2). However, the ACh-induced Ca<sup>2+</sup> sensitizing effect was completely blocked after the treatment with C3 both in the AHR and normal rats (Figure 2). These findings indicate that the ACh-induced Ca<sup>2+</sup> sensitization might be mediated by Rho proteins and that the Rho-mediated Ca<sup>2+</sup> sensitizing effect might be augmented at the AHR.

Although Rho family proteins comprise RhoA, RhoB, Rac, Cdc42, and so on, a strong involvement of RhoA in agonist-induced smooth muscle Ca<sup>2+</sup> sensitization has been reported (Otto *et al.*, 1996; Gong *et al.*, 1997). Therefore, we assessed the expression of RhoA in the homogenates of the intrapulmonary bronchi by using immunoblotting. As shown in Figure 3, immunoblotting with the antibody against RhoA gave a single 21 kD band, indicating the existence of RhoA protein in the intrapulmonary bronchi of rats. The level of RhoA in samples from the antigen-induced AHR rats was significantly increased compared with that from control rats (Figure 3). On the other hand, RhoB was not detected in the samples from either group (data not shown).


**Discussion** In the present study, the Ca<sup>2+</sup>-induced contractile responses (in the absence of ACh and GTP) of  $\beta$ -escin-permeabilized intrapulmonary bronchial smooth muscles from the AHR rats were normal (Table 1, Figure 1). This finding is in agreement with our previous study that the contractile



**Figure 1** Comparison of ACh-induced Ca<sup>2+</sup> sensitization in  $\beta$ -escin-permeabilized intrapulmonary bronchial smooth muscle from normal rats (Control; *n* = 6) and the antigen-induced airway hyperresponsiveness rats (AHR; *n* = 5). The contractile responses induced by 10<sup>-6</sup> M Ca<sup>2+</sup> in the presence (closed column) and absence (open column) of 100  $\mu$ M ACh and 100  $\mu$ M GTP are expressed as percentage of maximal contraction induced by 10<sup>-5</sup> M Ca<sup>2+</sup>. Data represent the mean  $\pm$  s.e.mean. \**P* < 0.05 and \*\**P* < 0.01 vs respective Ca<sup>2+</sup>-induced contraction in the absence of ACh and GTP. ###*P* < 0.01 vs ACh-induced Ca<sup>2+</sup> sensitization in the control group.



**Figure 2** Effect of *Clostridium botulinum* C3 exoenzyme on the ACh-induced Ca<sup>2+</sup> sensitization of the  $\beta$ -escin-permeabilized intrapulmonary bronchial smooth muscle from the antigen-induced airway hyperresponsive rats. Left panel: typical recordings. After the permeabilization with 10  $\mu$ M  $\beta$ -escin, the Ca<sup>2+</sup> (10<sup>-6</sup> M)-induced contractile responses in the presence and absence of 100  $\mu$ M ACh and 100  $\mu$ M GTP were observed as indicated (before C3). Then the strip was incubated with C3 exoenzyme (1  $\mu$ g ml<sup>-1</sup>, for 20 min) in relaxing solution (see Methods), and the contractile responses were re-estimated (After C3). Right panel: summary of the inhibition of ACh-induced Ca<sup>2+</sup> sensitization by C3 exoenzyme. The data are expressed as percentage increase in tension induced by ACh (in the presence of Ca<sup>2+</sup> and GTP) from the sustained contraction induced by Ca<sup>2+</sup> (10<sup>-6</sup> M). Data represent the mean  $\pm$  s.e. mean from four experiments. \*\**P* < 0.01 vs before C3.



**Figure 3** The levels of RhoA protein in intrapulmonary bronchi from normal rats (Control) and the antigen-induced airway hyperresponsive rats (AHR). Upper panel: typical immunoblot. Lane 1; control, Lane 2; AHR, markers; protein molecular weight markers, and GAPDH; glyceraldehyde-3-phosphate dehydrogenase as a tissue marker. The bands were analysed by a densitometer and normalized by loading protein, and the data are summarized as shown in the lower panel. The data represent the mean  $\pm$  s.e. mean from four individual experiments, respectively. \*\**P* < 0.01 vs control.

response of intact (non-permeabilized) bronchial smooth muscle induced by high K<sup>+</sup> depolarization was normal in AHR (Chiba & Misawa, 1995b). Taken together, the baseline Ca<sup>2+</sup> sensitivity (no receptor stimulation) of contractile elements is unlikely to change in AHR.

In the presence of GTP, ACh elicited a further contraction of the  $\beta$ -escin-permeabilized intrapulmonary bronchial smooth muscles from control rats, even though the Ca<sup>2+</sup> concentration was clamped (Figure 1). This effect was antagonized by atropine, indicating existence of a muscarinic receptor-mediated Ca<sup>2+</sup> sensitizing mechanism. Likewise, muscarinic receptor-mediated Ca<sup>2+</sup> sensitization has also been reported in permeabilized smooth muscles of the canine (Bremerich *et al.*, 1997) and porcine trachea (Croxtton *et al.*, 1998). So, it is possible that the agonist-induced Ca<sup>2+</sup> sensitization might occur in airway smooth muscles of other species. In the current study, the ACh-induced Ca<sup>2+</sup> sensitization was completely blocked by treatment with C3 exoenzyme, a selective inactivator of Rho family proteins (Fujita *et al.*, 1995). Although the signal transduction pathways involved in the regulation of Ca<sup>2+</sup> sensitization appear complex and, partly smooth muscle type specific, participation of Rho proteins has been reported in various types of smooth muscle including airways (Croxtton *et al.*, 1998). A part of the mechanism by which Rho mediated Ca<sup>2+</sup> sensitization has been demonstrated as the Rho-associated kinase phosphorylates the 20 kD myosin light chain (Amano *et al.*, 1996) and the application of Rho-associated kinase to permeabilized smooth muscle induces contraction (Kureishi *et al.*, 1997).

ACh-induced Ca<sup>2+</sup> sensitizing effect was augmented in bronchial smooth muscle from the AHR rats, the effect was blocked by C3 exoenzyme, and the protein level of RhoA in AHR was increased. This is the first study, to our knowledge, that demonstrates an augmentation of ACh-induced, Rho (probably RhoA)-mediated Ca<sup>2+</sup> sensitization, which coincides with enhanced RhoA protein expression. An increase in

responsiveness to muscarinic agonists of airway smooth muscle has been reported in animal models of AHR (Misawa & Chiba, 1993; Gavett *et al.*, 1993; Lee *et al.*, 1994; Chiba & Misawa, 1995a,b) and asthmatic patients (Roberts *et al.*, 1984), although no change in the levels of muscarinic receptors was observed (Gavett *et al.*, 1993; Lee *et al.*, 1994; Chiba & Misawa, 1995a). Thus, it is likely that the enhanced contractility to muscarinic agonists reflects the augmentation of muscarinic receptor- and Rho-mediated Ca<sup>2+</sup> sensitization. The mechanism(s) for the activation of Rho proteins by ACh is currently unclear. If Rho proteins are activated *via* receptors other than muscarinic receptor, it might account for the 'non-specific' AHR, which is a common feature of asthmatics.

## References

- AMANO, M., ITO, M., KIMURA, K., FUKATA, Y., CHIHARA, K., NAKANO, T., MATSUURA, Y. & KAIBUCHI, K. (1996). Phosphorylation and activation of myosin by Rho-associated kinase (Rho-kinase). *J. Biol. Chem.*, **271**, 20246–20249.
- BREMERICH, D.H., WARNER, D.O., LORENZ, R.R., SHUMWAY, R. & JONES, K.A. (1997). Role of protein kinase C in calcium sensitization during muscarinic stimulation in airway smooth muscle. *Am. J. Physiol.*, **273**, L775–L781.
- CHIBA, Y. & MISAWA, M. (1995a). Characteristics of muscarinic cholinergic receptors in airways of antigen-induced airway hyperresponsive rats. *Comp. Biochem. Physiol.*, **111C**, 351–357.
- CHIBA, Y. & MISAWA, M. (1995b). Alteration in Ca<sup>2+</sup> availability involved in antigen-induced airway hyperresponsiveness in rats. *Eur. J. Pharmacol.*, **278**, 79–82.
- CROXTON, T.L., LANDE, B. & HIRSHMAN, C.A. (1998). Role of G proteins in agonist-induced Ca<sup>2+</sup> sensitization of tracheal smooth muscle. *Am. J. Physiol.*, **275**, L748–L755.
- FUJITA, A., TAKEUCHI, T., NAKAJIMA, H., NISHIO, H. & HATA, F. (1995). Involvement of heterotrimeric GTP-binding protein and rho protein, but not protein kinase C, in agonist-induced Ca<sup>2+</sup> sensitization of skinned muscle of guinea pig vas deferens. *J. Pharmacol. Exp. Ther.*, **274**, 555–561.
- GAVETT, S.H. & WILLS-KARP, M. (1993). Elevated lung G protein levels and muscarinic receptor affinity in a mouse model of airway hyperreactivity. *Am. J. Physiol.*, **265**, L493–L500.
- GONG, M.C., FUJIHARA, H., SOMLYO, A.V. & SOMLYO, A.P. (1997). Translocation of rhoA associated with Ca<sup>2+</sup> sensitization of smooth muscle. *J. Biol. Chem.*, **272**, 10704–10709.
- HORI, M., SATO, K., MIYAMOTO, S., OZAKI, H. & KARAKI, H. (1993). Different pathways of calcium sensitization activated by receptor agonists and phorbol esters in vascular smooth muscle. *Br. J. Pharmacol.*, **110**, 1527–1531.
- KUREISHI, Y., KOBAYASHI, S., AMANO, M., KIMURA, K., KANAIDE, H., NAKANO, T., KAIBUCHI, K. & ITO, M. (1997). Rho-associated kinase directly induces smooth muscle contraction through myosin light chain phosphorylation. *J. Biol. Chem.*, **272**, 12257–12260.
- LEE, J.Y., UCHIDA, Y., SAKAMOTO, T., HIRATA, A., HASEGAWA, S. & HIRATA, F. (1994). Alteration of G protein levels in antigen-challenged guinea pigs. *J. Pharmacol. Exp. Ther.*, **271**, 1713–1720.
- MISAWA, M. & CHIBA, Y. (1993). Repeated antigenic challenge-induced airway hyperresponsiveness and airway inflammation in actively sensitized rats. *Jpn. J. Pharmacol.*, **61**, 41–50.
- OLENIC, C., BARTH, H., JUST, I., AKTORIES, K. & MEYER, D.K. (1997). Gene expression of the small GTP-binding proteins RhoA, RhoB, Rac1, and Cdc42 in adult rat brain. *Brain Res. Mol. Brain Res.*, **15**, 263–269.
- OTTO, B., STEUSLOFF, A., JUST, I., AKTORIES, K. & PFITZER, G. (1996). Role of Rho proteins in carbachol-induced contractions in intact and permeabilized guinea-pig intestinal smooth muscle. *J. Physiol.*, **496**, 317–329.
- OKAZI, H., KWON, S.-C., TAJIMI, M. & KARAKI, H. (1990). Changes in cytosolic Ca<sup>2+</sup> and contraction induced by various stimulants and relaxants in canine tracheal smooth muscle. *Pflügers Arch.*, **416**, 351–359.
- ROBERTS, J.A., RAEBURN, D., RODGER, I.W. & THOMSON, N.C. (1984). Comparison of in-vivo responsiveness and in-vitro smooth muscle sensitivity to methacholine. *Thorax*, **39**, 837–843.
- SATO, K., OZAKI, H. & KARAKI, H. (1988). Changes in cytosolic calcium level in vascular smooth muscle strip measured simultaneously with contraction using fluorescent calcium indicator Fura 2. *J. Pharmacol. Exp. Ther.*, **246**, 294–300.
- SATO, S., KREUTZ, R., WILM, C., GANTEN, D. & PFITZER, G. (1994). Augmented agonist-induced Ca<sup>2+</sup>-sensitization of coronary artery contraction in genetically hypertensive rats. Evidence for altered signal transduction in the coronary smooth muscle cells. *J. Clin. Invest.*, **94**, 1397–1403.

(Received January 28, 1999)

Revised March 4, 1999

Accepted March 9, 1999