

# Potential Mechanisms of Thyroid Disruption in Humans: Interaction of Organochlorine Compounds with Thyroid Receptor, Transthyretin, and Thyroid-binding Globulin

Ann Oliver Cheek,<sup>1,2</sup> Kelvin Kow,<sup>1</sup> Jian Chen,<sup>1</sup> and John A. McLachlan<sup>1,3</sup>

<sup>1</sup>Tulane/Xavier Center for Bioenvironmental Research; <sup>2</sup>Department of Ecology, Evolution, and Organismal Biology; <sup>3</sup>Department of Pharmacology, Tulane University, New Orleans, LA 70402 USA

Organochlorine compounds, particularly polychlorinated biphenyls (PCBs), alter serum thyroid hormone levels in humans. Hydroxylated organochlorines have relatively high affinities for the serum transport protein transthyretin, but the ability of these compounds to interact with the human thyroid receptor is unknown. Using a baculovirus expression system in insect cells (Sf9 cells), we produced recombinant human thyroid receptor  $\beta$  (hTR $\beta$ ). In competitive binding experiments, the recombinant receptor had the expected relative affinity for thyroid hormones and their analogs. In competitive inhibition experiments with PCBs, hydroxylated PCBs (OH-PCBs), DDT and its metabolites, and several organochlorine herbicides, only the OH-PCBs competed for binding. The affinity of hTR $\beta$  for OH-PCBs was 10,000-fold lower ( $K_i = 20\text{--}50\ \mu\text{M}$ ) than its affinity for thyroid hormone (3,3',5-triiodothyronine, T<sub>3</sub>;  $K_i = 10\ \text{nM}$ ). Because their relative affinity for the receptor was low, we tested the ability of OH-PCBs to interact with the serum transport proteins—transthyretin and thyroid-binding globulin (TBG). With the exception of one compound, the OH-PCBs had the same affinity ( $K_i = 10\text{--}80\ \text{nM}$ ) for transthyretin as thyroid hormone (thyroxine; T<sub>4</sub>). Only two of the OH-PCBs bound TBG ( $K_i = 3\text{--}7\ \mu\text{M}$ ), but with a 100-fold lower affinity than T<sub>4</sub>. Hydroxylated PCBs have relatively low affinities for the human thyroid receptor *in vitro*, but they have a thyroid hormonelike affinity for the serum transport protein transthyretin. Based on these results, OH-PCBs *in vivo* are more likely to compete for binding to serum transport proteins than for binding to the thyroid receptor. **Key words:** endocrine disruption, PCB, thyroid-binding globulin, thyroid receptor, transthyretin. *Environ Health Perspect* 107:273–278 (1999). [Online 9 March 1999] <http://ehpnet1.niehs.nih.gov/docs/1999/107p273-278cheek/abstract.html>

An important question in endocrine disruption is the mechanism by which a xenobiotic compound alters the action of endogenous hormones. One possible mechanism is direct interaction with the hormone receptor, either as an agonist or as an antagonist. In the case of thyroid hormone, a second important mechanism may be the ability of compounds to alter serum transport of thyroid hormones (TH). In nonmammalian vertebrates, the major transport protein is prealbumin (transthyretin); while some mammals, including humans, have a second binding protein, thyroid-binding globulin (TBG) (1). Assessing the relative affinity of the thyroid receptor and the serum transport proteins for xenobiotics should help clarify one of the mechanisms by which xenobiotics alter thyroid homeostasis.

Alterations in thyroid homeostasis by organochlorine compounds have been documented for many species, including humans. In most cases, exposure to organochlorine compounds is correlated with decreased serum levels of thyroid hormone, particularly thyroxine (T<sub>4</sub>). Exposure to polychlorinated biphenyls (PCBs) has been correlated with decreased serum T<sub>4</sub> concentrations in rats (2–10) and humans (6,11,12). Evidence from rat studies indicates that PCB-induced

decreases in serum T<sub>4</sub> are the result of increased metabolism by uridine diphosphate glucuronosyltransferase (UDPGT), a hepatic enzyme that glucuronidates T<sub>4</sub> (5,13–15). Another class of organochlorines, the chloroacetanilides acetochlor and alachlor, elevates UDPGT activity and concomitantly decreases serum T<sub>4</sub> levels in rats (16,17). Acetochlor also alters thyroid hormone (3,3',5-triiodothyronine; T<sub>3</sub>) action in amphibians, accelerating T<sub>3</sub>-induced metamorphosis (18). DDT and its metabolites alter serum T<sub>4</sub> levels in birds (19) and humans (20). DDT also alters thyroid metabolism in rats by increasing hepatic UDPGT activity (21).

Because of their physiological effects and their structural resemblance to thyroid hormones (Fig. 1), several studies have investigated the ability of PCBs to bind to the serum transport proteins transthyretin (3,8,22–24) and TBG (22) and to the rat thyroid receptor (23). Transthyretin (TTR) and TBG have similar affinities for the natural ligand, T<sub>4</sub> (50–90 nM) (22), but have different affinities for PCBs. Hydroxylated PCBs are potent ligands for TTR, having affinities in the 1 nM range, 50-fold greater than that of T<sub>4</sub> (8,22,24). Few hydroxylated PCBs bind TBG (22)

and few unmetabolized PCBs have strong affinities for either TTR or TBG (8,22,23). Like the transport proteins, the rat thyroid receptor appears to have a higher affinity for hydroxylated versus parent PCBs (23).

Although studies of binding affinity suggest that organochlorine compounds may alter thyroid homeostasis by interacting with thyroid hormone transport proteins in humans and animals, little is known about the ability of organochlorines to interact with the thyroid receptor, particularly in humans. We examined the ability of PCBs, DDTs, chloroacetanilides, and an isoprenoid to bind a recombinant human thyroid receptor (hTR $\beta$ 1). To evaluate the relative significance of receptor versus transport protein binding for disrupting thyroid homeostasis, compounds that bound the receptor were also tested for binding to human transthyretin and TBG.

## Methods


### Chemicals

T<sub>3</sub>, T<sub>4</sub>, 3,3',5-triiodothyroacetic acid (Triac), 3,3',5,5'-tetraiodothyroacetic acid (Tetrac), 3,3',5'-triiodo-L-thyronine (rT<sub>3</sub>), 2,2-bis(*p*-chlorophenyl)-ethanol (DDOH), human TBG, and human transthyretin (prealbumin) were purchased from Sigma Chemical Co. (St. Louis, MO). The PCBs—3,3',4,4',5-pentachlorobiphenyl (PCB 126), 3,3',4,4'-tetrachlorobiphenyl (PCB 77), 3-OH-2',4',6'-trichlorobiphenyl, 4-OH-3,5-dichlorobiphenyl, 4-OH-2',3',4',5'-tetrachlorobiphenyl, 4-OH-2',3',4',6'-tetrachlorobiphenyl, 4-OH-2',3',4',5'-pentachlorobiphenyl, and 4-OH-2',3',4',5,6'-pentachlorobiphenyl—were purchased from AccuStandard (New Haven, CT). The hydroxylated PCB, 4,4'-diOH-3,3',5,5'-tetrachlorobiphenyl, was purchased from

Address correspondence to A.O. Cheek, Department of Biology, Southeastern Louisiana University, SLU 10736, Room 205 Mead Hall, North Pine Extension, Hammond, LA 70402 USA.

Many thanks to P. Vonier for teaching techniques in cell culture and baculovirus expression and to A. Holstein for performing preliminary Western blot analysis. This study was supported by Department of Energy grant DE-FG07-96ER62308.

Received 21 September 1998; accepted 21 December 1998.



**Figure 1.** Structures of thyroid hormones and hydroxylated polychlorinated biphenyls (PCBs). CB, chlorinated biphenyl.

Ultra Scientific (North Kingstown, RI). The DDTs—1-(2-chlorophenyl)-1-(4-chlorophenyl)-2,2,2-trichloroethane (*o,p'*-DDT), 1-(2-chlorophenyl)-1-(4-chlorophenyl)-2,2-dichloroethane (*o,p'*-DDD), 1,1-bis(4-chlorophenyl)-2,2,2-trichloroethane (*p,p'*-DDT), and 2,2-bis(4-chlorophenyl)-1,1-dichloroethane (*p,p'*-DDE)—were purchased from Aldrich Chemical (Milwaukee,

WI). All chemicals were dissolved in DMSO. DMSO did not exceed 0.01% in the binding assays.

### Transfection

Sf9 insect cells (pupal ovarian cells from the fall armyworm) were purchased from Invitrogen (Carlsbad, CA) and cultured at 27°C in complete Grace's media (10% fetal bovine serum lactalbumin hydrolysate, tissue culture yeastolate, and glutamine; Invitrogen). A baculovirus phagemid containing the human thyroid receptor  $\beta 1$  cDNA was cloned into the multiple cloning site of the baculovirus phagemid pFastBac1 (Gibco BRL, Grand Island, NY) by DNA Technologies (Gaithersburg, MD). Confluent cells in a T-25 flask (Corning, Corning, NY) were rinsed with serum-free Grace's media and transfected with 1  $\mu$ g phagemid DNA and 2  $\mu$ l lipofectin in 2 ml serum-free Grace's media at 27°C. At the end of 5 hr, the transfection medium was replaced with complete Grace's media. After 7 days, medium containing baculovirus was harvested and stored at 4°C. To produce recombinant hTR $\beta$ , confluent Sf9 cells in a T-150 flask were incubated with 7 ml baculovirus-containing medium at 27°C for 1.5 hr. An additional 13 ml of Grace's media was added and cells were cultured for 5 days. Virus-containing medium was harvested and cell extracts were prepared.

### Preparation of Protein Extract

Protocols for preparing cell extracts were modified from Toscano (25), Bres and Eales (26), and Sullivan et al. (27). Transfected Sf9 cells were scraped from the flask and centrifuged at 5,000 rpm for 5 min. The supernatant was harvested and stored at 4°C. The cell pellet was resuspended in 2.5 ml buffer A [10 mM Tris-HCl (pH 7.6), 10% glycerol, 3 mM MgCl<sub>2</sub>, 2 mM CaCl<sub>2</sub>, 5 mM dithiothreitol (DTT), 1 mM Pefabloc, 1  $\mu$ g/ml aprotinin, and 20  $\mu$ M leupeptin], incubated for 20 min on ice, and homogenized in a glass dounce. KCl was added to a concentration of 0.4 M and

the homogenate was incubated on ice for 30 min, with shearing through a pasteur pipet every 10 min. The homogenate was then centrifuged at 25,000 rpm for 15 min at 4°C. The supernatant (cell extract) was aliquotted and stored at -80°C until use.

### Immunodetection of TR Protein

Cell extracts were heated (95°C) in sodium dodecyl sulfate (SDS) loading buffer and electrophoresed through 10% SDS-PAGE. Gels were electroblotted onto polyvinylidene difluoride membranes (Sigma) at 25 V overnight. Membranes were rinsed in Tris-buffered saline (TBS; 10 mM Tris-HCl, pH 7.6, and 150 mM NaCl) and blocked in TBS supplemented with 3% bovine serum albumin, fraction V (Sigma). Membranes were incubated with a polyclonal antiserum to amino acids 62–82 of human TR $\beta 1$  (1:1250; Affinity BioReagents, Golden, CO) in TBST (TBS + 0.1% Tween-20) for 2 hr, rinsed 3 times in TBST, incubated for 1 hr with HRP-conjugated goat antirabbit IgG (Kirkegaard-Perry, Gaithersburg, MD), rinsed 4 times with TBST, and visualized with enhanced chemiluminescence (Amersham, Arlington Heights, IL).

### TR Binding Assays

**Saturation analysis.** Cell extract containing hTR $\beta$  was added to assay buffer [10 mM Tris-HCl (pH 7.4), 10% glycerol, 5 mM DTT, and 0.5% CHAPS] to achieve a final concentration of 50  $\mu$ g protein/ml in 100  $\mu$ l total volume. Extracts were preincubated with 1,000-fold excess unlabeled T<sub>3</sub> for 15 min at 21°C, then increasing concentrations of [<sup>125</sup>I]T<sub>3</sub> (0.05, 0.1, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 5, and 10 nM) were added and extracts were incubated for an additional 60 min. The reaction was terminated on ice and unbound [<sup>125</sup>I]T<sub>3</sub> was separated from bound [<sup>125</sup>I]T<sub>3</sub> with the addition of hydroxylapatite (HAP) slurry (1:1 v/v in 0.1 M KCl, 10 mM Tris-HCl, pH 7.4). Extracts were incubated in HAP slurry for 40 min, with vortexing every 10 min. The slurry was centrifuged at 5,000 rpm for 3 min and buffer was aspirated. HAP pellets were then washed 3–4 times with 400  $\mu$ l wash buffer [0.1 M KCl, 10 mM Tris-HCl (pH 7.4), 0.5% CHAPS], resuspended in 200  $\mu$ l slurry buffer, and counted in ScintiVerse scintillation fluid (Fisher, Houston, TX).

**Competitive inhibition experiments.** Based on the saturation analysis, 2.5 nM [<sup>125</sup>I]T<sub>3</sub> was used in competitive inhibition experiments. Nonsaturable binding was estimated by preincubating cell extracts with 10,000-fold molar excess unlabeled T<sub>3</sub> for 15 min. Varying concentrations (10<sup>-8</sup>–10<sup>-4</sup> M) of unlabeled competitors were also



preincubated with cell extract before addition of [ $^{125}$ I]T $_3$ . All other conditions were as described above.

To characterize the specificity of the recombinant receptor, the known TR agonists T $_3$ , T $_4$ , Triac, and Tetrac, and the inactive T $_3$  metabolite, rT $_3$ , were tested in competitive inhibition experiments. We selected organochlorine compounds for testing based on their reported ability to alter *in vivo* responses or to bind serum transport proteins. PCB mixtures and some specific PCBs decrease serum T $_4$  levels, but PCB 126 and PCB 77 markedly enhance spatial learning in rats exposed *in utero*, while only slightly altering serum T $_4$  (2). Because the enhancement of learning is similar to that in hyperthyroid rat pups, Schantz et al. (2) proposed that PCBs 126 and 77 might be thyroid receptor agonists. We examined the interaction of these two compounds with the human thyroid receptor. Because several OH-PCBs bind TTR as effectively as T $_4$  (22), we also tested the ability of OH-PCBs to bind the receptor. DDTs and chloroacetanilides alter serum T $_4$  levels and catabolism, but their ability to bind TR is unknown.

### Thyroid-binding Protein and Transthyretin Binding Assays

Conditions for these binding assays were modified from Lans et al. (22). Briefly, purified TBG or TTR was added to 200  $\mu$ l assay buffer (TR assay buffer discussed previously) for a final concentration of 30 nM. We used 55 nM L-T $_4$  containing 100,000 cpm [ $^{125}$ I]L-T $_4$  to estimate total binding. Nonsaturable binding was estimated by preincubating protein solutions with 100-fold molar excess unlabeled L-T $_4$  for 15

min. Varying concentrations of unlabeled competitors were also pre-incubated with the protein before the addition of [ $^{125}$ I]L-T $_4$ . All other conditions were as described for TR saturation analysis, except the volumes of HAP slurry and of slurry buffer for resuspension were doubled to account for the twofold larger assay volume.

## Results

### Immunodetection

Infected Sf9 cells expressed a protein of approximately 51 kDa that cross-reacted with a polyclonal antibody specific to hTR $\beta$ 1 (Fig. 2). This protein is similar in size to the 52–55 kDa proteins recombinantly expressed in *Escherichia coli* (28,29).

### TR Binding


**Saturation analysis.** Preliminary experiments using 50, 100, 200, and 300  $\mu$ g protein/ml indicated that minimal nonsaturable binding ( $\leq 10\%$  of total binding) occurred at 50  $\mu$ g protein/ml. Similarly, varying incubation times (15, 30, 45, 60, 90, and 120 min) showed that a 60-min incubation period was sufficient for achieving equilibrium binding. Binding to recombinant hTR $\beta$  was saturable at 3 nM [ $^{125}$ I]T $_3$  with a  $K_d$  of  $1.37 \pm 0.24$  nM ( $n = 4$  experiments) and a  $B_{max}$  of  $0.30 \pm 0.09$  nM (Fig. 3). The binding affinity observed with recombinant hTR $\beta$  expressed in Sf9 cells is similar to that reported for other recombinant thyroid receptors (28,29), but is lower than that reported for thyroid receptors extracted from tissues (approximately 0.1 nM) (30).

**Competitive binding experiments.** Thyroid hormones and their analog showed the expected order of affinity for recombinant


hTR $\beta$ : Triac > T $_3$  = L-T $_4$  >> DL-T $_4$   $\approx$  Tetrac (Fig. 4 and Table 1). Of the xenobiotics tested, only the hydroxylated PCBs could inhibit 50% of [ $^{125}$ I]T $_3$  binding to hTR $\beta$  (Fig. 5 and Table 1), although with 10,000-fold lower potency than L-T $_4$ . The coplanar PCBs, PCB 77 and PCB 126, did not displace T $_3$  from the receptor, nor did *p,p'*-DDE, *p,p'*-DDT, acetochlor, or methoprene. Although some displacement (20%) was achieved by the highest concentrations (100  $\mu$ M) of *o,p'*-DDD, *o,p'*-DDT, DDOH, and alachlor, none of these compounds could inhibit 50% of [ $^{125}$ I]T $_3$  binding.

### TTR and TBG Binding

The binding affinity of unlabeled L-T $_4$  was  $62 \pm 12$  nM for TTR and  $76 \pm 15$  nM for TBG. These affinities are in agreement with those reported by Lans et al. (22): 88–138 nM for TTR and 52–85 nM for TBG. All



**Figure 2.** Western blot of cell extract from Sf9 cells infected with baculovirus containing the human thyroid receptor (hTR $\beta$ ) gene. A 51-kDa protein that cross-reacts specifically with a polyclonal antibody to hTR $\beta$ 1 is produced by the infected Sf9 cells.



**Figure 3.** (A) Saturation analysis and (B) Scatchard analysis of [ $^{125}$ I]3,3',5-triiodothyronine (T $_3$ ) binding to recombinant human thyroid receptor (hTR $\beta$ ). At each concentration of [ $^{125}$ I]T $_3$ , a 1,000-fold molar excess of unlabeled T $_3$  was used to estimate nonsaturable binding (NSB). Curves are the average  $\pm$  standard error of four experiments. Abbreviations: TB, total binding; SB, saturable binding; B/F, bound/free concentration.

of the 4-hydroxylated PCBs bound TTR with affinities (10–140 nM) similar to that of the natural ligand L-T<sub>4</sub> (62 nM) (Fig. 6 and Table 1). Hydroxylation in the *meta* position appeared to abolish TTR binding (Table 1). PCB 126 bound TTR weakly, with a 1,000-fold lower affinity than L-T<sub>4</sub> and the 4-hydroxylated PCBs (Table 1 and Fig. 6). DDOH was the only DDT metabolite with weak affinity for TTR (approximately 100 μM).

Few of the xenobiotics competitively bound TBG. Two of the hydroxylated PCBs competed for TBG binding—4-OH-2',3,4',6'-tetraCB and 3-OH-2',4',6'-triCB (Fig. 7 and Table 1)—but with affinities 30–100-fold lower than L-T<sub>4</sub>. Cl<sup>-</sup> ions in the 2',4', and 6' positions appeared to facilitate binding to TBG. Addition of an extra Cl<sup>-</sup> at the 5 position seemed to abolish TBG binding, as 4-OH-2',3,4',5,6'-pentaCB did not compete for binding, whereas 4-OH-2',3,4',6'-tetraCB did. None of the other hydroxylated PCBs bound TBG. *o,p'*-DDD and DDOH bound TBG, but with affinities 70–800-fold lower than L-T<sub>4</sub> (Fig. 7 and Table 1). Acetochlor, alachlor, and methoprene showed no affinity for TTR or TBG.

## Discussion

Of the four groups of compounds examined, only the hydroxylated PCBs bound to human TRβ1 with affinities ranging from 30–90 μM—affinities 10,000-fold lower than the natural ligand T<sub>3</sub>. The hydroxylated PCBs had 1,000-fold greater


affinities for TTR than for TR, making them competitors for the natural ligand L-T<sub>4</sub>. Half of the hydroxylated PCBs tested had higher affinities for TTR than did T<sub>4</sub>.

Only two of the hydroxylated PCBs bound TBG—3-OH-2',4',6'-triCB and 4-OH-2',3,4',6'-tetraCB. In fact, 3-OH-2',4',6'-triCB did not bind TTR, but had a 20-fold


**Table 1.** Inhibition constants (± standard error, *n* = 3 experiments in duplicate) for thyroid hormones and environmental chemicals interacting with recombinant hTRβ, hTTR, and hTBG

Compound	K <sub>i</sub> (μM)		
	TRβ	TTR	TBG
T <sub>3</sub>	0.020 ± 0.015	—	—
L-T <sub>4</sub>	0.024 ± 0.012	0.062 ± 0.012	0.076 ± 0.015
Dl-T <sub>4</sub>	0.135 ± 0.054	—	—
Triac	0.006 ± 0.005	—	—
Tetrac	0.075 ± 0.014	—	—
rT <sub>3</sub>	0.808 ± 0.151	—	—
3-OH-2',4',6'-trichlorobiphenyl	90.3 ± 23.7	>100	5.46
4-OH-3,5-dichlorobiphenyl	83.5 ± 18.7	0.016	>100
4-OH-2',3',4',5'-tetrachlorobiphenyl	37.7 ± 26.0	0.089 ± 0.009	>100
4-OH-2',3,4',6'-tetrachlorobiphenyl	43.0 ± 16.9	0.033 ± 0.002	2.34 ± 0.029
4-OH-2',3,3',4',5'-pentachlorobiphenyl	67.2 ± 17.1	0.141	>100
4-OH-2',3,4',5,6'-pentachlorobiphenyl	36.5 ± 6.65	0.040 ± 0.008	>100
4,4'-diOH-3,3',5,5'-tetrachlorobiphenyl	32.7 ± 12.1	0.011 ± 0.002	>100
PCB 77	>100	>100	>100
PCB 126	>100	140 ± 18.7	>100
<i>o,p'</i> -DDT	>100	>100	>100
<i>o,p'</i> -DDD	>100	>100	4.99 ± 4.13
DDOH	>100	89 ± 11	62.20 ± 40.9
<i>p,p'</i> -DDE	>100	>100	>100
<i>p,p'</i> -DDT	>100	>100	>100
Acetochlor	>100	>100	>100
Alachlor	>100	>100	>100
Methoprene	>100	>100	>100

Abbreviations: TRβ, thyroid receptor β; TTR, transthyretin; TBG, thyroid-binding globulin; T<sub>3</sub>, 3,3',5-triiodothyronine; T<sub>4</sub>, thyroxine; Triac, 3,3',5-triiodothyroacetic acid; Tetrac, 3,3',5,5'-tetraiodothyroacetic acid; rT<sub>3</sub>, 3,3',5'-triiodo-L-thyronine; PCB 77, 3,3',4,4'-tetrachlorobiphenyl; PCB 126, 3,3',4,4',5-pentachlorobiphenyl; *o,p'*-DDT, 1-(2-chlorophenyl)-1-(4-chlorophenyl)-2,2-dichloroethane; *o,p'*-DDD, 1-(2-chlorophenyl)-1-(4-chlorophenyl)-2,2-dichloroethane; DDOH, 2,2-bis(*p*-chlorophenyl)-ethanol; *p,p'*-DDE, 2,2-bis(4-chlorophenyl)-1,1-dichloroethane; *p,p'*-DDT, 1,1-bis(4-chlorophenyl)-2,2,2-trichloroethane.



**Figure 4.** Competitive binding of known thyroid receptor (TR) agonists to recombinant human thyroid receptor β. Increasing concentrations of agonist competed with 2.5-nM [<sup>125</sup>I]3,3',5-triiodothyronine (T<sub>3</sub>). Nonsaturable binding was estimated by incubation with 10,000-fold molar excess unlabeled T<sub>3</sub>. *n* = 3 experiments performed in duplicate, except *n* = 5 experiments for T<sub>3</sub>.



**Figure 5.** Competitive binding of hydroxylated polychlorinated biphenyls to recombinant human thyroid receptor β. Assay conditions are as described in Figure 4. *n* = 3 experiments performed in duplicate. Abbreviations: T<sub>3</sub>, 3,3',5-triiodothyronine; CB, chlorinated biphenyl.



greater affinity for TBG than for TR. PCBs and especially their hydroxylated metabolites interact with multiple components of the thyroid system, enhancing hepatic metabolism of thyroid hormones (5,13–15), competing for transport via serum proteins (21), especially TTR (3,8,22–24), and competing for receptor binding (this study). The physiological result is alteration in serum thyroid hormone levels. Of the known mechanisms, interaction with the thyroid receptor is likely to be less important than competition for serum transport proteins and induction of hepatic metabolism of  $T_4$ . Given  $K_i$  values of 30–90  $\mu\text{M}$  for xenobiotics that interacted with the receptor, high concentrations on the order of 20 ppm (micrograms per gram) would have to be achieved in target tissues for hydroxylated PCBs to significantly alter  $T_3$  binding to the receptor. Alternatively, the  $K_i$  values for TTR suggest that a concentration of only 0.017 ppm would have to be achieved for hydroxylated PCBs to significantly alter thyroid hormone transport via TTR. Hydroxylated PCB concentrations on the order of 0.36 ppm have been measured in human serum (31). Based on these data, *in vivo* disruption of TTR binding is more likely than disruption of receptor binding.

Neither of the coplanar dioxinlike PCBs, PCB 77 and PCB 126, bound the TR. Schantz et al. (2) observed that these PCBs accelerated spatial learning in rat pups


exposed *in utero* and during lactation, an effect observed in hyperthyroid neonatal rats. Because both coplanar PCBs caused slight decreases in serum  $T_3$  and  $T_4$  but accelerated spatial learning, Schantz et al. (2) suggested that these PCBs might directly activate the thyroid receptor. In this study, PCB 77 and PCB 126 did not bind the human TR $\beta$ 1, so direct interaction with the receptor probably does not explain the thyromimetic learning effect observed in the rats.

Previous work by McKinney et al. (23) indicated that two other coplanar PCBs, PCB 169 and PCB 80, bound to a rat nuclear extract with 100-fold lower affinities than  $L-T_4$ , whereas PCB 54, an *ortho*-substituted congener, did not bind at all. Although the present study did not examine the binding affinity of PCBs 169 and 80, reported differences in affinity for coplanar PCBs may be due to several factors. First, the current study used a protein extract of insect cells producing recombinant human TR $\beta$ 1. Therefore, only a single TR isoform was available to interact with compounds in competitive binding experiments. In contrast, the rat liver nuclear extract probably contained not only rat TR $\beta$ 1, but also TR $\alpha$ 1 and TR $\alpha$ 2 (30). Second, because only a single TR isoform was present in recombinant cell extracts, only TR homodimers could form, while in rat liver nuclear extracts, retinoid X receptors (RXRs), the heterodimeric partners of TR, were probably also present


(30,32). *In vitro*, TR-RXR heterodimers exhibit different affinities for ligands than do TR-TR homodimers, although both appear to form spontaneously in cells (30). Third, species-specific differences in TR affinity for coplanar PCBs may exist.

DDTs and chloroacetanilide herbicides cause hypothyroidlike effects in animals, decreasing serum  $T_4$  (16,17,19–21). None of these compounds bound to the thyroid receptor, indicating that they are unlikely to disrupt the thyroid axis via receptor interaction. DDOH bound to TTR and to TBG, but with such low affinity that concentrations in serum are unlikely to be high enough to compete for  $T_4$  binding. *o,p'*-DDD bound to TBG with a fairly low affinity (5  $\mu\text{M}$ ) and is unlikely to reach such high concentrations in serum because of environmental exposure. However, clinical treatment of adrenal carcinomas resulted in 100–600  $\mu\text{M}$  doses of *o,p'*-DDD (20). One consequence of *o,p'*-DDD treatment was decreased serum  $T_4$ , purportedly due to direct competition with *o,p'*-DDD for TBG binding (20). Our results support that hypothesis. Neither of the chloroacetanilides acetochlor nor alachlor bound TTR or TBG, but they enhance hepatic metabolism of  $T_4$  in rats (16,17). Alteration of metabolism is probably the major mechanism by which chloroacetanilides affect thyroid axis function.

Our results suggest that disruption of thyroid hormone transport is one of the



**Figure 6.** Competitive binding of OH-polychlorinated biphenyls (PCBs) with purified human transthyretin (TTR). Increasing concentrations of OH-PCBs competed with 55 nM  $L$ -thyroxine ( $T_4$ ) spiked with 100,000 cpm [ $^{125}\text{I}$ ] $L$ - $T_4$ . Nonsaturable binding was estimated by incubation with 100-fold molar excess unlabeled  $L$ - $T_4$ .  $n = 3$  experiments performed in duplicate. Abbreviations: DDOH, 2,2-bis(*p*-chlorophenyl)-ethanol; CB, chlorinated biphenyl.




**Figure 7.** Competitive binding of hydroxylated polychlorinated biphenyls with purified human thyroid-binding globulin. Assay conditions are as described in Figure 6.  $n = 3$  experiments performed in duplicate. Abbreviations:  $T_4$ , thyroxine; CB, chlorinated biphenyl; *o,p'*-DDD, 1-(2-chlorophenyl)-1-(4-chlorophenyl)-2,2-dichloroethane; DDOH, 2,2-bis(*p*-chlorophenyl)-ethanol.

mechanisms by which organochlorine compounds alter thyroid homeostasis. In particular, hydroxylated PCBs compete effectively for  $T_4$  binding to TTR, but few compounds compete for TBG binding, even at  $\mu\text{M}$  concentrations. TBG is found only in some mammals, including primates, ungulates (cattle, sheep, goats, pigs, water buffalo, and horses), and carnivores (dog), but not in rodents (rat) or lagomorphs (rabbit). Depending on species, TBG binds 60–90% of serum  $T_4$ . Interestingly, TBG deficiency in humans does not interfere with euthyroid status, suggesting that TTR is also important for  $T_4$  transport in humans (1). TTR is a highly conserved TH binding protein in all vertebrate species (1), so disruption of thyroid hormone transport by hydroxylated PCBs could potentially occur in all vertebrates, not only in humans.

#### REFERENCES AND NOTES

- Larsson M, Pettersson T, Carlstrom A. Thyroid hormone binding in serum of 15 vertebrate species: isolation of thyroxine-binding globulin and prealbumin analogs. *Gen Comp Endocrinol* 58:360–375 (1985).
- Schantz SL, Seo BW, Moshtaghian J, Amin S. Developmental exposure to polychlorinated biphenyls or dioxin: do changes in thyroid function mediate effects on spatial learning? *Am Zool* 37:399–408 (1997).
- Darnerud PO, Morse D, Klasson-Wehler E, Brouwer A. Binding of 3,3',4,4'-tetrachlorobiphenyl (CB-77) metabolite to fetal transthyretin and effects on fetal thyroid hormone levels in mice. *Toxicology* 106:105–114 (1996).
- Goldey ES, Kehn LS, Lau C, Rehnberg GL, Crofton KM. Developmental exposure to polychlorinated biphenyls (Aroclor 1254) reduces circulating thyroid hormone concentrations and causes hearing deficits in rats. *Toxicol Appl Pharmacol* 135:77–88 (1995).
- Van Birgelen AP, Smit EA, Kampen IM, Groeneveld CN, Fase KM, Van der Kolk J, Poiger H, VandenBerg M, Koeman JH, Brouwer A. Subchronic effects of 2,3,7,8-TCDD or PCBs on thyroid hormone metabolism: use in risk assessment. *Eur J Pharmacol* 293:77–85 (1995).
- Sauer PJJ, Huisman M, Koopman-Esseboom C, Morse DC, Prooije AESV, Berg KJVD, Tuinstra LGMT, Paauw CGVD, Boersma ER, Weisglas-Kuperus N, et al. Effects of polychlorinated biphenyls (PCBs) and dioxins on growth and development. *Hum Exp Toxicol* 13:900–906 (1994).
- Morse DC, Wehler EK, Wesseling W, Koeman JH, Brouwer A. Alterations in rat brain thyroid hormone status following pre- and postnatal exposure to polychlorinated biphenyls (Aroclor 1254). *Toxicol Appl Pharmacol* 136:269–279 (1996).
- Lans MC, Klasson-Wehler E, Willemssen M, Meussen E, Safe S, Brouwer A. Structure-dependent, competitive interaction of hydroxy-polychlorobiphenyls, -dibenzo-*p*-dioxins and -dibenzofurans with human transthyretin. *Chem Biol Interact* 88:7–21 (1993).
- Gray LEJ, Ostby J, Marshall R, Andrews J. Reproductive and thyroid effects of low-level polychlorinated biphenyl (Aroclor 1254) exposure. *Fundam Appl Toxicol* 20:288–294 (1993).
- Byrne JJ, Carbone JP, Hanson EA. Hypothyroidism and abnormalities in the kinetics of thyroid hormone metabolism in rats treated chronically with polychlorinated biphenyl and polybrominated biphenyl. *Endocrinology* 121:520–527 (1987).
- Koopman-Esseboom C, Huisman M, Touwen BC, Boersma ER, Brouwer A, Sauer PJ, Weisglas-Kuperus N. Newborn infants diagnosed as neurologically abnormal with relation to PCB and dioxin exposure and their thyroid-hormone status. *Dev Med Child Neurol* 39:785 (1997).
- Koopman-Esseboom C, Morse DC, Weisglas-Kuperus N, Lutkeschipholt IJ, Van Der Paauw CG, Tuinstra LGMT, Brouwer A, Sauer PJ. Effects of dioxins and polychlorinated biphenyls on thyroid hormone status of pregnant women and their infants. *Pediatr Res* 36:468–473 (1994).
- Beetstra JB, van Engelen JGM, Karels P, van der Hoek HJ, deJong M, Doctor R, Krenning EP, Hennemann G, Brouwer A, Visser TJ. Thyroxine and 3,3',5-triiodothyronine are glucuronidated in rat liver by different uridine diphosphate-glucuronyltransferases. *Endocrinology* 128:741–746 (1991).
- Barter RA, Klaassen CD. UDP-glucuronyltransferase inducers reduce thyroid hormone levels in rats by an extrathyroidal mechanism. *Toxicol Appl Pharmacol* 113:36–42 (1992).
- Barter RA, Klaassen CD. Reduction of thyroid hormone levels and alteration of thyroid function by four representative UDP-glucuronosyltransferase inducers in rats. *Toxicol Appl Pharmacol* 128:9–17 (1994).
- Wilson AGE, Thake DC, Heydens WE, Brewster DW, Hotz KJ. Mode of action of thyroid tumor formation in the male Long-Evans rat administered high doses of alachlor. *Fundam Appl Toxicol* 33:16–23 (1996).
- Ashby J, Kier L, Wilson AGE, Green T, Lefevre PA, Tinwell H, Willis GA, Heydens WF, Clapp MJL. Evaluation of the potential carcinogenicity and genetic toxicity to humans of the herbicide alachlor. *Hum Exp Toxicol* 15:702–735 (1996).
- Cheek AD, Ide CF, Bollinger JE, Rider CV, McLachlan JA. Alteration of leopard frog (*Rana pipiens*) metamorphosis by the herbicide acetochlor. *Arch Environ Contam Toxicol* (in press).
- Grässle B, Biessmann A. Effects of DDT, polychlorinated biphenyls and thiouracil on circulating thyroid hormones, thyroid histology and eggshell quality in Japanese quail (*Coturnix coturnix japonica*). *Chem Biol Interact* 42:371–377 (1982).
- Marshall JS, Tompkins L. Effect of *o,p'*-DDD and similar compounds on thyroxine binding globulin. *J Clin Endocrinol* 28:386–392 (1968).
- Bastomsky CH. Effects of a polychlorinated biphenyl mixture (Aroclor 1254) and DDT on biliary thyroxine excretion in rats. *Endocrinology* 95:1150–1155 (1974).
- Lans MC, Spiertz C, Brouwer A, Koeman JH. Different competition of thyroxine binding to transthyretin and thyroxine-binding globulin by hydroxy-PCBs, PCDDs, PCDFs. *Eur J Pharmacol* 270:129–136 (1994).
- McKinney J, Fannin R, Jordan S, Chae K, Rickenbacher U, Pedersen L. Polychlorinated biphenyls and related compound interactions with specific binding sites for thyroxine in rat liver nuclear extracts. *J Med Chem* 80:79–86 (1987).
- Rickenbacher U, McKinney JD, Oatley SJ, Blake CCF. Structurally specific binding of halogenated biphenyls to thyroxine transport protein. *J Med Chem* 29:641–648 (1986).
- Toscano D. Personal communication.
- Bres O, Eales J. Thyroid hormone binding to isolated trout (*Salmo gairdneri*) liver nuclei *in vitro*: binding affinity, capacity, and chemical specificity. *Gen Comp Endocrinol* 61:29–39 (1986).
- Sullivan C, Darling D, Dickhoff W. Nuclear receptors for L-triiodothyronine in trout erythrocytes. *Gen Comp Endocrinol* 65:149–160 (1987).
- Park J-B, Ashizawa K, Parkison C, Cheng S-Y. One-step immunoaffinity purification of human  $\beta 1$  thyroid hormone receptor with DNA and hormone binding affinity. *J Biochem Biophys Meth* 27:95–103 (1993).
- Falcone M, Miyamoto T, Fierro-Renoy F, Macchia E, DeGroot LJ. Antipeptide polyclonal antibodies specifically recognize each human thyroid hormone receptor isoform. *Endocrinology* 131:2419–2429 (1992).
- Lazar MA. Thyroid hormone receptors: multiple forms, multiple possibilities. *Endocrinol Rev* 14:184–193 (1993).
- Bergman A, Klasson-Wehler E, Kuroki H. Selective retention of hydroxylated PCB metabolites in blood. *Environ Health Perspect* 102:464–469 (1994).
- Glass CK. Some new twists in the regulation of gene expression by thyroid hormone and retinoic acid receptors. *J Endocrinol* 150:349–357 (1996).



## EHP PUTS EVEN MORE ENVIRONMENTAL HEALTH INFORMATION RIGHT AT YOUR FINGERTIPS!

EHP online articles contain convenient links to PubMed—the National Library of Medicine's free online search service of more than 9 million citations! Search MEDLINE and Pre-MEDLINE (including links to other online journals and databases) for information directly related to each EHP article's topic!

Subscribe to EHP today at <http://ehp.niehs.nih.gov>