

Hemopoietic Stem Cells: Stochastic Differentiation and Humoral Control of Proliferation

by Makio Ogawa*

The central feature of hemopoiesis is life-long, stable cell renewal. This process is supported by hemopoietic stem cells which, in the steady state, appear to be dormant in cell cycling. The entry into cell cycle of the dormant stem cells may be promoted by such factors as interleukin-1, interleukin-6 (IL-6), and granulocyte colony-stimulating factor (G-CSF). Once the stem cells leave G_0 and begin proliferation, the subsequent process is characterized by continued proliferation and differentiation. While several models of stem cell differentiation have been proposed, micromanipulation studies of individual progenitors suggest that the commitment of multipotential progenitors to single lineages is a random (stochastic) process. The proliferation of early hemopoietic progenitors requires the presence of interleukin-3 (IL-3), and the intermediate process appears to be supported by granulocyte/macrophage colony-stimulating factor (GM-CSF). Once the progenitors are committed to individual lineages, the subsequent maturation process appears to be supported by late-acting, lineage-specific factors such as erythropoietin and G-CSF. Synthesis of a hemopoietic factor may take place in different cell types and is regulated by multiple factors. The physiological regulator of erythropoiesis is erythropoietin, which, by a feedback mechanism, provides fine control of erythrocyte production. Feedback mechanisms for leukocyte production have not been identified. It is possible that there is no feedback regulator of leukopoiesis. In this model, leukocyte production in the steady state is maintained at a genetically determined level. When an infection occurs, the bacterial lipopolysaccharides may augment the production of interleukin 1α and β , tumor necrosis factor, macrophage colony-stimulating factor, etc. These factors circulate and stimulate the production in the marrow of G-CSF, GM-CSF, IL-3, IL-6, etc., with resultant leukocytosis. Once the infection subsides, the production of the factors and leukopoiesis return to the steady-state levels.

Introduction

The central feature of the hemopoietic system is its continuous process of cell turnover. It is estimated that approximately 200 billion erythrocytes (1) and 60 billion neutrophilic leukocytes (2) are produced and destroyed daily in a man weighing 70 kg. This life-long cell renewal process is supported by stem cells that possess the ability to self-renew and to produce progenitors that are committed to differentiation in single lineages. The mechanisms regulating stem cell functions have intrigued not only hematologists but also biologists interested in cellular differentiation and have been the object of active investigations and model building. During the last three decades, studies using functional assays for hemopoietic progenitors, i.e., the *in vivo* spleen colony assay for murine stem cells (3) and clonal cell culture assays for progenitors at various stages of development (4), have resulted in sig-

nificant elucidation of the mechanisms of hemopoiesis. In addition, these assays allowed identification of several types of hemopoietic factors. Recent clinical trials indicate that some of the recombinant proteins promise to be effective in the treatment with patients with hemopoietic disorders. In this review, I will summarize first the current understanding of the stem cell kinetics and differentiation and second the mechanisms of the hemopoietic factors regulating the dynamics of cell turnover.

Progenitors

General Consideration

Available evidence suggests that at a given time, hemopoietic cells occupying the entire hemopoietic tissue may be supplied by a small number of stem cell clones. Mulligan and his associates (5,6) reconstituted the hemopoietic system of mice by transferring a retrovirally labeled clone of stem cells, thereby demonstrating that the descendants of a single stem cell can generate enough

*Veterans Administration Medical Center and Department of Medicine, Medical University of South Carolina, Charleston, SC 29403.

cells to occupy the entire lympho-hemopoietic system. Follow-up studies of such animals indicated that hemopoiesis may be supported by sequential activation of different stem cell clones (6). This clonal succession of hemopoietic stem cells *in vivo* had also been evinced by Mintz et al. (7) who injected suspensions of liver cells from 13-day mouse fetuses into the placental circulation of *W/W* or *W^f/W^f* mice and obtained mice whose hemopoiesis was supported by more than one clone of stem cells. Serial observations of the erythrocyte types up to 60 weeks after transplantation revealed a complementary rise and fall in the proportion of the cells of different genotypes. These observations clearly demonstrated major proliferative potentials of a single stem cell and the exponential increase in the incidences of the progenitors as they progress through the developmental process.

Cell-Cycle Dormancy of Stem Cells

It is generally held that in the steady state, most hemopoietic stem cells are not engaged in active proliferation. Experimental evidence for this concept has been provided by investigators in several laboratories. Becker et al. (8) demonstrated that short-term exposure to ³H-thymidine with high specific activity does not kill colony-forming units in spleen (CFU-S). Similarly, multilineage colony formation in culture was not affected by the brief exposure to ³H-thymidine with high specific activity (9). Hodgson and Bradley (10) reported that high-dose 5-fluorouracil (5-FU), which preferentially kills cells in the cell cycle, does not affect cells that form late-appearing spleen colonies termed pre-CFU-S. In our laboratory, sequential observations of the growth of individual multipotential progenitors in culture also suggested that most primitive multipotential progenitors are dormant in cell proliferation (11). The concept of a true resting state was originally proposed by Lajtha (12) and was based on studies of liver cell regeneration. He envisioned that this state is metabolically distinct from the other phases of the cell cycle, and coined the term *G*₀ state. Lajtha proposed that hemopoietic stem cells are in *G*₀ since cell-cycle dormancy confers the stem cells time to repair DNA damage and thus allows maintenance of the genetic integrity of the stem cell populations (13). The mechanisms that initiate the cell division of the stem cells are not known. Lajtha proposed that this is a random process (13). As will be discussed later in this review, there is evidence that interleukin-1, interleukin-6, and granulocyte colony-stimulating factors appear to shorten the *G*₀ period and may induce the stem cells to begin active proliferation. It is possible that these factors, like the other hemopoietic factors, act as survival and/or growth factors on the most primitive hemopoietic stem cells and that the survival of stem cells and the duration of *G*₀ state may depend on the ambient levels of these factors.

Self-Renewal of Stem Cells

When a stem cell divides, its progeny can self-renew or differentiate. Following the development of the spleen

colony assay, Till and his associates (14) transplanted single-cell suspensions of individual spleen colonies into secondary recipient mice and observed a very heterogeneous distribution of the incidences of CFU-S in individual spleen colonies. The frequency distribution could not be fitted by a Poisson distribution but could be approximated by a γ distribution (14). Simultaneously, they carried out a computer simulation of a birth and death model in which they envisioned that self-renewal of CFU-S is a birth process and that loss of colony-forming ability associated with differentiation is a death process. This simulation of the model based on generation of random numbers also resulted in a γ distribution of CFU-S incidences. Based on these data, they proposed that self-renewal or differentiation is a stochastic (random) process.

In our laboratory, we analyzed distributions of colony-forming cells for blast cell colonies and those of multilineage colonies by replating individual blast cell colonies. We assumed that the blast cell colony formation and multilineage colony formation represent the culture equivalent of the birth and death processes, respectively (15). The frequency distributions of the secondary blast cell colonies and multilineage colonies could be approximated by variates of γ distributions. Therefore, the studies *in vivo* and in culture mutually corroborated and were consistent with the concept that self-renewal and commitment to differentiation of an individual stem cell is a stochastic process.

We also carried out serial observations (mapping) of the development multi-potential blast cell colonies (11) from spleen cells of 5-FU-treated mice (10). Blast cell colonies emerged after variable lag times and once identified, grew with relatively fixed cell doubling times and developed into multilineage colonies. These observations were consistent with the notion that in the steady-state the dormant stem cells enter cell-cycle randomly and that once committed to differentiation, their progeny proliferate at a relatively constant cell doubling time.

Differentiation of Stem Cells

Several models for stem cell differentiation have been proposed (16). Two deterministic models propose that external factors instruct individual stem cells regarding their directions of differentiation. The hemopoietic inductive microenvironment (HIM) model of Trentin and his associates (17,18) features small anatomical niches that are specific for individual lineages and determine the commitment of stem cells. The stem cell competition model envisions control of stem cell differentiation by lineage-specific humoral factors. For example, VanZant and Goldwasser (19,20) proposed that exposure of a stem cell in a certain cell cycle phase to erythropoietin or colony-stimulating factors determines the commitment to the erythroid or granulocytic lineage. In addition, a model of predetermined, sequential loss of lineage potentials has been proposed (21). As discussed previously in our review (16), specific criticisms may be raised against the data supporting these models. Most importantly, these models are based on studies of populations of colony-forming cells

rather than on individual progenitors. Identification of murine blast cell colonies with high replating efficiencies (22) provided us with a unique opportunity to micro-manipulate single progenitors and study differentiation potentials of single stem cells.

When we carried out cytological analysis of multilineage colonies that were derived from single progenitor cells and were cultured under identical conditions, we observed varying lineage combinations expressed in individual colonies (23). Subsequently, studies of paired progenitors revealed heterologous pairs of single and multilineage colonies expressing diverse lineage combinations (24). These results were consistent with the concept that the lineage selection by the progenitors at the time of cell division is a random process. Sequential manipulation of paired progenitors and analysis of the differentials of colonies derived from these progenitors indicated that the apparent random commitment takes place sequentially during stem cell differentiation (25). Analysis of single and paired human progenitors also revealed similar results and were consistent with the notion that the principle of human stem cell differentiation is a random restriction in the types and number of lineage potentials (26,27).

In the analysis of human multilineage colonies of single cell origin, we observed that individual lineages were represented by a variable number of cells. For example, a mixed colony consisted of 1340 erythrocytes and 4 eosinophilic leukocytes (26). This observation suggested that the number of times that a progenitor divides after commitment to a single lineage is variable. This observation is also in agreement with earlier observations in replating analysis of single lineage colonies. Investigators had observed variable replating efficiencies and heterogeneous size distributions of secondary colonies. For example, Wu (28) reported that the cumulative frequency distribution of the numbers of secondary T-lymphocyte colonies per primary T-lymphocyte colony could be approximated by a γ distribution. Analysis of the size of secondary mast cell colonies and computer simulation of a modification of the birth and death model of Till et al. (14) also suggested that mast cell proliferation in culture is a stochastic process (29). Together these data are consistent with the notion that both differentiation and proliferation of hemopoietic progenitors are separate stochastic processes.

Humoral Factors

General Considerations

While differentiation may be a stochastic process, proliferation and survival of progenitors appear to be under the control of humoral factors. Recently, a number of hemopoietic factors have been identified, purified, and molecularly cloned. The major hemopoietic factors include erythropoietin (Ep) (30-32), a physiological regulator of erythropoiesis; granulocyte/macrophage colony-stimulating factor (GM-CSF) (33-36); granulocyte (G)-CSF (37-39); interleukin-3 (IL-3) (40-43); and macro-

phage/monocyte (M)-CSF (44-46). Except for Ep, the physiological roles of these factors have not yet been clearly established. However, recent clinical trials of some of the factors indicate that they may possess significant therapeutic potential in the treatment of some of the hemopoietic disorders. Readers are referred to recent reviews (47-49) for detailed information on the biosynthesis, biochemistry and molecular biology of these proteins.

G-CSF was initially identified as a differentiation-inducing factor for leukemic cell lines (37). Some investigators have proposed that hemopoietic factors induce differentiation of progenitors in normal hemopoiesis. As reviewed earlier, our micromanipulation studies of differentiation of single and paired progenitors suggested that stem cell differentiation is not a directed process. The primary function of the humoral factors is to support the survival and/or proliferation of hemopoietic progenitors. In this model, the apparent induction of differentiation is a mere reflection of survival/proliferation of a population of progenitors that are supported by a humoral factor and the death of populations of progenitors that are not supported by the factor. Based on these concepts, I have summarized the recent evidence on the targets of the individual hemopoietic progenitors. The factors are arbitrarily divided into late-acting, lineage-specific factors, early-acting, lineage-nonspecific factors and those affecting dormant stem cells. The proliferative kinetics of the stem cells and the targets of the major hemopoietic factors are schematically presented in Figure 1.

Late-Acting, Lineage-Specific Factors

Ep is a physiological regulator of erythropoiesis and is secreted by the kidney. The recombinant protein has been given to patients with chronic renal failure and intractable anemia with remarkable therapeutic success (50). While some controversy still exists, the cells that are the most sensitive to Ep appear to be late erythroid progenitors that include erythroid colony-forming units (CFU-E) (51), erythroid cluster-forming units (52), and pronormoblasts. Since these cells are present in large numbers, are close to terminal maturation and are actively engaged




FIGURE 1. The model depicts the cell cycle dormancy of the stem cells and active proliferation of their progenies once they begin differentiation. It also illustrates the principle targets of the major hemopoietic factors.

in cell proliferation, a slight fluctuation in the ambient Ep levels would result in rapid changes in the production and release of reticulocytes. Experimental data suggest that multipotential stem cells (53) and early erythroid progenitors (54-56) are not under the control of Ep.

G-CSF appears to be a late-acting, neutrophil-specific factor. It has been demonstrated that in serum-containing cultures G-CSF supports the formation of neutrophil/macrophage colonies and pure neutrophil colonies (57), whereas in serum-free culture, it supports formation of only neutrophil colonies (58). In our laboratory, we studied the effects of G-CSF alone and in combination with other factors on colony formation by purified human progenitors in serum-free culture (59). G-CSF revealed significant synergism with earlier-acting factors such as IL-3 and GM-CSF in support of the formation of neutrophil colonies. When G-CSF was injected into animals, prompt and dose-dependent neutrophilic granulocytosis was observed (60-62). In addition, the differential revealed predominance of mature neutrophils (61). Together, these observations support the concept that G-CSF is a late-acting, neutrophil-specific factor. Interestingly, G-CSF appears also to play a role in stimulation of active proliferation of dormant stem cells. This will be elaborated in more detail later.

M-CSF has long been considered as a late-acting macrophage/monocyte-specific factor. Recently, CSF-HU that is abundant in human urine (63) was proved to be human M-CSF (46). A clinical trial of this material produced modest elevation in the granulocyte and macrophage counts in patients (63). It was also reported that during pregnancy, there is concomitant increase in the serum levels of M-CSF and blood monocyte counts (64). In serum-free culture of purified human progenitors, M-CSF alone or in combination with GM-CSF, G-CSF, IL-3, and Ep was an ineffective factor for macrophage colony formation (59). It is possible that M-CSF needs to interact with other factors in serum in macrophage colony formation.

Early-Acting, Lineage-Nonspecific Factors

IL-3 appears to be an important factor for proliferation of early hemopoietic progenitors. Purified murine IL-3 supports the growth of various types of colonies, including multilineage colonies (65-67) and blast cell colonies (66,68). Gibbon IL-3, which is 93% homologous with human IL-3, supported the formation of human multilineage and blast cell colonies (69). We also observed that delayed addition of murine IL-3 to cultures 7 days after cell plating decreases the number of multipotential blast cell colonies to one-half the number in cultures with IL-3 added on day 0. It did not, however, alter the proliferative and differentiative characteristics of late-emerging, multipotential blast cell colonies (66).

Based on these observations, we proposed that IL-3 does not trigger stem cells into active proliferation but is required for the continued proliferation of early multipotential progenitors. Subsequently, we observed that the development of multipotential blast cell colonies requires less IL-3 than the process of multilineage colony forma-

tion from blast cell colonies (68). These observations suggested that during stem cell development, the early multipotential progenitors are sensitive to IL-3, whereas as they gradually differentiate, the sensitivity to IL-3 slowly declines.

More recently, we examined the effects of human IL-3 on colony formation by purified hemopoietic progenitors in serum-free culture. IL-3 alone was not effective in support of colony formation except for a few small eosinophil colonies (59). Serial observation of the serum-free culture dishes containing human IL-3 revealed that blast cell clusters appear after varying lag time, reach approximately the 50 cell stage, and degenerate in the absence of late-acting, lineage-specific factors. IL-3 revealed significant synergism with Ep and G-CSF in support of erythroid bursts and neutrophil colonies, respectively. These observations indicated that IL-3 supports progenitors that are in the early stages of hemopoietic development, but it does not support the terminal maturation process (59).

The primary function of GM-CSF may also be to support the intermediate stages of hemopoietic development. It may not support the terminal neutrophil/macrophage maturation process effectively. Although it was originally identified as the factor that supported neutrophil/macrophage colonies in culture, Metcalf et al. (70) in 1980 observed that the few cell divisions of multipotential progenitors are supported by murine GM-CSF. Using serial transfer of multipotential blast cell colonies, we demonstrated that a subpopulation of the multipotential progenitors that respond to IL-3 also respond to GM-CSF (71). Investigators in several laboratories (72-74) have reported that human GM-CSF supports multilineage colony formation in serum-containing culture and possesses significant burst-promoting activity (BPA). In the studies of human enriched progenitors in serum-free culture, we observed that GM-CSF alone did not support colony formation except for a few small eosinophil colonies (59). However, in combination with G-CSF and Ep, GM-CSF effectively supported a large number of neutrophil colonies and erythroid bursts. Recently, administration of the recombinant GM-CSF to primates (75,76) and patients with acquired immunodeficiency syndrome (77) raised the levels of circulating neutrophils, eosinophils, and monocytes. These observations indicate that the primary target of GM-CSF is a population of multipotential progenitors that are intermediate between those responding to IL-3 and those sensitive to late-acting, lineage-specific factors.

Factors That Affect Proliferation of Dormant Hemopoietic Stem Cells

Very recently, several factors have been identified that appear to induce proliferation of hemopoietic stem cells in G₀. Stanley and his co-workers (78) reported that hemopoietin-1 (H-1), which was purified from human bladder carcinoma cell line, 5637 (79), acts synergistically with IL-3 in support of proliferation of murine hemopoietic stem cells. The recent reports by Mochizuki and her as-

sociates (80) and Moore and Warren (81) suggest that IL-1, which is abundant in the supernatant of this cell line, may account for the H-1 activity.

In our laboratory, we have found IL-6 [also called interferon β 2 (82) and B-cell stimulatory factor-2 (83)] to possess synergistic activity with IL-3 in the support of the active proliferation of murine hemopoietic stem cells (84). In the presence of IL-3, multipotential blast cell colonies emerged after varying lag times from spleen cells of 5-FU-treated mice (66,84). When IL-6 is also present, the appearance of multipotential blast cell colonies is significantly hastened, although the speed of the development of the blast cell colonies was not affected (84). These data suggested that part of the H-1 effect of IL-6 is the apparent shortening of the G_0 period of hemopoietic stem cells.

Human IL-6 also works synergistically with human IL-3 in stimulation of the proliferation of early human progenitors (85). Most recently, we have identified human G-CSF as an additional synergistic factor for the IL-3-dependent proliferation of murine hemopoietic stem cells (86). Both G-CSF and IL-6 appear to possess stronger synergistic effects than IL-1. It is of interest that there appears to be structural similarity between IL-6 and G-CSF including the positions of the cysteine residues, indicating similarity in three-dimensional structures of the two proteins. It is possible that the genes for the two proteins may share a common ancestral gene.

There are experimental data to indicate that production of IL-1, IL-6, and G-CSF may be significantly augmented immediately after invasion by microorganisms. For example, bacterial lipopolysaccharides induce synthesis of IL-1 in macrophages (87). IL-1 then circulates in the body and may augment synthesis of G-CSF (88,89) and IL-6 (90,91) by fibroblasts and other cell types. G-CSF stimulates the production of neutrophils, and IL-6 enhances maturation of B-cells and thereby the synthesis of immunoglobulins. The additional effects of these factors on hemopoietic stem cells in G_0 may assure an uninterrupted supply of the hemopoietic and possibly lymphoid progenitors for an integrated and effective defense against bacterial infections.

Factors Regulating Other Lineages

Regulation of megakaryopoiesis and thrombopoiesis remains unclear. While several factors such as megakaryocyte-CSF (92) and megakaryocyte stimulatory factor, which enhances synthesis of platelet-specific proteins by a rat megakaryoblast cell line (93), have been purified, their roles in the physiology of megakaryopoiesis and thrombopoiesis *in vivo* have not been established.

The proliferation of mast cells appears to be regulated by IL-3 and IL-4. Several years ago, investigators in a number of laboratories succeeded in growing murine mast cells in suspension culture in the presence of medium conditioned by pokeweed mitogen-stimulated spleen cells (PWM-SCM) (94-97). IL-3 was identified to be the main factor in PWM-SCM. However, IL-3 could not support the continued proliferation of mast cells in suspension cultures. In addition, it was found that IL-3 supports primar-

ily proliferation of mucosal-type mast cells but not connective tissue-type mast cells. More recently, IL-4 (98) [also known as B-cell stimulatory factor-1 (99)] was found to be an additional mast cell stimulating factor and was identified as the factor that is needed for the proliferation of connective tissue-type mast cells (100). The roles of human IL-3 and IL-4 in the proliferation of human mast cells have not been established.

Several factors have been shown to stimulate proliferation of eosinophils recently. Previously, several investigators documented that the human and murine GM-CSF and IL-3 support eosinophil proliferation in methylcellulose culture (66,68,69,72-74). In addition, *in vivo* administration of human GM-CSF in monkeys and men (75-77) revealed significant augmentation of the number of circulating eosinophils. Recently, Sanderson and his colleagues (101) identified eosinophil differentiation factor in the supernatant of cultures of alloreactive T-cell clones. This factor proved to be identical to T-cell replating factor (102) and is generally referred to as interleukin-5 (IL-5). The exact relationship between IL-5, GM-CSF, and IL-3 in eosinophilopoiesis remains yet to be determined.

Control of Hemopoiesis—A Model of Granulopoiesis

In addition to the factors already described, there are a number of factors known to affect hemopoiesis indirectly. Foremost, IL-1 α and β appear to play major roles in granulocyte/macrophage production. The synthesis of IL-1 by macrophages may be augmented by bacterial lipopolysaccharides (87). IL-1 then augments the production and the release of G-CSF (88,89) and GM-CSF (88,103-106) by a variety of cells, including fibroblasts, endothelial cells, and smooth muscle cells. Tumor necrosis factors (TNF α and TNF β) and interferons are known to inhibit myeloid proliferation in culture. A confusing aspect is that TNF α can stimulate the mesenchymal cells to synthesize and release G-CSF (107) and GM-CSF (108,109). It was reported that GM-CSF may in turn augment the synthesis and release of TNF α by monocytic cells (110). IL-4 may indirectly influence proliferation of progenitors supported by other cytokines (111,112). These observations indicate that a complex network of the cellular and molecular interactions regulate the continuous cellular production in hemopoiesis.

It is generally held that the feedback mechanisms exist for the control of individual lineages. In erythropoiesis, tissue oxygenation and Ep provide the feedback fine control of cell production. Regarding other lineages, no feedback system has been identified. It is possible that there is no such feedback regulation existing for granulocyte production. In this model, the production of granulocyte/macrophages in the steady state is maintained at genetically determined levels and is augmented only by external stimuli such as infections. In the presence of infections, bacterial lipopolysaccharides, IL-1 and TNF α will be involved in the augmentation of the synthesis and release of G-CSF and GM-CSF by various types of mesenchymal cells as outlined above. These factors, produced lo-

cally in the bone marrow, in turn augment the production of neutrophils, macrophages, and eosinophils. Once the external stimuli are removed, the production of the leukocytes reverts back to the steady-state level. Since the lifespans of leukocytes are brief, only coarse regulation of the cellular proliferation may be necessary. The wide normal ranges of granulocyte counts may attest to this hypothesis.

This work was supported by NIH grant AM32294 and the Veterans Administration. M. Ogawa is a Veterans Administration Medical Investigator. The author thanks Dr. Pamela N. Pharr and Linda S. Vann for their assistance in the preparation of this review.

REFERENCES

- Erslev, A. J. Production of erythrocytes. In: Hematology (W. J. Williams, E. Beutler, A. J. Erslev, M. A. Lichtman, Eds.), McGraw-Hill, New York, 1983, pp. 365-376.
- Dancey, J. T., Deubelbeiss, K. A., Harker, L. A., and Finch, C. A. Neutrophil kinetics in man. *J. Clin. Invest.* 58: 705-715 (1976).
- Till, J. E., and McCulloch, E. A. A direct measurement of the radiation sensitivity of normal mouse bone marrow cells. *Radiat. Res.* 14: 213-222 (1961).
- Golde, D. W. Methods in Hematology. Hematopoiesis. Churchill Livingstone, New York, 1984.
- Williams, D. A., Lemischka, I. R., Nathan, D. G., and Mulligan, R. C. Introduction of new genetic material into pluripotent hematopoietic stem cells of the mouse. *Nature* 310: 476-480.
- Lemischka, I. R., Raulet, D. H., and Mulligan, R. C. Developmental potential and dynamic behavior of hematopoietic stem cells. *Cell* 45: 917-927 (1986).
- Mintz, B., Anthony, K., and Litwin, S. Monoclonal derivation of mouse myeloid and lymphoid lineages from totipotent hematopoietic stem cells experimentally engrafted from fetal hosts. *Proc. Natl. Acad. Sci. (U.S.)* 81: 7835-7839 (1984).
- Becker, A. J., McCulloch, E. A., Siminovitch, L., and Till, J. E. The effect of differing demands for blood cell production on DNA synthesis by hemopoietic colony forming cells of mice. *Blood* 26: 296-308 (1965).
- Hara, H., and Ogawa, M. Murine hemopoietic colonies in culture containing normoblasts, macrophages, and megakaryocytes. *Am. J. Hematol.* 4: 23-34 (1978).
- Hodgson, G. S., and Bradley, T. R. Properties of hematopoietic stem cells surviving 5-fluorouracil treatment: Evidence for a pre-CFU-S cell? *Nature* 281: 381-382 (1979).
- Suda, T., Suda, J., and Ogawa, M. Proliferative kinetics and differentiation of murine blast cell colonies in culture: Evidence for variable G₀ periods and constant doubling rates of early pluripotent hemopoietic progenitors. *J. Cell. Physiol.* 117: 308-318 (1983).
- Lajtha, L. G. On the concept of the cell cycle. *J. Cell. Comp. Physiol.* 62(suppl. 1): 143-144 (1963).
- Lajtha, L. G. Stem cell concepts. *Differentiation* 14: 23-34 (1979).
- Till, J. E., McCulloch, E. A., and Siminovitch, L. A stochastic model of stem cell proliferation, based on the growth of spleen colony forming cells. *Proc. Natl. Acad. Sci. (U.S.)* 51: 29-36 (1964).
- Nakahata, T., Gross, A. J., and Ogawa, M. A stochastic model of self-renewal and commitment to differentiation of the primitive hemopoietic stem cells in culture. *J. Cell. Physiol.* 113: 455-458 (1982).
- Ogawa, M., Porter, P. N., and Nakahata, T. Renewal and commitment to differentiation of hemopoietic stem cells: An interpretive review. *Blood* 61: 823-829 (1983).
- Curry, J. L., and Trentin, J. J. Hemopoietic spleen colony studies: I. Growth and differentiation. *Dev. Biol.* 15: 395-413 (1967).
- Trentin, J. J. Influence of hematopoietic organ stroma (hematopoietic inductive microenvironments) on stem cell differentiation. In: Regulation of Hematopoiesis (A. S. Gordon, Ed.), Appleton-Century-Crofts, New York, 1970, pp. 161-186.
- VanZant, G., and Goldwasser, E. Simultaneous effects of erythropoietin and colony-stimulating factor on bone marrow cells. *Science* 198: 733-735 (1977).
- Goldwasser, E. Erythropoietin and differentiation of red blood cells. *Fed. Proc.* 34: 2285-2292 (1975).
- Johnson, G. R. Is erythropoiesis an obligatory step in the commitment of multipotential hematopoietic stem cells? In: Experimental Hematology Today (S. J. Baum, G. D. Ledney, and A. Kahn, Eds.), Springer-Verlag, New York, 1981, pp. 13-20.
- Nakahata, T., and Ogawa, M. Identification in culture of a class of hemopoietic colony-forming units with extensive capability to self-renew and generate multipotential colonies. *Proc. Natl. Acad. Sci. (U.S.)* 79: 3843-3847 (1982).
- Suda, T., Suda, J., and Ogawa, M. Single-cell origin of mouse hemopoietic colonies expressing multiple lineages in variable combinations. *Proc. Natl. Acad. Sci. (U.S.)* 80: 6689-6693 (1983).
- Suda, T., Suda, J., and Ogawa, M. Disparate differentiation in mouse hemopoietic colonies derived from paired progenitors. *Proc. Natl. Acad. Sci. (U.S.)* 81: 2520-2524 (1984).
- Suda, J., Suda, T., and Ogawa, M. Analysis of differentiation of mouse hemopoietic stem cells in culture by sequential replating of paired progenitors. *Blood* 64: 393-399 (1984).
- Leary, A. G., Ogawa, M., Strauss, L. C., and Civin, C. I. Single cell origin of multilineage colonies in culture: Evidence that differentiation of multipotent progenitors and restriction of proliferative potential of monopotent progenitors are stochastic processes. *J. Clin. Invest.* 74: 2193-2197 (1984).
- Leary, A. G., Strauss, L. C., Civin, C. I., and Ogawa, M. Disparate differentiation in hemopoietic colonies derived from human paired progenitors. *Blood* 66: 327-332 (1985).
- Wu, A. M. Regulation of self-renewal on human T lymphocyte colony-forming units (TL-CFUs). *J. Cell. Physiol.* 117: 101-108 (1983).
- Pharr, P. N., Nedelman, J., Downs, H. P., Ogawa, M., and Gross, A. J. A stochastic model for mast cell proliferation in culture. *J. Cell. Physiol.* 125: 379-386 (1985).
- Miyake, T., Kung, C. H., and Goldwasser, E. Purification of human erythropoietin. *J. Biol. Chem.* 252: 5558-5564 (1977).
- Lin, F.-K., Suggs, S., Lin, C.-H., Browne, J. K., Smalling, R., Egrie, J. C., Chen, K. K., Fox, G. M., Martin, F., Stabinsky, Z., Badrawi, S. M., Lai, P.-H., and Goldwasser, E. Cloning and expression of the human erythropoietin gene. *Proc. Natl. Acad. Sci. (U.S.)* 82: 7580-7584 (1985).
- Jacobs, K., Schoemaker, C., Rudersdorf, R., Neill, S. D., Kaufman, R. J., Mufson, A., Sehra, J., Jones, S. S., Hewick, R., Fritch, E. F., Kawakita, M., Shimizu, T., and Miyake, T. Isolation and characterization of genomic and cDNA clones of human erythropoietin. *Nature* 313: 806-810 (1985).
- Burgess, A. W., Camakaris, J., and Metcalf, D. Purification and properties of colony-stimulating factor from mouse lung-conditioned medium. *J. Biol. Chem.* 252: 1998-2003 (1977).
- Gough, N. M., Gough, J., Metcalf, D., Kelso, A., Grail, D., Nicola, N. A., Burgess, A. W., and Dunn, A. R. Molecular cloning of cDNA encoding a murine hematopoietic growth regulator, granulocyte-macrophage colony stimulating factor. *Nature* 309: 763-767 (1984).
- Gasson, J. C., Weisbart, R. H., Kaufman, S. E., Clark, S. C., Hewick, R. M., Wong, G. G., and Golde, D. W. Purified human granulocyte-macrophage colony-stimulating factor: Direct action on neutrophils. *Science* 226: 1339-1342 (1984).
- Wong, G. G., Witek, J. S., Temple, P. A., Wilkens, K. M., Leary, A. C., Luxenburg, D. P., Jones, S. S., Brown, E. L., Kay, R. M., Orr, E. C., Shoemaker, C., Golde, D. W., Kaufman, R. J., and Hewick, R. M. Human GM-CSF: Molecular cloning of cDNA and purification of the natural and recombinant proteins. *Science* 228: 810-815 (1985).
- Nicola, N. A., Metcalf, D., Matsumoto, M., and Johnson, G. R. Purification of a factor inducing differentiation in murine myelomonocytic leukemia cells. Identification as a granulocyte colony-stimulating factor. *J. Biol. Chem.* 258: 9017-9023 (1983).
- Welte, K., Platzer, E., Lu, L., Gabilove, J. L., Levi, E., Martelsmann, R., and Moore, M. A. Purification and biochemical characterization of human pluripotent hematopoietic colony-stimulating factor. *Proc. Natl. Acad. Sci. (U.S.)* 82: 1526-1530 (1985).

39. Nagata, S., Tsuchiya, M., Asano, S., Kaziro, Y., Yamazaki, T., Yamamoto, O., Hirata, Y., Kubota, N., Oheda, M., Nomme, H., and Ono, M. Molecular cloning and expression of cDNA for human granulocyte colony-stimulating factor. *Nature* 319: 415-417 (1986).
40. Ihle, J. N., Keller, J., Henderson, L., Klein, F., and Palaszynski, E. Procedures for the purification of interleukin 3 to homogeneity. *J. Immunol.* 129: 2431-2436 (1982).
41. Yokota, T., Lee, F., Rennick, D., Hall, C., Arai, N., Mosmann, T., Nabel, G., Cantor, H., and Arai, K. Isolation and characterization of a mouse cDNA clone that expresses mast-cell growth-factor activity in monkey cells. *Proc. Natl. Acad. Sci. (U.S.)* 81: 1070-1074 (1984).
42. Fung, M. C., Hapel, A. J., Ymer, S., Cohen, D. R., Johnson, R. M., Campbell, H. D., and Young, I. G. Molecular cloning of cDNA for murine interleukin-3. *Nature* 307: 233-237 (1984).
43. Yang Y.-C., Ciarletta, A. B., Temple, P. A., Chung, M. P., Kovacic, S., Witek-Giannotti, J. S., Leary, A. C., Kriz, R., Donahue, R. E., Wong, G. G., and Clark, S. C. Human IL-3 (multi CSF): Identification by expression cloning of a novel hemopoietic growth factor related to murine IL-3. *Cell* 47: 3-10 (1986).
44. Stanley, E. R., and Heard, P. M. Factors regulating macrophage production and growth. Purification and some properties of the colony stimulating factor from medium conditioned by mouse L cells. *J. Biol. Chem.* 252: 4305-4312 (1977).
45. Kawasaki, E. S., Ledner, M. B., Wang, A. M., Arsdell, J. V., Warren, M. K., Cayne, M. Y., Schiveickart, V. L., Lee, M.-T., Wilson, K. J., Boosman, A., Stanley, E. R., Ralph, P., and Mark, D. F. Molecular cloning of a complementary DNA encoding human macrophage-specific colony-stimulating factor (CSF-1). *Science* 230: 291-296 (1985).
46. Wong, G. G., Temple, P. A., Leary, A. C., Witek-Giannotti, J. S., Yang, Y.-C., Ciarletta, A. B., Chung, M., Murtha, P., Kriz, R., Kaufman, R. J., Ferenz, C. R., Sibley, B. S., Turner, K. J., Hewick, R. M., Clark, S. C., Yanai, N., Yokota, H., Yamada, M., Saito, M., Motoyoshi, K., and Takaku, F. Human CSF-1: Molecular cloning and expression of 4-kb cDNA encoding the human urinary protein. *Science* 235: 1504-1508 (1987).
47. Metcalf, D. The granulocyte-macrophage colony-stimulating factor. *Science* 229: 16-22 (1985).
48. Metcalf, D. The molecular biology and functions of the granulocyte-macrophage colony-stimulating factors. *Blood* 67: 257-267 (1986).
49. Clark, S. C., and Kamen, R. The human hemopoietic colony-stimulating factors. *Science* 236: 1229-1237 (1987).
50. Eschbach, J. W., Egrie, J. C., Downing, M. R., Browne, J. K., and Adamson, J. W. Correction of the anemia of end-stage renal disease with recombinant human erythropoietin. *N. Engl. J. Med.* 316: 73-78 (1987).
51. Gregory, C. J. Erythropoietin sensitivity as a differentiation marker in the hemopoietic system: Studies of three erythropoietic colony responses in culture. *J. Cell. Physiol.* 89: 289-302 (1976).
52. Ouellette, P. L., and Monette, F. C. Erythroid progenitors forming clusters *in vitro* demonstrate high erythropoietin sensitivity. *J. Cell. Physiol.* 105: 181-184 (1980).
53. Bruce, W. R., and McCulloch, E. A. The effect of erythropoietic stimulation on the hemopoietic colony-forming cells of mice. *Blood* 23: 216-232 (1964).
54. Hara, H., and Ogawa, M. Erythropoietic precursors in mice under erythropoietic stimulation and suppression. *Exp. Hematol.* 5: 141-148 (1977).
55. Iscove, N. N. The role of erythropoietin in the regulation of population size and cell cycling of early and late erythroid precursors in mouse bone marrow. *Cell Tissue Kinet.* 10: 323-334 (1977).
56. Kimura, H., Finch, C. A., and Adamson, J. W. Hematopoiesis in the rat: Quantitation of hemopoietic progenitors and the response to iron deficiency anemia. *J. Cell. Physiol.* 126: 298-306 (1986).
57. Metcalf, D., and Nicola, N. A. Proliferative effects of purified granulocyte colony-stimulating factor (G-CSF) on normal mouse hemopoietic cells. *J. Cell. Physiol.* 116: 198-206 (1983).
58. Eliason, J. F. Granulocyte-macrophage colony formation in serum-free culture: Effects of purified colony-stimulating factors and modulation by hydrocortisone. *J. Cell. Physiol.* 128: 231-238 (1986).
59. Sonoda, Y., Yang, Y.-C., Wong, G. G., Clark, S. C., and Ogawa, M. Analysis in serum-free culture of the targets of recombinant human hemopoietic growth factors: Interleukin-3 and granulocyte/macrophage colony-stimulating factors are specific for early developmental stages. *Proc. Natl. Acad. Sci. (U.S.)* 85: 4360-4364 (1988).
60. Shimamura, M., Kobayashi, Y., Yuo, A., Urabe, A., Okabe, T., Komatsu, Y., Itoh, S., and Takaku, F. Effect of human recombinant granulocyte colony-stimulating factor on hematopoietic injury in mice induced by 5-fluorouracil. *Blood* 69: 353-355 (1987).
61. Welte, K., Bonilla, A. M., Gillio, A. P., Boone, T. C., Potter, G. K., Gabrilove, J. L., Moore, M. A. S., O'Reilly, R. J., and Souza, L. M. Recombinant human granulocyte colony-stimulating factor: Effects on hematopoiesis in normal and cyclophosphamide-treated primates. *J. Exp. Med.* 165: 941-948 (1987).
62. Cohen, A. M., Zsebo, K. M., Inoue, H., Hines, D., Boone, T. C., Chazin, V. R., Tsai, L., Ritch, T., and Souza, L. M. *In vivo* stimulation of granulopoiesis by recombinant granulocyte colony-stimulating factor. *Proc. Natl. Acad. Sci. (U.S.)* 84: 2484-2488 (1987).
63. Motoyoshi, K., Takaku, F., and Miura, Y. High serum colony-stimulating activity of leukocytopenic patients after intravenous infusions of human urinary colony-stimulating factor. *Blood* 62: 685-688 (1983).
64. Bartocci, A., Pollard, J. W., and Stanley, E. R. Regulation of colony-stimulating factor 1 during pregnancy. *J. Exp. Med.* 164: 956-961 (1986).
65. Ihle, J. N., Keller, J., Oroszlan, S., Henderson, L. E., Copeland, T. D., Fitch, F., Prystowsky, M. B., Goldwasser, E., Schrader, J. W., Palaszynski, E., Dy, M., and Lebel, B. Biologic properties of homogeneous interleukin 3. I. Demonstration of WEHI-3 growth factor activity, mast cell growth factor activity, P cell-stimulating factor activity, colony-stimulating factor activity and histamine-producing cell-stimulating factor activity. *J. Immunol.* 131: 282-287 (1983).
66. Suda, T., Suda, J., Ogawa, M., and Ihle, J. N. Permissive role of interleukin 3 (IL-3) in proliferation and differentiation of multipotential hemopoietic progenitors in culture. *J. Cell. Physiol.* 124: 182-190 (1985).
67. Rennick, D. M., Lee, F. D., Yokota, T., Arai, K., Cantor, H., and Nabel, G. J. A cloned MCGF cDNA encodes a multilineage hemopoietic growth factor: Multiple activities of interleukin 3. *J. Immunol.* 134: 910-914 (1985).
68. Koike, K., Ihle, J. N., and Ogawa, M. Declining sensitivity to interleukin 3 of murine multipotential hemopoietic progenitors during their development: Application to a culture system that favors blast cell colony formation. *J. Clin. Invest.* 77: 894-899 (1986).
69. Leary, A. G., Yang, Y.-C., Clark, S. C., Gasson, J. C., Golde, D. W., and Ogawa, M. Recombinant gibbon interleukin-3 (IL-3) supports formation of human multilineage colonies and blast cell colonies in culture: Comparison with recombinant human granulocyte-macrophage colony-stimulating factor (GM-CSF). *Blood* 70: 1343-1348 (1987).
70. Metcalf, D., Johnson, G. R., and Burgess, A. W. Direct stimulation by purified GM-CSF of the proliferation of multipotential and erythroid precursors. *Blood* 55: 138-147 (1980).
71. Koike, K., Ogawa, M., Ihle, J. N., Miyake, T., Shimizu, T., Miyajima, A., Yokota, T., and Arai, K. Recombinant murine granulocyte-macrophage (GM) colony-stimulating factor supports formation of GM and multipotential blast cell colonies in culture: Comparison with the effects of interleukin-3. *J. Cell. Physiol.* 131: 458-464 (1987).
72. Sieff, C. A., Emerson, S. G., Donahue, R. E., Nathan, D. G., Wang, E. A., Wong, G. G., and Clark, S. C. Human recombinant granulocyte-macrophage colony-stimulating factor: A multilineage hemopoietin. *Science* 230: 1171-1173 (1985).
73. Gabrilove, J. L., Welte, K., Harris, P., Platzer, E., Lu, L., Levi, E., Mertelsmann, R., and Moore, M. A. S. Pluripoietin α : A second human hemopoietic colony-stimulating factor produced by the human bladder carcinoma cell line 5637. *Proc. Natl. Acad. Sci. (U.S.)* 83: 2478-2482 (1986).
74. Kaushansky, K., O'Hara, P. J., Berkner, K., Segel, G. M., Hagen, F. S., and Adamson, J. W. Genomic cloning, characterization, and multilineage growth-promoting activity of human granulocyte-macrophage colony-stimulating factor. *Proc. Natl. Acad. Sci. (U.S.)* 83: 3101-3105 (1986).

75. Donahue, R. E., Wang, E. A., Stone, D. K., Kamen, R., Wong, G. G., Sehgal, P. K., Nathan, D. G., and Clark, S. C. Stimulation of haematopoiesis in primates by continuous infusion of recombinant human GM-CSF. *Nature* 321: 872-875 (1986).
76. Nienhuis, A. W., Donahue, R. E., Karlsson, S., Clark, S. C., Agricola, B., Antinoff, N., Pierce, J. E., Turner, P., Anderson, W. F., and Nathan, D. G. Recombinant human granulocyte-macrophage colony-stimulating factor (GM-CSF) shortens the period of neutropenia after autologous bone marrow transplantation in a primate model. *J. Clin. Invest.* 80: 573-577 (1987).
77. Groopman, J. E., Mitsuyasu, R. T., DeLeo, M. J., Oette, D. H., and Golde, D. W. Effect of recombinant human granulocyte-macrophage colony-stimulating factor on myelopoiesis in the acquired immunodeficiency syndrome. *N. Engl. J. Med.* 317: 593-598 (1987).
78. Stanley, E. R., Bartocci, A., Patinkin, D., Rosendaal, M., and Bradley, T. R. Regulation of very primitive, multipotent, hemopoietic cells by hemopoietin-1. *Cell* 45: 667-674 (1986).
79. Jubinsky, P. T., and Stanley, E. R. Purification of hemopoietin 1: A multilineage hemopoietic growth factor. *Proc. Natl. Acad. Sci. (U.S.)* 82: 2764-2768 (1985).
80. Mochizuki, D. Y., Eisenman, J. A., Conlon, P. J., Larsen, A. D., and Tushinski, R. J. Interleukin 1 regulates hematopoietic activity, a role previously ascribed to hemopoietin 1. *Proc. Natl. Acad. Sci. (U.S.)* 84: 5267-5271 (1987).
81. Moore, M. A. S., and Warren, D. J. Synergy of interleukin-1 and granulocyte colony-stimulating factor: *In vivo* stimulation of stem-cell recovery and hematopoietic regeneration following 5-fluorouracil treatment of mice. *Proc. Natl. Acad. Sci. (U.S.)* 84: 7134-7138 (1987).
82. Zilberstein, A., Ruggieri, R., Korn, J. H., and Revel, M. Structure and expression of cDNA and genes for human interferon-beta-2, a distinct species inducible by growth stimulatory cytokines. *EMBO J.* 5: 2529-2537 (1986).
83. Hirano, T., Yasukawa, K., Harada, H., Taga, T., Watanabe, Y., Matsuda, T., Kashiwamura, S., Nakajima, K., Koyama, K., Iwamatsu, A., Tsunashima, S., Sakiyama, F., Matsui, H., Takahara, Y., Taniguchi, T., and Kishimoto, T. Complementary DNA for a novel human interleukin (BSF-2) that induces B lymphocytes to produce immunoglobulin. *Nature* 324: 73-77 (1986).
84. Ikebuchi, K., Wong, G. G., Clark, S. C., Ihle, J. N., Hirai, Y., and Ogawa, M. Interleukin-6 enhancement of interleukin-3-dependent proliferation of multipotential hemopoietic progenitors. *Proc. Natl. Acad. Sci. (U.S.)* 84: 9035-9039 (1987).
85. Leary, A. G., Ikebuchi, K., Wong, G. G., Yang, Y.-C., Clark, S. C., and Ogawa, M. Interleukin-6 is a synergistic factor for human hemopoietic stem cells. *Blood* 70(suppl. 1): 176a (1987).
86. Ikebuchi, K., Clark, S. C., Ihle, J. N., Souza, L. M., and Ogawa, M. Granulocyte colony-stimulating factor enhances interleukin-3-dependent proliferation of multipotential hemopoietic progenitors. *Proc. Natl. Acad. Sci. (U.S.)* 85: 3445-3449 (1988).
87. Dinarello, C. A. Interleukin-1. *Rev. Infect. Dis.* 6: 51-95 (1984).
88. Rennick, D., Yang, G., Gemmell, L., and Lee, F. Control of hemopoiesis by a bone marrow stromal cell clone: Lipopolysaccharide- and interleukin-1-inducible production of colony-stimulating factors. *Blood* 69: 682-691 (1987).
89. Broudy, V. C., Kaushansky, K., Harlan, J. M., and Adamson, J. W. Interleukin 1 stimulates human endothelial cells to produce granulocyte-macrophage colony-stimulating factor and granulocyte colony-stimulating factor. *J. Immunol.* 139: 464-468 (1987).
90. Yang, Y.-C., Tsai, S., Wong, G. G., and Clark, S. C. Interleukin-1 regulation of hematopoietic growth factor production by human stromal fibroblasts. *J. Cell. Physiol.* 134: 292-296 (1988).
91. Yasukawa, K., Hirano, T., Watanabe, Y., Muratani, K., Matsuda, T., Nakai, S., and Kishimoto, T. Structure and gene expression of human B cell stimulatory factor-2 (BSF-2/IL-6) gene. *EMBO J.* 5: 2529-2537 (1987).
92. Hoffman, R., Yang, H. H., Bruno, E., and Straneva, J. E. Purification and partial characterization of a megakaryocyte colony-stimulating factor from human plasma. *J. Clin. Invest.* 75: 1174-1182 (1985).
93. Tayrien, G., and Rosenberg, R. D. Purification and properties of a megakaryocyte stimulatory factor present both in the serum-free conditioned medium of human embryonic kidney cells and in thrombocytopenic plasma. *J. Biol. Chem.* 262: 3262-3268 (1987).
94. Schrader, J. W., Lewis, S. J., Clark-Lewis, I., and Culvenor, J. G. The persisting (P) cell: Histamine content, regulation by a T cell-derived factor, origin from a bone marrow precursor, and relationship to mast cells. *Proc. Natl. Acad. Sci. (U.S.)* 78: 323-327 (1981).
95. Nagao, K., Yokoro, K., and Aaronson, S. A. Continuous lines of basophil/mast cells derived from normal mouse bone marrow. *Science* 212: 333-335 (1981).
96. Tertian, G., Yung, Y. P., Guy-Grand, D., and Moore, M. A. S. Long-term *in vitro* culture of murine mast cells. 1. Description of a growth factor-dependent culture technique. *J. Immunol.* 127: 788-794 (1981).
97. Razin, E., Cordon-Cordo, C., and Good, R. A. Selective growth of a population of human basophil cells *in vitro*. *Proc. Natl. Acad. Sci. (U.S.)* 78: 5793-5796 (1981).
98. Lee, F., Yokota, T., Otsuka, T., Meyerson, P., Villaret, D., Coffman, R., Mosmann, T., Rennick, D., Roehm, N., Smith, C., Zlotnik, A., and Arai, K. Isolation and characterization of a mouse interleukin cDNA clone that expresses B-cell stimulatory factor 1 activities and T-cell- and mast-cell-stimulating activities. *Proc. Natl. Acad. Sci. (U.S.)* 83: 2061-2065 (1986).
99. Howard, M., Farrar, J., Hilfiker, M., Johnson, B., Takatsu, K., Hamaoka, T., and Paul, W. E. Identification of a T cell-derived B cell growth factor distinct from interleukin-2. *J. Exp. Med.* 155: 914-923 (1982).
100. Hamaguchi, Y., Kanakura, Y., Fujita, J., Takeda, S., Nakano, T., Tarui, S., Honjo, T., and Kitamura, Y. Interleukin 4 as an essential factor for *in vitro* clonal growth of murine connective tissue-type mast cells. *J. Exp. Med.* 165: 268-273 (1987).
101. Sanderson, C. J., O'Garra, A., Warren, D. J., and Klaus, G. G. B. Eosinophil differentiation factor also has B-cell growth factor activity: Proposed name interleukin 4. *Proc. Natl. Acad. Sci. (U.S.)* 83: 437-440 (1986).
102. Kinashi, T., Harada, N., Severinson, E., Tanabe, T., Sideras, P., Tonishi, M., Azuma, C., Tominaga, A., Bergstedt-Lindqvist, S., Takahashi, M., Matsuda, F., Yaoita, Y., Takatsu, K., and Honjo, T. Cloning of complementary DNA encoding T-cell replacing factor and identity with B-cell growth factor II. *Nature* 324: 70-73 (1986).
103. Zucali, J. R., Dinarello, C. A., Oblon, D. J., Gross, M. A., Anderson, L., and Weiner, R. S. Interleukin 1 stimulates fibroblasts to produce granulocyte-macrophage colony-stimulating activity and prostaglandin E₂. *J. Clin. Invest.* 77: 1857-1863 (1986).
104. Bagby, G. C. Jr., Dinarello, C. A., Wallace, P., Wagner, C., Hefeneider, S., and McCall, E. Interleukin 1 stimulates granulocyte macrophage colony-stimulating activity release by vascular endothelial cells. *J. Clin. Invest.* 78: 1316-1323 (1986).
105. Fibbe, W. E., vanDamme, J., Billiau, A., Voogt, P. J., Duinkerken, N., Kluck, P. M. C., and Falkenburg, J. H. F. Interleukin-1 (22-K factor) induces release of granulocyte-macrophage colony-stimulating activity from human mononuclear phagocytes. *Blood* 68: 1316-1321 (1986).
106. Sieff, C. A., Tsai, S., and Faller, D. V. Interleukin 1 induces cultured human endothelial cell production of granulocyte-macrophage colony-stimulating factor. *J. Clin. Invest.* 79: 48-51 (1987).
107. Koeffler, H. P., Gasson, J., Ranyard, J., Souza, L., Shepard, M., and Munker, R. Recombinant human TNF α stimulates production of granulocyte colony-stimulating factor. *Blood* 70: 55-59 (1987).
108. Munker, R., Gasson, J., Ogawa, M., and Koeffler, H. P. Recombinant human TNF induces production of granulocyte-macrophage colony-stimulating factor. *Nature* 323: 79-82 (1986).
109. Broudy, V. C., Kaushansky, K., Segal, G. M., Harlan, J. M., and Adamson, J. W. Tumor necrosis factor type α stimulates human endothelial cells to produce granulocyte-macrophage colony-stimulating factor. *Proc. Natl. Acad. Sci. (U.S.)* 83: 7467-7471 (1986).
110. Cannistra, S. A., Rambaldi, A., Spriggs, D. R., Herrmann, F., Kufe, D., and Griffin, J. D. Human granulocyte-macrophage colony-stimulating factor induces expression of the tumor necrosis factor

- gene by the U937 cell line and by normal human monocytes. *J. Clin. Invest.* 79: 1720-1728 (1987).
111. Peschel, C., Paul, W. E., Ohara, J., and Green, I. Effects of B cell stimulatory factor-1/interleukin 4 on hematopoietic progenitor cells. *Blood* 70: 254-263 (1987).
112. Rennick, D., Yang, G., Muller-Sieburg, C., Smith, C., Arai, N., Takabe, Y., and Gemmell, L. Interleukin 4 (B-cell stimulatory factor 1) can enhance or antagonize the factor-dependent growth of hemopoietic progenitor cells. *Proc. Natl. Acad. Sci. (U.S.)* 84: 6889-6893 (1987).