

Computational Estimation and Experimental Verification of Off-Target Silencing during Posttranscriptional Gene Silencing in Plants^{1[W][OA]}

Ping Xu², Yuanji Zhang², Li Kang, Marilyn J. Roossinck, and Kirankumar S. Mysore*

Plant Biology Division, The Samuel Roberts Noble Foundation, Ardmore, Oklahoma 73401

Successful application of posttranscriptional gene silencing (PTGS) for gene function study in both plants and animals depends on high target specificity and silencing efficiency. By computational analysis with genome and/or transcriptome sequences of 25 plant species, we predicted that about 50% to 70% of gene transcripts in plants have potential off-targets when used for PTGS that could obscure experimental results. We have developed a publicly available Web-based computational tool called siRNA Scan to identify potential off-targets during PTGS. Some of the potential off-targets obtained from this tool were tested by measuring the amount of off-target transcripts using quantitative reverse transcription-PCR. Up to 50% of the predicted off-target genes tested in plants were actually silenced when tested experimentally. Our results suggest that a high risk of off-target gene silencing exists during PTGS in plants. Our siRNA Scan tool is useful to design better constructs for PTGS by minimizing off-target gene silencing in both plants and animals.

Posttranscriptional gene silencing (PTGS), also known as RNA interference (RNAi) in animals, cosuppression in plants, and RNA quelling in fungi, is an epigenetic phenomenon that results in sequence-specific degradation of endogenous mRNAs (Cogoni and Macino, 2000). PTGS is mediated by 21- to 24-nucleotide (nt) double-strand RNA (dsRNA) molecules, termed short/small interference RNAs (siRNAs). The siRNAs are incorporated into a multisubunit protein complex, the RNAi-induced silencing complex, which directs the siRNA to and degrades the complementary mRNA (Baulcombe, 2004). Because PTGS allows silencing of a specific target gene, it has become a popular tool to study gene function in plants and animals (Kamath et al., 2003; Baulcombe, 2004; Hannon and Rossi, 2004; Kuttenkeuler and Boutros, 2004).

In plants, PTGS can be induced by antisense and sense transgenic technology and it is also achieved by expressing dsRNA through stable or transient transformation with RNAi constructs (binary vectors) to knock down the expression of target genes (Wesley et al., 2001; Miki and Shimamoto, 2004). In addition, a transient PTGS of plant genes by recombinant viruses

carrying a near-identical sequence was adapted by a process called virus-induced gene silencing (VIGS; Baulcombe, 1999; Burch-Smith et al., 2004). Both PTGS approaches, RNAi and VIGS, are becoming powerful tools in functional genomic studies of plants. However, to infer gene function through PTGS, it is essential to determine the specificity of mRNAs that are targeted for silencing.

Theoretically, PTGS functions in a siRNA-specific rather than a target-specific manner. However, analyses of mammalian cells transfected with different siRNAs against a target gene by two different research groups led to contradictory conclusions about silencing unintended genes (Chi et al., 2003; Jackson et al., 2003; Jackson and Linsley, 2004). The reasons for these contradictory results are probably due to the differences in experimental designs or microarray analyses. In plants, several RNAi and VIGS studies successfully targeted specific members of gene families for silencing without affecting the transcript level of the most closely related family members or simultaneously silenced a few family members to overcome functional redundancy (Burch-Smith et al., 2004; Hwang and Gelvin, 2004). However, PTGS relies upon sequences of contiguous nucleotide identity and does not target genes simply based on family relationships. A dsRNA expressed in plants from a binary or virus vector is usually chosen to be identical to a partial or full-length sequence of the target gene. Many distinct siRNAs of 21 to 24 nt can be derived from the cleavage of the dsRNA by Dicer. This may improve the efficiency of RNA silencing, but could also increase the opportunity to suppress unintended genes, the off-targets, containing sequences identical to some of these siRNAs.

Investigating off-target gene silencing is crucial for accurate interpretation of gene function by PTGS and

¹ This work was supported by The Samuel Roberts Noble Foundation.

² These authors contributed equally to the paper.

* Corresponding author; e-mail ksmysore@noble.org; fax 580-224-6692.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Kirankumar S. Mysore (ksmysore@noble.org).

[W] The online version of this article contains Web-only data.

[OA] Open Access articles can be viewed online without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.106.083295

for use of PTGS application in agriculture. In this study, we estimate potential off-targets based on sequence identity for 25 plant species whose genomic or expressed sequence tag (EST) sequences are publicly available. Experimentally, we investigated the expression level of several potential off-target genes in an RNAi transgenic Arabidopsis (*Arabidopsis thaliana*) line and in gene-silenced (by VIGS) *Nicotiana benthamiana* plants. To assist the design of PTGS constructs to minimize off-target gene silencing or to identify potential off-targets from a particular PTGS construct, we have developed a Web-based computational tool. The tool provides an integrated sequence similarity search environment for plant and animal species, identifies potential off-targets, and predicts putatively effective siRNAs from the target query sequences.

RESULTS

Computational Analysis of Potential Off-Targets during PTGS

PTGS in plants is mediated by siRNAs derived from the cleavage of dsRNA produced through binary vector-based RNAi or VIGS constructs. The siRNAs recognize their targets by base pairing. Therefore, the potential off-targets are identified by a direct sequence identity and reverse complementary sequence identity search. In this study, we define the cDNA producing dsRNA for silencing of the intended target gene (from which the cDNA comes) as the trigger and the unintended genes that share a contiguous ≥ 21 -nt region of identity or reverse complementary identity to the trigger as off-targets. The continuous identical or reverse complementary sequence to a trigger region in an off-target is defined as an off-target region.

Using the full-length cDNA sequence of every transcript in the Arabidopsis genome mRNA dataset (ATH1) as a trigger, we estimated the patterns of potential off-targets by searching for identical or reverse complementary regions of ≥ 21 nt between a trigger sequence and all the other genes, except the target itself. Differently spliced transcripts from the same locus in the genome were not counted as additional off-targets. The potential off-target region was scanned along the full length of all released transcript sequences. Off-target regions can be in the 5' and 3' untranslated regions or coding regions because all these regions have been shown to be targeted by siRNAs (McManus et al., 2002; Yu et al., 2002; Yokota et al., 2003; Shirane et al., 2004). By using results obtained from Arabidopsis data analyses, we compared the distributions of off-target/trigger, off-target site/off-target, and length of off-target site/off-target site (Fig. 1) with their statistical averages (Table I). The histograms shown in Figure 1 have skewed distributions, with a majority of the trigger genes having similar potential off-targets as the average (Fig. 1A), a majority of the off-targets having approximately the

same number of off-target sites as the average (Fig. 1B), and a majority of off-target sites having almost the same length as the average (Fig. 1C). The average values for off-targets, off-target sites, and lengths of off-target sites are also close to the peak values (medians).

On average, 68.7% of Arabidopsis transcripts can potentially silence 3.9 off-target genes (Table I). Each candidate off-target has an average of three off-target regions with an average length of over 40 nt (Table I). However, this could be an overestimation because not all siRNAs derived from the cleavage of dsRNAs are efficient at silencing (Khvorova et al., 2003; Schwarz et al., 2003). Because little is known about the efficacy of specific siRNAs in plants, we adapted the rules developed by Ui-Tei et al. (2004) to predict whether and how many of the trigger sequences mentioned above produce efficient siRNAs to silence their potential off-targets. Specifically, the rules for a 21-nt siRNA duplex with a 2-nt overhang include: (1) antisense strand starts with A/U; (2) sense strand starts with G/C; (3) at least five of the first seven residues at the 5' terminus of the antisense strand should be A/U; and (4) there is no G/C stretch of more than 9 nt in length (Ui-Tei et al., 2004). As shown in Table I, 68.7% of all trigger sequences in the ATH1 database are predicted to generate an average of 35 efficient 21-nt siRNAs to cause potential off-target silencing.

In addition to Arabidopsis, we also estimated the potential off-targets in 24 other plant species, using Gene Indices assembled from ESTs and other expressed transcripts provided by The Institute for Genomic Research (TIGR; Quackenbush et al., 2001). We first compared the results from Arabidopsis Gene Index version 11 (AGI 11) with the results from the genome mRNAs described above (ATH1). Slightly more trigger sequences were predicted to have potential off-targets when the AGI 11 dataset was analyzed (Table I). This is because numerous unique gene transcripts in AGI 11 are not represented in the current genome sequence. Estimations of off-targets in other plant species also indicated high likelihood of off-target silencing (Table I). The potential off-target effects for plants like rye (*Secale cereale*), pepper (*Capiscum annuum*), and Lotus japonicus may be underestimated in this analysis because of limited sequence availability. For example, the percentage of target sequences predicted to silence off-targets is approximately 26% for these species. When those species with <20,000 EST sequences in their gene indices are excluded, 50% to 70% of all gene transcripts in each of 17 plant species used as silencing triggers can cause potential off-target silencing. The average numbers of off-targets for each trigger sequence range from 2.4 in tomato (*Lycopersicon esculentum*) to 24 in lettuce (*Lactuca sativa*) and sugarcane (*Saccharum officinarum*; Table I).

Off-target silencing, however, may be desirable in PTGS applications to generate a loss-of-function phenotype if the target and off-targets are functionally redundant. For example, the same family members or




Figure 1. Histograms showing distributions of the numbers of silencing triggers and off-targets in Arabidopsis. A, Number of silencing triggers versus the number of off-targets. B, Number of off-targets versus the number of off-target sites. C, Number of off-target sites versus the length of off-target sites. y axis is in log₁₀ scale.

genes encoding proteins that share conserved functional protein domains may be simultaneously silenced. Arabidopsis gene family information at The Arabidopsis Information Resource (TAIR) Web site (<http://www.arabidopsis.org>) was used to evaluate the chance that potential off-targets belong to the same gene family as the target gene. The data from the TAIR ftp site, after removing ambiguous entries and single-member families, contain 5,842 genes from 674 families (see Supplemental Table S1), with family size ranging from two to 307 members. Our analysis indicates that 4,677 gene sequences (79.9% of 5,842 genes) of these families may trigger off-target silencing of other members of the same family when a full-length

sequence is used as a trigger. Among the families with different numbers of gene members, the number of off-targets was different but generally fewer than six (Fig. 2; Supplemental Table S1). Moreover, only 418 genes from 192 families containing fewer than 10 members share an identical region of ≥ 21 nt with all other members in the same gene family, so only these transcripts used as trigger sequences can silence all family members. No single trigger sequence from any family with more than 10 members was predicted to silence all members of the family. In addition, the percentage of gene members to be silenced decreases from approximately 79% to $< 9\%$ as family size increases from two to > 35 . The family CYP705A in the

Table 1. Estimated off-target patterns during PTGS in plants

Species	Source Sequence Set	Release Date	Total No. Unique Gene Transcripts	No. Triggers Predicted to Silence Off-Targets (% Total)	Average No. Predicted Off-Targets/Trigger	Average No. Predicted Off-Target Regions/Off-Target	Average Length (nt)/Off-Target Region	Average No. Efficient siRNAs/Trigger
<i>Arabidopsis</i>	ATH1	6/10/04	28,952	19,882 (68.7)	3.9	2.9	41.2	35.2
<i>Arabidopsis</i>	AGI 11	1/29/04	45,683	37,250 (81.5)	4.3	3	99.2	147.5
<i>Capsicum annuum</i>	CaGI 1	6/4/04	10,712	27,88 (26.0)	1.9	2.2	40.7	12.5
<i>Gossypium</i> spp.	CGI 5	9/16/03	24,350	14,270 (58.6)	6.9	2.3	74	87.5
<i>Chlamydomonas reinhardtii</i>	ChrGI 4	1/31/04	30,339	16,579 (54.6)	6.1	1.9	40.2	12.5
<i>Glycine max</i>	GmGI 11	9/16/03	67,826	49,658 (73.2)	9.4	2.1	77.9	108.4
<i>Helianthus annuus</i>	HaGI 3	9/16/03	20,520	11,961 (58.3)	15.8	1.4	34.6	38.8
<i>Hordeum vulgare</i>	HvGI 8	1/30/04	49,190	32,297 (65.6)	16.3	2.6	99.4	205.9
Tomato	LeGI 9	5/19/03	31,012	15,340 (49.5)	2.4	2.3	60.8	30.9
<i>Lotus japonicus</i>	LjGI 2	5/16/03	11,025	2,951 (26.8)	2.6	1.7	35.6	10
Lettuce	LsGI 2	2/3/04	22,185	12,345 (55.6)	24.1	1.4	30.4	26.8
<i>Mesembryanthemum crystallinum</i>	McGI 4	5/15/03	8,455	3,459 (42)	13.3	2	33	29.4
<i>Medicago truncatula</i>	MtGI 7	5/19/03	36,976	19,468 (52.7)	5.6	2.2	49.4	50.6
<i>Nicotiana benthamiana</i>	NbGI 1	1/28/04	6,118	31,99 (52.3)	2.8	3.8	53.4	42.1
<i>Nicotiana tabacum</i>	NtGI 1	6/4/04	10,232	2,184 (21.3)	1.8	3.1	38.3	15.2
<i>Allium cepa</i>	OnGI 1	9/17/03	11,726	3,735 (31.9)	1.7	2.3	54.2	20.2
Rice	OsGI 15	5/27/04	88,765	45,406 (51.2)	22.5	4.4	41.7	174.3
<i>Pinus</i> spp.	PGI 4	1/28/04	31,771	16,487 (51.9)	11.7	5.5	38.4	27.5
Rye	RyeGI 3	1/29/04	5,347	1,472 (27.5)	2.7	6	32.4	10.7
<i>Sorghum bicolor</i>	SbGI 8	5/26/04	39,148	22,039 (56.3)	6.4	1.9	70.3	56.9
Sugarcane	SoGI 1	1/30/04	95,884	55,269 (57.6)	24.1	3.5	69.1	205.4
<i>Solanum tuberosum</i>	StGI 9	5/26/04	32,553	20,469 (62.9)	4.4	2.9	46.2	42.2
<i>Triticum aestivum</i>	TaGI 8	1/29/04	12,3807	89,862 (72.6)	19.5	2.7	52.4	102.2
<i>Theobroma cacao</i>	TcaGI 1	6/4/04	2,539	531 (20.9)	2.6	1.9	37.4	13
<i>Vitis vinifera</i>	VvGI 3	11/14/03	23,109	10,548 (45.6)	8.8	1.8	40.3	37.1
<i>Zea mays</i>	ZmGI 14	1/2/04	56,364	40,028 (71)	8.9	3	63.1	96.9

P450 superfamily is an exception. This family has 24 gene members, and >75% of them were predicted to be potential off-targets when any one of its members is used as an RNAi trigger (Fig. 2).

An Integrated Search Environment for PTGS

An online tool (available at <http://bioinfo2.noble.org/RNAiScan.htm>) has been developed to evaluate and minimize the risk of off-target effects during

PTGS. The tool is a search environment with several integrated components, including a sequence similarity search to identify potential off-targets, efficiency estimation of siRNAs, and functional analysis of off-target genes. In the sequence similarity search, the query sequence is compared with a gene transcript dataset using the BLASTn program (Altschul et al., 1997) with user-specified mismatches that may be tolerable in siRNA target recognition (Saxena et al., 2003). The datasets include *Arabidopsis* and rice




Figure 2. Average number of off-target gene family members predicted to be silenced by a trigger from the same family in *Arabidopsis*. Data for gene families with the same size are combined and averaged.

(*Oryza sativa*) genome mRNAs and all TIGR gene indices for plants and animals, and they are updated regularly. All possible siRNAs derived from the query are then evaluated for efficiency using Ui-Tei's rule (Ui-Tei et al., 2004). The predicted off-target genes can be subject to further functional analysis by identifying Pfam domains (Sonnhammer et al., 1998) and associated gene ontology assignments. The tool can also be used to estimate off-targets during siRNA-induced gene silencing in animals. In addition to displaying the potential siRNAs and off-target candidates, the results page also displays all the potential siRNAs with no sequence similarity to any other genes in the searched database. Therefore, the RNAi scan tool can be used to design siRNAs with minimal off-targets.

Experimental Verification of Off-Target Silencing in Plants

Expression of at Least Three Off-Target Genes Was Knocked Down in *BT11* RNAi Transgenic Lines

BT11 is an *Arabidopsis* protein that interacts with the *Agrobacterium tumefaciens* VirB2 protein (Hwang and Gelvin, 2004). A functional study of *BT11* through RNAi transgenic *Arabidopsis* lines has been published, where dsRNA covering the coding region of *BT11* was produced from the RNAi vector pFGC5941 (Hwang and Gelvin, 2004). We used the coding region of *BT11*

as a trigger sequence to search for potential off-targets against the TIGR AGI 12.1 database with the Web-based tool siRNA Scan described above. We further confirmed the annotations of these off-targets by BLAST search in the National Center for Biotechnology (NCBI). In addition to one family member, *BT12*, 13 other genes were found that share at least 21-nt continuous direct identity or reverse complementary identity to the *BT11* coding sequence (Table II). These 14 genes have the potential to be targeted by the siRNAs generated from *BT11*-derived dsRNA. Twelve of them were selected for the analysis of their expression levels in *BT11* RNAi transgenic lines (Table II). They are TC251703 (*BT12*), TC251496 (putative 3-isopropylmalate dehydratase large subunit), TC256637 (encodes a probable Ser/Thr kinase), TC258543 (RTNLB6), TC255665 (encodes a zinc-finger motif protein), TC262843 (encodes a hypothetical protein), TC263798 (encodes a protein of unknown function), TC269146 (encodes a hypothetical protein), TC275407 (encodes a putative Ser/Thr kinase), TC265975 (encodes a probable membrane protein), TC275528 (encodes a maturase-related protein), and TC275625 (encodes protein of unknown function). In addition, expression of the other 22 genes that share almost 21- or 22-nt continuous identity to the *BT11* coding region, but with one mismatch, was also analyzed. The expression levels of these genes, as well as the target gene *BT11*, were measured by real-time quantitative reverse transcription-PCR

Table II. Result from the siRNA Scan search with *Arabidopsis BT11* coding sequence as a trigger

TC numbers in bold are the ones used for quantitative RT-PCR.

Off-Target ID	Identical (I) or Reverse Complementary (RC) Sequence	BLAST Search
NP167859	I, AGCCTGTTTCATAAGGTTCTCGG	<i>Arabidopsis</i> reticulon family protein (RTNLB6) (At3g61560).
NP302761	I, AGAAGAAGAAGACTAAGAAGC I, AGCCTGTTTCATAAGGTTCTCGG	Contains similarity to DnaJ gene YM8520.10 gb 825566 from <i>Saccharomyces cerevisiae</i> cosmid gb Z49705. ESTs gb Z47720 and gb Z29879 come from this gene.
TC251496	RC, TGATTCTTCTTCGTTCTTCTC	GB AAM51226 Unknown protein { <i>Arabidopsis</i> }; similar to UP Q6Z702 (Q6Z702) putative 3-isopropylmalate dehydratase large subunit.
TC251703	I, CCTGTTTCATAAGGTTCTCGGCG; I, TGGTCTAATGCCACTATGTTTCATT; I, CCAAAGATTCTGAAGTTCATATCCCT- GAAGAACCT; I, TTCTTGACATTGGCATAACATAGCTCT	GB AAP47461.1 32331867 AY164887 RTNLB2 { <i>Arabidopsis</i> }, complete. Recently identified as <i>BT12</i> .
TC255665	I, TCTTCGTTCTTTCATCTTCT	GB AAD31078.1 4850408 F3F19 Contains PF 00097 zinc finger (C3HC4) ring finger motif. { <i>Arabidopsis</i> }, complete.
TC256637	RC, TTCTTCTTCGTTCTTCTTCATC	PIR B96716 Probable Ser/Thr kinase F23O10.20 [imported]— <i>Arabidopsis</i> .
TC258543	I, AGCCTGTTTCATAAGGTTCTCGG	GB AAP47457.1 32331859 AY164883 RTNLB6 { <i>Arabidopsis</i> }, partial (90%).
TC262843	I, CTTCTTCGTTCTTTCATCTTCT	UPI O49467 (O49467) Hypothetical protein F24J7.50 (hypothetical protein AT4g19490).
TC263798	RC, TTCTTCTTCGTTCTTCTTCATC	GB AAO50470 Unknown protein { <i>Arabidopsis</i> }.
TC265975	I, AGAAGAAGAAGACTAAGAAGC	UP 080799 (080799) T8F5.5 protein; weakly similar to PIR S64314 probable membrane protein YGR023w—yeast (<i>S. cerevisiae</i>).
TC269146	I, CTTCTTCGTTCTTTCATCTTCT	UPI O49467 (O49467) Hypothetical protein F24J7.50 (hypothetical protein AT4g19490).
TC275407	RC, TTCTTCTTCGTTCTTCTTCATC	UPI Q8VYC1 (Q8VYC1) Putative Ser/Thr kinase.
TC275528	I, GAGAAGAAGAAGACTAAGAAG	UPI Q9FJR9 (Q9FJR9) Similarity to maturase-related protein, complete.
TC275625	RC, TTCTTCTTCGTTCTTTCATCTTCT	GB AAM51439 Unknown protein { <i>Arabidopsis</i> }; weakly similar to UP 081812(081812) auxilin-like protein, partial (13%).

(qRT-PCR) and compared with their expression in non-transgenic plants. The Arabidopsis elongation factor-1 α (*EF-1 α*) gene was used as the endogenous control to normalize the relative transcripts in the reactions. The results showed about 90% down-regulation of target gene *BT11* in one RNAi transgenic line (Fig. 3). Down-regulation of some of the selected potential off-targets, TC262843 and TC269146, and a family member, TC251703 (*BT12*), was detected in the same RNAi transgenic line. They share at least 23 contiguous nucleotide identity with the *BT11* coding region (Table II). About 75% and 50% down-regulation occurred to TC262843 and TC251703, respectively (Fig. 3). Surprisingly, expression of TC269146 is about 95% down-regulated, a little more than that of the target gene. TC262843 and TC269146 share the same 23-nt identity with the *BT11* gene. An oligo DNA with this 23-nt sequence was synthesized and labeled for detection of the potential siRNA involved, but it was not detectable. Nevertheless, a low amount of siRNA could be detected when a 200-bp DNA probe of *BT11* containing this 23-nt sequence was used for hybridization (data not shown). No significant difference was seen in the expression of other analyzed genes with at least 21-nt identity or reverse complementarity to *BT11*, whereas very low (unquantifiable) amounts of transcripts were detected for TC255665 in all the tested leaf tissues (data not shown). Also, no significant differences were seen in the expression of the selected 22 genes that share almost 21- or 22-nt continuous identity to the *BT11* coding region, but with one mismatch (Supplemental Fig. S1). Similar results were seen for all the above-mentioned experiments from two other independent *BT11* RNAi transgenic lines tested (data not shown).

To rule out the possibility that the lowered expression of some of these selected genes might be due to the loss of function of the *BT11* protein instead of off-target silencing, we analyzed the expression levels of all the above investigated genes in an Arabidopsis *bti1-2* null mutant line (T-DNA knockout line; Salk-032220). As expected, the expression of *BT11* was

below the detectable levels in the mutant plants, but the expression levels of all the other genes tested were similar to those in nontransgenic Arabidopsis plants (Fig. 3). Expression of two additional housekeeping genes, *eIF4* and *actin II*, was also investigated for the confirmation of equal RNA amounts used for real-time qRT-PCR. Expression levels were similar among the samples from the three groups, further confirming the equal loading of total RNA in the reactions (Fig. 3). Thus, it is off-target silencing caused by *BT11*-derived dsRNA that down-regulated the expression of three of the five selected potential off-targets in *BT11* RNAi transgenic plants. In addition, there is a striking phenotypic difference between *BT11* RNAi transgenic lines and the *bti1-2* mutant line. Plants from four independent *BT11* RNAi transgenic lines flowered earlier than the wild-type and *bti1-2* mutant plants (Fig. 4) and this is probably due to off-target gene silencing in the RNAi plants.

Expression of Some Potential Off-Targets Was Down-Regulated in N. benthamiana Plants Infiltrated with VIGS Constructs

Computational analysis showed that more than one-half of the EST contigs (TCs) in the *N. benthamiana* database have multiple potential off-targets during PTGS (Table I). We randomly chose two genes for experimental analyses. They were TC381 and TC1146 encoding the U2 small nuclear ribonucleoprotein A (snRNAP A) and pyruvate decarboxylase, respectively. Partial fragments from both genes were separately cloned into the tobacco rattle virus (TRV) RNA2 vector (Liu et al., 2002) for VIGS analysis (their sequences can be found in Supplemental Fig. S2). The search with siRNA Scan using both fragments as trigger sequences revealed several potential off-targets (Table III). Clone TC381 shares contiguous identity of at least 21-nt identity to the following unique sequence entries in the current *N. benthamiana* database: CK286172, CK288691, CK289650, TC10748, and TC7796. These genes encode proteins that are unrelated to

Figure 3. Comparison of the expression levels of *BT11* and some of its potential off-targets in wild-type, *bti1-2* mutant, and *BT11* RNAi transgenic Arabidopsis plants. Initiation factor-4 (*eIF4*) and *actin II* genes were used as controls to show equal loading. Error bars are the sds of three biological replicates of quantitative RT-PCR. *, Expression level of the gene in the *BT11* RNAi transgenic line is significantly lower than that in the wild-type or *bti1-2* mutant plants with 98% confidence by *t* test.






Figure 4. Comparison of phenotypes exhibited by *BT11* RNAi transgenic lines, *bti1-2* mutant plants, and wild-type nontransgenic plants. All plants were grown in short-day conditions for 5 weeks and moved to a growth chamber in long-day conditions for 1 week. A, Plants from four independent *BT11* RNAi transgenic lines. B, *bti1-2* mutant plants. C, Wild-type plants.

snRNAP A. Three of these potential off-targets, CK286172, CK288691, and TC10748, were selected for expression analysis by real-time qRT-PCR in TC381 VIGS-silenced plants. Expression levels of these genes, as well as that of the target gene TC381, were compared with their expression in the plants infiltrated with TRV1 (RNA1) + TRV2::00 (RNA2 empty vector). The *EF-1 α* gene was used as the endogenous reaction control to normalize the relative quantity and expression of the β -*tubulin* gene was investigated to confirm the equal loading of total RNA. The results showed a greater silencing of TC10748 than the target gene TC381 and very little silencing (not statistically significant) for CK286172 and CK288691 genes (Fig. 5A). The analysis was repeated twice with two different groups of plants and similar results were observed.

A similar scenario was found in TRV2::TC1146-infiltrated plants. Five unique EST entries share at least 21 nt of contiguous reverse complementary sequences to the cloned TC1146 fragment (Table III). They are CK282591 encoding a protein similar to the VPI/ABI3 family regulatory protein, CK292351 encoding a Ser carboxypeptidase II-2 precursor, and CK287535, CK296810, and TC8666 encoding three different members of the auxin efflux protein family. The expression levels of these genes and TC1146 in silenced plants were compared between the silenced plants and TRV-RNA1 + TRV2:00-infiltrated plants. The expression of target gene TC1146 was decreased to about 50% in TC1146-silenced plants, whereas among the five analyzed potential off-targets, CK287535, CK296810, and TC8666 were down-regulated to a greater extent than TC1146 (Fig. 5B). These three genes all belong to the auxin efflux protein family and have the same 22-nt reverse complementary sequences to TC1146. The expression level of CK282591 was slightly reduced, whereas there was no significant change in the expression of CK292351 in TC1146-silenced plants. These experiments were repeated twice with two individual silenced plants and the results were similar. In conclusion, some of the potential off-target genes were silenced to different degrees when VIGS was used to silence target genes of

interest in *N. benthamiana*. Because we do not have null mutations of these target genes in *N. benthamiana*, we cannot rule out the possibility that the reduced transcripts of the off-target genes result from the reduction in the amount of target protein.

DISCUSSION

The use of RNA silencing/interference for suppressing gene expression has become a powerful and promising approach in gene function exploration and disease treatment in both plants and animals. Its successful application relies on specific and efficient silencing of particular genes or gene families. Exquisite specificity of RNAi through siRNA in animal cells has been supported by several studies (Tuschl et al., 1999; Elbashir et al., 2001; Chi et al., 2003; Semizarov et al., 2003). However, some contradictory reports indicated that siRNA used for RNA silencing can cause off-target suppression at both transcriptional and translational levels in animals. For example, siRNA causes silencing of unintended genes that lack complete sequence identity and sometimes induces nonspecific interferon responses in animals (Holen et al., 2002; Amarzguoui et al., 2003; Bridge et al., 2003; Saxena et al., 2003; Sledz et al., 2003; Jackson and Linsley, 2004; Scacheri et al., 2004). Recently, a computational study using the genome and transcriptome sequence data of *Homo sapiens*, *Caenorhabditis elegans*, and *Schizosaccharomyces pombe* suggested that the risk of transcriptional off-target silencing by siRNA is considerable in all of these organisms (Qiu et al., 2005). However, to date there is no experimental evidence in plants to show that off-target silencing of unrelated genes can occur as a result of nucleotide sequence similarity with siRNA.

Direct introduction of siRNA into plant protoplasts for PTGS was reported, but it is difficult to apply in intact plant tissues (Vanitharani et al., 2003). The functions of various plant genes revealed by PTGS have been studied mostly through VIGS or by generating stable transgenic lines that express antisense RNA or dsRNA that is identical or complementary to the partial or full-length sequences of target genes. Cleavage of the expressed dsRNAs by Dicer produces many siRNAs. Because siRNAs of 21 to 26 nt have been reported in plants (Hamilton et al., 2002; Llave et al., 2002; Tang et al., 2003; Qi et al., 2005), theoretically, 21-nt identity or reverse complementarity between a trigger sequence and a target could be the minimal requirement for successful RNAi. Our computational analyses indicate a high risk of off-target gene silencing among different plant species (Table I). Results from our experimental analyses with both an RNAi transgenic line and VIGS further confirmed the silencing of some potential off-targets that share at least 22 nt of direct identity or reverse complementary identity to the trigger sequences (Figs. 3 and 5). We were not able to detect the particular siRNAs that have

Table III. Result from the siRNA Scan search for potential off-targets with the cloned sequences of *N. benthamiana* TC381 and TC1146 (Supplemental Fig. S1) as trigger sequences

TC numbers in bold are the ones used for quantitative RT-PCR.

Target ID	Annotation of the Target	Potential Off-Target	Identical (I) or Reverse Complementary (RC) Sequence	Annotation of the Potential Off-Target
TC381	Similar to TIGR_Osa1 9630.m01295 U2 snRNAP protein, Arabidopsis, partial 79%	CK286172	RC, GGGTGTATTCTGGCCCCGGGCC TGRC, TATCTTGATAGTAGTTATTAGTATAGT	Similar to UP O93419 (O93419) collagen XVIII precursor, partial (1%).
		CK288691	RC, TATCTTGATAGTAGTTATTAG-TATAGT	Annotation not available.
		CK289650	I, TTCGCGGTCCCCGGGCTTCGTG	Similar to TIGR_Osa1 9631.m00421 CYFIP2, partial (9%).
		TC10748	I, TATCTTGATAGTAGTTATTA GTATA	Similar to TIGR_Ath1 At5g13210.1 68418.m01516 expressed protein, partial (17%).
		TC7796	I, TATCTTGATAGTAGTTATTA TGTATAG	Weakly similar to UP Q75NB3 (Q75NB3) Cys proteinase, partial (34%).
TC1146	Similar to UP Q8H9C6 (Q8H9C6) pyruvate decarboxylase (fragment), partial (33%)	CK282591	RC, ATCGCTTTGCGAACCCGACTAG	Similar to UP Q6Z3U3 (Q6Z3U3) VP1/ABL3 family regulatory protein-like, partial (5%).
		CK287535	RC, ATCGCTTTGCGAACCCGACTAG	Similar to TIGR_Ath1 At5g65980.1 68418.m08307 auxin efflux carrier family protein contains auxin efflux carrier domain, Pfam:PF03547, partial (14%).
		CK292351	RC, TTTGTTCTCGGGCCTTACCAG	Similar to UP CP22_HORVU (P55748) Ser carboxypeptidase II-2 precursor (CP-MII.2) (fragment), partial (30%).
		CK296810	RC, ATCGCTTTGCGAACCCGACTAG	Similar to TIGR_Ath1 At5g65980.1 68418.m08307 auxin efflux carrier family protein contains auxin efflux carrier domain, Pfam:PF03547, partial (11%).
		TC8666	RC, ATCGCTTTGCGAACCCGACTAG	Similar to TIGR_Ath1 At2g17500.1 68415.m02022 auxin efflux carrier family protein contains auxin efflux carrier domain, Pfam:PF03547, partial (41%).

caused off-target silencing through normal northern-blot analysis. This may be due to low abundance or rapid degradation of one particular siRNA molecule. Nevertheless, the obvious down-regulation of some predicted off-targets that only occurred in *BTI* RNAi transgenic lines, but not in *bt1* mutant plants, is compelling. In other studies, a stretch ≥ 23 nt of perfect sequence identity was found necessary to silence a green fluorescent protein transgene (Thomas et al., 2001) and heterologous silencing occurred when at least 23- or 24-nt identity existed between the RNAi trigger sequence and the intended gene (Ekengren et al., 2003; Liu et al., 2004). In our analysis, 22-nt sequence identity was sufficient to cause off-target silencing. None of the investigated genes with 21- or 22-nt identity, but containing one mismatch to the trigger sequence, was silenced. Thus, at least 22-nt identity may be required for off-target silencing to

occur. However, we only analyzed 19 of 70 predicted off-target candidates with 22-nt identity, but containing one mismatch identified by siRNA Scan. Hopefully, a systematic experimental analysis to determine the minimal sequence identity for PTGS between trigger and target sequences will help to set a baseline for potential off-target searching with a particular trigger sequence in the future.

Statistically, longer siRNAs should be less likely to silence unintended genes by chance. Therefore, predicted off-target risk might be overestimated for siRNAs longer than 21 nt. However, because on average each predicted off-target for all plant species analyzed here has multiple off-target regions and each average off-target region is longer than the longest siRNAs reported in plants (Table I), the overall trend of off-target risk predicted from this study should be valid for longer siRNAs such as 22 to 26 nt. More importantly,




Figure 5. Off-target gene silencing in *N. benthamiana* by VIGS. A, Comparison of the expression levels of TC381, CK286172, TC10748, CK288691, and β -tubulin in TC381 silenced and nonsilenced *N. benthamiana*. B, Comparison of the expression levels of TC1146, CK282591, CK287535, CK292351, CK296810, TC8666, and β -tubulin in the *N. benthamiana* plants silenced with the TRV2::00 or TRV2::TC1146 construct. Error bars are the SDs among three technical replicates of real-time qRT-PCR. *, Expression level of the gene in the TRV2::TC381- or TRV2::TC1146-infiltrated plant is significantly lower than that in the TRV2::00-infiltrated plant with 98% confidence by *t* test. Both of the experiments in A and B were repeated twice with two individual silenced plants and the results were similar.

because siRNAs of various lengths simultaneously coexist within the plant cell (Hamilton et al., 2002), off-target risk should be estimated with the shortest siRNAs. In addition, mismatches in siRNA target recognition are tolerable and these siRNAs can cause RNA degradation and translational repression in animals (Saxena et al., 2003). It is not known whether this applies to plants. In the case of *BT11* RNAi transgenic lines, transcript accumulation of all the investigated potential off-targets with contiguous ≥ 22 -nt identity or with 21- or 22-nt identity, but containing one mismatch to the expressed dsRNA, was not affected (Supplemental Fig. S1). However, this is just one particular case. If mismatches of siRNAs can be tolerated in plant PTGS, our computational analysis results may represent the lower limit of off-target effects as only 21-nt identical regions were counted as potential off-target sites in this study. Nevertheless, our siRNA Scan tool includes the option of searching potential off-targets with complete sequence identity or reverse complementary identity of 18 to 29 nt to the trigger sequence, as well as allowing a few mismatches to the potential siRNAs.

Although our computational analysis showed a high risk of off-target gene silencing during PTGS in plants, the efficiency of off-target gene silencing should be the main factor that affects the functional analysis of a particular target gene. Results from our experimental analyses showed a varied reduction of expression levels of the potential off-targets in both

Arabidopsis and *N. benthamiana*. The reduction ranges from none to a greater reduction than that of the target gene expression. The underlying mechanisms for this variation in expression levels of off-targets are not yet clear. Gene silencing efficiency is correlated with siRNA sequence-specific features (Khvorova et al., 2003; Schwarz et al., 2003; Amarzquioui and Prydz, 2004; Reynolds et al., 2004; Ui-Tei et al., 2004), location of the complementary sequence of siRNA in the target (Birmingham et al., 2006), and target accessibility (Luo and Chang, 2004; Pancoska et al., 2004; Brown et al., 2005). At present, target accessibility cannot be reliably predicted because sequences may be partial for most datasets, which are assembled from ESTs and other transcripts. Besides, the *in vivo* mRNA secondary structure is unknown for the datasets in our analysis. Therefore, we only used siRNA sequence properties to assess gene silencing efficiency. Specifically, siRNA efficacy was predicted according to the rules by Ui-Tei et al. (2004). A recent comparison study indicates that, among currently available siRNA efficacy predictors, Ui-Tei's rules are stable and high performance (Saetrom and Snove, 2004). Although Ui-Tei's rules have not been experimentally tested in plants, they agree with the requirement for thermodynamic features of an efficient siRNA duplex, which favors siRNA unwinding by helicases. However, the siRNAs predicted to cause off-target gene silencing had different efficiencies in silencing the target and off-target genes in our experimental analysis. This could be due

to the different accessibility of the mRNAs to the siRNA. Earlier, it was shown that siRNAs against different regions of genes display marked variation in their potency in mediating mRNA degradation (Thomas et al., 2001). In addition, strong RNAi effects, as assessed by phenotypic analysis, were found to correlate with high expression levels of the targeted genes or higher expression of RNAi trigger sequences in *C. elegans* and plants (Chuang and Meyerowitz, 2000; Cutter et al., 2003; Hu et al., 2004; Kerschen et al., 2004). The structure of siRNA may also have an effect on its efficiency to induce gene silencing. These complex factors together may result in different silencing efficiencies among target and off-target genes.

Short reverse complementary sequences (over 21 nt) were found in some completely unrelated genes in our computational analyses. In animals, siRNAs can simultaneously induce sequence-specific degradation of two endogenous mammalian transcripts oriented in opposite directions (Hu et al., 2004). Our experimental results indicate that either strand of the siRNA duplex may silence the genes that contain identical sequences. It is not known whether these endogenous short reverse complementary sequences in different genes have any biological function.

CONCLUSION

The understanding of off-target silencing is crucial for accurate interpretation of gene function by PTGS. Our computational analyses with the genome and transcriptome sequences from 25 plant species showed a high risk of off-target gene silencing when a full-length sequence of each transcript entry in the datasets was used as an RNAi trigger. This off-target gene silencing risk was confirmed by our experimental analysis with both RNAi transgenic *Arabidopsis* lines and *N. benthamiana* plants infected by VIGS constructs. So far, the contribution of silenced off-targets to the silencing phenotype is not known and needs to be analyzed in exploring target gene function. For example, although the *BT11* RNAi lines had off-target gene silencing, the transformation recalcitrant phenotype of these lines is not in question because it was confirmed by null mutant lines of the target gene (Hwang and Gelvin, 2004). Nevertheless, no approach used to inactivate gene function is free from potential problems, and the reality of off-target silencing does not override the enormous potential of RNAi as a tool for individual or high-throughput studies of gene function. On the other hand, off-target silencing in PTGS provides an advantage in overcoming gene functional redundancy and in its potential to be applied in heterologous gene silencing across species. Hopefully, a further understanding of the molecular mechanisms of RNAi will add more restraining rules for off-target prediction and reveal possible approaches to overcome off-target effects for target specificity. At this point, our siRNA Scan should provide an extremely

useful tool in searching for the potential off-targets of an RNAi trigger of interest and will also help to design more specific RNAi triggers and appropriate controls in experiments.

MATERIALS AND METHODS

Datasets

Arabidopsis (*Arabidopsis thaliana*) and rice (*Oryza sativa* subsp. *japonica*) genome mRNAs were downloaded from TIGR (<http://www.tigr.org>). Gene Indices for 25 plant organisms were downloaded from TIGR. The Gene Ontology (GO) database was provided by the GO Consortium (<http://geneontology.org>). *Arabidopsis* gene family information and GO annotation were downloaded from TAIR (<http://www.arabidopsis.org>). Protein domain data (<http://pfam.wustl.edu>) was also downloaded for off-target function analysis.

Computational Off-Target Estimation

The gene transcript sequence dataset for each organism was searched against itself for contiguous ≥ 21 -nt identical or reverse complementary regions using the BLAST algorithm (Altschul et al., 1997). The BLAST output was loaded into a local data warehouse into which GO and gene family databases were also integrated. Patterns of potential off-targets were obtained by querying the database. Possible distinct siRNAs were enumerated from each identical region between a query and an off-target, and efficient siRNAs were also predicted according to the rules by (Ui-Tei et al. 2004). The rules include a 5' antisense strand, starting with an A or U base, a 5' sense strand, starting with G or C, and the first seven bases from a 5' antisense strand end containing at least three to five A/U bases. The procedures of the above analyses were modified and implemented in an online tool that is publicly accessible (<http://bioinfo2.noble.org/RNAiScan.htm>). The input parameters of the tool for analysis of siRNAs and off-targets can be adjusted by users. Additionally, gene function prediction is provided for off-targets by aligning them to Pfam-A seed sequences using the BLASTx program. The E-value cutoff is set to 10^{-5} . When a domain is identified, a local GO database is searched for its available GO annotations in molecular function, biological process, and cellular component.

Plant Materials and Growth Conditions

Arabidopsis (ecotype Columbia) and *BT11* RNAi transgenic lines (Hwang and Gelvin, 2004) and *BT11* T-DNA insertion mutants (Salk-032220, *bti1-2*, ecotype Columbia) were grown in a growth chamber at 22°C with 1 h daylight. *Nicotiana benthamiana* and the *Agrobacterium tumefaciens* strain GV2260 were used for VIGS analysis. The planting conditions were the same as described previously (Ryu et al., 2004).

Plasmid Construction

The sequences of TC381 and TC1146 were obtained from the TIGR database of *N. benthamiana*. cDNA fragments, including the sequences with >20 -nt continuous identity or complementarity to their potential off-target genes, were amplified by RT-PCR. Primers for the amplification of specific TC381 and TC1146 fragments are shown in Supplemental Table S2. About 5 μ g total RNA from *N. benthamiana* leaves were used for RT at 42°C with NNpoly(dT)₂₀ as the primer for 2 h. The RT product was used for PCR amplification for TC381 and TC1146 fragments, respectively, in a PTC-100 Peltier thermal cycler (M.J. Research). The PCR products were cloned into the pGEM-T-easy vector (Promega). Clones with inserts that are identical to the area in the target genes were amplified by PCR with primers that contain attB recombination sequences adapted to the previous primer pairs. PCR products were purified, sequenced, and cloned into the *pTRV2* VIGS vector using the GATEWAY cloning system with the protocol from the manufacturer (Invitrogen). The recombinant plasmids were named TRV2::TC381 and TRV2::TC1146.

VIGS

Agrobacterium strain GV2260 containing *pTRV1*, TRV2::00, TRV2::TC381, and TRV2::TC1146 were grown in an incubator at 28°C on Luria-Bertani broth with 10 μ g L⁻¹ rifampicin and 50 μ g L⁻¹ kanamycin for 2 d. Inoculum was

prepared with the protocol published previously (Ryu et al., 2004). Leaves of 2-week-old *N. benthamiana* plants (two- to three-leaf stage) were infiltrated with a 1:1 *Agrobacterium* mixture of either TRV1+TRV2::00 or TRV1+TRV2::TC381 or TRV1+TRV2::TC1146 as described earlier (Liu et al., 2002). The infiltrated plants were grown for 14 to 18 d for silencing to occur. Leaf samples were then collected for RT-PCR analysis.

RNA Extraction and Real-Time qRT-PCR Analysis

For *Arabidopsis*, total RNA was isolated from leaf tissues using TRIzol reagent (Invitrogen), followed by RNase-free DNase treatment (Promega). First-strand cDNA was synthesized with the Omniscript RT kit (Qiagen) using oligo(dT)₁₅ according to the manufacturer's instructions. For qRT-PCR, real-time experiments were conducted in an ABI PRISM 7000 sequence detection system (Applied Biosystems), using the intercalation dye SYBR Green I as a fluorescent reporter. Quantification of PCR products was performed via a calibration curve procedure using *EF-1α* as an endogenous control. The ratio of gene-specific expression to the expression level of the designated calibrator was defined as relative expression using a standard curve method described in User Bulletin Number 2 (Applied Biosystems).

For *N. benthamiana*, total RNA from each plant at 15 d postinfiltration was extracted using TRIzol reagent with the protocol provided by the manufacturer (Molecular Research Center). First-strand cDNA was synthesized with 2 μg of total RNA using primer NNpoly(dT)₂₀ as described above. Semiquantitative RT-PCR was performed with the modified program from the one described in "Plasmid Construction" with annealing temperature at 60°C and 20 cycles for TC381, 33 cycles for TC1146. The primer sequences are listed in Supplemental Table S2. Further real-time qRT-PCR analysis for each comparison was done with the total RNA of one TRV2::00-infiltrated and one recombinant clone-infiltrated plant, as described above for *Arabidopsis* samples. The analysis was repeated twice with two different groups of plants for each combination. The primer sequences used for real-time qRT-PCR are listed in Supplemental Table S3.

Data Analysis

Data were subjected to ANOVA using Student's *t* test software of Excel 2003 with 98% confidence.

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Figure S1. Comparison of the expression levels of three genes with 21- or 22-nt identity but containing one mismatch to *BTII* gene sequence in wild-type, *bti1-2* mutant, and *BTII* RNAi transgenic *Arabidopsis* plants.

Supplemental Figure S2. List of the sequences that were cloned for VIGS analysis.

Supplemental Table S1. Gene family members as triggers and their off-targets in the same family in *Arabidopsis*.

Supplemental Table S2. Primers for cDNA cloning and semiquantitative RT-PCR.

Supplemental Table S3. Primers for real-time qRT-PCR assay.

ACKNOWLEDGMENTS

We thank Dr. Stan Gelvin and Dr. Hau-Hsuan Hwang for sending us the *Arabidopsis* seeds of *BTII* RNAi lines and the *bti1-2* T-DNA mutant; Dr. S.P. Dinesh-Kumar for providing GATEWAY-ready TRV-VIGS vectors; Dr. Choong-Min Ryu for help with infiltrations and discussions; and Dr. Alison Blancaflor, Dr. Stan Gelvin, and Dr. Jianzhong Liu for reviewing this manuscript.

Received May 8, 2006; accepted August 7, 2006; published August 18, 2006.

LITERATURE CITED

Altschul SE, Madden TL, Schäffer A, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**: 3389–3402

- Amarzguioui M, Holen T, Babaie E, Prydz H (2003) Tolerance for mutations and chemical modifications in a siRNA. *Nucleic Acids Res* **31**: 589–595
- Amarzguioui M, Prydz H (2004) An algorithm for selection of functional siRNA sequences. *Biochem Biophys Res Commun* **316**: 1050–1058
- Baulcombe D (1999) Fast forward genetics based on virus-induced gene silencing. *Curr Opin Plant Biol* **2**: 109–130
- Baulcombe D (2004) RNA silencing in plants. *Nature* **431**: 356–363
- Birmingham A, Anderson EM, Reynolds A, Ilsley-Tyree D, Leake D, Fedorov Y, Baskerville S, Maksimova E, Robinson K, Karpilow J, et al (2006) 3' UTR seed matches, but not overall identity, are associated with RNAi off-targets. *Nat Methods* **3**: 199–204
- Bridge AJ, Pebernard S, Ducraux A, Nicoulaz A-L, Iggo R (2003) Induction of an interferon response by RNAi vectors in mammalian cells. *Nat Genet* **34**: 263–264
- Brown KM, Chu CY, Rana TM (2005) Target accessibility dictates the potency of human RISC. *Nat Struct Mol Biol* **12**: 469–470
- Burch-Smith TM, Anderson JC, Martin GB, Dinesh-Kumar SP (2004) Applications and advantages of virus-induced gene silencing for gene function studies in plants. *Plant J* **39**: 734–746
- Chi J-T, Chang HY, Wang NN, Chang DS, Dunphy N, Brown PO (2003) Genomewide view of gene silencing by small interfering RNAs. *Proc Natl Acad Sci USA* **100**: 6343–6346
- Chuang C-F, Meyerowitz EM (2000) Specific and heritable genetic interference by double-stranded RNA in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* **97**: 4985–4990
- Cogoni C, Macino G (2000) Post-transcriptional gene silencing across kingdoms. *Curr Opin Genet Dev* **10**: 638–643
- Cutter AD, Payseur BA, Salcedo T, Estes AM, Good JM, Wood E, Hartl T, Maughan H, Strempel J, Wang B, et al (2003) Molecular correlates of genes exhibiting RNAi phenotypes in *Caenorhabditis elegans*. *Genome Res* **13**: 2651–2657
- Ekengren SK, Liu Y, Schiff M, Dinesh-Kumar SP, Martin GB (2003) Two MAPK cascades, NPR1, and TGA transcription factors play a role in Pto-mediated disease resistance in tomato. *Plant J* **36**: 905–917
- Elbashir SM, Garborth J, Lendeckel W, Yalcin A, Weber K, Tuschl T (2001) Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* **411**: 494–498
- Hamilton A, Voinnet O, Chappell L, Baulcombe D (2002) Two classes of short interfering RNA in RNA silencing. *EMBO J* **21**: 4671–4679
- Hannon GJ, Rossi JJ (2004) Unlocking the potential of the human genome with RNA interference. *Nature* **431**: 371–378
- Holen T, Amarzguioui M, Wiiger MT, Babaie E, Prydz H (2002) Positional effects of short interfering RNAs targeting the human coagulation trigger tissue factor. *Nucleic Acids Res* **30**: 1757–1766
- Hu X, Hipolito S, Lynn R, Abraham V, Ramos S, Wong-Staal F (2004) Relative gene-silencing efficiencies of small interfering RNAs targeting sense and antisense transcripts from the same genetic locus. *Nucleic Acids Res* **32**: 4609–4617
- Hwang H-H, Gelvin SB (2004) Plant proteins that interact with VirB2, the *Agrobacterium tumefaciens* pilin protein, mediate plant transformation. *Plant Cell* **16**: 3148–3167
- Jackson AL, Bartz SR, Schelter J, Kobayashi SV, Burchard J, Mao M, Li B, Cavet G, Linsley PS (2003) Expression profiling reveals off-target gene regulation by RNAi. *Nat Biotechnol* **21**: 635–637
- Jackson AL, Linsley PS (2004) Noise amidst the silence: off-target effects of siRNAs? *Trends Genet* **20**: 521–524
- Kamath RS, Fraser AG, Dong Y, Poulin R, Gotta M, Kanapin A, Le Bot N, Moreno S, Sohrmann M, Welchman DP, et al (2003) Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* **421**: 231–237
- Kerschen A, Napoli CA, Jorgensen RA, Muller AE (2004) Effectiveness of RNA interference in transgenic plants. *FEBS Lett* **466**: 223–228
- Khvorova A, Reynolds A, Jayasena SD (2003) Functional siRNAs and miRNAs exhibit strand bias. *Cell* **115**: 209–216
- Kuttenkeuler D, Boutros M (2004) Genome-wide RNAi as a route to gene function in *Drosophila*. *Brief Func Genomics Proteomics* **3**: 168–176
- Liu Y, Schiff M, Dinesh-Kumar SP (2002) Virus-induced gene silencing in tomato. *Plant J* **31**: 777–786
- Liu Y, Schiff M, Dinesh-Kumar SP (2004) Involvement of MEK1 MAPKK, NTF6 MAPK, WRKY/MYB transcription factors, *COI1* and *CTR1* in N-mediated resistance to tobacco mosaic virus. *Plant J* **38**: 800–809

- Llave C, Kasschau KD, Rector MA, Carrington JC (2002) Endogenous and silencing-associated small RNAs in plants. *Plant Cell* **14**: 1605–1619
- Luo KQ, Chang DC (2004) The gene-silencing efficiency of siRNA is strongly dependent on the local structure of mRNA at the targeted region. *Biochem Biophys Res Commun* **318**: 303–310
- McManus MT, Petersen CP, Haines BB, Chen J, Sharp PA (2002) Gene silencing using micro-RNA designed hairpins. *RNA* **8**: 842–850
- Miki D, Shimamoto K (2004) Simple RNAi vectors for stable and transient suppression of gene function in rice. *Plant Cell Physiol* **45**: 490–495
- Pancoska P, Moravek Z, Moll UM (2004) Efficient RNA interference depends on global context of the target sequence: quantitative analysis of silencing efficiency using Eulerian graph representation of siRNA. *Nucleic Acids Res* **32**: 1469–1479
- Qi Y, Denli AM, Hannon GJ (2005) Biochemical specialization within Arabidopsis RNA silencing pathways. *Mol Cell* **19**: 421–428
- Qiu S, Adema CM, Lane T (2005) A computational study of off-target effects of RNA interference. *Nucleic Acids Res* **33**: 1834–1847
- Quackenbush J, Cho J, Lee D, Liang F, Holt I, Karamycheva S, Parvizi B, Pertea G, Sultana R, White J (2001) The TIGR gene indices: analysis of gene transcript sequences in highly sampled eukaryotic species. *Nucleic Acids Res* **29**: 159–164
- Reynolds A, Leake D, Boese Q, Scaringe S, Marshall WS, Khvorova A (2004) Rational siRNA design for RNA interference. *Nat Biotechnol* **22**: 326–330
- Ryu C-M, Anand A, Kang L, Mysore KS (2004) Agrodrench: a novel and effective agroinoculation method for virus-induced gene silencing in roots and diverse Solanaceous species. *Plant J* **40**: 322–331
- Saetrom P, Snove O Jr (2004) A comparison of siRNA efficacy predictors. *Biochem Biophys Res Commun* **321**: 247–253
- Saxena S, Jónsson SO, Dutta Z (2003) Small RNAs with imperfect match to endogenous mRNA repress translation. *J Biol Chem* **278**: 44312–44319
- Scacheri PC, Rozenblatt-Rosen O, Caplen NJ, Wolfsberg TG, Umayam L, Lee JC, Hughes CM, Shanmugam KS, Bhattacharjee A, Meyerson M, et al (2004) Short interfering RNAs can induce unexpected and divergent changes in the levels of untargeted proteins in mammalian cells. *Proc Natl Acad Sci USA* **101**: 1892–1897
- Schwarz DS, Hutvagner G, Du T, Xu Z, Aronin N, Zamore PD (2003) Asymmetry in the assembly of the RNAi enzyme complex. *Cell* **115**: 199–208
- Semizarov D, Frost L, Sarthy A, Kroeger P, Halbert DN, Fesik SW (2003) Specificity of short interfering RNA determined through gene expression signatures. *Proc Natl Acad Sci USA* **100**: 6347–6352
- Shirane D, Sugao K, Namiki S, Tanabe M, Iino M, Hirose K (2004) Enzymatic production of RNAi libraries from cDNAs. *Nat Genet* **36**: 190–196
- Sledz CA, Holko M, de Veer MJ, Silverman RH, Williams BRG (2003) Activation of the interferon system by short-interfering RNAs. *Nat Cell Biol* **5**: 834–839
- Sonnhammer EL, Eddy SR, Birney E, Bateman A, Durbin R (1998) Pfam: multiple sequence alignments and HMM-profiles of protein domains. *Nucleic Acids Res* **26**: 320–322
- Tang G, Reinhart BJ, Bartel DP, Zamore PD (2003) A biochemical framework for RNA silencing in plants. *Genes Dev* **17**: 49–63
- Thomas CL, Jones L, Baulcombe DC, Maule AJ (2001) Size constraints for targeting post-transcriptional gene silencing and for RNA-directed methylation in *Nicotiana benthamiana* using a potato virus X vector. *Plant J* **25**: 417–425
- Tuschl T, Zamore PD, Lehmann R, Bartel DP, Sharp PA (1999) Targeted mRNA degradation by double-stranded RNA in vitro. *Genes Dev* **13**: 3191–3197
- Ui-Tei K, Naito Y, Takahashi F, Haraguchi T, Ohki-Hamazaki H, Juni A, Ueda R, Saigo K (2004) Guidelines for the selection of highly effective siRNA sequences for mammalian and chick RNA interference. *Nucleic Acids Res* **32**: 936–948
- Vanitharani R, Chellappan P, Fauquet CM (2003) Short interfering RNA-mediated interference of gene expression and viral DNA accumulation in cultured plant cells. *Proc Natl Acad Sci USA* **100**: 9632–9636
- Wesley SV, Helliwell CA, Smith NA, Wang M, Rouse DT, Liu Q, Gooding PS, Singh SP, Abbott D, Stoutjesdijk PA, et al (2001) Construct design for efficient, effective and high-throughput gene silencing in plants. *Plant J* **27**: 581–590
- Yokota T, Sakamoto N, Enomoto N, Tanabe Y, Miyagishi M, Maekawa S, Yi L, Kurosaki M, Taira K, Watanabe M, et al (2003) Inhibition of intracellular hepatitis C virus replication by synthetic and vector-derived small interfering RNAs. *EMBO Rep* **4**: 602–608
- Yu JY, DeRuiter SL, Turner DL (2002) RNA interference by expression of short-interfering RNAs and hairpin RNAs in mammalian cells. *Proc Natl Acad Sci USA* **99**: 6047–6052