

Plant Gene Register

Sequence of a *psaC* Gene from the Cyanobacterium *Synechococcus* sp. PCC 6301¹

Patricia L. Herman*, Kartika Adiwilaga², John H. Golbeck, and Donald P. Weeks

Department of Biochemistry, University of Nebraska-Lincoln, Lincoln, Nebraska 68583-0718

The *psaC* gene encodes PsaC, the apoprotein for the terminal iron-sulfur clusters, F_A and F_B, in the PSI reaction center of cyanobacteria, algae, and higher plants. PsaC functions as a membrane-bound oxidoreductase, accepting electrons from the F_x iron-sulfur cluster located on the PsaA/PsaB heterodimer and donating them to the soluble electron transfer proteins Fd and flavodoxin (reviewed in Bryant, 1992). The objective of our work is to clarify the role of PsaC in linear and cyclic electron transfer. The experimental organism is *Synechococcus* sp. PCC 6301, a unicellular, freshwater cyanobacterium that is used extensively for physiological, biochemical, and spectroscopic studies of photosynthesis. The PSI complex of *Synechococcus* sp. PCC 6301 is widely used in structural and functional studies of the F_A and F_B iron-sulfur clusters (Parrett et al., 1990).

We became interested in the *psaC* sequence of *Synechococcus* sp. PCC 6301 because there are subtle differences in the electron paramagnetic resonance spectra of the F_A and F_B iron-sulfur clusters between cyanobacteria and higher plants. This suggests that there may be a correlation with structure: in all organisms studied thus far, the higher-plant PsaC protein differs from the cyanobacterial PsaC protein in two regions: at position 37, higher plants contain a Lys whereas cyanobacteria contain a neutral amino acid (Gly or Ala), and at positions 70 and 71, dicots contain a Trp-His pair, monocots contain a Gly-Pro pair, and cyanobacteria contain a Gly-Ala pair. No sequence for the PsaC protein of *Synechococcus* sp. PCC 6301 is available, even though entire sequences are available for PsaD and PsaE and partial sequences are available for PsaL and PsaF. This information is relevant because it has been reported that two *psaC* genes exist in the cyanobacterium *Synechocystis* sp. PCC 6803, one with a higher-plant sequence (Anderson and McIntosh, 1991) and the other with a cyanobacterial sequence (Steinmüller, 1992). We were interested in determining whether the *psaC* gene of *Synechococcus* sp. PCC 6301 was higher plant-like or cyanobacterial-like and whether the gene was flanked by the mitochondrial *ndh* genes that encode subunits of the NADH dehydrogenase complex (Schantz and Bogorad, 1988).

We report here the isolation and nucleotide sequence of a *psaC* gene from *Synechococcus* sp. PCC 6301 (Table I). Oligonucleotide primers based on conserved sequences in the 5' and 3' coding regions of previously sequenced cyanobacteria *psaC* genes were used to generate a fragment from *Synechococcus* sp. PCC 6301 genomic DNA by PCR amplification. The PCR product was cloned into pBluescript II KS+ (Stratagene), sequenced to confirm its identity, and labeled with digoxigenin by the random-prime method (Genius System, Boehringer). The labeled probe was then used in Southern hybridization experiments (using chemiluminescent detection) to map the position of a number of restriction sites relative to an *AccI* site located 50 bp from the 3' end of the *psaC* coding region. Southern analysis showed that the PCR product hybridized to a single 4.2-kb *XhoI/HindIII* genomic fragment. A size-fractionated genomic library consisting of *XhoI/HindIII* fragments from 3.5 to 5 kb in length was constructed in pBluescript II KS+ and transformed into DH5 α cells. Ninety-seven bacterial colonies were screened with the digoxigenin-labeled PCR product, and two putative positive colonies were identified. Restriction enzyme analysis confirmed that both colonies contained clones with the expected genomic fragment. When several restriction enzyme sites in the 4.2-kb clone were mapped, the putative *psaC* gene was localized to a 327-bp *EagI/BamHI* fragment. This fragment was subcloned into pBluescript II KS+, and both strands were sequenced. To sequence the regions flanking the *psaC* gene, a set of unidirectional nested deletions was generated for each strand of the 4.2-kb genomic clone using the Erase-a-Base system (Promega).

The nucleotide sequence of the 246-bp ORF that encodes the *psaC* gene of *Synechococcus* sp. PCC 6301 is highly homologous to previously reported *psaC* genes from six other cyanobacteria. A sequence comparison using the FastA program of the Genetics Computer Group (1991) showed the following results: 82.1% identity to *Synechococcus* sp. PCC 7002 (Rhiel et al., 1992), 81.2% identity to *Synechococcus vulcanus* (Shimizu et al., 1990), 80.8% identity to a *Synechococcus* species (GenBank, X63767), 80.5% identity to an *Anabaena* species (GenBank, X57153), 80.1% identity to *Anabaena* sp. PCC 7120 (Mulligan and Jackman, 1992), and 80.1% identity to *Synechosystis* sp. PCC 6803 (Steinmüller, 1992). The identity of the *psaC* coding sequence from *Synechococcus* sp. PCC 6301 to previously sequenced *psaC* genes

¹ This research was supported by a grant from the National Science Foundation (MCB-9205756) to J.H.G. This is Journal Series No. 10540 of the Agricultural Research Division, University of Nebraska.

² Present address: Institute of Molecular and Cell Biology, National University of Singapore, Singapore 0511.

* Corresponding author; fax 1-402-472-7842.

Abbreviation: ORF, open reading frame.

Table I. Characteristics of a *psaC* gene from *Synechococcus* sp. PCC 6301

Organism:	<i>Synechococcus</i> sp. PCC 6301.
Function of Gene:	Encodes PsaC, a protein that binds the iron-sulfur clusters that are the terminal electron acceptors in PSI.
Type of Clone:	Genomic.
Techniques:	A genomic fragment generated by PCR amplification was used as a probe to isolate a 4.2-kb clone from a size-fractionated genomic library of <i>XhoI/HindIII</i> fragments. Restriction fragment subclones and nested deletion subclones were generated. Both strands of the <i>psaC</i> gene and the flanking regions were sequenced on a Li-Cor 4000 automated DNA sequencer (Brumbaugh et al., 1988) using a double-stranded method that employs dideoxy chain termination reactions and Sequenase (United States Biochemical).
Method of Identification:	Sequence comparison to <i>psaC</i> genes isolated from other photosynthetic organisms.
Features of Gene Structure:	The <i>psaC</i> gene is contained on an <i>EagI/BamHI</i> fragment with a length of 327 bp. The ATG translational start codon begins at position 59 and is preceded 11 bp upstream by a putative Shine-Dalgarno ribosome binding site (5'-GGAG-3'). The coding sequence of the <i>psaC</i> gene is 246 nucleotides in length (including the stop codon that begins at position 302) and has a (C + G) content of 57.3%.
Features of the Deduced Amino Acid Sequence:	The predicted sequence of the PsaC polypeptide contains 81 codons and has 98.8% identity to the deduced sequence of the PsaC protein from <i>Synechococcus</i> sp. PCC 7002 (Rhiel et al., 1992). The only difference between these two sequences is at position 41, where a Ser residue in the <i>Synechococcus</i> sp. PCC 7002 sequence is replaced by an Ala residue in the <i>Synechococcus</i> sp. PCC 6301 sequence. The Ala residue at position 41 is unique because all of the PsaC proteins that have been sequenced have a Ser residue at this position.
Antibodies:	The highly homologous PsaC protein from <i>Synechococcus</i> sp. PCC 7002 has been overexpressed in <i>Escherichia coli</i> and antibodies have been generated by the University of Nebraska Antibody Core Facility.

from 12 other photosynthetic organisms is approximately 10% lower, ranging from 69.1% for the liverwort *Marchantia polymorpha* to 73.6% for maize. Interestingly, there is only 72% identity to another *psaC* sequence from the cyanobacterium *Synechosystis* sp. PCC 6803 (Anderson and McIntosh, 1991). This result supports the conclusion of Steinmüller (1992), who suggested that the *psaC* gene isolated by Anderson and McIntosh (1991) was the result of a rare transformation event by chloroplast DNA from another plant species.

The predicted sequence of the PsaC polypeptide encoded by the *psaC* gene from *Synechococcus* sp. PCC 6301 consists of 81 codons and has a high degree of sequence similarity to the previously reported PsaC proteins from six other cyanobacteria. The Gly residue at position 37 and the Gly-Ala pair at positions 70 and 71 are indicative of a cyanobacterial PsaC protein. A sequence comparison using the TFASTA program of the Genetics Computer Group (1991) showed the following results: 98.8% identity to *Synechococcus* sp. PCC 7002 (Rhiel et al., 1992), 97.5% identity to *Synechocystis* sp. PCC 6803 (Steinmüller, 1992), 95.1% identity to *S. vulcanus* (Shimizu et al., 1990) and another *Synechococcus* species (GenBank, X63767), and 93.8% identity to *Anabaena* sp. PCC 7120 (Mulligan and Jackman, 1992) and to another *Anabaena* species (GenBank, X57153). As expected, the identity of the deduced PsaC sequence from *Synechococcus* sp. PCC 6301 to

the PsaC proteins from 13 other photosynthetic organisms is lower, ranging from 85.2% for the liverwort *M. polymorpha* to 88.9% for tobacco. The only difference between the predicted PsaC sequences from *Synechococcus* sp. PCC 7002 and *Synechococcus* sp. PCC 6301 is at position 41, where an Ala residue in the *Synechococcus* sp. PCC 6301 sequence replaces a Ser residue in the *Synechococcus* sp. PCC 7002 sequence. The substitution of an Ala for a Ser at position 41 is unique because this Ser residue is conserved in the 19 PsaC proteins whose genes have previously been isolated and sequenced.

In the chloroplast genome of a number of plants, the *psaC* gene is located between sequences with homology to two mitochondrial *ndh* genes that encode subunits of the NADH dehydrogenase complex (Schantz and Bogorad, 1988). The first homolog, *ndhE*, is upstream from the *psaC* gene. The second homolog, *ndhD*, is downstream and actually co-transcribed with the *psaC* gene from maize (Schantz and Bogorad, 1988). Anderson and McIntosh (1991) reported that this gene arrangement is conserved in the cyanobacterium *Synechocystis* sp. PCC 6803. The Compare and DotPlot programs of the Genetics Computer Group (1991) were used to compare the nucleotide sequence of the region surrounding the *psaC* gene of *Synechococcus* sp. PCC 6301 to the published nucleotide sequence of the region flanking 19 *psaC* genes from both cyanobacteria and other photosynthetic organisms. This

analysis revealed no significant similarity between the sequence flanking the *Synechococcus* sp. PCC 6301 *psaC* gene and the sequence surrounding any of the previously reported *psaC* genes. In particular, the *psaC* gene of *Synechococcus* sp. PCC 6301 is not flanked by sequences with homology to the *ndhE* or *ndhD* genes. This result agrees with that of Rhiel et al. (1992), who found no evidence of *ndhE* or *ndhD* sequences immediately adjacent to the coding region of the *psaC* gene from the cyanobacterium *Synechococcus* sp. PCC 7002. Ellersiek and Steinmüller (1992) reported a sequence with homology to *ndhD* located 368 bp upstream from another *psaC* gene isolated from the cyanobacterium *Synechocystis* sp. PCC 6803. However, this *ndhD* gene is transcribed on the opposite DNA strand. It can be concluded that most of the available evidence suggests that cyanobacteria do not have the same gene arrangement as plant chloroplasts in the region of the genome that contains the *psaC* gene.

A sequence analysis using the FastA program of the Genetics Computer Group (1991) detected five ORFs not previously reported for cyanobacteria in the 4.2-kb clone containing the *psaC* gene of *Synechococcus* sp. PCC 6301 (P.L. Herman and D.P. Weeks, unpublished data). There is a cluster of four consecutive ORFs starting 459 bp upstream of the *psaC* gene encoded on the opposite strand that is highly homologous to previously sequenced genes that function in polyketide biosynthesis. In addition, there is an ORF that begins 111 bp downstream from the *psaC* coding region on the same strand. This ORF has a high degree of sequence similarity to *nodM*, a gene found in the nitrogen-fixing *Rhizobium* bacterium that is involved in the production of a host-specific nodulation signal.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the DNA Sequencing Facility supported by the Center for Biotechnology at the University of Nebraska-Lincoln for their help in sequencing the clones reported in this study. The authors also thank Drs. Sylvia Darr and Robert Spreitzer (Department of Biochemistry, University of Nebraska-Lincoln) for their critical review of the manuscript and Dr. Don Bryant

(Pennsylvania State University) for advice on designing the PCR primers.

Received November 9, 1993; accepted November 22, 1993.

Copyright Clearance Center: 0032-0889/94/104/1459/03.

The GenBank accession number for the sequence reported in this article is U01536.

LITERATURE CITED

- Anderson SL, McIntosh L** (1991) Partial conservation of the 5' *ndhE-psaC-ndhD* 3' gene arrangement of chloroplasts in the cyanobacterium *Synechocystis* sp. PCC 6803: implications for NDH-D function in cyanobacteria and chloroplasts. *Plant Mol Biol* **16**: 487-499
- Brumbaugh JA, Middendorf LR, Grone DL, Ruth JL** (1988) Continuous, online sequencing using oligonucleotide primers with multiple fluorophores. *Proc Natl Acad Sci USA* **85**: 5610-5614
- Bryant DA** (1992) Molecular biology of photosystem I. In J Barber, ed, *The Photosystems: Structure, Function and Molecular Biology*. Elsevier, Amsterdam, pp 501-549
- Ellersiek U, Steinmüller K** (1992) Cloning and transcription analysis of the *ndh(A-I-G-E)* gene cluster and the *ndhD* gene of the cyanobacterium *Synechocystis* sp. PCC6803. *Plant Mol Biol* **20**: 1097-1110
- Genetics Computer Group** (1991) Program Manual for the GCG Package, Version 7, 1991
- Mulligan ME, Jackman DM** (1992) Nucleotide sequence and expression of the gene for the 9 kDa FA/FB component of photosystem I from the cyanobacterium *Anabaena* sp. strain PCC 7120. *Plant Mol Biol* **18**: 803-808
- Parrett KG, Mehari T, Golbeck JH** (1990) Resolution and reconstitution of the cyanobacterial photosystem I complex. *Biochim Biophys Acta* **1015**: 341-352
- Rhiel E, Stirewalt VL, Gasparich GE, Bryant DA** (1992) The *psaC* genes of *Synechococcus* sp. PCC7002 and *Cyanophora paradoxa*: cloning and sequence analysis. *Gene* **112**: 123-128
- Schantz R, Bogorad L** (1988) Maize chloroplast genes *ndhD*, *ndhE*, and *psaC*. Sequences, transcripts and transcript pools. *Plant Mol Biol* **11**: 239-247
- Shimizu T, Hiyama T, Ikeuchi M, Koike H, Inoue Y** (1990) Nucleotide sequence of the *psaC* gene of the cyanobacterium *Synechococcus vulcanus*. *Nucleic Acids Res* **18**: 3644
- Steinmüller K** (1992) Identification of a second *psaC* gene in the cyanobacterium *Synechocystis* sp. PCC6803. *Plant Mol Biol* **20**: 997-1001