

# Germination and Dormancy of Abscisic Acid- and Gibberellin-Deficient Mutant Tomato (*Lycopersicon esculentum*) Seeds<sup>1</sup>

## Sensitivity of Germination to Abscisic Acid, Gibberellin, and Water Potential

Bing-Rui Ni and Kent J. Bradford\*

Department of Vegetable Crops, University of California, Davis, California 95616–8631

Germination responses of wild-type (*MM*), abscisic acid (ABA)-deficient (*sit<sup>v</sup>*), and gibberellin (GA)-deficient (*gib-1*) mutant tomato (*Lycopersicon esculentum* Mill. cv MoneyMaker) seeds to ABA, GA<sub>4+7</sub>, reduced water potential ( $\psi$ ), and their combinations were analyzed using a population-based threshold model (B.R. Ni and K.J. Bradford [1992] Plant Physiol 98: 1057–1068). Among the three genotypes, *sit<sup>v</sup>* seeds germinated rapidly and completely in water, *MM* seeds germinated more slowly and were partially dormant, and *gib-1* seeds did not germinate without exogenous GA<sub>4+7</sub>. Times to germination were inversely proportional to the differences between the external osmoticum, ABA, or GA<sub>4+7</sub> concentrations and the corresponding threshold levels that would either prevent ( $\psi_b$ ,  $\log[ABA_b]$ ) or promote ( $\log[GA_b]$ ) germination. The sensitivity of germination to ABA, GA<sub>4+7</sub>, and  $\psi$  varied widely among individual seeds in the population, resulting in a distribution of germination times. The rapid germination rate of *sit<sup>v</sup>* seeds was attributable to their low mean  $\psi_b$  (–1.17 MPa). Postharvest dormancy in *MM* seeds was due to a high mean  $\psi_b$  (–0.35 MPa) and a distribution of  $\psi_b$  among seeds such that some seeds were unable to germinate even on water. GA<sub>4+7</sub> (100  $\mu$ M) stimulated germination of *MM* and *gib-1* seeds by lowering the mean  $\psi_b$  to –0.75 MPa, whereas ABA inhibited germination of *MM* and *sit<sup>v</sup>* seeds by increasing the mean  $\psi_b$ . The changes in  $\psi_b$  were not due to changes in embryo osmotic potential. Rather, hormonal effects on endosperm weakening opposite the radicle tip apparently determine the threshold  $\psi$  for germination. The analysis demonstrates that ABA- and GA-dependent changes in seed dormancy and germination rates, whether due to endogenous or exogenous growth regulators, are based primarily upon corresponding shifts in the  $\psi$  thresholds for radicle emergence. The  $\psi$  thresholds, in turn, determine both the rate and final extent of germination within the seed population.

Welbaum et al., 1990; Ni and Bradford, 1992). In addition, high ABA concentrations caused an increase in the sensitivity of germination to reduced  $\psi$  (Ni and Bradford, 1992). GAs are also involved in the regulation of seed development and germination but with opposite effects to those of ABA. In general, ABA delays or prevents seed germination and determines the depth of dormancy during development, whereas GA breaks dormancy and promotes germination upon imbibition by the mature seeds (Karssen et al., 1983, 1987; Koornneef et al., 1989; Groot and Karssen, 1992). The quantitative amounts of ABA during development result in different depths of dormancy, which in turn require different amounts of GA to stimulate germination. Hence, ABA-deficient mutants produce nondormant seeds that germinate rapidly, whereas GA-deficient mutant seeds will not germinate without additional GA (Karssen et al., 1983, 1987; Groot and Karssen, 1992).

The responses of germination to ABA, osmoticum, and their combination have been quantitatively characterized using a population-based threshold model (Ni and Bradford, 1992). Seed germination rates and percentages can be predicted with reasonable accuracy at any given ABA concentration or  $\psi$  on the basis of the derived threshold values for each factor and their respective time constants. The basic assumption of the model is that the rate of approach of a seed toward germination is proportional to the difference between the factor level and the seed's threshold value for the regulatory factor. The response of germination to reduced  $\psi$ , for example, can be expressed as

$$\theta_H = [\psi - \psi_b(g)]t_g \quad (1)$$

or

$$GR_g = 1/t_g = [\psi - \psi_b(g)]/\theta_H \quad (2)$$

where  $\theta_H$  is the "hydrotime constant,"  $\psi$  is the water potential of the imbibition medium,  $\psi_b(g)$  is the base (threshold or minimum)  $\psi$  allowing germination of percentage  $g$ ,  $t_g$  is the time required for germination of percentage  $g$ , and  $GR_g$  is the germination rate of the seeds of fraction or percentage  $g$  (Gummerson, 1986; Bradford, 1990). The normal distribution

Seed germination is sensitive to both endogenous plant growth regulators and environmental factors. Both ABA and reduced  $\psi$  (Table I) inhibit seed germination, and their interactions have been analyzed for several species (Schopfer and Plachy, 1985; Kermodé, 1990; Welbaum et al., 1990; Ni and Bradford, 1992). The inhibitory effect of exogenous ABA on germination is empirically similar to, and quantitatively additive with, that of reduced  $\psi$  (Schopfer and Plachy, 1985;

<sup>1</sup> Supported by National Science Foundation grants DCB-8817758 and IBN-9204782 to K.J.B.

\* Corresponding author; fax 1-916-752-4554.

The abbreviations used in this article are listed in Table I.

**Table I.** Abbreviations used in this paper

$ABA_0$	the highest ABA concentration having no effect on germination
$ABA_{endo}$	endogenous ABA
$ABA_{exo}$	exogenous ABA
$ABA_b(g)$	base ABA concentration inhibiting germination of percentage $g$
$ABA_b(50)$	mean base ABA concentration reducing germination to 50%
$GR_g$	germination rate (inverse of time to germination) for percentage $g$
$GA_0$	GA concentration stimulating maximum germination rates
$GA_{endo}$	endogenous GA
$GA_{exo}$	exogenous GA
$GA_b(g)$	base GA concentration stimulating germination of percentage $g$
$GA_b(50)$	mean base GA concentration stimulating 50% germination
$gib-1$	GA-deficient mutant tomato
$MM$	wild type cv MoneyMaker tomato
$\psi$	water potential
$\psi_b(g)$	base water potential inhibiting germination of percentage $g$
$\psi_b(50)$	mean base $\psi$ reducing germination to 50%
$\psi_\pi$	osmotic potential
RWC	relative water content
$\sigma_{\psi_b}$	standard deviation of $\psi_b(g)$
$\sigma_{ABA_b}$	standard deviation of $ABA_b(g)$
$\sigma_{GA_b}$	standard deviation of $GA_b(g)$
$sit^w$	ABA-deficient mutant <i>sitiens</i> tomato
$t_g$	time to radicle emergence of percentage $g$
$t_g(0)$	time to radicle emergence of percentage $g$ in water
$t_g(ABA)$	time to radicle emergence of percentage $g$ in ABA
$\theta_H$	hydrotime constant
$\theta_{ABA}$	ABA time constant
$\theta_{GA}$	GA time constant

of  $\psi_b(g)$  values among seeds in the population can be characterized by its mean [ $\psi_b(50)$ ] and SD ( $\sigma_{\psi_b}$ ), derived using repeated probit analysis (Bradford, 1990). The resulting probit equation is

$$\text{probit}(g) = [\psi - (\theta_H/t_g) - \psi_b(50)]/\sigma_{\psi_b} \quad (3)$$

which relates  $g$  to  $t_g$  at any constant  $\psi$ . Seed germination time courses can then be predicted at any  $\psi$  on the basis of the three basic parameters  $\theta_H$ ,  $\psi_b(50)$ , and  $\sigma_{\psi_b}$ .

The same formal equations also can be used to quantitate and predict seed germination behavior in the presence of exogenous ABA (Ni and Bradford, 1992). The defining model is

$$\theta_{ABA} = (\log[ABA] - \log[ABA_b(g)])t_g \quad (4)$$

and the probit equation relating germination percentage to ABA concentration and imbibition time is

$$\text{probit}(g) = \{\log[ABA] - (\theta_{ABA}/t_g) - \log[ABA_b(50)]\}/\sigma_{ABA_b} \quad (5)$$

In this case, the relevant parameters are the "ABA time constant" ( $\theta_{ABA}$ ), the log of the exogenous ABA concentration ( $\log[ABA]$ ), the normally distributed log threshold or base ABA concentrations inhibiting germination of percentage  $g$  ( $\log[ABA_b(g)]$ ), the log of the mean base or threshold ABA concentration ( $\log[ABA_b(50)]$ ), and the SD of  $ABA_b$  values ( $\sigma_{ABA_b}$ ). To determine whether endogenous ABA influences germination in a manner consistent with this model, we have examined the sensitivity to both ABA and  $\psi$  of *sit<sup>w</sup>* seeds (Karssen et al., 1987; Groot and Karssen, 1992) and its wild-type parent, *MM*.

Seeds from a tomato mutant deficient in GA (*gib-1*) require applied GA to germinate, and germination percentage increases as the GA concentration increases (Groot, 1987; Groot and Karssen, 1987; Groot et al., 1988). We have examined whether the responses of seed germination to GA can be analyzed using the same model used for ABA responses but based upon a threshold concentration of GA permitting, rather than inhibiting, germination. The defining model would be

$$\theta_{GA} = (\log[GA] - \log[GA_b(g)])t_g \quad (6)$$

and the relationship among germination percentage, imbibition time, and GA concentration is

$$\text{probit}(g) = \{\log[GA] - (\theta_{GA}/t_g) - \log[GA_b(50)]\}/\sigma_{GA_b} \quad (7)$$

where  $\theta_{GA}$  is the "GA time constant,"  $\log[GA]$  is the GA concentration of the imbibition medium,  $\log[GA_b(g)]$  is the normally distributed threshold or base GA concentration permitting germination of percentage  $g$ , and  $\log[GA_b(50)]$  and  $\sigma_{GA_b}$  are the mean and SD of  $GA_b$  values, respectively. The model predicts that, as the GA concentration increases above the threshold level for a given seed, the time to germination will be shortened proportionately, because  $\theta_{GA}$  is a constant (Eq. 6).

We show here that these models adequately describe the germination characteristics of wild-type, *sit<sup>w</sup>*, and *gib-1* seeds and their responses to applied hormones and reduced  $\psi$ . The influence of both endogenous and exogenous growth regulators on germination behavior and water relations can be quantitatively evaluated using this approach. We further demonstrate that the effects of ABA and GA on germination and dormancy are mediated primarily through changes in the threshold  $\psi$ s permitting radicle emergence.

## MATERIALS AND METHODS

### Plant Material

Tomato (*Lycopersicon esculentum* Mill.) seeds of the homozygous *gib-1* and *sit<sup>w</sup>* mutants and their wild-type isogenic parent (cv *MM*) were obtained from Dr. C.M. Karssen, Agricultural University, Wageningen, The Netherlands. Plants of all genotypes were grown in potting soil in a greenhouse. The *gib-1* plants were sprayed with 10  $\mu\text{M}$  GA<sub>4+7</sub> once or twice each week to stimulate shoot growth and flower development (Groot, 1987). Seeds were extracted from mature fruit, fermented in the juice for 1 d, rinsed, dried, and stored at 4°C.

## Seed Germination

Four replicates of 50 seeds each were placed on two 4.5-cm filter papers in 5-cm Petri dishes moistened with 4 mL of water, ABA, GA<sub>4+7</sub>, or PEG solutions. The dishes were covered with tightly fitting lids, placed in a covered polystyrene box lined with water-saturated tissue paper, and incubated in the dark at 25 ± 1°C. The PEG solutions were prepared according to the procedure of Michel (1983) using PEG 8000 (Carbowax PEG 8000; Fisher Scientific Co., Fair Lawn, NJ), and the  $\psi_s$  of the solutions were verified with a vapor pressure osmometer (model 5100C; Wescor Inc., Logan, UT) calibrated with NaCl standards. It has been reported that the exclusion of PEG from the filter paper matrix can cause errors in  $\psi$  determinations (Hardegee and Emmerich, 1990). However, a similar solution volume to filter paper weight ratio was present in both the osmometer measurement and in the germination dishes. The  $\psi$  measured by the osmometer should, therefore, correspond to that in the incubation dishes. The ABA and GA<sub>4+7</sub> solutions were prepared by dissolving ABA (Sigma, St. Louis, MO) and GA<sub>4+7</sub> (Abbott Laboratories, North Chicago, IL) in 1 M KOH and diluting with distilled water. Solutions containing PEG were changed after the first 24 h and then weekly thereafter, and ABA and GA solutions were changed every 2 weeks. Clearly visible radicle protrusion was used as the criterion for the completion of germination.

## Measurement of Embryonic RWC and $\psi_\pi$

Embryo water contents (dry weight basis) were measured by oven drying at 130°C for 1 h. Embryo  $\psi_\pi$  was determined by thermocouple psychrometry on frozen and thawed tissue using the vapor pressure osmometer. For each genotype, the embryonic water content of seeds that had fully imbibed on water was taken as 100% RWC. The  $\psi_\pi$  values determined for each genotype that had imbibed in PEG, ABA, or GA<sub>4+7</sub> were corrected for water content changes as described previously (Ni and Bradford, 1992).


## Data Analysis

Data analyses to determine the values of the model parameters were conducted using repeated probit regression as described previously (Bradford, 1990; Dahal and Bradford, 1990; Ni and Bradford, 1992).

## RESULTS

### Germination of *MM* Seeds in ABA, GA, or Reduced $\psi$

Groot and Karssen (1992) reported that the occurrence of postharvest dormancy in *MM* seeds is sensitive to environmental conditions during seed development. The final germination of our *MM* seed lot was only about 75% in water because of partial dormancy. Although 1  $\mu$ M ABA had no effect on germination in comparison with the water control, higher concentrations progressively delayed and inhibited germination (Fig. 1A). The threshold model fit the data well ( $r^2 = 0.93$ ; Table II), and the predicted curves are shown in Figure 1A. The low  $ABA_b(50)$  of 6  $\mu$ M (Table II) indicates the relatively high sensitivity of this seed lot to ABA. The  $\sigma_{ABA_b}$



**Figure 1.** Cumulative germination time courses of *MM* seeds incubated at a range of ABA concentrations (A),  $\psi$  (B), and  $\psi + 100 \mu\text{M}$  GA<sub>4+7</sub> (C). The symbols represent the actual data, and the solid curves are predicted by the threshold models (Eqs. 3 and 5) using the parameters in Table II. The time axis is plotted on a logarithmic scale to separate the curves for clarity.

value (0.84 log[M]) reflects wide variation in ABA sensitivity among individual seeds (Table II).


Germination of *MM* seeds was markedly delayed and inhibited by small reductions in  $\psi$  (Fig. 1B). This sensitivity to  $\psi$  was evident in the high  $\psi_b(50)$  value of  $-0.35$  MPa (Table II). With the addition of 100  $\mu\text{M}$  GA<sub>4+7</sub>, however, germination rate was greatly enhanced, and seed dormancy was completely broken (Fig. 1C). Germination was complete within 100 h on 100  $\mu\text{M}$  GA<sub>4+7</sub>, a time when <50% germination had occurred on water (Fig. 1, A and B). Germination was also much less sensitive to reduced  $\psi$  in the presence of GA<sub>4+7</sub> (Fig. 1C). This increased tolerance to low  $\psi$  was mainly attributable to a GA-induced lowering of  $\psi_b(50)$  from  $-0.35$  to  $-0.75$  MPa, although the  $\theta_H$  was also reduced by 20% (Table II).

The distributions of  $\psi_b(g)$  for *MM* seeds in the presence and absence of GA<sub>4+7</sub> are shown diagrammatically in Figure 2A. The relative frequency of a particular  $\psi_b$  value within the

**Table II.** Parameters of water relations, ABA, and GA models characterizing seed germination time courses

The parameters were derived using probit analysis of data from all factor levels. The time constants include  $\theta_H$  (MPa h),  $\theta_{ABA}$ , and  $\theta_{GA}$  (log[M] h). The mean base values are  $\psi_b(50)$  (MPa),  $ABA_b(50)$  and  $GA_b(50)$  (log[M] or  $\mu\text{M}$ ). The corresponding SD ( $\sigma$ ) of each threshold distribution and coefficients of determination ( $r^2$ ) of the probit regressions are also shown.

Genotype	Factor	Time Constants	Mean Base Values	$\sigma$	$r^2$
MM	ABA	-108 log[M] h	-5.2 (6 $\mu\text{M}$ )	-0.84 log[M]	0.93
MM	$\psi$	42 MPa h	-0.35 MPa	0.28 MPa	0.95
MM	$\psi + 100 \mu\text{M GA}_{4+7}$	34 MPa h	-0.75 MPa	0.15 MPa	0.86
<i>sit<sup>w</sup></i>	ABA	-48 log[M] h	-4.9 (13 $\mu\text{M}$ )	-0.54 log[M]	0.91
<i>sit<sup>w</sup></i>	$\psi$	45 MPa h	-1.17 MPa	0.33 MPa	0.96
<i>sit<sup>w</sup></i>	$\psi + 4.5 \mu\text{M ABA}$	72 MPa h	-0.90 MPa	0.43 MPa	0.91
<i>gib-1</i>	$GA_{4+7}$	99 log[M] h	-5.9 (1.3 $\mu\text{M}$ )	1.01 log[M]	0.97
<i>gib-1</i>	$\psi + 10 \mu\text{M GA}_{4+7}$	40 MPa h	-0.41 MPa	0.40 MPa	0.96
<i>gib-1</i>	$\psi + 100 \mu\text{M GA}_{4+7}$	41 MPa h	-0.71 MPa	0.23 MPa	0.97




**Figure 2.** Relative frequency distributions of  $\psi_b(g)$  values within seed populations of MM (A), *sit<sup>w</sup>* (B), and *gib-1* (C). The ordinate indicates the relative frequency within the seed population of the possible values of  $\psi_b$  indicated on the abscissa. The vertical dotted line at 0 MPa shows the highest possible  $\psi_b(g)$  at which germination will occur in water. The shaded areas indicate the fraction of dormant or inhibited seeds in each case.

seed population is illustrated as a normal distribution based upon the  $\psi_b(50)$  and  $\sigma_{\psi_b}$  values in Table II. The high  $\psi_b(50)$  and relatively large  $\sigma_{\psi_b}$  for MM seeds result in a distribution that extends above 0 MPa (shaded region in Fig. 2A). Because  $\psi$  cannot exceed 0 MPa, the fraction of the seed population with an estimated  $\psi_b(g) > 0$  MPa would be unable to germinate even on water, accounting for the percentage of dormant seeds. In addition, as the external  $\psi$  is reduced, the  $\psi_b$  of an increasing fraction of the population would exceed the current  $\psi$  level and be prevented from germinating. The time to completion of germination of the remaining seeds would also be delayed, because the time to germination is inversely proportional to the difference between the seed  $\psi$  and its own  $\psi_b$  threshold, generating the  $\psi$  response curves shown in Figure 1B. In the presence of 100  $\mu\text{M GA}_{4+7}$ , the  $\psi_b(g)$  distribution is shifted to lower values and is narrower because of the smaller  $\sigma_{\psi_b}$  (Fig. 2A). The distribution does not extend above 0 MPa, in agreement with 100% germination on water or breaking of dormancy (Fig. 1C). The greater tolerance to reduced  $\psi$  in the presence of  $GA_{4+7}$  (Fig. 1C) is also evident from the position of the  $\psi_b(g)$  distribution, because  $\psi$  would have to be reduced considerably before it would be below the threshold of a significant fraction of seeds in the population. We can conclude, therefore, that both dormancy and its release by  $GA_{4+7}$  in MM seeds are based upon the values of the  $\psi_b(g)$  distributions within the seed populations and that the time to germination, final germination percentage, and sensitivity of germination to reduced  $\psi$  are all linked manifestations of this underlying  $\psi$  threshold distribution.

#### Germination of *sit<sup>w</sup>* Seeds in ABA or Reduced $\psi$

In contrast to its wild-type parent, all homozygous *sit<sup>w</sup>* seeds germinated rapidly in water, and higher concentrations of ABA were required to inhibit germination of this genotype in comparison with MM seeds (Fig. 3A). The more rapid germination of *sit<sup>w</sup>* seeds resulted in a  $\theta_{ABA}$  of -48 log[M] h, as compared to -108 log[M] h for MM, and the  $ABA_b(50)$  was 13  $\mu\text{M}$ , double that of MM seeds (Table II).

Bradford (1990) derived a factor  $[1 - \psi/\psi_b(g)]$  that could be used to normalize the times to completion of germination



**Figure 3.** Cumulative germination time courses of *sit<sup>w</sup>* seeds incubated at a range of ABA concentrations (A),  $\psi$  (B), and  $\psi + 4.5 \mu\text{M}$  ABA (C). The symbols represent the actual data, and the solid curves are predicted by the threshold models (Eqs. 3 and 5) using the parameters in Table II. The time axis is plotted on a logarithmic scale to separate the curves for clarity.


at different  $\psi$  values to a common time course equal to that in water. The same procedure was used here to normalize the germination time courses at different ABA concentrations to a common time course equal to that in water. For the *sit<sup>w</sup>* seeds, we can calculate from Table II that the highest ABA concentration that would have no effect on germination ( $ABA_0$ ) would be  $0.3 \mu\text{M}$  (i.e. 99.9% germination = 3 SD units above the mean;  $\log[ABA_b(99.9)] = -4.9 - 3(0.54) = -6.5 \log[M]$ , or  $0.3 \mu\text{M}$ ). The maximum ABA concentration permitting germination of any individual seed would be  $\log[ABA_b(g)]$ , so  $\log[ABA_0] - \log[ABA_b(g)]$  represents the maximum range of ABA concentrations allowing germination of percentage  $g$ . Normalizing the ABA model (Eq. 4) by dividing both sides of the equation by  $\log[ABA_0] - \log[ABA_b(g)]$ , the resulting equation is

$$t_g(0) = \left( \frac{\log[ABA] - \log[ABA_b(g)]}{\log[ABA_0] - \log[ABA_b(g)]} \right) t_g(ABA) \quad (8)$$

where  $t_g(0)$  is the time to germination at the minimum thresh-

old ABA concentration (equivalent to that in water) and  $t_g(ABA)$  is the time to germination at a given higher ABA concentration. Figure 4 shows the germination time courses of *sit<sup>w</sup>* seeds at various ABA concentrations (from Fig. 3A) normalized to a single curve approximating the time course in water (closed symbols and solid line). The ability of this normalization equation to collapse the array of curves in Figure 3A to a single curve in Figure 4 is an indication of the degree to which parameters derived from the model describe the actual germination responses.

The rapid germination of *sit<sup>w</sup>* seeds was particularly evident at reduced  $\psi$ . Germination of *sit<sup>w</sup>* seeds was complete within 200 h at  $-0.45 \text{ MPa}$  (Fig. 3B), whereas only 30% of *MM* seeds were able to germinate at this  $\psi$  (Fig. 1B). The rapid germination observed in this genotype resulted entirely from its low  $\psi_b(50)$  ( $-1.17 \text{ MPa}$ ), because its  $\theta_H$  value was essentially the same as that of *MM* seeds (Table II). Using the ABA model, we can calculate that the *sit<sup>w</sup>* seeds on  $4.5 \mu\text{M}$  ABA should germinate at approximately the same rate as *MM* seeds on water. We tested this prediction by germinating *sit<sup>w</sup>* seeds in water or osmotic solutions containing  $4.5 \mu\text{M}$  ABA (Fig. 3C). The addition of ABA did delay germination to approximate that of the *MM* seeds in water, but a higher concentration of ABA was apparently required to completely reproduce the wild-type behavior at all  $\psi$  values (compare with Fig. 1B). The  $\psi_b(50)$  value of *sit<sup>w</sup>* seeds increased from  $-1.17$  to  $-0.90 \text{ MPa}$  in the presence of  $4.5 \mu\text{M}$  ABA, which is still much lower than the  $\psi_b(50)$  of  $-0.35 \text{ MPa}$  of *MM* seeds (Table II). This change in the  $\psi_b(g)$  distribution of *sit<sup>w</sup>* seeds due to ABA is shown in Figure 2B, which illustrates that,




**Figure 4.** Cumulative germination time courses for *sit<sup>w</sup>* (from Fig. 3A; closed symbols) and for *gib-1* seeds (from Fig. 5A; open symbols) at a range of ABA and  $\text{GA}_{4+7}$  concentrations, respectively, plotted on a normalized time scale. The data for *sit<sup>w</sup>* seeds are normalized to the predicted time in water (solid line) using Equation 8 and the values in Table II, and those for *gib-1* seeds are normalized to the predicted time course occurring in  $100 \mu\text{M}$   $\text{GA}_{4+7}$  (dashed curve) using Equation 8 (with GA substituted for ABA) and the values in Table II. The dotted line represents the theoretical germination time course predicted by the model for *gib-1* seeds germinating in  $1 \text{ mM}$   $\text{GA}_{4+7}$ .

although ABA increased the mean  $\psi_b$  value of the population, the majority of seeds still had thresholds considerably below those of *MM* seeds (Fig. 2A). Previous work with a different genotype showed that the mean  $\psi_b$  value would increase further at higher ABA concentrations (Ni and Bradford, 1992). The  $\theta_H$  also increased from 45 to 72 MPa h in the presence of 4.5  $\mu\text{M}$  ABA (Table II). Thus, ABA delayed germination by a dual effect on the  $\theta_H$  and on the  $\psi_b(g)$  distribution.

#### Germination of *gib-1* Seeds in GA or Reduced $\psi$

Seeds of the *gib-1* mutant showed a complete and quantitative dependence of germination on  $\text{GA}_{4+7}$ . No germination was observed without applied  $\text{GA}_{4+7}$ , and the germination percentage and rate progressively increased with increasing  $\text{GA}_{4+7}$  concentrations (Fig. 5A). The models of Equations 6 and 7 fit the GA response of *gib-1* seeds very well ( $r^2 = 0.97$ ; Table II; predicted curves in Fig. 5A). The very low value of  $\text{GA}_{b(50)}$  (1.3  $\mu\text{M}$ ) indicates the high sensitivity of germination



**Figure 5.** Cumulative germination time courses of *gib-1* seeds incubated at a range of  $\text{GA}_{4+7}$  concentrations (A),  $\psi + 10 \mu\text{M}$   $\text{GA}_{4+7}$  (B), and  $\psi + 100 \mu\text{M}$   $\text{GA}_{4+7}$  (C). The symbols represent the actual data, and the solid curves are predicted by the threshold models (Eqs. 3 and 7) using the parameters in Table II. The time axis is plotted on a logarithmic scale to separate the curves for clarity.

to GA in these GA-deficient mutant seeds. As for ABA, a function can be developed to normalize germination time courses at any GA concentration to that predicted for a saturating dose of GA. The function will be the same as Equation 8 with GA values substituted for ABA, except that  $\text{GA}_0$  will be the saturating dose of  $\text{GA}_{4+7}$  (100  $\mu\text{M}$ ) rather than the minimum threshold dose. This function worked well to collapse the array of time courses in Figure 5A to a single normalized time course (Fig. 4, open symbols and dashed line).

When incubated in osmoticum containing 10  $\mu\text{M}$   $\text{GA}_{4+7}$ , the germination time courses and water relations parameters of *gib-1* seeds were similar to those of *MM* seeds in the absence of GA (compare Figs. 1B and 5B; Table II). When both genotypes were supplied with 100  $\mu\text{M}$   $\text{GA}_{4+7}$ , the germination behavior and water relations parameters of *gib-1* and *MM* seeds were again very similar (compare Figs. 1C and 5C; Table II).  $\text{GA}_{4+7}$  could, therefore, restore both the germination rate and water potential sensitivity of *gib-1* seeds to equal that of its wild-type parent. This is evident in the  $\psi_b(g)$  distributions of *gib-1* seeds in either 10 or 100  $\mu\text{M}$   $\text{GA}_{4+7}$  (Fig. 2C), which correspond closely to those of *MM* seeds in water or 100  $\mu\text{M}$   $\text{GA}_{4+7}$  (Fig. 2A). Projecting this trend to lower  $\text{GA}_{4+7}$  concentrations, we can suggest that the threshold distribution would continue to increase, resulting in the progressive delay and reduction in germination evident in Figure 5A. On the other hand, when  $\text{GA}_{4+7}$  stimulates germination of dormant *gib-1* seeds, the entire population distribution of  $\psi_b(g)$  is shifted to lower values, accounting for both the increased rate and final percentage of germination.

Even in 100  $\mu\text{M}$   $\text{GA}_{4+7}$ , germination of *gib-1* or *MM* seeds was not as rapid as that of *sit<sup>w</sup>* seeds (Figs. 1C, 3A, 4, and 5C). It is interesting that inserting a value of 1 mM  $\text{GA}_{4+7}$  into the GA model (Eq. 7) using the parameters for the GA response in Table II resulted in a predicted time course for *gib-1* seeds that was essentially identical with that of *sit<sup>w</sup>* seeds in water (dotted curve in Fig. 4). We tested this prediction experimentally by germinating *gib-1* and *MM* seeds in 1 mM  $\text{GA}_{4+7}$ , but the time courses were not significantly more rapid than those in 100  $\mu\text{M}$   $\text{GA}_{4+7}$  (data not shown). Although experiments at these very high hormone concentrations must be interpreted with caution, this result suggests that, although the model predicts that *sit<sup>w</sup>* seeds germinate as if they were in the presence of 1 mM  $\text{GA}_{4+7}$ , germination of *gib-1* or *MM* seeds cannot be accelerated to an equivalent degree by supraoptimal GA. Low concentrations of ABA during development may precondition the seeds for a subsequent germination rate that cannot be duplicated by excess GA applied to seeds that had developed with normal ABA concentrations.

#### Effect of Reduced $\psi$ , ABA, and GA on Embryo $\psi_\pi$

The changes in  $\psi_b(g)$  values due to genotype, ABA, and GA described above could result from accumulation or loss of solutes in the embryos, because embryo  $\psi_\pi$  determines the maximum turgor force that can be generated by the embryo to penetrate through the enclosing endosperm tissues. To test this possibility, embryo  $\psi_\pi$  was measured in embryos of seeds from which the radicle had not protruded at a time when some seeds had just completed germination under each im-

bibition condition (Table III). Among the three genotypes, *sit<sup>w</sup>* embryos had the lowest  $\psi_{\pi}$  (-1.41 MPa), 0.14 and 0.19 MPa lower than that of *MM* and *gib-1* embryos, respectively (Table III). This pattern is consistent with the lower  $\psi_b(50)$  of *sit<sup>w</sup>* seeds, but the differences in  $\psi_{\pi}$  are far too small to account for the differences in  $\psi_b(50)$  (Table II). Similar results were also found by H.W.M. Hilhorst at Wageningen Agricultural University (personal communication). The presence of ABA caused a small increase in embryonic RWC but did not cause any change in  $\psi_{\pi}$  that could not be accounted for by the increased water content (compare observed and predicted values in Table III). When seeds were incubated on 100  $\mu\text{M}$  GA<sub>4+7</sub> solutions, however, the predicted embryo  $\psi_{\pi}$  values for both *MM* and *gib-1* seeds were about 0.1 MPa lower than the corresponding measured values, indicating a net accumulation of solutes equivalent to about 0.1 MPa (Table III). Overall, the changes in embryonic  $\psi_{\pi}$  due to genotype or hormones were small compared to the corresponding differences in  $\psi_b(g)$ .

## DISCUSSION

### Population-Based Threshold Model

The population-based threshold model worked well to describe germination of *MM*, *sit<sup>w</sup>*, and *gib-1* seeds in response to reduced  $\psi$  and a range of exogenous ABA and GA<sub>4+7</sub> concentrations, with >86% of the variation in germination times accounted for by the model in all cases (Table II). A good match between the actual data and the curves predicted by the models is evident for all three genotypes across all  $\psi$  and hormone levels (Figs. 1, 3, and 5). It should be emphasized that the parameters of the models and the predicted curves are not empirical fits to each individual time course. Rather, as discussed previously (Ni and Bradford, 1992), the parameters derived from simultaneous analysis of data at all factor levels are estimators of the mean sensitivity of a seed population to  $\psi$  [ $\psi_b(50)$ ], ABA [ $ABA_b(50)$ ], or GA [ $GA_b(50)$ ], the

corresponding sd values of the threshold distributions ( $\sigma_{\psi_b}$ ,  $\sigma_{ABA_b}$ , and  $\sigma_{GA_b}$ , respectively), and the proportionality or time constants that relate the difference between the external level and the threshold value of a factor to a corresponding time to completion of germination ( $\theta_H$ ,  $\theta_{ABA}$ , and  $\theta_{GA}$ ). When the three basic parameters for each model are determined, germination time courses of the entire population in response to  $\psi$ , ABA, or GA can be predicted by changing only the factor level in Equations 3, 5, or 7. In addition to the factors described here, the same basic model is likely to be applicable to other agents that affect germination, including ethylene, light, nitrate, or aging (Abeles, 1986; Hilhorst and Karssen, 1989; Tarquis et al., 1992).

A novel feature of this model is its ability to simultaneously account for both the timing and the final extent of germination at different factor levels. This is a result of the distribution of different factor threshold values among individual seeds and the constant proportionality between the germination response rate and the difference between the current factor level and its threshold. Other approaches to quantitating the sensitivity of germination to environmental or physiological factors from germination response curves, such as those derived from the Michaelis-Menten enzyme kinetic equation (Weyers et al., 1987; Hilhorst and Karssen, 1989) or from fitted monomolecular, logistic, or Gompertz functions (Tipton, 1984), require the estimation of separate values for maximum response rate and other "constants" for each individual curve and do not explicitly incorporate the population aspects of germination behavior into the analysis. The models developed here characterize the physiological sensitivity and variation in sensitivity of the seed population to a given factor and then use this distribution to generate the predicted response at any factor level.

We have also demonstrated previously how the apparently additive or synergistic action of multiple factors (e.g. ABA and  $\psi$ ) can be predicted by mathematically combining the separate response functions for each factor (Ni and Bradford, 1992). The model also predicts that equivalent effects on

**Table III.** Water content, RWC, and  $\psi_{\pi}$  of embryos after imbibition by tomato seeds of water, 33  $\mu\text{M}$  ABA, or 100  $\mu\text{M}$  GA<sub>4+7</sub>

Seeds imbibed for the period indicated (just before radicle emergence), the embryos were excised, and the embryonic water contents (dry weight basis) and  $\psi_{\pi}$  were measured. The water content of the embryos of each genotype that had imbibed water was taken as 100% RWC, and the RWC of the embryos that had imbibed ABA or GA are calculated as a percentage of this value. The predicted values of  $\psi_{\pi}$  corrected for changes in embryo RWC were calculated by dividing the measured  $\psi_{\pi}$  of the water-imbibing embryos of each genotype by the factor  $[1 - (100 - \text{RWC})/83]$  using a constant nonosmotic volume of 17% (Ni and Bradford, 1992). Means  $\pm$  SE are shown ( $n = 4$ ).

Genotype	Imbibition Condition	Embryonic Water Content		$\psi_{\pi}$	
		Percentage	RWC	Observed	Predicted
				MPa	
<i>MM</i>	H <sub>2</sub> O (45 h)	49.9 $\pm$ 1.3	100	-1.27 $\pm$ 0.03	-1.27
<i>MM</i>	33 $\mu\text{M}$ ABA (7 d)	52.6 $\pm$ 2.2	105	-1.22 $\pm$ 0.04	-1.20
<i>MM</i>	100 $\mu\text{M}$ GA <sub>4+7</sub> (32 h)	49.4 $\pm$ 1.6	99	-1.38 $\pm$ 0.02	-1.29
<i>sit<sup>w</sup></i>	H <sub>2</sub> O (24 h)	50.3 $\pm$ 3.2	100	-1.41 $\pm$ 0.03	-1.41
<i>sit<sup>w</sup></i>	33 $\mu\text{M}$ ABA (36 h)	53.8 $\pm$ 2.4	107	-1.32 $\pm$ 0.06	-1.30
<i>gib-1</i>	H <sub>2</sub> O (45 h)	56.5 $\pm$ 2.4	100	-1.22 $\pm$ 0.06	-1.22
<i>gib-1</i>	100 $\mu\text{M}$ GA <sub>4+7</sub> (36 h)	59.8 $\pm$ 1.7	106	-1.22 $\pm$ 0.02	-1.14

germination would result from a given change in either the factor level or in the threshold value, because the response is proportional only to the difference between them (Bradford et al., 1992). The model, therefore, can be used to characterize responses due to changes in either factor levels or tissue sensitivity and can discriminate between the two possibilities.

Application of a population-based model is not limited to seed germination, because it is general in plants that individual tissues or cells within the same tissue may differ in their hormonal sensitivity, resulting in wide dose-response curves and variable responses to a given dosage among cells or tissues (Trewavas, 1991). In fact, the concept of target tissues or cells that are particularly responsive to a given factor is virtually axiomatic in our understanding of hormonal regulation of growth and development. Although this variation in sensitivity has been recognized, it has seldom been quantified as a significant physiological parameter in its own right.

Seed germination is perhaps only a more obvious example of this fundamental physiological feature because a seed population is a collection of individual seeds that do not germinate simultaneously but, rather, show a distribution of germination events over time. This distribution in time is mainly attributable to the differences in sensitivity to environmental or hormonal factors among individual seeds in the population. We can calculate, for example, that the differences in the sensitivity to ABA [ $ABA_b(g)$  values] between the first 10% and the last 10% of germinating seeds ( $\pm 1.28 \sigma_{ABA_b}$  around the mean) were as large as 140-fold for *MM* seeds and 25-fold for *sit<sup>w</sup>* seeds. Differences were even greater among *gib-1* seeds in response to  $GA_{4+7}$  (390-fold). The analytical methods developed here using seeds as a model system should be applicable to quantifying and understanding the variations in sensitivity that occur in other systems as well, such as, for example, nonuniform stomatal closure in response to ABA (Terashima et al., 1988) and mild water stress (Sharkey and Seemann, 1989; Ni and Pallardy, 1992), variable changes in cytosolic calcium concentration among epidermal guard cells in response to a single ABA concentration (McAinsh et al., 1992), tissue-specific gene expression patterns in response to hormones or water stress (Cohen et al., 1991; Li et al., 1991; van Loon and Bruinsma, 1992), and differences in  $\alpha$ -amylase production among individual aleurone cells in response to GA (Jacobsen and Knox, 1973).

A final implication of the present analysis is that, in analogy to thermal time, the concepts of "hydrottime," "ABA time," and "GA time" also have physiological meaning for seed germination. The application of thermal time, or a heat-units or degree-days analysis, to normalize the rates of biological processes at different temperatures is well established in modeling of crop growth or insect development (Fry, 1983). Germination behavior at different  $\psi$ , ABA, or GA levels can be normalized using hydrottime, ABA time, and GA time in analogy to the use of thermal time to normalize germination rates at different temperatures (Fig. 4; Bradford, 1990; Dahal and Bradford, 1990; Dahal et al., 1990). That is, progress toward germination is proportional to the prevailing level of  $\psi$ , ABA, or GA in much the same way that biological developmental rates are proportional to temperature. An alternative way to view this is to consider that, as  $\psi$  is reduced or ABA concentration is increased, biological time "stretches" or

slows down so that more clock time is required to reach the same physiological state (i.e. radicle emergence in this case).

In contrast, as GA concentration or  $\psi$  is increased, biological time "contracts" so that more development is achieved per unit of clock time, as would occur if the temperature were increased. There is also evidence that biological or metabolic time can accumulate under conditions in which radicle emergence is prevented by low  $\psi$  or ABA, shortening subsequent times to germination when the inhibitor is removed (Finch-Savage and McQuistan, 1991; Dahal et al., 1992). This is not to say that the underlying mechanisms of thermal time, hydrottime, ABA time, and GA time are necessarily identical. These rather novel concepts of biological time are, however, a well-supported consequence of our analysis of seed germination, and this alternative viewpoint may have more general implications for understanding the regulation of plant development.

### Regulation of Germination and Dormancy in Tomato Seeds


It has been well documented in studies using ABA- and GA-deficient mutants of *Arabidopsis thaliana* and tomato that ABA determines the depth of dormancy during development and the amount of GA required to break dormancy during imbibition (Karssen et al., 1983, 1987; Groot, 1987; Koornneef et al., 1989; Groot and Karssen, 1992). Our results are in excellent qualitative agreement with those recently reported by Groot and Karssen (1992) using the same tomato genotypes. Our work extends their analysis to provide quantitative measures of  $\psi$ , ABA, and GA sensitivity for each genotype and develops a comprehensive model explaining the underlying linkage among germination rates, sensitivity to  $\psi$ , and dormancy.

In our analysis, the influence of both endogenous and exogenous ABA and GA on dormancy and germination can be explained largely on the basis of their effects on the  $\psi_b(g)$  distributions within the seed population. When GA was absent, as in the *gib-1* mutant, the  $\psi_b(g)$  distribution was entirely above 0 MPa so that no germination occurred even on water. As the exogenous  $GA_{4+7}$  concentration increased, the  $\psi_b(g)$  distribution shifted to lower values (Fig. 2C), allowing a larger percentage of seeds to germinate and reducing the time to germination (Fig. 5A).

In contrast, low ABA levels during development of *sit<sup>w</sup>* seeds resulted in very low  $\psi_b(g)$  values, absence of dormancy, and rapid germination, and exogenous ABA delayed and inhibited germination by causing an increase in both  $\psi_b(g)$  and  $\theta_H$  (Fig. 2B; Table II; Ni and Bradford, 1992). Partial dormancy in *MM* seeds was due to a high  $\psi_b(g)$  distribution that extended above 0 MPa; that fraction of the seed population for which  $\psi_b > 0$  MPa was unable to germinate even on water or was dormant (Fig. 2A).

If the distribution of  $\psi_b(g)$  determines when and if a given seed will germinate, what determines the values of  $\psi_b(g)$ ? A scheme summarizing our current understanding of the factors controlling tomato seed germination is shown in Figure 6. Whether radicle emergence occurs or not under a given condition is determined by the balance of forces between the growth potential or turgor of the embryo (essentially the difference in  $\psi_x$  between the embryo and the external me-





**Figure 6.** A diagram summarizing the water and hormonal relations in the embryo and the endosperm cap that influence radicle emergence of tomato seeds. The inset illustrates the enclosure of the embryo by the endosperm and the thin-walled endosperm cap region opposite the radicle tip that weakens to allow radicle emergence (Haigh, 1988; modified from Groot, 1987).

dium) and the yield threshold or mechanical resistance of the endosperm cap. The value of  $\psi_b(g)$  is the net result of these opposing forces. The primary control of germination timing seems to reside in the endosperm cap, whereas secondary effects on cell wall loosening of the embryo are expressed after the endosperm has weakened sufficiently to allow embryo expansion (Haigh, 1988; Karssen et al., 1989). High  $\psi$  increases the growth potential of the embryo and allows the weakening process in the endosperm to proceed. Low  $\psi$  decreases embryo growth potential directly by lowering turgor and also delays or inhibits endosperm weakening (Dahal and Bradford, 1990).

On the other hand, prolonged incubation at low  $\psi$  ( $\leq -0.5$  MPa) results in solute accumulation in the embryo, increasing the embryo growth potential, decreasing  $\psi_b(g)$  by as much as 0.3 MPa, and allowing germination at  $\psi$  values that were initially inhibitory (Ni and Bradford, 1992). However, the small differences in embryonic  $\psi_\pi$  detected here among genotypes or hormone treatments in water (Table III) were far too small to account for the observed differences in  $\psi_b(g)$  (Table II; Fig. 2). For example,  $\psi_\pi$  of *MM* and *sit<sup>w</sup>* embryos differed by only 0.14 MPa, and the difference in  $\psi_b(50)$  was 0.82 MPa. Imbibition in ABA or  $GA_{4+7}$  caused large changes in  $\psi_b(g)$ , but ABA had no effect on  $\psi_\pi$  and  $GA_{4+7}$  lowered  $\psi_\pi$  by only 0.1 MPa. In addition, there was no significant difference in  $\psi_\pi$  between *MM* and *gib-1* embryos when both seeds imbibed on water (Table III), indicating that the inability of *gib-1* seeds to germinate was not due to insufficient embryo turgor. It can be concluded that the differences in  $\psi_b(g)$  among the three genotypes and the shifts in  $\psi_b(50)$  caused by addition

of  $GA_{4+7}$  or ABA are not due to large changes in embryonic  $\psi_\pi$ .

Alternatively, the changes in  $\psi_b(g)$  could be dependent on the rate or extent of the weakening of the endosperm tissue directly opposite the radicle tip, because the endosperm completely encloses the embryo and limits its water uptake (Fig. 6; Groot, 1987; Haigh, 1988; Karssen et al., 1989; Dahal and Bradford, 1990). It has been suggested that values of  $\psi_b(g)$  are related to the resistance or yield threshold of the endosperm during a final weakening stage just before radicle emergence (Karssen et al., 1989; Dahal and Bradford, 1990). Linear relationships have been found between germination rate and endosperm puncture force (Karssen et al., 1989) and between germination rate and  $\psi_b(g)$  (Dahal and Bradford, 1990; Ni and Bradford, 1992). Hence, it is very likely that  $\psi_b(g)$  and endosperm resistance to penetration are closely correlated.

There is good evidence that ABA inhibits and GA promotes tomato seed germination by affecting the rate and extent of endosperm weakening, with lesser effects on embryo growth capacity (Groot and Karssen, 1987, 1992; Groot et al., 1988). Tomato endosperm cell walls are composed primarily of a mannan polymer with smaller quantities of Glc and Gal (Groot et al., 1988).  $GA_{4+7}$  induced endo- $\beta$ -mannanase and increased mannohydrolase activities in the endosperm of *gib-1* seeds, and the puncture force required to penetrate the endosperm cap was significantly reduced coincident with the initiation of radicle emergence (Groot and Karssen, 1987; Groot et al., 1988).

When ABA was added to the incubation solution, however,

the GA-induced endosperm weakening was completely inhibited and the puncture force remained high (Groot and Karssen, 1992). The *sit<sup>o</sup>* mutation that lowers endogenous ABA contents during seed development prevents the induction of dormancy, speeds germination, and dramatically lowers the  $\psi_w(g)$  for germination (Figs. 2 and 3; Table II; Groot and Karssen, 1992). To our knowledge, it has not been tested experimentally in tomato seeds whether ABA and/or low  $\psi$  act by inhibiting the synthesis or activity of the same set of cell wall hydrolases that are stimulated by GA, although it is a likely hypothesis. In isolated lettuce endosperms, ABA does inhibit the appearance of mannanase activity, and it has been argued that endogenous ABA is an in vivo regulator of the synthesis of this enzyme (Dulson et al., 1988). However, the effects of low  $\psi$  on tomato seed germination are not mediated through increases in endogenous ABA (Ni and Bradford, 1992). The changes in endosperm strength due to ABA and GA correspond well with the effects of these hormones on  $\psi_w(g)$  (Fig. 2), indicating that  $\psi_w(g)$  is determined mainly by modulation of cell wall hydrolase activity leading to endosperm weakening (Fig. 6).

Received August 24, 1992; accepted October 29, 1992.

Copyright Clearance Center: 0032-0889/93/101/0607/11.

#### LITERATURE CITED

- Abeles FB** (1986) Role of ethylene in *Lactuca sativa* cv. 'Grand Rapids' seed germination. *Plant Physiol* **81**: 780-787
- Bradford KJ** (1990) A water relations analysis of seed germination rates. *Plant Physiol* **94**: 840-849
- Bradford KJ, Dahal P, Ni B-R** (1992) Quantitative models describing germination responses to temperature, water potential, and growth regulators. In D Côme, F Corbineau, eds, *Proceedings of the Fourth International Workshop on Seeds: Basic and Applied Aspects of Seed Biology*. Association pour la Formation Professionnelle de l'Interprofession Semences, Paris, France (in press)
- Cohen A, Plant AL, Moses MS, Bray EA** (1991) Organ-specific and environmentally regulated expression of two abscisic acid-induced genes of tomato. Nucleotide sequence and analysis of the corresponding cDNAs. *Plant Physiol* **97**: 1367-1374
- Dahal P, Bradford KJ** (1990) Effects of priming and endosperm integrity on seed germination rates of tomato genotypes. II. Germination at reduced water potential. *J Exp Bot* **41**: 1441-1453
- Dahal P, Bradford KJ, Haigh AM** (1992) The concept of hydrothermal time in seed germination and priming. In D Côme, F Corbineau, eds, *Proceedings of the Fourth International Workshop on Seeds: Basic and Applied Aspects of Seed Biology*. Association pour la Formation Professionnelle de l'Interprofession Semences, Paris, France (in press)
- Dahal P, Bradford KJ, Jones RA** (1990) Effects of priming and endosperm integrity on seed germination rates of tomato genotypes. I. Germination at suboptimal temperature. *J Exp Bot* **41**: 1431-1439
- Dulson J, Bewley JD, Johnson RN** (1988) Abscisic acid is an endogenous inhibitor in the regulation of mannanase production by isolated lettuce (*Lactuca sativa* cv. Grand Rapids) endosperms. *Plant Physiol* **87**: 660-665
- Finch-Savage WE, McQuistan CI** (1991) Abscisic acid: an agent to advance and synchronize germination for tomato (*Lycopersicon esculentum* Mill.) seeds. *Seed Sci Technol* **19**: 537-544
- Fry KE** (1983) Heat-Unit Calculations in Cotton Crop and Insect Models. U.S. Department of Agriculture/Agricultural Research Service Advances in Agricultural Technology AAT-W-23
- Groot SPC** (1987) Hormonal regulation of seed development and germination in tomato. Studies on abscisic acid- and gibberellin-deficient mutants. PhD dissertation, Agricultural University, Wageningen, The Netherlands
- Groot SPC, Karssen CM** (1987) Gibberellins regulate seed germination in tomato by endosperm weakening: a study with gibberellin-deficient mutants. *Planta* **171**: 525-531
- Groot SPC, Karssen CM** (1992) Dormancy and germination of abscisic acid-deficient tomato seeds. Studies with the *sitiens* mutant. *Plant Physiol* **99**: 952-958
- Groot SPC, Kieliszewska-Rockika B, Vermeer E, Karssen CM** (1988) Gibberellin-induced hydrolysis of endosperm cell walls in gibberellin-deficient tomato seeds prior to radicle protrusion. *Planta* **174**: 500-504
- Gummerson RJ** (1986) The effect of constant temperatures and osmotic potential on the germination of sugar beet. *J Exp Bot* **37**: 729-741
- Haigh AM** (1988) Why do tomato seeds prime? Physiological investigations into the control of tomato seed germination and priming. PhD thesis. Macquarie University, Sydney, Australia
- Hardegree SP, Emmerich WE** (1990) Effect of polyethylene glycol exclusion on the water potential of solution-saturated filter paper. *Plant Physiol* **92**: 462-466
- Hilhorst HWM, Karssen CM** (1989) The role of light and nitrate in seed germination. In RB Taylorson, ed, *Recent Advances in the Development and Germination of Seeds*. Plenum Press, New York, pp 191-205
- Jacobsen JV, Knox RB** (1973) Cytochemical localization and antigenicity of  $\alpha$ -amylase in barley aleurone tissue. *Planta* **112**: 213-224
- Karssen CM, Brinkhorst-van der Swan DLC, Breekland AE, Koornneef M** (1983) Induction of dormancy during seed development by endogenous abscisic acid: studies on abscisic acid deficient genotypes of *Arabidopsis thaliana*. *Planta* **157**: 158-165
- Karssen CM, Groot SPC, Koornneef M** (1987) Hormone mutants and seed dormancy in *Arabidopsis* and tomato. In H Thomas, D Grierson, eds, *Developmental Mutants in Higher Plants*. Cambridge University Press, Cambridge, UK, pp 119-133
- Karssen CM, Haigh A, van der Toorn P, Weges R** (1989) Physiological mechanisms involved in seed priming. In RB Taylorson, ed, *Recent Advances in the Development and Germination of Seeds*. Plenum Press, New York, pp 269-280
- Kermode AR** (1990) Regulatory mechanisms involved in the transition from seed development to germination. *CRC Crit Rev Plant Sci* **9**: 155-195
- Koornneef M, Hanhart CJ, Hilhorst HWM, Karssen CM** (1989) In vivo inhibition of seed development and reserve protein accumulation in recombinants of abscisic acid biosynthesis and responsiveness mutants in *Arabidopsis thaliana*. *Plant Physiol* **90**: 463-469
- Li Y, Hagen G, Guilfoyle TJ** (1991) An auxin-responsive promoter is differentially induced by auxin gradients during tropisms. *Plant Cell* **3**: 1167-1175
- McAinsh MR, Brownlee C, Hetherington AM** (1992) Visualizing changes in cytosolic-free  $Ca^{2+}$  during the response of stomatal guard cells to abscisic acid. *Plant Cell* **4**: 1113-1122
- Michel BE** (1983) Evaluation of the water potentials of solutions of polyethylene glycol 8000 both in the absence and presence of other solutes. *Plant Physiol* **72**: 66-70
- Ni B-R, Bradford KJ** (1992) Quantitative models describing the sensitivity of tomato seed germination to abscisic acid and osmoticum. *Plant Physiol* **98**: 1057-1068
- Ni B-R, Pallardy SG** (1992) Stomatal and nonstomatal limitations to net photosynthesis in seedlings of woody angiosperms. *Plant Physiol* **99**: 1502-1508
- Schopfer P, Plachy C** (1985) Control of seed germination by abscisic acid. II. Effect on embryo growth potential (minimum turgor pressure) and growth coefficient (cell wall extensibility) in *Brassica napus* L. *Plant Physiol* **77**: 676-686

- Sharkey TD, Seemann JR** (1989) Mild water stress effects on carbon-reduction-cycle intermediates, RuBP carboxylase activity, and spatial homogeneity of photosynthesis in intact leaves. *Plant Physiol* **89**: 1060–1065
- Tarquis AM, Bradford KJ, Durán JM** (1992) A threshold model relating seed germination rates to ageing period during storage. In D Côme, F Corbineau, eds, Proceedings of the Fourth International Workshop on Seeds: Basic and Applied Aspects of Seed Biology. Association pour la Formation Professionnelle de l'Interprofession Semences, Paris, France (in press)
- Terashima I, Wong S-C, Osmond CB, Farquhar GD** (1988) Characterization of non-uniform photosynthesis induced by abscisic acid in leaves having different mesophyll anatomies. *Plant Cell Physiol* **29**: 385–394
- Tipton JL** (1984) Evaluation of three growth curve models for germination data analysis. *J Am Soc Hortic Sci* **109**: 451–454
- Trewavas A** (1991) How do plant growth substances work? II. *Plant Cell Environ* **14**: 1–12
- Van Loon LC, Bruinsma J** (1992) The new plant physiology—molecular approaches to studying hormonal regulation of plant development. *Acta Bot Neerl* **41**: 1–23
- Welbaum GE, Tissaoui T, Bradford KJ** (1990) Water relations of seed germination in muskmelon (*Cucumis melo* L.). III. Sensitivity of germination to water potential and abscisic acid during development. *Plant Physiol* **92**: 1029–1037
- Weyers JDB, Paterson NW, A'Brook R** (1987) Towards a quantitative definition of plant hormone sensitivity. *Plant Cell Environ* **10**: 1–10