

Luminal NaCl delivery regulates basolateral PGE₂ release from macula densa cells

János Peti-Peterdi,^{1,2} Peter Komlosi,^{1,2} Amanda L. Fuson,¹ Youfei Guan,³ André Schneider,³ Zhonghua Qi,³ Reyadh Redha,³ Laszlo Rosivall,² Matthew D. Breyer,³ and P. Darwin Bell¹

¹Department of Medicine and Department of Physiology, Division of Nephrology, University of Alabama at Birmingham, Birmingham, Alabama, USA

²Institute of Pathophysiology, Hungarian Academy of Sciences, and Semmelweis University Nephrology Research Group, Budapest, Hungary

³Department of Medicine, Molecular Physiology and Biophysics, Division of Nephrology, Vanderbilt University, Nashville, Tennessee, USA

Macula densa (MD) cells express COX-2 and COX-2-derived PGs appear to signal the release of renin from the renal juxtaglomerular apparatus, especially during volume depletion. However, the synthetic machinery and identity of the specific prostanoid released from intact MD cells remains uncertain. In the present studies, a novel biosensor tool was engineered to directly determine whether MD cells release PGE₂ in response to low luminal NaCl concentration ([NaCl]_L). HEK293 cells were transfected with the Ca²⁺-coupled E-prostanoid receptor EP₁ (HEK/EP₁) and loaded with fura-2. HEK/EP₁ cells produced a significant elevation in intracellular [Ca²⁺]_i ([Ca²⁺]_i) by 29.6 ± 12.8 nM (*n* = 6) when positioned at the basolateral surface of isolated perfused MD cells and [NaCl]_L was reduced from 150 mM to zero. HEK/EP₁ [Ca²⁺]_i responses were observed mainly in preparations from rabbits on a low-salt diet and were completely inhibited by either a selective COX-2 inhibitor or an EP₁ antagonist, and also by 100 μM luminal furosemide. Also, 20-mM graduated reductions in [NaCl]_L between 80 and 0 mM caused step-by-step increases in HEK/EP₁ [Ca²⁺]_i. Low-salt diet greatly increased the expression of both COX-2 and microsome-associated PGE synthase (mPGES) in the MD. These studies provide the first direct evidence that intact MD cells synthesize and release PGE₂ during reduced luminal salt content and suggest that this response is important in the control of renin release and renal vascular resistance during salt deprivation.

J. Clin. Invest. 112:76–82 (2003). doi:10.1172/JCI200318018.

Introduction

PGE₂ is a major product of PGH₂ derived from COX metabolism and is an important paracrine regulator of salt and water homeostasis in the kidney. In the renal cortex, COX-1 expression predominates in collecting duct, vascular tissue, and glomerular mesangial cells (1). In contrast, COX-2 is expressed and presumably mediates PG production in the macula densa (MD) and surrounding cortical thick ascending limb (cTAL) cells (2–5). MD cells are in direct contact with the vascular pole of the same glomerulus from which the filtrate originates. These cells sense changes in tubular [NaCl] and send signals that control preglomerular

vascular resistance and glomerular filtration rate in a process known as tubuloglomerular feedback. MD cells also control the release of renin from juxtaglomerular granular cells (3–6). COX-2-derived PGs may participate in MD-mediated control of juxtaglomerular function, particularly in high renin states such as low salt intake, loop diuretic treatment, and renovascular hypertension (4, 7). In particular, PGE₂ produced by MD cells has been suggested as the mediator of renin release induced by low luminal [NaCl] ([NaCl]_L) (3–10). In addition, PGE₂, as a potent vasodilator, may also modulate preglomerular vascular resistance (11–13) and tubuloglomerular feedback (14).

Although COX activity has been considered the key step in PG synthesis, metabolism of arachidonate by either COX-1 or COX-2 yields only the unstable intermediary PGH₂ (1). The subsequent fate of PGH₂ is dictated by coexpression of a PG synthase, which is the other key enzyme of PG synthesis. This enzyme is capable of converting PGH₂ to one of the prostanoid end products including PGE₂, PGF_{2α}, PGD₂, PGI₂, and thromboxane A₂ (11). Because of the limited number and inaccessibility of MD cells, as opposed to other nephron segments in the kidney (15), direct evidence for the expression of a PGE synthase in the MD and release of PGE₂ from intact MD cells is lacking. Inves-

Received for publication February 3, 2003, and accepted in revised form April 29, 2003.

Address correspondence to: János Peti-Peterdi, 865 Sparks Center, 1720 Seventh Avenue South, Birmingham, Alabama 35294, USA. Phone: (205) 934-5783; Fax: (205) 934-1147; E-mail: petjan@uab.edu.

Conflict of interest: The authors have declared that no conflict of interest exists.

Nonstandard abbreviations used: macula densa (MD); cortical thick ascending limb (cTAL); luminal NaCl concentration ([NaCl]_L); E-prostanoid receptor-1 (EP₁); EP₁-transfected HEK293 cells (HEK/EP₁ cells); intracellular [Ca²⁺]_i ([Ca²⁺]_i); microsome-associated PGE synthase (mPGES).

titating PGE₂ release from MD cells is important since a recent work actually questioned whether MD cells are equipped with PGE₂ synthetic machinery in the normal kidney (16).

The present studies used a cloned PG receptor (E-prostanoid receptor-1, or EP₁) to engineer a novel biosensor tool that could be used to detect local PGE₂ production. PGE₂ interacts with four different G protein-coupled E-prostanoid receptors designated EP₁, EP₂, EP₃, and EP₄ (11, 12). EP₂, EP₃, and EP₄ couple to adenylyl cyclase-associated G proteins to increase or decrease cAMP generation. In contrast, EP₁ is coupled to a Ca²⁺-signaling mechanism that is thought to involve inositol 1,4,5-triphosphate and diacylglycerol formation (11, 12). Therefore, EP₁-mediated effects of PGE₂ can be detected using fluorescence microscopy and a biosensor cell expressing EP₁ and loaded with a calcium fluorophore. The present studies were undertaken to determine, for the first time, whether the enzymatic machinery necessary for PGE₂ synthesis is expressed in MD cells from rabbits on a standard or low-salt diet, and to provide direct functional evidence for the control of MD PGE₂ release in response to low [NaCl]_L.

Methods

Materials. All materials were purchased from Sigma-Aldrich (St. Louis, Missouri, USA) unless otherwise stated. The COX-2 inhibitor SC58236 was generously provided by Peter Isakson and Karen Siebert, and the EP₁ blocker SC51322 was provided by Ed Drower (Pharmacia Research and Development, St. Louis, Missouri, USA). These compounds were dissolved in DMSO with a final DMSO concentration of below 0.1% (vol/vol).

Salt diet. Separate groups of New Zealand white rabbits (0.5–1.0 kg) were fed standard chow (8630, 0.3% sodium) or low salt (TD 90188, 0.01% sodium; both from Harlan Teklad, Madison, Wisconsin, USA) rabbit chow for a minimum of 1 week.

Tubule perfusion. Individual cTALs containing the MD segment with attached glomeruli were dissected from rabbit kidneys and perfused in vitro using methods similar to those described previously (17). For these biosensor studies, the cTAL containing the MD plaque was dissected away from the glomerulus, so that the basolateral surface of MD cells was accessible from the bath (Figure 1). The dissection solution was an isosmotic, low NaCl-containing Ringer's solution consisting of (in mM): 25 NaCl, 120 N-methyl-D-glucamine cyclamate (NMDG cyclamate), 5 KCl, 1 MgSO₄, 1.6 Na₂HPO₄, 0.4 NaH₂PO₄, 1.5 CaCl₂, 5 D-glucose, and 10 HEPES. Dissection was performed at 4°C. An individual cTAL was transferred to a chamber mounted on an inverted microscope. The tubule was kept in the low [NaCl] solution until it was cannulated and perfused with the same Ringer's solution except containing approximately 150 mM NaCl and no NMDG cyclamate. The bathing solution was the same 150 mM NaCl Ringer's solution and temperature was maintained at 37°C. When removing

NaCl from the tubular perfusate, NaCl was isosmotically substituted with NMDG cyclamate, KCl with potassium gluconate, and CaCl₂ with calcium gluconate to achieve an [NaCl] of 0 mM. Graduated changes in [NaCl]_L between 80 and 0 mM were achieved by isosmotically substituting NaCl with NMDG cyclamate.

Engineering the PGE₂ biosensor cells. HEK293 cells were stably transfected with the full-length mouse E-prostanoid receptor EP₁. EP₁ is activated specifically by nanomolar concentrations of PGE₂ and not by other prostanoids, and produces increases in intracellular calcium (18). HEK293 cells were transfected with a full-length mouse EP₁ cDNA cloned into pcDNA3.1 using SuperFect reagent (QIAGEN Inc., Valencia, California, USA) according to the manufacturer's instructions. Following transfection, cells were grown and maintained in media containing 400 µg/ml G418. Preliminary studies in monolayers of EP₁-transfected HEK293 cells (HEK/EP₁) exhibited increases in intracellular [Ca²⁺]_i ([Ca²⁺]_i) in response to nanomolar concentrations of PGE₂.

Following cannulation and perfusion of a microdissected cTAL/MD segment (Figure 1), a single fura-2-loaded HEK/EP₁ cell was gently positioned at the basolateral membrane surface of the MD segment using a separate holding pipette. Based on preliminary experiments, HEK/EP₁ [Ca²⁺]_i also appears to be responsive to ATP released by MD cells via endogenous purinergic receptors (19). To exclude the effect of purinergic receptors on the biosensor cell calcium signaling, 100 µM gadolinium, which not only inhibits MD ATP release (19, 20) but also blocks P₂X receptors (21), was always present in the bath.

Fluorescence microscopy. The [Ca²⁺]_i of HEK/EP₁ biosensor cells was measured with dual excitation wavelength fluorescence microscopy (Photon Technology Interna-




Figure 1

Photomicrograph of the biosensor technique for assessing PGE₂ release at the basolateral membrane of MD cells. The glomerulus (G) has been dissected away from the MD plaque. A single HEK/EP₁ cell is loaded with fura-2 and held with a pipette (arrowhead) positioned at the MD basolateral membrane surface while perfusing the intact cTAL.




Figure 2

Effects of exogenous PGE₂ (added to the bath) on [Ca²⁺]_i of an HEK/EP₁ biosensor cell. (a) Recordings of five different HEK/EP₁ cells demonstrate almost identical sensitivities of cells to 1 μM PGE₂. HEK/EP₁ [Ca²⁺]_i responses were prevented by the coadministration of an EP₁ blocker (SC51322). (b) Dose-response curve. Each data point represents the average of four cells studied.

tional, Lawrenceville, New Jersey, USA) using the fluorescent probe fura-2 (TEF LABS Inc., Austin, Texas, USA) as described previously (22). Fura-2 fluorescence was measured at an emission wavelength of 510 nm in response to excitation wavelengths of 340 and 380 nm alternated at a rate of 50 Hz by a computer-controlled chopper assembly. An adjustable photometer window was positioned over the single biosensor cell and emitted photons were detected by a Leitz photometer (Vashaw Scientific, Atlanta, Georgia, USA) that was modified for photon counting. Magnification was ×1,000 using an Olympus 100× UVFL lens. Autofluorescence-corrected ratios (340 nm/380 nm) were calculated at a rate of 5 points/s using Photon Technology International software. Biosensor cells were loaded with the dye by adding 10 μM fura-2 AM dissolved in dimethyl sulfoxide to the culture medium. Loading required approximately 60 minutes, after which fura-2 AM was removed. The 340/380 ratios were converted into [Ca²⁺]_i values as described before (22) using the methods and equation of Grynkiewicz et al. (23).

In situ hybridization. A 290-bp fragment of rabbit microsome-associated PGE synthase (mPGES) cDNA was obtained using RT-PCR with mPGES-selective primers based on mouse and human conserved sequences (5′-GCT GGT CAT CAA GAT GTA CG-3′ for sense; 5′-CCA GGT AGG CCA CGG TGT GT-3′ for antisense). This cDNA fragment was used to determine the expression of this isoform in MD. For in situ hybridization studies, tissues were fixed in 4% paraformaldehyde. Tissues were embedded in paraffin and 7-μm sections were cut. Prior to hybridization, sections were deparaffinized, refixed in paraformaldehyde, treated with proteinase K (20 μg/ml), washed with PBS, and treated with triethanolamine plus acetic anhydride (0.25% vol/vol). Finally sections were dehydrated in 100% ethanol. Antisense RNA was hybridized to the sections at 50–55°C for approximately 18 hours as described by Pelton et al. (24). Following hybridization, sections were washed at 50°C in 5× SSC plus 10 mM β-mercaptoethanol for 30 minutes. This was followed by a wash in 50% formamide,

2× SSC, and 100 mM β-mercaptoethanol for 60 minutes. Following additional washes in 10 mM Tris, 5 mM EDTA, and 500 mM NaCl, sections were treated with 10 μg/ml RNase at 37°C for 30 minutes, followed by another wash in 10 mM Tris, 5 mM EDTA, and 500 mM NaCl at 37°C. Sections were then washed twice in 2× SSC and twice in 0.1× SSC at 50°C. Slides were dehydrated with a series of graded ethanol containing 300 mM ammonium acetate. For detection of the hybridized probe, slides were dipped in photoemulsion (K5; Ilford Imaging UK Ltd., Knutsford, United Kingdom) diluted 1:1 with 2% glycerol/water and exposed for 7 days at 4°C. After development in Kodak D19 (Eastman Kodak Co. Scientific Imaging Systems, New Haven, Connecticut, USA), slides were counterstained with hematoxylin and eosin. Photomicrographs were taken using a Zeiss Axioskop (Carl Zeiss Inc., Thornwood, New York, USA) using both bright-field and dark-field optics.

Immunofluorescence. Rabbit kidneys were perfusion-fixed with paraformaldehyde and tissue sections were processed as described earlier (17). Sections were subjected to microwave antigen retrieval before staining and were blocked for 40 minutes with PBS-Tween containing 2% BSA to lower background fluorescence. Subsequent blocking with goat anti-rabbit Fab IgG (1:100; Jackson ImmunoResearch Laboratories Inc., West Grove, Pennsylvania, USA) was carried out for 40 minutes to reduce nonspecific binding when a rabbit polyclonal antibody (anti-mPGES) was used on rabbit tissue. After subsequent washings in PBS, tissues were treated overnight with either a goat polyclonal COX-2 antibody (C-20, 1:100; Santa Cruz Biotechnology Inc., Santa Cruz, California, USA) that recognizes COX-2 in various species (25) or with the affinity-purified rabbit polyclonal mPGES antibody (1:50; Cayman Chemical Co., Ann Arbor, Michigan, USA). After washing, there was a 40-minute incubation with Alexa 594-conjugated donkey anti-goat IgG (1:500; Molecular Probes Inc., Eugene, Oregon, USA) for COX-2 sections or with




Figure 3

Representative recordings of changes in HEK/EP₁ [Ca²⁺]_i/MD PGE₂ release. (a) Luminal NaCl removal in the low-salt group (in contrast to normal diet) induced significant PGE₂ release from MD, but not from distant cTAL cells, as evidenced by increases in HEK/EP₁ [Ca²⁺]_i. (b) HEK/EP₁ calcium responses (low-salt group) were inhibited by either luminal addition of furosemide, an Na:2Cl:K cotransport blocker, or by a selective COX-2 inhibitor (SC58236, 100 nM) or bath SC51322 (10 μM), an EP₁ blocker.




Figure 4 Biosensor HEK/EP₁ cell baseline [Ca²⁺]_i (a) and tubular NaCl removal-induced increases in [Ca²⁺]_i (Δ[Ca²⁺]_i) (b). A low-salt diet significantly elevated biosensor cell baseline [Ca²⁺]_i and Δ[Ca²⁺]_i upon luminal NaCl removal, compared with a normal diet. Biosensor cell calcium responses were inhibited by luminal addition of a selective COX-2 inhibitor (SC58236, 100 nM) or bath SC51322 (10 μM), an EP₁ blocker. (*n* = 6 in each group, **P* < 0.01 compared with normal diet).

Alexa 488–conjugated goat anti-rabbit IgG (1:500; Molecular Probes Inc.) for mPGES sections. Sections were mounted with VECTASHIELD medium containing DAPI for nuclear staining (Vector Laboratories Inc., Burlingame, California, USA). Tissue sections were examined with an Olympus IX70 inverted epifluorescence microscope using a 40× objective. Images were captured using a SenSys digital camera and IPLab Spectrum software equipped with a power microtome (Signal Analytics Corp., Vienna, Virginia, USA).

Statistical analysis. Data are expressed as mean ± SE. Statistical significance was tested using ANOVA. Significance was accepted at *P* < 0.05.

Results

Effects of exogenous PGE₂. Initial studies were performed to assess the sensitivity of individual HEK/EP₁ cells to exogenous PGE₂. Representative recordings (Figure 2a) demonstrate that individual HEK/EP₁ biosensor cells produced very similar, reversible, and dose-dependent (Figure 2b) elevations in [Ca²⁺]_i in response to exogenous PGE₂. Calcium signals were completely abolished (Figure 2a) when HEK/EP₁ cells were preincubated with the EP₁ blocker SC51322 (10 μM).

Effects of luminal NaCl removal. Since decreasing [NaCl]_L produces MD signals that stimulate renin release, initial studies were performed in which [NaCl]_L was decreased from 150 mM to 0 mM. Removing luminal NaCl resulted in a significant increase in [Ca²⁺]_i of single HEK/EP₁ cells positioned next to the MD basolateral membrane (Δ = 29.6 ± 12.8 nM, *n* = 6; Figure 3a and Figure 4b). This response was observed in preparations from animals kept on a low-salt diet for 1 week. In contrast, we were unable to detect significant elevations in HEK/EP₁ cell [Ca²⁺]_i (i.e., PGE₂ release) from MD plaques obtained from rabbits kept on a standard salt diet (Δ = 2.8 ± 2.8 nM, *n* = 6) (Figure 3a and Figure 4b). Also, this response was MD cell-specific, since the same maneuver caused no detectable change in [Ca²⁺]_i of

HEK/EP₁ cells placed next to distant cTAL cells (Figure 3a). Luminal NaCl removal-induced HEK/EP₁ calcium responses in the low-salt group were completely inhibited by luminal addition of a selective COX-2 inhibitor (SC58236) or bath SC51322, a blocker of the receptor EP₁ (Figure 3b and Figure 4b). It should also be noted that in low-salt diet studies there was a significant elevation in biosensor cell baseline [Ca²⁺]_i (Figure 4a).

Effects of various [NaCl]_L's and furosemide. Next we tested whether our model is sensitive in the physiological range of [NaCl]_L, and if more subtle changes in [NaCl]_L can cause HEK/EP₁ calcium responses. Figure 5 demonstrates that graduated changes in [NaCl]_L between 80 mM and 0 mM caused step-by-step increases in HEK/EP₁ cell fura-2 ratio. The most sensitive range of [NaCl]_L where the most significant changes occurred was between 40 mM and 20 mM (Figure 5a). The [NaCl]_L producing the half-maximal effect was around 30 mM (Figure 5b). Additional experiments examined whether HEK/EP₁ cell calcium signals are related to changes in MD NaCl transport. Furosemide (100 μM), a Na:2Cl:K cotransport blocker and loop diuretic, added to the luminal perfusate significantly inhibited the magnitude of HEK/EP₁ [Ca²⁺]_i increases (by 81.0% ± 7.6% compared with control; *n* = 5, *P* < 0.01) in response to 150 mM [NaCl]_L removal, as illustrated by the representative recording in Figure 3b. In the presence of furosemide, HEK/EP₁ cells were still sensitive to exogenous PGE₂ added to the bathing solution (Figure 3b). Similar confirmation of HEK/EP₁ biosensor cell reactivity was always performed when detecting no calcium response (such as in the normal salt group and with COX-2 inhibitor, not shown).

Expression of COX-2 and mPGES in rabbit MD. As illustrated in Figure 6a, there was very little staining of the rabbit MD with an antibody directed toward a C-terminal COX-2 peptide in animals on a normal salt diet. In contrast, the low-salt diet greatly increased the number of COX-2-immunoreactive cells in the MD (Figure 6, b and c). As with COX-2, there was a low level of mPGES expression in the MD in the normal kidney, but mPGES expression greatly increased in response to




Figure 5 Luminal [NaCl] dependency of MD PGE₂ release in cell preparations from rabbits fed a low-salt diet. (a) Representative recording shows step-by-step increases in HEK/EP₁ cell fura-2 ratio in response to graduated reductions in luminal [NaCl] from 80 to 0 mM. (b) Relationship between biosensor cell responses and luminal [NaCl]. Data points represent mean ± SE, *n* = 4, all low-salt preparations.




Figure 6 COX-2 (a–c) and mPGES (d and e) immunofluorescence of kidney cortex in control rabbits (a and d) and rabbits fed a low-salt diet (b and e). Number of COX-2-positive cells (c, $n = 6$ in each group, $*P < 0.05$ compared with normal salt diet) and mPGES expression in the MD significantly increased in response to the low-salt diet (LS) compared with the normal salt diet (NS). Nuclei are blue. In situ hybridization clearly demonstrated mPGES mRNA expression in MD cells (f).

the low-salt diet (Figure 6e). In addition to localizing the mPGES protein, in situ hybridization clearly demonstrated the expression of mPGES mRNA in rabbit MD cells (Figure 6f). These findings indicate that the complete enzymatic machinery necessary for PGH₂ synthesis and its conversion to PGE₂ is indeed present in MD cells, and more importantly, both COX-2 and mPGES expression are upregulated by low salt intake.

Discussion

The present studies used a modification of a recently established biosensor technique (26) to provide direct functional evidence, for the first time, that PGE₂ is released from the basolateral membrane of intact, perfused MD cells in a setting of reduced luminal salt delivery. This bioassay was originally used to measure cellular ATP release from the MD (19) using a rat pheochromocytoma clonal cell line (PC12) that endogenously expresses purinergic P₂X receptors. Instead of using a native cell line, we engineered a novel biosensor tool, HEK293 cells transfected with the mouse receptor EP₁, to selectively detect PGE₂. The fura-2 fluorescence ratio, and consequently [Ca²⁺]_i of individual HEK/EP₁ cells, increased in response to exogenous PGE₂ added to the bathing solution (Figure 2). The calcium responses of several individual HEK/EP₁ cells were almost identical, indicating similar EP₁ density and PGE₂ sensitivity (Figure 2a). Also, calcium responses were dose-dependent and were completely abolished by preincubation with the EP₁ blocker SC51322, which is consistent with EP₁-mediated calcium signaling (11, 12). It is important to note that nontransfected HEK293 cells did not increase [Ca²⁺]_i in response to PGE₂ but did respond to exoge-

nous ATP (data not shown), presumably through endogenously expressed purinergic receptors. This latter effect was completely blocked by gadolinium, which not only blocks P₂X receptors (21) but also inhibits ATP release (19, 20). The experiments measuring PGE₂ release were therefore carried out in the presence of gadolinium to exclude ATP purinergic signaling and its effects on cell [Ca²⁺]_i. Gadolinium did not affect PGE₂ detection by HEK/EP₁ cells (similar responses in Figure 2a and Figure 3b).

Because of the limited number and inaccessibility of MD cells (only 15–20 cells per nephron), it has not been possible to directly test for the release of prostanoids by MD cells. The HEK/EP₁ biosensor technique provides a unique tool allowing the detection of PGE₂ release from single MD plaques. As shown in Figure 1, a fura-2-loaded HEK/EP₁ cell can be positioned at the MD basolateral membrane surface, allowing real-time determination of basolateral PGE₂ release from an intact microperfused cTAL-MD segment.

Removing NaCl from the luminal perfusate significantly elevated [Ca²⁺]_i in HEK/EP₁ biosensor cells, indicating PGE₂ release from MD cells. Importantly, not only a single large change in luminal NaCl delivery (Figure 3a), but also more subtle reductions in [NaCl]_L in the physiological range (Figure 5a), caused detectable PGE₂ release from MD cells. The largest change in biosensor cell responses were observed between 40 mM and 20 mM [NaCl]_L, and the half-maximal effect was around 30 mM [NaCl]_L (Figure 5b). These findings are not only consistent with earlier work (27), but the data are essentially identical to the previously established [Cl]_{1/2}, the luminal chloride concentration producing a half-maximal renin secretory effect (27).

Additional studies using furosemide provided further evidence that biosensor cell calcium responses (and therefore MD PGE₂ release) were related to MD NaCl transport rather than a nonspecific effect of luminal NaCl. Inhibition of MD Na:2Cl:K cotransport with furosemide prevented [NaCl]_L-dependent increases in HEK/EP₁ [Ca²⁺]_i (Figure 3b), consistent with earlier findings that furosemide blocks MD NaCl transport dependency of renin release (27). Low distal tubular [NaCl] is associated with states of volume depletion and is a stimulus for MD signaling that increases synthesis and release of renin from juxtaglomerular granular cells (3–6). Basolateral MD PGE₂ release in response to decreased luminal salt delivery represents the physiological direction of signal transmission from MD cells into the juxtaglomerular apparatus area. Recent evidence has accumulated that this signaling mechanism involves MD-derived PGs produced by COX-2 in these cells (3–12). Consistent with this, we found that HEK/EP₁ calcium responses were inhibited by luminal addition of a selective COX-2 inhibitor (SC58236) or bath SC51322, a blocker of the PGE₂ receptor EP₁. These findings provide direct evidence that intact MD cells release PGE₂, consistent with PGE₂ synthesis reported in a recently developed putative MD

cell line (25). In our experiments, PGE₂ release appeared to be MD cell-specific, since applying this biosensor approach upstream using HEK/EP₁ cells placed next to distant cTAL cells produced no detectable signals (Figure 3a). However, it is possible that cTAL cells adjacent to the MD can also synthesize and release PGE₂ and that this contributed to the detected biosensor signal.

Interestingly, we were unable to detect [NaCl]_L-dependent PGE₂ release from MD cells obtained from animals kept on a standard salt diet. Only the low-salt diet caused consistent and significant [NaCl]_L-dependent increases in HEK/EP₁ cell [Ca²⁺]_i. A low-salt diet produces an experimental model characterized by a high renin state and is also known to increase the expression of COX-2 in MD cells (2). Upregulation of COX-2 in kidneys obtained from rabbits on a low-salt diet was indeed confirmed in the present studies by immunohistochemistry (Figure 6, a–c). Thus, similar to findings observed in other species (2, 28), the number of COX-2-immunoreactive cells in the MD plaque was substantially increased (by about sixfold) when animals were subjected to a low-salt diet for 1 week. This number actually nicely correlates to findings of biosensor studies in that only one of six normal salt preparations produced a biosensor calcium signal. We believe that the failure to show MD PGE₂ release in the majority of normal salt preparations is mainly technical: the chance that the biosensor cell is positioned next to a COX-2- and mPGES-expressing (and consequently PGE₂-producing) MD cell is much higher in the low-salt group. In other words, the sensitivity of this biosensor technique is probably limited to those MD cells in direct contact with the biosensor cell. Still, average MD PGE₂ release was much higher on the low-salt diet. If we compare Figure 4b with Figure 2b, the estimated amount of PGE₂ released from MD cells is around 50–60 nM on the low-salt diet and only 1–2 nM on normal salt intake.

In addition to COX-2, the present studies also demonstrated that mPGES, an enzyme capable of converting COX-2-derived PGH₂ to PGE₂ (29), was also present in the rabbit MD (Figure 6, d and f), and that mPGES expression was greatly upregulated in response to low salt intake (Figure 6e). Previous studies performed in rats failed to detect mPGES in microdissected MD (16), but whether this represents a species difference or a difference in the physiologic status of the animals studied remains unclear.

Not only did MD cells obtained from animals on a low-salt diet exhibit increased PGE₂ production in response to reduced [NaCl]_L, but baseline [Ca²⁺]_i was also increased in HEK/EP₁ cells placed adjacent to the MD (Figure 5a). Increased baseline HEK/EP₁ cell [Ca²⁺]_i is consistent with increased COX-2 and mPGES expression in MD cells and higher baseline PGE₂ production in these MD. NaCl_L removal caused an almost immediate elevation in biosensor cell [Ca²⁺]_i (Figure 3), suggesting a rapid activation of COX-2 and mPGES, with synthesis and release of PGE₂ from MD

cells. The rapidity of PGE₂ synthesis in response to low luminal Cl⁻ suggests rapid activation of an intracellular signaling cascade. At present the nature of this immediate signaling cascade, i.e., the link between reduced [NaCl]_L and COX-2 activation, remains uncertain. Cl⁻ transport-dependent activation of the MAPKs ERK and p38 has recently been suggested in cultured cells (25). However, tyrosine phosphorylation (30) could also stimulate MD COX-2 as well as activation of phospholipase A₂, and increased substrate availability could be the cause of acute PGE₂ generation (31). The biosensor technique should provide a novel tool that will help to clarify in detail the intracellular signaling mechanisms involved in MD NaCl transport and related PGE₂ production in response to low [NaCl]_L during salt deprivation.

In summary, these studies provided direct evidence for the synthesis and release of PGE₂ from intact, perfused MD cells. Our findings are consistent with the recent view (9) that renin release and renal vascular resistance are critically dependent on MD COX-2- and mPGES-derived PGE₂, particularly during volume depletion.

Acknowledgments

This work was supported by an NIH grant (5RO1 DK-32032) to P.D. Bell; an American Heart Association Scientist Development Grant (0230074N); an American Society of Nephrology Carl W. Gottschalk Research Scholar grant to J. Peti-Peterdi; a Hungarian Ministry of Welfare (ETT) grant and NATO Science Fellowships to J. Peti-Peterdi and L. Rosivall; and a Veterans Merit Award and NIH grant (RO1 DK-37097) to M.D. Breyer.

1. Smith, W.L., and Bell, T.G. 1978. Immunohistochemical localization of the prostaglandin-forming cyclooxygenase in renal cortex. *Am. J. Physiol.* **235**:F451–F457.
2. Harris, R.C., et al. 1994. Cyclooxygenase-2 is associated with the macula densa of rat kidney and increases with salt restriction. *J. Clin. Invest.* **94**:2504–2510.
3. Schnermann, J. 2001. Cyclooxygenase-2 and macula densa control of renin secretion. *Nephrol. Dial. Transplant.* **16**:1735–1738.
4. Harris, R.C., Wang, J.L., Cheng, H.F., Zhang, M.Z., and McKanna, J.A. 1998. Prostaglandins in macula densa function. *Kidney Int.* **67**:S49–S52.
5. Harris, R.C. 2000. Macula densa signalling—a potential role of cyclooxygenase-2 (COX-2)? *Nephrol. Dial. Transplant.* **15**:1504–1506.
6. Schnermann, J., and Briggs, J. 1985. Function of the juxtaglomerular apparatus: local control of glomerular hemodynamics. In *The kidney*. D.W. Seldin and G. Giebisch, editors. Raven Press. New York, New York, USA. 669–697.
7. Harris, R.C., and Breyer, M.D. 2001. Physiological regulation of cyclooxygenase-2 in the kidney. *Am. J. Physiol. Renal Physiol.* **281**:F1–F11.
8. Yang, T., et al. 2000. Renin expression in COX-2-knockout mice on normal or low-salt diets. *Am. J. Physiol. Renal Physiol.* **279**:F819–F825.
9. Castrop, H., Schweda, F., Schumacher, K., Wolf, K., and Kurtz, A. 2001. Role of renocortical cyclooxygenase-2 for renal vascular resistance and macula densa control of renin secretion. *J. Am. Soc. Nephrol.* **12**:867–874.
10. Traynor, T.R., Smart, A., Briggs, J.P., and Schnermann, J. 1999. Inhibition of macula densa-stimulated renin secretion by pharmacological blockade of cyclooxygenase-2. *Am. J. Physiol.* **277**:F706–F710.
11. Breyer, M.D., and Breyer, R.M. 2001. G protein-coupled prostanoid receptors and the kidney. *Ann. Rev. Physiol.* **63**:579–605.
12. Breyer, M.D., and Breyer, R.M. 2000. Prostaglandin E receptors and the kidney. *Am. J. Physiol. Renal Physiol.* **279**:F12–F23.
13. Chaudhari, A., Gupta, S., and Kirschenbaum, M.A. 1990. Biochemical evidence for PGI₂ and PGE₂ receptors in the rabbit renal preglomerular microvasculature. *Biochim. Biophys. Acta.* **1053**:156–161.
14. Schnermann, J., and Weber, P.C. 1982. Reversal of indomethacin-induced inhibition of tubuloglomerular feedback by prostaglandin

- infusion. *Prostaglandins*. **24**:351–361.
15. Farman, N., Pradelles, P., Bonvalet, J.P. 1987. PGE₂, PGF₂ alpha, 6-keto-PGF₁ alpha, and Tx_{B2} synthesis along the rabbit nephron. *Am. J. Physiol.* **252**:F53–F59.
 16. Vitzthum, H., Abt, I., Einhellig, S., and Kurtz, A. 2002. Gene expression of prostanoid forming enzymes along the rat nephron. *Kidney Int.* **62**:1570–1581.
 17. Peti-Peterdi, J., et al. 2000. Macula densa Na:H exchange activities mediated by apical NHE2 and basolateral NHE4 isoforms. *Am. J. Physiol.* **278**:F452–F463.
 18. Narumiya, S., Sugimoto, Y., and Ushikubi, F. 1999. Prostanoid receptors: structures, properties, and functions. *Physiol. Rev.* **79**:1193–1226.
 19. Bell, P.D., Lapointe, J.Y., Sabirov, R., Hayashi, S., and Okada, Y. 2000. Maxi-chloride channel in macula densa cells: possible pathway for ATP release. *FASEB J.* **14**:A134. (Abstr.)
 20. Sabirov, R.Z., Dutta, A.K., and Okada, Y. 2001. Volume-dependent ATP-conductive large-conductance anion channel as a pathway for swelling-induced ATP release. *J. Gen. Physiol.* **118**:251–266.
 21. Roman, R.M., Feranchak, A.P., Davison, A.K., Schwiebert, E.M., and Fitz, J.G. 1999. Evidence for Gd(3+) inhibition of membrane ATP permeability and purinergic signaling. *Am. J. Physiol.* **277**:G1222–G1230.
 22. Peti-Peterdi, J., and Bell, P.D. 1999. Cytosolic [Ca²⁺] signaling pathway in macula densa cells. *Am. J. Physiol.* **277**:F472–F476.
 23. Grynkiewicz, G., Ponie, M., and Tsien, R.Y. 1985. A new generation of Ca²⁺ indicators with greatly improved fluorescence properties. *J. Biol. Chem.* **260**:3440–3450.
 24. Pelton, R.W., Johnson, M.D., Perrett, E.A., Gold, L.I., and Moses, H.L. 1991. Expression of transforming growth factor-beta 1, -beta 2, and -beta 3 mRNA and protein in the murine lung. *Am. J. Respir. Cell Mol. Biol.* **5**:522–530.
 25. Yang, T., et al. 2000. Low chloride stimulation of prostaglandin E₂ release and cyclooxygenase-2 expression in a mouse macula densa cell line. *J. Biol. Chem.* **275**:37922–37929.
 26. Hazama, A., Hayashi, S., and Okada, Y. 1998. Cell surface measurements of ATP release from single pancreatic beta cells using a novel biosensor technique. *Pflugers Arch.* **437**:31–35.
 27. He, X.R., Greenberg, S.G., Briggs, J.P., and Schnermann, J. 1995. Effects of furosemide and verapamil on the NaCl dependency of macula densa-mediated renin secretion. *Hypertension*. **26**:137–142.
 28. Theilig, F., et al. 2002. Epithelial COX-2 expression is not regulated by nitric oxide in rodent renal cortex. *Hypertension*. **39**:848–853.
 29. Jakobsson, P.J., Thoren, S., Morgenstern, R., and Samuelsson, B. 1999. Identification of human prostaglandin E synthase: a microsomal, glutathione-dependent, inducible enzyme, constituting a potential novel drug target. *Proc. Natl. Acad. Sci. U. S. A.* **96**:7220–7225.
 30. Parfenova, H., Balabanova, L., and Leffler, C.W. 1998. Posttranslational regulation of cyclooxygenase by tyrosine phosphorylation in cerebral endothelial cells. *Am. J. Physiol.* **274**:C72–C81.
 31. Mangat, H., Peterson, L.N., and Burns, K.D. 1997. Hypercalcemia stimulates expression of intrarenal phospholipase A₂ and prostaglandin H synthase-2 in rats. Role of angiotensin II AT₁ receptors. *J. Clin. Invest.* **100**:1941–1950.