

Research

Open Access

Ethnomedicines used in Trinidad and Tobago for urinary problems and diabetes mellitus

Cheryl A Lans*

Address: BCICS, University of Victoria, British Columbia, V8W 2Y2, Canada

Email: Cheryl A Lans* - cher2lans@netscape.net

* Corresponding author

Published: 13 October 2006

Received: 07 August 2006

Journal of Ethnobiology and Ethnomedicine 2006, **2**:45 doi:10.1186/1746-4269-2-45

Accepted: 13 October 2006

This article is available from: <http://www.ethnobiomed.com/content/2/1/45>

© 2006 Lans; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: This paper is based on ethnobotanical interviews conducted from 1996–2000 in Trinidad and Tobago with thirty male and female respondents.

Methods: A non-experimental validation was conducted on the plants used for urinary problems and diabetes mellitus: This is a preliminary step to establish that the plants used are safe or effective, to help direct clinical trials, and to inform Caribbean physicians of the plants' known properties to avoid counter-prescribing.

Results: The following plants are used to treat diabetes: *Antigonon leptopus*, *Bidens alba*, *Bidens pilosa*, *Bixa orellana*, *Bontia daphnoides*, *Carica papaya*, *Catharanthus roseus*, *Cocos nucifera*, *Gomphrena globosa*, *Laportea aestuans*, *Momordica charantia*, *Morus alba*, *Phyllanthus urinaria* and *Spiranthes acaulis*. *Apium graveolens* is used as a heart tonic and for low blood pressure. *Bixa orellana*, *Bontia daphnoides*, *Cuscuta americana* and *Gomphrena globosa* are used for jaundice. The following plants are used for hypertension: *Aloe vera*, *Annona muricata*, *Artocarpus atilis*, *Bixa orellana*, *Bidens alba*, *Bidens pilosa*, *Bontia daphnoides*, *Carica papaya*, *Cecropia peltata*, *Citrus paradisi*, *Cola nitida*, *Crescentia cujete*, *Gomphrena globosa*, *Hibiscus sabdariffa*, *Kalanchoe pinnata*, *Morus alba*, *Nopalea cochinellifera*, *Ocimum campechianum*, *Passiflora quadrangularis*, *Persea americana* and *Tamarindus indicus*.

The plants used for kidney problems are *Theobroma cacao*, *Chamaesyce hirta*, *Flemingia strobilifera*, *Peperomia rotundifolia*, *Petiveria alliacea*, *Nopalea cochinellifera*, *Apium graveolens*, *Cynodon dactylon*, *Eleusine indica*, *Gomphrena globosa*, *Pityrogramma calomelanos* and *Vetiveria zizanioides*. Plants are also used for gall stones and for cooling.

Conclusion: *Chamaesyce hirta*, *Cissus verticillata*, *Kalanchoe pinnata*, *Peperomia* spp., *Portulaca oleraceae*, *Scoparia dulcis*, and *Zea mays* have sufficient evidence to support their traditional use for urinary problems, "cooling" and high cholesterol.

Eggplant extract as a hypocholesterolemic agent has some support but needs more study. The plants used for hypertension, jaundice and diabetes that may be safe and justify more formal evaluation are *Annona squamosa*, *Aloe vera*, *Apium graveolens*, *Bidens alba*, *Carica papaya*, *Catharanthus roseus*, *Cecropia peltata*, *Citrus paradisi*, *Hibiscus sabdariffa*, *Momordica charantia*, *Morus alba*, *Persea americana*, *Phyllanthus urinaria*, *Tamarindus indicus* and *Tournefortia hirsutissima*. Several of the plants are used for more than one condition and further trials should take this into account.

Background

Latin American and Caribbean policy makers have recognised that many problems in primary health care are due to lack of knowledge and sensitivity to local health practices, and to the economic and cultural factors associated with these practices [1]. However available herbal products have no clear statement of content; or medically related information on the package labels, and have not been validated or certified by any recognised body. This concerns consumers (potential and actual) and medical practitioners who may unknowingly counter-prescribe these herbal products [1]. Not all of the plants reported to be useful are harmless. These considerations (among others) underlie the study of ethnomedicinal plants used in Trinidad and Tobago from 1995 to 2000 [1]. This paper focuses on the plants used for diabetes, urinary problems and related problems. The World Health Organization has projected that the incidence of type 2 diabetes in Trinidad and Tobago will increase from 70,000 in the year 2000 to 89,000 by the year 2010 [2]. Type 1 diabetes typically affects less than 10% of patients [3]. Both types of diabetes involve hyperglycemia and several grave complications resulting from long-term hyperglycemia [3]. Control of the glucose levels in the blood is the most effective treatment. Long term reductions in hyperglycemia reduce the likelihood of developing microvascular and macrovascular complications [3]. Type 2 diabetes mellitus has both genetic and environmental components [2]. Previous studies (including Caribbean studies) of offspring or first-degree relatives of patients with type 2 diabetes have confirmed that insulin resistance and/or hyperinsulinaemia is the antecedent biochemical marker for developing diabetes in middle age [2]. First-degree relatives of diabetic patients are more likely to develop diabetes and also have an increased cluster of biochemical metabolic syndrome risk factors [2]. Although insulin resistance is independently associated with obesity, it is more severe in obese patients (80% of type 2 diabetic patients in the West) [3]. Diabetic patients typically have delayed or impaired wound healing, and may develop chronic ulcers [4]. Diabetic ulcers of the lower limbs and feet are associated with high morbidity and amputation is a common treatment. Peripheral neuropathy and peripheral vascular disease are possible underlying factors in diabetic wound formation, but dermal atrophy is a possible contributing factor [4]. Reduced fibroblast growth, increased expression of matrix-degrading matrix metalloproteinases (MMPs), and decreased matrix synthesis are consequences of chronic vascular disease in diabetic skin as well as in other situations characterised by peripheral vascular disease [4]. Atrophic skin is more likely to develop wounds than healthy skin. In addition, insufficient extracellular matrix production during the proliferative phase of wound repair contributes to poor healing [4].

A 1990 Caribbean study on folk medicine used for diabetes mellitus found no difference between non-users of informal medication and those who used it in addition to, or in replacement of, formal medication to control diabetes mellitus in Jamaica [5]. The authors found that teas made from periwinkle (*Catharantus* species) and rice biters (*Andrographis paniculata*) interfered with the control of diabetes mellitus. In a previous study by one of the authors, with more severe cases of diabetes, formal medication gave better control of the diabetes than folk medicine teas [6].

A more recent study conducted in Trinidad and Tobago examined 622 people with diabetes mellitus. Herbal remedies for diabetes were used by 152 (24%) of patients [7]. Caraailli (*Momordica charantia*), aloes (*Aloe vera*), olive bush (*Bontia daphnoides*) and seed-under-leaf (*Phyllanthus urinaria*) were the plants most frequently used. Patients who reported burning or numbness in the feet or feelings of tiredness, weakness, giddiness or dizziness used folk medicines more frequently than those who reported other symptoms [7]. Subsequent studies by the same authors have discarded folk medicine and focussed on the standards of conventional medical care for diabetes [8-10]. In a 2004 study they found that diabetes was associated with low income status and worse health status and more frequent expenditure on medical services [9]. Although the authors of this study found that the control group had access to health insurance this access is not common in Trinidad. In an earlier paper the same authors estimated the annual cost of hospital admissions with diabetes at TT10.66 million (UK 1.24 million pounds) [10], type 2 diabetes is a costly and largely avoidable problem.

In the late 1980s an IDRC-funded study examined the prevalence of diabetes and hypertension in women in Trinidad [11]. They screened 4013 women older than 25 years in 3152 households and 769 had been diagnosed with diabetes mellitus or hypertension. They concluded that the women's conditions were not managed adequately by the health system and that they should assume greater responsibility for managing their own health [11]. It is therefore puzzling that little research has been conducted on the medicinal plants used for diabetes and their efficacy.

Instead studies have examined ethnic or gender differences in prevalence of diabetes mellitus or compliance with conventional treatment protocols [12]. Finding alternatives is important because 10 – 20% of the population is affected. The leading causes of death and illness for the past thirty years in Trinidad and Tobago have been cardiovascular disease, cerebrovascular disease, cancer, hypertension and diabetes mellitus [13]. In a 2002 study, 2117 randomly selected patients from 35 randomly selected

government health centres in Trinidad were found to be suffering from diabetes-related foot problems. Of these, 49% had burning or numbness in the feet, 1241 (59%) self-reported visual impairment and 12% had foot ulceration of whom 92 (4%) had major or minor amputation [14].

This author conducted ethnoveterinary research in Trinidad and Tobago from 1995 to 2000. During this research many ethnomedicinal plant uses were discovered that did not have comparative uses for animals. This paper deals with the twenty-two plants used for hypertension, the four plants used for jaundice and the fourteen used for diabetes. It also presents the forty-five plants for urinary problems, "cooling" and high cholesterol levels.

A substantial body of research published outside the Caribbean since 2000 has provided sufficient data to conduct a preliminary evaluation of the researched plants and this evaluation is presented in the discussion section of this paper [15]. Several (approx. 30) of the plant uses have already been recorded in an ethnomedicinal study in one rural community in the northern range of Trinidad [16] and in the study on diabetes mellitus mentioned previously [7]. Several older studies referred to in the discussion section of the paper were reviewed previously and their original references will not be given for space reasons [1].

The Trinidadian hot/cold system

Caribbean folk medicine is a marriage- α -cinq:European folkmedicine, scientific medicine, African-based practices, Amerindianmedicine and Indian-based medicine, a product of inter-group borrowing or medical syncretism [1]. Activities, food and medicines are classified in various ethnomedicinal systems as hot or cold. The Trinidadian hot/cold system is not humoral in the sense that balance must be established between hot and cold, it is cathartic in that remedies are taken to remove heat from the system [1]. Some researchers consider that the hot-cold concept of health and illness is absent in Spanish folk medicine and did not exist at the folk level in the past [1]. These beliefs are now widespread in Latin America and the Caribbean, and according to G.M. Foster were derived from the *élite* and scholarly Hippocratic-galenic traditions that were brought to the Spanish colonies by Spanish physicians and clergy [17,18]. The hot-cold valence refers to the traditional belief that heat opens the body and facilitates the blood's free flow, whereas cold causes the blood to stop flowing and clog the arteries, veins and womb [1,17,18]. Heat comes from the sun, work, sleeping, burns, cooking, and reproductive activities. Linked to the hot/cold dichotomy is a system of blood beliefs where an excess or lack of cold or heat in the body through exposure or diet causes illness. Blood then becomes 'bad' or dirty [1]. Female

Warao herbalists in Eastern Venezuela also use the concept of bad blood [19].

Teas are used for 'cooling' if there is too much 'heat' in the body. Cooling teas are used prophylactically when they are taken to keep the body healthy by cooling the 'system', or the bladder, meaning that they remove the 'heat' or impurities in the system [1,17-20]. Cooling teas become treatment when they are taken for undiagnosed or unspecified illnesses or when feeling unwell. Purges reduce the heat further and 'clean the blood'. There are also 'hot' plants to stimulate the blood or to treat 'cold' illnesses, and 'hot' external applications like 'soft candle' 'grated nutmeg' and hot poultices [1,17-20]. Medicines are administered in accordance with the identity between cause and effect. Expulsion of disease-causing impurities is the primary mechanism by which bodily equilibrium is restored. Folk medicines achieve cures through 'bitterness', 'cutting', 'cooling', 'building', 'purging or washing out', and 'drawing out' [1,17-20].

Methods

This study adhered to the research guidelines and ethical protocols of Wageningen University in the Netherlands. Thirty respondents, ten of whom were male were interviewed from September 1996 to September 2000. The respondents were obtained by snowball sampling, and were found in thirteen different sites, 12 in Trinidad and one in Tobago. Snowball sampling was used because there was no other means of identifying respondents and because the research covered the entire island-area rather than concentrating on one village. The chief objective of the sampling method was to identify knowledgeable respondents; no priority was given to extrapolating the data to the wider population to establish prevalence of use. No statistical analysis was applied to the data.

Twenty respondents were interviewed once, the other ten (who were healers) were interviewed three or four times. Healers were also asked to reconstruct the circumstances and contexts of the plant uses so that the means of administration of the plants could be identified. No interview schedule of questions was used but a more qualitative, conversational technique. Plants were collected when available to verify that the common names used by each respondent were the same in each ethnic group as those recorded in the literature. The majority of the plants were identified at the Herbarium of the University of the West Indies but voucher samples were not deposited. This ethnomedicinal study was part of a larger research project on ethnoveterinary medicine; other data collecting techniques were used in the larger study [1].

Validation of practices

A preliminary validation of ethnomedicinal practices is considered a preliminary step to establish which plants are safe or effective and which uses should be discontinued. It also ensures that clinical trials are not wasted on plants that are used for cultural or religious reasons. This is important not only because of the waste of time, money and energy, but because negative results can lead to the discrediting of further effort [1].

Anthropologists like Posey [21] and Hastrup and Elsass [22] claim that anthropologists should not decide whether indigenous beliefs and practices are or are not scientific as this has colonial overtones. Other anthropologists claim that indigenous knowledge systems represent the cultural dimension of development and cannot be reduced to the empirical knowledge that they contain [23]. These anthropological reservations have some value; however validation of traditional medicines is important since most scientists will not use local medicines without some form of validation and they have discouraged the use of these medicines leading to their disuse and eventual loss [1]. A framework of indigenous/ethnoveterinary knowledge that can interface with science and technology is more likely to influence scientific research agendas and development work [1,24].

The validation of the remedies was conducted with a non-experimental method [25]. This method consists of:

1. obtaining an accurate botanical identification,
2. determining whether the folk data can be understood in terms of bioscientific concepts and methods,
3. searching the chemical/pharmaceutical/pharmacological literature for the plant's known chemical constituents and to determine the known physiological effects of either the crude plant, related species, or isolated chemical compounds that the plant is known to contain.

This information is used to assess whether the plant use is based on empirically verifiable principles or whether symbolic aspects of healing are of greater relevance. If ethnobotanical data, phytochemical and pharmacological information supports the folk use of a plant species it can be grouped into the validation level with the highest degree of confidence.

Four levels of validity were established [25]:

1. If no information supports the use it indicates that the plant may be inactive; or no research has been done on the plant.

2. A plant (or closely related species of the same genus), which is used in geographically or temporally distinct areas in the treatment of similar illnesses, attains the lowest level of validity, if no further phytochemical or pharmacological information validates the popular use. Use in other areas increases the likelihood that the plant is active against the illness.

3. If in addition to the ethnobotanical data, phytochemical or pharmacological information also validates the use in Trinidad and Tobago, the plant may exert a physiological action on the patient and is more likely to be effective than those at the lowest level of validity.

4. If ethnobotanical, phytochemical and pharmacological data supports the folk use of the plant, it is grouped in the highest level of validity and is most likely an effective remedy.

Results

Plants used for urinary problems

Forty-five plants were used for urinary problems, "cooling" and high cholesterol levels. The term "stoppage of water" means urinary retention. Four plants were used for bladder problems: *Costus scaber*, *Pilea microphylla*, *Kalanchoe pinnata* and *Cocos nucifera*. Two plants were used for high cholesterol levels: *Solanum melongena* and *Portulaca oleraceae*. *Bauhinia cumanensis*/*Bauhinia excisa* and *Capraria biflora* were used for gall stones.

Twenty-four plants were used for "cooling": *Musa* species, *Begonia humilis*, *Bontia daphnoides*, *Cissus verticillata*, *Coleus aromaticus*, *Commelina elegans*, *Cuscuta americana*, *Cyperus rotundus*, *Desmodium canum*, *Entada polystachya*, *Justicia pectoralis*, *Momordica charantia*, *Peperomia pellucida*, *Ruellia tuberosa*, *Sansevieria guineensis*, *Stachytarpheta jamaicensis*, *Scoparia dulcis*, *Cassia alata*, *Capraria biflora*, *Kalachoe pinnata*, *Mimosa pudica*, *Lepianthes peltata*, *Tournefortia hirsutissima* and *Solanum americanum*. *Bauhinia cumanensis*, *Bauhinia excisa* and *Capraria biflora* were used for gall stones, while *Hibiscus sabdariffa* was used to "clean the liver and blood".

The following fourteen plants were used for kidney and other urinary problems: *Kalachoe pinnata*, *Mimosa pudica*, *Chamaesyce hirta*, *Flemingia strobilifera*, *Peperomia rotundifolia*, *Petiveria alliacea*, *Nopalea cochinellifera*, *Apium graveolens*, *Cynodon dactylon*, *Zea mays*, *Theobroma cacao*, *Lepianthes peltata*, *Eleusine indica*, *Gomphrena globosa*, *Ptyrogramma calomelanos* and *Vetiveria zizanioides*.

The results are summarized in Table 1.

Table 1: Ethnomedicinal plants used for "cooling", high cholesterol and urinary problems in Trinidad and Tobago

Scientific name	Family	Common name	Part used	Use
<i>Apium graveolens</i>	Apiaceae	Celery		Kidney tonic
<i>Bauhinia cumanensis /excisa</i>	Fabaceae	Monkey step	Bark	Gall stones
<i>Begonia humilis</i>	Begoniaceae	Lozeille		Cooling
<i>Bontia daphnoides</i>	Myoporaceae	Olive bush	Leaves	Cooling
<i>Capraria biflora</i>	Scrophulariaceae	Du thé pays	Leaves	Gall stones, cooling
<i>Cassia alata</i>	Fabaceae	Senna	Leaves	Cooling with cloves & ginger
<i>Chamaesyce hirta</i>	Euphorbiaceae	Mal nommée		Kidney problems
<i>Cissus verticillata</i>	Vitaceae	Blister bush	Vine	Cooling
<i>Cocos nucifera</i>	Arecaceae	Coconut	Root	Bladder stones
<i>Coleus aromaticus</i>	Lamiaceae	Spanish thyme	Leaves	Cooling
<i>Commelina elegans</i>	Commelinaceae	Water grass	Plant	Cooling
<i>Costus scaber</i>	Zingiberaceae	Wild cane		Cleans bladder
<i>Cuscuta americana</i>	Convolvulaceae	Love vine	Vine	Cooling
<i>Cynodon dactylon</i>	Poaceae	Dube		Stoppage of water
<i>Cyperus rotundus</i>	Cyperaceae	Nut grass		Cooling
<i>Desmodium canum</i>	Fabaceae	Sweet heart bush	Plant	Cooling
<i>Eleusine indica</i>	Poaceae	Dead man's grass	Root, Leaves	Urinary
<i>Entada polystachya</i>	Fabaceae	Mayoc chapelle		Cooling
<i>Flemingia strobilifera</i>	Fabaceae	Kidney bush		Kidney problems
<i>Gomphrena globosa</i>	Amaranthaceae	Bachelor button	Leaves	Urinary problems
<i>Hibiscus sabdariffa</i>	Malvaceae	Sorrel	Flower & seed	Cleans liver and blood
<i>Justicia pectoralis</i>	Acanthaceae	Carpenter grass	Leaves	Cooling
<i>Kalachoe pinnata</i>	Crassulaceae	Wonder of the world	Leaves	Cooling, Bladder stones
<i>Lepianthes peltata</i>	Piperaceae	Lani bois	Leaves	Tea
<i>Mimosa pudica</i>	Fabaceae	Ti marie, mese marie		Cooling, Kidney problems
<i>Momordica charantia</i>	Cucurbitaceae	Caraaili	Vine	Cooling
<i>Musa species</i>	Musaceae	Banana	Dry leaf	Boil for cooling
<i>Nopalea cochinellifera</i>	Cactaceae	Rachette	Joint	Kidney stones
<i>Peperomia rotundifolia</i>	Piperaceae	Giron fleur, mowon		Kidney problems
<i>Peperomia pellucida</i>	Piperaceae	Shining bush		Cooling
<i>Petiveria alliacea</i>	Phytolaccaceae	Mapourite, kudjuruk		Kidney problems
<i>Pilea microphylla</i>	Urticaceae	Du thé bethelmay	Leaves	Bladder cleanser
<i>Pityrogramma calomelanos</i>	Pteridaceae	Fern		Urinary problems
<i>Portulaca oleraceae</i>	Portulacaceae	Pussley	Plant	Cholesterol, shortness of breath
<i>Ruellia tuberosa</i>	Acanthaceae	Miny root	Root	Cooling
<i>Sansevieria guineensis</i>	Agavaceae	Langue bouef, lash	Leaves	Cooling
<i>Scoparia dulcis</i>	Scrophulariaceae	Sweet broom	Plant	Cooling for babies
<i>Solanum americanum</i>	Solanaceae	Agouma, gouma	Plant	Cooling, provides iron
<i>Solanum melongena</i>	Solanaceae	Melongene	Fruit	Cholesterol
<i>Stachytarpheta jamaicensis</i>	Verbenaceae	Vervine	Leaves	Cooling
<i>Theobroma cacao</i>	Sterculiaceae	Cocoa	Core	Eat for urinary problems
<i>Tournefortia hirsutissima</i>	Boraginaceae	Chigger bush	Leaves	Cooling
<i>Vetiveria zizanioides</i>	Poaceae	Vetivert		Urinary problems
<i>Zea mays</i>	Poaceae	Corn silk	Stigma	Diuretic

Plants used for high blood pressure, diabetes and jaundice

Twenty-nine plants are used for diabetes and hypertension including four for jaundice.

Multiple-plant remedies are used for several conditions including one used for jaundice which combined white bachelor button (*Gomphrena globosa*), olive bush (*Bontia*

daphnoides), small white vere michelle (unidentified), and fine-stemmed rather than thick-stemmed love vine (*Cuscuta americana*).

The following plants are used to treat diabetes: *Antigonon leptopus*, *Bidens alba*, *Bidens pilosa*, *Bontia daphnoides*, *Carica papaya*, *Gomphrena globosa*, *Bixa orellana*, *Catharanthus roseus*, *Cocos nucifera*, *Laportea aestuans*, *Momordica charantia*, *Morus alba*, *Phyllanthus urinaria* and *Spiranthes acaulis*.

Apium graveolens is used as a heart tonic and the following plants are used for hypertension: *Aloe vera*, *Annona muricata*, *Artocarpus altilis*, *Bixa orellana*, *Bidens alba*, *Bidens pilosa*, *Bontia daphnoides*, *Carica papaya*, *Cecropia peltata*, *Citrus paradisi*, *Cola nitida*, *Crescentia cujete*, *Gomphrena globosa*, *Hibiscus sabdariffa*, *Kalanchoe pinnata*, *Nopalea cochinellifera*, *Morus alba*, *Ocimum campechianum*, *Passiflora quadrangularis*, *Persea americana* and *Tamarindus indicus*. Low blood pressure is treated with *Apium graveolens*.

Jaundice is treated with the following plants (many of which are also listed above): *Bixa orellana*, *Bontia daphnoides*, *Gomphrena globosa* and *Cuscuta americana*.

The ethnomedicinal plants used in Trinidad and Tobago for diabetes are summarised in Table 2.

Discussion

Non-experimental validation

For each species or genus the ethnomedicinal uses in other countries was examined but will not be presented in the table due to space constraints. Only the most relevant clinical trials are presented. The non-experimental validation is summarised in Tables 3 and 4. Table 4 is an additional file (see Additional file 1).

Conclusion

From the 1930s the impact of western medicine in the Caribbean has been that of a dominant paradigm that was not totally accepted, but which offered elements that were selectively appropriated in a process of indigenisation [117,118]. The concepts of 'structural superiority' and 'functional strength' imply that western medicine acquired elite status because of its ability to control diseases (or suppress symptoms), while the folk medicinal system retained functional strength because it was more accessible and available to those isolated communities that existed well into the twentieth century [1,117,118]. However this explanatory model, western medicine, has become the dominant medical system; the main means of

Table 2: Ethnomedicinal plants used for high blood pressure, jaundice and diabetes in Trinidad and Tobago

Scientific name	Family	Common name	Plant part used	Use
<i>Aloe vera</i>	Liliaceae	Aloes	Leaf gel	Hypertension
<i>Annona muricata</i>	Annonaceae	Soursop	Leaves	Hypertension
<i>Antigonon leptopus</i>	Polygonaceae	Coralita	Vine	Diabetes
<i>Apium graveolens</i>	Apiaceae	Celery		Heart tonic, Low blood pressure
<i>Artocarpus altilis</i>	Moraceae	Breadfruit	Leaves	Hypertension
<i>Bidens alba/Bidens pilosa</i>	Asteraceae	Needle grass	Leafy branch	Hypertension, Diabetes
<i>Bixa orellana</i>	Bixaceae	Roucou	Leaves, root	Hypertension, Diabetes, Jaundice
<i>Bontia daphnoides</i>	Myoporaceae	Olive bush	Leaves	Diabetes, Jaundice, Hypertension
<i>Carica papaya</i>	Caricaceae	Papaya	Green fruit	Hypertension, Diabetes
<i>Catharanthus roseus</i>	Apocynaceae	White Periwinkle		Diabetes
<i>Cecropia peltata</i>	Cecropiaceae	Bois canôt	Leaves	Hypertension
<i>Citrus paradisi</i>	Rutaceae	Grapefruit	Peel	Hypertension
<i>Cocos nucifera</i>	Arecaceae	Coconut	Shell, flower	Diabetes
<i>Cola nitida</i>	Sterculiaceae	Obie seed	Seed	Hypertension
<i>Crescentia cujete</i>	Bignoniaceae	Calabash	Leaves	Hypertension
<i>Cuscuta americana</i>	Convolvulaceae	Love vine	Vine	Jaundice
<i>Gomphrena globosa</i>	Amaranthaceae	Bachelor button	Leaves	Jaundice, Diabetes, Hypertension
<i>Hibiscus sabdariffa</i>	Malvaceae	Sorrel	Leaf	Hypertension
<i>Kalanchoe pinnata</i>	Crassulaceae	W/world	Leaf	Hypertension
<i>Laportea aestuans</i>	Urticaceae	Red stinging nettle	Leaves	Diabetes
<i>Momordica charantia</i>	Cucurbitaceae	Carailli		Diabetes
<i>Morus alba</i>	Moraceae	Pawi bush		Diabetes, Hypertension
<i>Nopalea cochinellifera</i>	Cactaceae	Rachette	Joint	Hypertension
<i>Ocimum campechianum</i>	Lamiaceae	Ti bom	Leaves	Hypertension
<i>Passiflora quadrangularis</i>	Passifloraceae	Barbadine	Leaves	Hypertension
<i>Persea americana</i>	Lauraceae	Avocado	Leaf	Hypertension
<i>Phyllanthus urinaria</i>	Euphorbiaceae	Red seed under leaf		Diabetes
<i>Spiranthes acaulis</i>	Orchidaceae	Lapsogen		Early-stage diabetes
<i>Tamarindus indicus</i>	Fabaceae	Tamarind	Seed	Hypertension

Table 3: Validation of plants used for "cooling", high cholesterol and urinary problems in Trinidad and Tobago

Scientific name	Common name	Use	Validation score
<i>Apium graveolens</i>	Celery	Kidney tonic	3 for pain
<i>Bauhinia cumanensis /excisa</i>	Monkey step	Gall stones	3 for pain
<i>Begonia humilis</i>	Lozeille	Cooling	2
<i>Bontia daphnoides</i>	Olive bush	Cooling	2
<i>Capraria biflora</i>	Du thé pays	Gall stones, cooling	3 for pain
<i>Cassia alata</i>	Senna	Cooling with cloves and ginger	2
<i>Chamaesyce hirta</i>	Malomay, Mal nommée	Kidney problems	3 diuretic; 3 sedative
<i>Cissus verticillata</i>	Blister bush	Cooling	3 diarrhoea; 3 gastric ulcers
<i>Cocos nucifera</i>	Coconut	Bladder stones	2
<i>Coleus aromaticus</i>	Spanish thyme	Cooling	2
<i>Commelina elegans</i>	Water grass	Cooling	2
<i>Costus scaber</i>	Wild cane	Cleans bladder	data needed
<i>Cuscuta americana</i>	Love vine	Cooling	2
<i>Cynodon dactylon</i>	Dube	Stoppage of water	2
<i>Cyperus rotundus</i>	Nut grass	Cooling	3
<i>Desmodium canum</i>	Sweet heart bush	Cooling	3
<i>Eleusine indica</i>	Dead man's grass	Urinary	2
<i>Entada polystachya</i>	Mayoc chapelle	Cooling	2 data needed
<i>Flemingia strobilifera</i>	Kidney bush	Kidney problems	2 data needed
<i>Gomphrena globosa</i>	Bachelor button	Urinary problems	2 data needed
<i>Hibiscus sabdariffa</i>	Sorrel	Cleans liver and blood	3 hypertension; 3 liver problems
<i>Justicia pectoralis</i>	Carpenter grass	Cooling	Data needed
<i>Kalachoe pinnata</i>	Wonder of the world	Cooling, Bladder stones	3 jaundice
<i>Lepianthes peltata</i>	Lani bois	Tea	Data needed
<i>Mimosa pudica</i>	Ti marie, mese marie	Cooling, Kidney problems	2
<i>Momordica charantia</i>	Caraaili	Cooling	Data needed
<i>Musa species</i>	Banana	Boil for cooling	2
<i>Nopalea cochinellifera</i>	Rachette	Kidney stones	2 data needed
<i>Peperomia rotundifolia</i>	Giron fleur, mowon	Kidney problems	3
<i>Peperomia pellucida</i>	Shining bush	Cooling	3
<i>Petiveria alliacea</i>	Mapourite, kudjuruk	Kidney problems	Data needed
<i>Pilea microphylla</i>	Du thé bethelmay	Bladder cleanser	2
<i>Pityrogramma calomelanos</i>	Fern	Urinary problems	Data needed
<i>Portulaca oleraceae</i>	Pussley	Cholesterol, short breath	3 pain; 3 gastroprotective
<i>Ruellia tuberosa</i>	Minny root	Cooling	Data needed
<i>Sansevieria guineensis</i>	Langue bouef, lash	Cooling	Data needed
<i>Scoparia dulcis</i>	Sweet broom	Cooling for babies	3 pain; 3 diuretic
<i>Solanum americanum</i>	Agouma, gouma	Cooling, provides iron	2 data needed
<i>Solanum melongena</i>	Melongene	Cholesterol	Not proven
<i>Stachytarpheta jamaicensis</i>	Vervine	Cooling	Data needed
<i>Theobroma cacao</i>	Cocoa	Eat for urinary problems	Data needed
<i>Vetiveria zizanioides</i>	Vetivert	Urinary problems	Data needed
<i>Zea mays</i>	Corn silk	Diuretic	3 diuretic

establishing whether a technology works and how. The non-experimental validation of the ethnoveterinary medicines was undertaken in recognition of that fact. Validation of traditional medicines is important since most scientists will not use local medicines without some form of validation and they have discouraged the use of these medicines leading to their disuse and eventual loss [1]. This loss is exacerbated by the death of knowledgeable elders. A framework of indigenous knowledge that can interface with science and technology is more likely to

influence scientific research agendas and development work, leading to the integration of indigenous knowledge into modern life [1,24].

More data is necessary to evaluate the safety of the following plants used to treat urinary problems, "cooling" and high cholesterol levels: *Costus scaber*, *Cynodon dactylon*, *Entada polystachya*, *Flemingia strobilifera*, *Gomphrena globosa*, *Justicia pectoralis*, *Lepianthes peltata*, *Momordica charantia*, *Nopalea cochinellifera*, *Petiveria alliacea*, *Pityrogramma*

calomelanos, *Ruellia tuberosa*, *Sansevieria guineensis*, *Stachytarpheta jamaicensis*, *Theobroma cacao* and *Vetiveria zizanioides*.

Little data was found to support the use of the following plants to treat urinary problems: *Justicia pectoralis*, *Lepianthes pelata*, *Momordica charantia*, *Petiveria alliacea*, *Pityrogramma calomelanos*, *Ruellia tuberosa*, *Sansevieria guineensis*, *Stachytarpheta jamaicensis*, *Theobroma cacao* and *Vetiveria zizanioides*.

The following plants have established analgesic or sedative effects: *Apium graveolens*, *Bauhinia cumanensis*, *Capraria biflora*, *Chamaesyce hirta* and *Portulaca oleraceae*.

Chamaesyce hirta, *Cissus verticillata*, *Kalanchoe pinnata*, *Peperomia* spp.,

Chamaesyce hirta, *Cissus verticillata*, *Kalanchoe pinnata*, *Peperomia* spp., *Portulaca oleraceae*, *Scoparia dulcis*, and *Zea mays* have sufficient evidence to support their traditional use for urinary problems, "cooling" and high cholesterol. *Cuscuta americana* also merits more study.

The plants used for hypertension, jaundice and diabetes that may be safe and justify more formal evaluation are *Annona squamosa*, *Aloe vera*, *Apium graveolens*, *Bidens alba*, *Carica papaya*, *Catharanthus roseus*, *Cecropia peltata*, *Citrus paradisi*, *Hibiscus sabdariffa*, *Momordica charantia*, *Morus alba*, *Persea americana*, *Phyllanthus urinaria*, *Tamarindus indicus* and *Tournefortia hirsutissima*. Several of the plants are used for more than one condition and further trials should take this into account. The use of eggplant extract as a hypocholesterolemic agent has some support but needs more study.

It is likely that medical practitioners in Trinidad and Tobago are counter-prescribing the majority of these remedies and should learn more about them.

Competing interests

The author(s) declare that they have no competing interests.

Additional material

Additional File 1

Table 4. Non-experimental validation of plants used in Trinidad and Tobago for diabetes and urinary problems. The data provided represents the chemical/pharmaceutical/pharmacological literature of the plant's known chemical constituents to determine the known physiological effects of either the crude plant, related species, or isolated chemical compounds that the plant is known to contain, and also any relevant clinical trials.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1746-4269-2-45-S1.pdf>]

Acknowledgements

This data collection was part of a larger study for a Ph.D. at Wageningen UR, the Netherlands. The fellowship support provided is appreciated. The Herbarium staff of the University of the West Indies provided essential plant identification. Dr. Lionel Robineau helped with the database searches. Dr. A.J.J. van den Berg provided invaluable editorial and pharmacological help.

References

- Lans C: *Creole remedies of Trinidad and Tobago* book self-published on Lulu.com; 2006.
- Ezenwaka CE, Kallou R, Uhlig M, Eckel J: **Relationship between adiponectin and metabolic variables in Caribbean offspring of patients with type 2 diabetes mellitus.** *Horm Metab Res* 2004, **36(4)**:238-42.
- Attele AS, Zhou YP, Xie JT, Wu JA, Zhang L, Dey L, Pugh W, Rue PA, Polonsky KS, Yuan CS: **Antidiabetic effects of Panax ginseng berry extract and the identification of an effective component.** *Diabetes* 2002, **51(6)**:1851-8.
- Lateef H, Abatan OI, Aslam MN, Stevens MJ, Varani J: **Topical pre-treatment of diabetic rats with all-trans retinoic acid improves healing of subsequently induced abrasion wounds.** *Diabetes* 2005, **54(3)**:855-61.
- Alleyne S, Cruickshank JK: **The use of informal medication – particularly bush teas – in Jamaican patients with diabetes mellitus.** *Cajanus* 1990, **23(1)**:57-67.
- Alleyne SI, Morrison EY, St A, Richard RR: **Some social factors related to control of diabetes mellitus in adult Jamaican patients.** *Diabetes Care* 1979, **2**:401-408.
- Mahabir D, Gulliford MC: **Use of medicinal plants for diabetes in Trinidad and Tobago.** *Revista Panamericana de Salud Pública* 1997, **1(3)**:174-179.
- Mahabir D, Gulliford MC: **Changing patterns of primary care for diabetes in Trinidad and Tobago over 10 years.** *Diabet Med* 2005, **22(5)**:619-24.
- Gulliford MC, Mahabir D, Rocke B: **Diabetes-related inequalities in health status and financial barriers to health care access in a population-based study.** *Diabet Med* 2004, **21(1)**:45-51.
- Gulliford MC, Ariyanayagam-Baksh SM, Bickram L, Picou D, Mahabir D: **Counting the cost of diabetic hospital admissions from a multi-ethnic population in Trinidad.** *Diabet Med* 1995, **12(12)**:1077-85.
- Zarowsky C, Stansfield S: *Diabetes and hypertension in women (Trinidad)* IDRC, Ottawa, Canada.
- Babwah F, Baksh S, Blake L, Cupid-Thuesday J, Hosein I, Sookhai A, Poon-King C, Hutchinson G: **The role of gender in compliance and attendance at an outpatient clinic for type 2 diabetes mellitus in Trinidad.** *Rev Panam Salud Publica* 2006, **19(2)**:79-84.
- Carr B-A, Alleyne L, Renaud D: **Report on the results of the Global Youth Tobacco survey in Trinidad and Tobago – 2000.** [[http://www.cdc.gov/tobacco/Global/gyts/reports/paho/2000/Trinidad and Tobago 2000 Paho01.htm](http://www.cdc.gov/tobacco/Global/gyts/reports/paho/2000/Trinidad%20and%20Tobago%2000/Paho01.htm)]. National Center for Chronic Disease Prevention and Health Promotion. PAHO. WHO Canadian Public Health Association

14. Gulliford MC, Mahabir D: **Diabetic foot disease and foot care in a Caribbean community.** *Diabetes Res Clin Pract* 2002, **56(1)**:35-40.
15. Grover JK, Yadav S, Vats V: **Medicinal plants of India with anti-diabetic potential.** *J Ethnopharmacol* 2002, **81(1)**:81-100.
16. Wong W: **Some folk medicinal plants from Trinidad.** *Economic Botany* 1976, **30**:103-142.
17. Foster GM: **Relationships between Spanish and Spanish-American folk medicine.** *Journal of American Folklore* 1953, **66**:201-217.
18. Aho WR, Minott K: **Creole and doctor medicine: folk beliefs, practices, and orientations to modern medicine in a rural and an industrial suburban setting in Trinidad and Tobago, the West Indies.** *Social Science and Medicine* 1977, **11(5)**:349-355.
19. Wilbert W: **The Pneumatic theory of female Warao herbalists.** *Social Science and Medicine* 1983, **25(10)**:1139-1146.
20. Littlewood R: **From vice to madness: the semantics of naturalistic and personalistic understandings in Trinidadian local medicine.** *Social Science and Medicine* 1988, **27(2)**:129-148.
21. Posey DA: **Changing fortunes: biodiversity and peasant livelihood in the Peruvian Andes.** *Journal of Latin American Studies* 1998, **30(3)**:682-683.
22. Hastrup K, Elsass P: **Anthropological advocacy.** *Current Anthropology* 1990, **31(3)**:301-311.
23. Warren DM, Slikkerveer LJ, Brokensha DW, Eds: *The cultural dimension of development: Indigenous knowledge systems* Intermediate Technology Publications, London; 1995.
24. Sillitoe P: **The development of indigenous knowledge.** *Current Anthropology* 1998, **39(2)**:223-252.
25. Heinrich N, Rimpler H, Antonio-Barrerra N: **Indigenous phytotherapy of gastrointestinal disorders in a lowland Mixe community (Oaxaca, Mexico): ethnopharmacological evaluation.** *J Ethnopharmacol* 1992, **36(1)**:63-80.
26. Chithra P, Sajithlal GB, Chanrakasan G: **Influence of Aloe vera on the glycosaminoglycans in the matrix of healing dermal wounds in rats.** *Journal of Ethnopharmacology* 1998, **59(3)**:179-186.
27. Shirwaikar A, Rajendran K, Dinesh Kumar C, Bodla R: **Antidiabetic activity of aqueous leaf extract of Annona squamosa in streptozotocin-nicotinamide type 2 diabetic rats.** *J Ethnopharmacol* 2004, **91(1)**:171-5.
28. Chistokhodova N, Nguyen C, Calvino T, Kachirskaia I, Cunningham G, Howard Miles D: **Antithrombin activity of medicinal plants from central Florida.** *J Ethnopharmacol* 2002, **81(2)**:277-80.
29. Atta AH, Alkofahi A: **Anti-nociceptive and anti-inflammatory effects of some Jordanian medicinal plant extracts.** *J Ethnopharmacol* 1998, **60(2)**:117-124.
30. Yadava RN, Reddy VM: **Anti-inflammatory activity of a novel flavonol glycoside from the Bauhinia variegata Linn.** *Nat Prod Res* 2003, **17(3)**:165-9.
31. Meyre-Silva C, Yunes RA, Delle Monache F, Santos AR, Schmeling LO, Gadotti VM, Liz F, Cechinel-Filho V: **Phytochemical and pharmacological analysis of Bauhinia microstachya (Raddi) Macbr. (Leguminosae).** *Z Naturforsch [C]* 2001, **56(11-12)**:939-42.
32. Ramesh N, Viswanathan MB, Saraswathy A, Balakrishna K, Brindha P, Lakshmanaperumalsamy P: **Phytochemical and antimicrobial studies of Begonia malabarica.** *J Ethnopharmacol* 1998, **79(1)**:129-132.
33. Chang SL, Chang CL, Chiang YM, Hsieh RH, Tzeng CR, Wu TK, Sytwu HK, Shyur LF, Yang WC: **Polyacetylenic compounds and butanol fraction from Bidens pilosa can modulate the differentiation of helper T cells and prevent autoimmune diabetes in non-obese diabetic mice.** *Planta Med* 2004, **70(11)**:1045-51.
34. Russell KR, Morrison EY, Ragoobirsingh D: **The effect of annatto on insulin binding properties in the dog.** *Phytother Res* 2005, **19(5)**:433-6.
35. De-Oliveira AC, Silva IB, Manhaes-Rocha DA, Paumgartten FJ: **Induction of liver monooxygenases by annatto and bixin in female rats.** *Braz J Med Biol Res* 2003, **36(1)**:113-8.
36. Liojier HA: *Plantas medicinales de Puerto Rico y del Caribe* Iberoamericana de Ediciones, Inc., San Juan, PR; 1990:566.
37. Kelcher S, West J, Crisp M, Chinnock R: **The Caribbean Bontia daphnoides and its Australian family Myoporaceae (Lamiaceae): evidence of an extreme dispersal event from morphological data and rp116 intron sequences.** [<http://www.botany2001.org/sympos12/abstracts/4.shtml>].
38. von Reis S, Lipp FJ: *New Plant Sources for Drugs and Foods from The New York Botanical Garden Herbarium* Cambridge, MA: Harvard University Press; 1982.
39. Acosta SL, Muro LV, Sacerio AL, Pena AR, Okwei SN: **Analgesic properties of Capraria biflora leaves aqueous extract.** *Fitoterapia* 2003, **74(7-8)**:686-8.
40. Collins DO, Gallimore WA, Reynolds WF, Williams LA, Reese PB: **New skeletal sesquiterpenoids, caprariolides A-D, from Capraria biflora and their insecticidal activity.** *J Nat Prod* 2000, **63(11)**:1515-1518.
41. Savickiene N, Dagilyte A, Lukosius A, Zitkevicius V: **Importance of biologically active components and plants in the prevention of complications of diabetes mellitus.** *Medicina (Kaunas)* 2002, **38(10)**:970-5. Article in Lithuanian
42. Eno AE, Owo OI, Itam EH, Konya RS: **Blood pressure depression by the fruit juice of Carica papaya (L.) in renal and DOCA-induced hypertension in the rat.** *Phytother Res* 2000, **14(4)**:235-9.
43. Palanichamy S, Nagarajan S, Devasagayam M: **Effect of Cassia alata leaf extract on hyperglycemic rats.** *J Ethnopharmacol* 1988, **22(1)**:81-90.
44. Nammi S, Boini MK, Lodagala SD, Behara RB: **The juice of fresh leaves of Catharanthus roseus Linn. reduces blood glucose in normal and alloxan diabetic rabbits.** *BMC Complement Altern Med* 2003, **3(1)**:4.
45. Perez-Guerrero C, Herrera MD, Ortiz R, Alvarez de Sotomayor M, Fernandez MA: **A pharmacological study of Cecropia obtusifolia Bertol aqueous extract.** *J Ethnopharmacol* 2001, **76(3)**:279-84.
46. Rocha FF, Lapa AJ, De Lima TC: **Evaluation of the anxiolytic-like effects of Cecropia glazouii Sneth in mice.** *Pharmacol Biochem Behav* 2002, **71(1-2)**:183-90.
47. Johnson PB, Abdurahman EM, Tiam EA, Abdu-Aguye I, Hussaini IM: **Euphorbia hirta leaf extracts increase urine output and electrolytes in rats.** *J Ethnopharmacol* 1999, **65(1)**:63-9.
48. Lanhers MC, Fleurentin J, Dorfman P, Mortier F, Pelt JM: **Analgesic, antipyretic and anti-inflammatory properties of Euphorbia hirta.** *Planta Med* 1991, **57(3)**:225-31.
49. Garcia MD, Saenz MT, Puerta R, Quilez A, Fernandez MA: **Antibacterial activity of Agave intermixta and Cissus sicyoides.** *Fitoterapia* 1999, **70(1)**:71-73.
50. Jainu M, Devi CS: **Effect of Cissus quadrangularis on gastric mucosal defensive factors in experimentally induced gastric ulcer-a comparative study with sucralfate.** *J Med Food* 2004, **7(3)**:372-6.
51. Longanga Otshudi AL, Vercruyse A, Foriers A: **Contribution to the ethnobotanical, phytochemical and pharmacological studies of traditionally used medicinal plants in the treatment of dysentery and diarrhoea in Lomela area, Democratic Republic of Congo (DRC).** *J Ethnopharmacol* 2000, **71(3)**:411-423.
52. Gorinstein S, Caspi A, Libman I, Katrich E, Lerner HT, Trakhtenberg S: **Fresh israeli jaffa sweetie juice consumption improves lipid metabolism and increases antioxidant capacity in hypercholesterolemic patients suffering from coronary artery disease: studies in vitro and in humans and positive changes in albumin and fibrinogen fractions.** *J Agric Food Chem* 2004, **52(16)**:5215-22.
53. Kirszberg C, Esquenazi D, Alviano CS, Rumjanek VM: **The effect of a catechin-rich extract of Cocos nucifera on lymphocytes proliferation.** *Phytother Res* 2003, **17(9)**:1054-8.
54. Alanis AD, Calzada F, Cervantes JA, Torres J, Ceballos GM: **Antibacterial properties of some plants used in Mexican traditional medicine for the treatment of gastrointestinal disorders.** *J Ethnopharmacol* 2005, **100(1-2)**:153-7.
55. Alleyne T, Roache S, Thomas C, Shirley A: **The control of hypertension by use of coconut water and mauby: two tropical food drinks.** *West Indian Med J* 2005, **54(1)**:3-8.
56. Kumaran A, Joel karunakaran R: **Antioxidant and free radical scavenging activity of an aqueous extract of Coleus aromaticus.** *Food Chemistry* 2006, **97(1)**:109-114.
57. Youn JY, Park HY, Cho KH: **Anti-hyperglycemic activity of Commelina communis L.: inhibition of alpha-glucosidase.** *Diabetes Res Clin Pract* 2004, **66(Suppl 1)**:S149-55.
58. Kala Prakash Chandra, Farooque Dhar Nehal A, Uppeandra : **Prioritization of medicinal plants on the basis of available knowl-**

- edge, existing practices and use value status in Uttaranchal, India. *Biodiversity and Conservation* 2004, **13**(2):453-469.
59. Mazumder UK, Gupta M, Pal D, Bhattacharya S: **Chemical and toxicological evaluation of methanol extract of *Cuscuta reflexa* Roxb. stem and *Corchorus olitorius* Linn. seed on hematological parameters and hepatorenal functions in mice.** *Acta Pol Pharm* 2003, **60**(4):317-23.
 60. Gupta M, Mazumder UK, Pal DK, Bhattacharya S: **Onset of puberty and ovarian steroidogenesis following administration of methanolic extract of *Cuscuta reflexa* Roxb. stem and *Corchorus olitorius* Linn. seed in mice.** *Journal of Ethnopharmacology* 2003, **89**(1):55-59.
 61. Pal D, Panda C, Sinhababu S, Dutta A, Bhattacharya S: **Evaluation of psychopharmacological effects of petroleum ether extract of *Cuscuta reflexa* Roxb. stem in mice.** *Acta Pol Pharm* 2003, **60**(6):481-6.
 62. Shinwari MI, Khan MA: **Folk use of medicinal herbs of Margalla Hills National Park, Islamabad.** *J Ethnopharmacol* 2000, **69**(1):45-56.
 63. Roman Ramos R, Alarcon-Aguilar F, Lara-Lemus A, Flores-Saenz JL: **Hypoglycemic effect of plants used in Mexico as antidiabetics.** *Arch Med Res* 1992, **23**(1):59-64.
 64. Auddy B, Ferreira M, Blasina F, Lafon L, Arredondo F, Dajas F, Tripathi PC, Seal T, Mukherjee B: **Screening of antioxidant activity of three Indian medicinal plants, traditionally used for the management of neurodegenerative diseases.** *J Ethnopharmacol* 2003, **84**(2-3):131-138.
 65. Jagtap AG, Shirke SS, Phadke AS: **Effect of polyherbal formulation on experimental models of inflammatory bowel diseases.** *J Ethnopharmacol* 2004, **90**(2-3):195-204.
 66. Rao M, Rao MNA: **Protective effects of cystone, a polyherbal ayurvedic preparation, on cisplatin-induced renal toxicity in rats.** *J Ethnopharmacol* 1998, **62**(1):1-6.
 67. Barreto GS: **Effect of butanolic fraction of *Desmodium adscendens* on the anococcygeus of the rat.** *Braz J Biol* 2002, **62**(2):223-30.
 68. Sugimoto K, Sakurai N, Shirasawa H, Fujise Y, Shibata K, Shimodo K, Sakata J: **Bovine cases of urolithiasis treated with traditional herbal medicine, P-3.** *J Vet Med Sci* 1992, **54**(3):579-582.
 69. Monache GD, Botta B, Vinciguerra V, de Mello JF, Andrade Chiapetta de A: **Antimicrobial isoflavonones from *Desmodium canum*.** *Phytochemistry* 1996, **41**(2):537-544.
 70. Nyarko AK, Asare-Anane H, Ofosuhene M, Addy ME: **Extract of *Ocimum canum* lowers blood glucose and facilitates insulin release by isolated pancreatic beta-islet cells.** *Phytomedicine* 2002, **9**(4):346-51.
 71. Ho CS, Wong YH, Chiu KW: **The hypotensive action of *Desmodium styracifolium* and *Clematis chinensis*.** *Am J Chin Med* 1989, **17**(3-4):189-202.
 72. Lakshmi Kumari P, Sumathi S: **Effect of consumption of finger millet on hyperglycemia in non-insulin dependent diabetes mellitus (NIDDM) subjects.** *Plant Foods Hum Nutr* 2002, **57**(3-4):205-13.
 73. Cos P, Hermans N, De Bruyne T, Apers S, Sindambiwe JB, Vanden Berghhe D, Pieters L, Vlietinck AJ: **Further evaluation of Rwandan medicinal plant extracts for their antimicrobial and antiviral activities.** *J Ethnopharmacol* 2002, **79**(2):155-63.
 74. Fabry W, Okemo P, Ansorg R: **Fungistatic and fungicidal activity of east African medicinal plants.** *Mycoses* 1996, **39**(1-2):67-70.
 75. Akinsinde KA, Olukoya DK: **Vibriocidal activities of some local herbs.** *J Diarrhoeal Dis Res* 1995, **13**(2):127-9.
 76. Rahman MM, Sarker SD, Byres M, Gray AI: **New salicylic acid and isoflavone derivatives from *Flemingia paniculata*.** *J Nat Prod* 2004, **67**(3):402-6.
 77. Das B, Tandon V, Saha N: **Effects of phytochemicals of *Flemingia vestita* (Fabaceae) on glucose 6-phosphate dehydrogenase and enzymes of gluconeogenesis in a cestode (*Railiellina echinobothrida*).** *Comp Biochem Physiol C Toxicol Pharmacol* 2004, **139**(1-3):141-6.
 78. de Moura RM, Pereira PS, Januario AH, Franca Sde C, Dias DA: **Antimicrobial screening and quantitative determination of benzoic acid derivative of *Gomphrena celosoides* by TLC-densitometry.** *Chem Pharm Bull (Tokyo)* 2004, **52**(11):1342-4.
 79. Herrera-Arellano A, Flores-Romero S, Chavez-Soto MA, Tortoriello J: **Effectiveness and tolerability of a standardized extract from *Hibiscus sabdariffa* in patients with mild to moderate hypertension: a controlled and randomized clinical trial.** *Phytomedicine* 2004, **11**(5):375-82.
 80. Akindahunsi AA, Olaleye MT: **Toxicological investigation of aqueous-methanolic extract of the calyces of *Hibiscus sabdariffa* L.** *J Ethnopharmacol* 2003, **89**(1):161-4.
 81. Ali BH, Mousa HM, El-Mougy S: **The effect of a water extract and anthocyanins of *Hibiscus sabdariffa* L on paracetamol-induced hepatotoxicity in rats.** *Phytother Res* 2003, **17**(1):56-9.
 82. MacRae WD, Towers GH: ***Justicia pectoralis*: a study of the basis for its use as a hallucinogenic snuff ingredient.** *J Ethnopharmacol* 1984, **12**(1):93-111.
 83. Yadav NP, Dixit VK: **Hepatoprotective activity of leaves of *Kalanchoe pinnata* Pers.** *J Ethnopharmacol* 2003, **86**(2-3):197-202.
 84. Momo CE, Oben JE, Tazoo D, Dongo E: **Antidiabetic and hypolipidaemic effects of a methanol/methylene-chloride extract of *Laportea ovalifolia* (Urticaceae), measured in rats with alloxan-induced diabetes.** *Ann Trop Med Parasitol* 2006, **100**(1):69-74.
 85. Joyamma V, Rao SG, Hrishikeshavan HJ, Aroor AR, Kulkarni DR: **Biochemical mechanisms and effects of *Mimosa pudica* (Linn) on experimental urolithiasis in rats.** *Indian J Exp Biol* 1990, **28**(3):237-40.
 86. Tongia A, Tongia SK, Dave M: **Phytochemical determination and extraction of *Momordica charantia* fruit and its hypoglycemic potentiation of oral hypoglycemic drugs in diabetes mellitus (NIDDM).** *Indian J Physiol Pharmacol* 2004, **48**(2):241-4.
 87. Cummings E, Hundal HS, Wackerhage H, Hope M, Belle M, Adeghate E, Singh J: ***Momordica charantia* fruit juice stimulates glucose and amino acid uptakes in L6 myotubes.** *Mol Cell Biochem* 2004, **261**(1-2):99-104.
 88. Lemus I, Garcia R, Delvillar E, Knop G: **Hypoglycaemic activity of four plants used in Chilean popular medicine.** *Phytother Res* 1999, **13**(2):91-4.
 89. Oku T, Yamada M, Nakamura M, Sadamori N, Nakamura S: **Inhibitory effects of extractives from leaves of *Morus alba* on human and rat small intestinal disaccharidase activity.** *Br J Nutr* 2006, **95**(5):933-8.
 90. Singab AN, El-Beshbishy HA, Yonekawa M, Nomura T, Fukai T: **Hypoglycemic effect of Egyptian *Morus alba* root bark extract: effect on diabetes and lipid peroxidation of streptozotocin-induced diabetic rats.** *J Ethnopharmacol* 2005, **100**(3):333-8.
 91. Dhanabal SP, Sureshkumar M, Ramanathan M, Suresh B: **Hypoglycemic effect of ethanolic extract of *Musa sapientum* on alloxan induced diabetes mellitus in rats and its relation with antioxidant potential.** *J Herb Pharmacother* 2005, **5**(2):7-19.
 92. Ojewole JA, Adewunmi CO: **Hypoglycemic effect of methanolic extract of *Musa paradisiaca* (Musaceae) green fruits in normal and diabetic mice.** *Methods Find Exp Clin Pharmacol* 2003, **25**(6):453-6.
 93. Wolfram R, Budinsky A, Efthimiou Y, Stomatopoulos J, Oguogho A, Sinzinger H: **Daily prickly pear consumption improves platelet function.** *Prostaglandins Leukot Essent Fatty Acids* 2003, **69**(1):61-6.
 94. Villasenor IM, Lamadrid MR: **Comparative anti-hyperglycemic potential of medicinal plants.** *J Ethnopharmacol* 2006, **104**(1-2):129-31.
 95. Singh S, Rehan HM, Majumdar DK: **Effect of *Ocimum sanctum* fixed oil on blood pressure, blood clotting time and pentobarbitone-induced sleeping time.** *J Ethnopharmacol* 2001, **78**(2-3):139-43.
 96. Hannan JM, Marenah L, Ali L, Rokeya B, Flatt PR, Abdel-Wahab YH: ***Ocimum sanctum* leaf extracts stimulate insulin secretion from perfused pancreas, isolated islets and clonal pancreatic beta-cells.** *J Endocrinol* 2006, **189**(1):127-36.
 97. Dhawan K, Dhawan S, Sharma A: **Passiflora: a review update.** *J Ethnopharmacol* 2004, **94**:1-23.
 98. Aziba PI, Adedeji A, Ekor M, Adeyemi O: **Analgesic activity of *Peperomia pellucida* aerial parts in mice.** *Fitoterapia* 2001, **72**(1):57-58.
 99. de Fatima Arrigoni-Blank M, Dmitrieva EG, Franzotti EM, Antonioli AR, Andrade MR, Marchioro M: **Anti-inflammatory and analgesic activity of *Peperomia pellucida* (L.) HBK (Piperaceae).** *J Ethnopharmacol* 2004, **91**(2-3):215-8.
 100. Khan MR, Omoloso AD: **Antibacterial activity of *Hygrophila stricta* and *Peperomia pellucida*.** *Fitoterapia* 2002, **73**(3):251-4.

101. Joshi A, Joshi K: **Indigenous knowledge and uses of medicinal plants by local communities of the Kali Gandaki Watershed area, Nepal.** *J Ethnopharmacol* 2000, **73(1-2)**:175-183.
102. de Fatima Arrigoni-Blank M, Dmitrieva EG, Franzotti EM, Antonioli AR, Andrade MR, Marchioro M: **Anti-inflammatory and analgesic activity of *Peperomia pellucida* (L.) HBK (Piperaceae).** *J Ethnopharmacol* 2004, **91(2-3)**:215-8.
103. Aziba PI, Adedeji A, Ekor M, Adeyemi O: **Analgesic activity of *Peperomia pellucida* aerial parts in mice.** *Fitoterapia* 2001, **72(1)**:57-58.
104. Raphael KR, Sabu MC, Kuttan R: **Hypoglycemic effect of methanol extract of *Phyllanthus amarus* Schum & Thonn on alloxan induced diabetes mellitus in rats and its relation with antioxidant potential.** *Indian J Exp Biol* 2002, **40(8)**:905-9.
105. Rocha MJ, Fulgencio SF, Rabetti AC, Nicolau M, Poli A, Simoes CM, Ribeiro-do-Valle RM: **Effects of hydroalcoholic extracts of *Portulaca pilosa* and *Achyrocline satureioides* on urinary sodium and potassium excretion.** *J Ethnopharmacol* 1994, **43(3)**:179-83.
106. Chan K, Islam MVW, Kamil M, Radhakrishnan R, Zakaria MNM, Habibullah M, Attas A: **The analgesic and anti-inflammatory effects of *Portulaca oleracea* L. subsp. *sativa* (Haw.) Celak.** *J Ethnopharmacol* 2000, **73(3)**:445-451.
107. Karimi G, Hosseinzadeh H, Eftehad N: **Evaluation of the gastric antiulcerogenic effects of *Portulaca oleracea* L. extracts in mice.** *Phytother Res* 2004, **18(6)**:484-7.
108. Pari L, Latha M: **Antihyperlipidemic effect of *Scoparia dulcis* (sweet broomweed) in streptozotocin diabetic rats.** *J Med Food* 2006, **9(1)**:102-7.
109. Guimaraes PR, Galvao AM, Batista CM, Azevedo GS, Oliveira RD, Lamounier RP, Freire N, Barros AM, Sakurai E, Oliveira JP, Vieira EC, Alvarez-Leite JI: **Eggplant (*Solanum melongena*) infusion has a modest and transitory effect on hypercholesterolemic subjects.** *Braz J Med Biol Res* 2000, **33(9)**:1027-36.
110. Botelho FV, Eneas LR, Cesar GC, Bizzotto CS, Tavares E, Oliveira FA, Gloria MB, Silvestre MP, Arantes RM, Alvarez-Leite JI: **Effects of eggplant (*Solanum melongena*) on the atherogenesis and oxidative stress in LDL receptor knock out mice (LDLR^{-/-}).** *Food Chem Toxicol* 2004, **42(8)**:1259-67.
111. Maiti R, Jana D, Das UK, Ghosh D: **Antidiabetic effect of aqueous extract of seed of *Tamarindus indica* in streptozotocin-induced diabetic rats.** *J Ethnopharmacol* 2004, **92(1)**:85-91.
112. Ruzaidi A, Amin I, Nawalyah AG, Hamid M, Faizul HA: **The effect of Malaysian cocoa extract on glucose levels and lipid profiles in diabetic rats.** *J Ethnopharmacol* 2005, **98(1-2)**:55-60.
113. Osakabe N, Yamagishi M, Natsume M, Yasuda A, Osawa T: **Ingestion of proanthocyanidins derived from cacao inhibits diabetes-induced cataract formation in rats.** *Exp Biol Med (Maywood)* 2004, **229(1)**:33-9.
114. Alarcon-Aguilara FJ, Roman-Ramos R, Perez-Gutierrez S, Aguilar-Contreras A, Contreras-Weber CC, Flores-Saenz JL: **Study of the anti-hyperglycemic effect of plants used as antidiabetics.** *J Ethnopharmacol* 1998, **61(2)**:101-10.
115. Jagtap AG, Shirke SS, Phadke AS: **Effect of polyherbal formulation on experimental models of inflammatory bowel diseases.** *J Ethnopharmacol* 2004, **90(2-3)**:195-204.
116. Habtemariam S: **Extract of corn silk (stigma of *Zea mays*) inhibits the tumour necrosis factor- α – and bacterial lipopolysaccharide-induced cell adhesion and ICAM-1 expression.** *Planta Med* 1998, **64(4)**:314-318.
117. Brereton B: *A history of modern Trinidad 1783–1962* Kingston, Jamaica: Heineman; 1981.
118. Buchler IR: **Caymanian folk medicine: A problem in applied anthropology.** *Human Organisation* 1964, **23(1)**:48-49.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

