



Published in final edited form as:
Caries Res. 2006 ; 40(6): 473–480.

DIETARY PATTERNS RELATED TO CARIES IN A LOW-INCOME ADULT POPULATION

Brian A. Burt^{*}, Justine L. Kolker^{**}, Anita M. Sandretto^{***}, Ying Yuan^{****}, Woosung Sohn^{**}, and Amid I. Ismail^{**}

^{*}*Department of Epidemiology, School of Public Health, University of Michigan*

^{**}*Department of Cariology, Restorative Sciences and Endodontics, School of Dentistry, University of Michigan*

^{***}*Department of Environmental Health Sciences, School of Public Health, University of Michigan*

^{****}*Department of Biostatistics, School of Public Health, University of Michigan*

Abstract

The aim of this study was to examine the relationship between dietary patterns and caries experience in a representative group of low-income African-American adults. Participants were residents of Detroit, Michigan, with household incomes below 250% of the federally-established poverty level (n = 1,021). Dietary histories were obtained by trained interviewers in face-to-face interviews with the adult participants, using the Block 98.2® food frequency questionnaire developed by Block Dietary Systems, Berkeley, CA. Caries was measured by the ICDAS criteria (International Caries Detection and Assessment System). There were 200 dietary records whose data were judged to be invalid; these participants were omitted from the dietary analyses to leave n = 821. Analyses were conducted using software from SAS and SUDAAN. Factor analysis identified patterns of liquid and solid food consumption, and the resulting factor scores were used as covariates in multivariable linear regression. Caries was extensive, with 82.3% of the 1,021 participants (n=839) having at least one cavitated lesion. Nearly three-quarters of the adult participants were overweight or obese. This population has severe caries, poor oral hygiene, and diets that are high in sugars and fats and low in fruits and vegetables. Apart from tapwater, the most frequently consumed food item by adults of all ages was soft drinks; 19% of all energy from sugar came from soft drinks alone. In both the bivariate analyses and in the regression model, frequency of soft drink consumption and the presence of gingival plaque deposits were significantly associated with caries. Interventions to promote oral health are unlikely to be successful without improvements in the social and physical environment.

Keywords

diet; dietary patterns; soft drinks; sugars; fat consumption; caries; plaque deposits; oral hygiene; adults; disparities; low-income

Introduction

Sugars are an integral part of caries etiology [Gustafsson et al.,1954; Sheiham, 2001]. Even though the sugars/caries relationship in the modern era of widespread fluoride exposure is not

Author for correspondence: Dr. Brian A. Burt Department of Epidemiology School of Public Health University of Michigan 109 Observatory Street Ann Arbor, MI 48109-2029.

Declaration of Interest: No author has any conflict of interest in the conduct of this study or in its findings.

as clear and direct as was previously assumed [Burt and Pai, 2001], sugars' role in caries development is still not seriously disputed.

There is far less certainty, however, about the relationship between overall dietary patterns and caries: i.e., does a poor-quality diet predispose to caries? By the term “poor-quality diet,” we mean one that provides adequate energy but has an imbalance in nutrient intake when measured against recommended levels: too much fat and sugars and not enough fruits and vegetables. Such diets in high-income countries have been linked to a number of disease states [Dreeben, 2001; Kayrooz et al., 1998]. Poor-quality diets have perhaps been linked most clearly to obesity, and the increase in obesity over recent years in the United States [Flegal et al., 2002] is a serious public health problem. This is true for adults as well as for children, and obesity-related problems are found at all socioeconomic levels [Ogden et al., 2002]. From the dental viewpoint, we might intuitively expect that poor-quality diets promote caries, but apart from sugars' role there is little firm evidence to support such a relationship. Caries is certainly higher in children who have been high sucrose consumers all their lives when compared to children with lower lifetime sucrose intake [Ruottinen et al., 2004].

This paper examines the relationship between dietary patterns and caries experience in a representative group of low-income African-American adults in Detroit, Michigan, USA. The study was conducted by the Detroit Center for Research in Oral Health Disparities. The Center has the overall goal of identifying reasons for the community-level and individual-level disparities in oral health found among this population, where families appear to live under much the same social and economic circumstances.

Materials and Methods

The study was conducted in accordance with the principles of the Declaration of Helsinki. All participants gave written consent to taking part. The study was approved by the Institutional Review Board (IRB) for Health Sciences at the University of Michigan, Ann Arbor.

Study Population

The participants were African-Americans in Detroit with household incomes below 250% of the poverty level as defined by the federal government. Participants were selected using a multistage area probability sample design based on data from the US census of 2000. The 39 census tracts within Wayne County (where Detroit is located) with the highest proportion of low-income African-American households were selected, and a random sample of 565 blocks divided into 118 neighborhood segments was drawn from them. A household listing of these blocks was conducted from June to August 2002. From that household list, a number of housing units were chosen to yield the target sample of 1,000 persons for the first wave of data collection. Between September 2002 and August 2003, a field team visited the selected housing units and successfully screened and recruited 1,021 families to take part in the study. A family, for the purposes of this study, was defined as a primary caregiver and one African-American child who had not yet attained his/her sixth birthday.

Data Collection

Data were collected by interview and clinical dental examination at the same facility in Detroit. Questionnaires, completed by trained interviewers working with the adult participants, were designed to explore the sociodemographic and lifestyle characteristics of each adult and child, as well as their general and oral health history. Dietary histories were obtained by trained interviewers from adult participants, using the the Block 98.2® food frequency questionnaire (FFQ) developed by Block Dietary Systems, Berkeley, CA. This FFQ asks the frequency of consumption of a wide variety of food items over the last year, plus the number of servings

per ingestion and the portion size (judged by the respondent from photographs and food models of various portion sizes). Algorithms used by Block Dietary Systems, based on data from several National Food Consumption Surveys conducted by the U.S. Department of Agriculture, permit the calculation of energy and nutrient intake as well as the frequency of consumption of food items.

Measurement of Body Mass Index

Participants were weighed on a calibrated digital scale and had their height measured from a wall-mounted tape measure, both from standard protocols. Body Mass Index (BMI), a broad measure of adiposity, is computed as weight in kilograms divided by the square of height in meters [National Institutes of Health, 2005]. Staff were trained over a period of one month, and the quality of the interviews was regularly checked throughout the 11-month data collection period by reviewing videotapes of the interviews. The investigators also conducted periodic face-to-face reviews with the interviewers.

Measurement of Dental Caries and Plaque Deposits

Caries was measured by the ICDAS criteria (International Caries Detection and Assessment System) applied by four trained dentists. ICDAS is a system of caries detection and assessment that was developed by an international group of caries researchers over the period 2002-05 [Pitts, 2004]. It was intended that the criteria (a) reflect the current scientific understanding of the process of dental caries; (b) be adopted for use in dental surveillance, research, practice, and education; and (c) provide a foundation for incorporating and validating new caries diagnostic tools.

In the Detroit study, four dentists carried out the scoring for dental caries, periodontal conditions, and plaque deposits. Participants were examined in a standard dental operatory in the reclining position with a standard light source. The teeth were cleaned of loose debris with a scaler, where necessary, and plaque deposits assessed. Teeth were then examined for caries using the ICDAS criteria. Assessment of caries was primarily visual. Caries criteria necessarily varied slightly from pit and fissures to proximal to smooth surfaces. To illustrate the ICDAS criteria, the scoring system for pit and fissure caries was as follows:

- 0- Sound tooth surface: no loss of integrity after air-drying for 5 s.
- 1- First visual change in enamel.
- 2- Distinct visual change in enamel.
- 3- Non-cavitated surface with underlying dark shadow from dentine.
- 4- Localized enamel breakdown, with no dentine visible.
- 5- Distinct cavity with dentine visible.
- 6- Extensive distinct cavity with dentine visible.

The caries outcome measure in data analysis was D₁MFS, where “D” included both noncavitated and cavitated lesions, i.e., ICDAS categories 1-6 above. A second measure, D₂MFS, recorded just cavitated lesions (ICDAS categories 4-6). However, these two measures were tightly correlated (Pearson's $r = 0.95$, $p < 0.001$), so we used just D₁MFS as the more complete outcome measure of caries.

Plaque deposits were scored by the Patient Hygiene Performance (PHP) index [Podshadley and Haley, 1968] applied to the “Ramfjord” index teeth [Ramfjord, 1959]. These teeth are the maxillary right first molar; maxillary left central incisor; maxillary left second premolar; mandibular left first molar; mandibular right central incisor; and mandibular right second premolar. The PHP index scores the degree of visible plaque deposits, after a disclosing rinse, as 0 (no plaque), 1 (recent plaque deposit), or 2 (long-term plaque deposit) in each of five zones of the tooth surface.

Statistical Analyses

Statistical analyses were conducted using software from SAS® [SAS Institute, 2005], and SUDAAN® [Research Triangle Institute, 2005] to estimate variances adjusted for the design effect. Significance was set at the conventional level of $p < 0.05$, and the Wald F test was used to derive p-values.

Completed FFQs were mailed to Block Dietary Systems for analysis. All other data were entered twice using a custom-designed Access program. Data checking for error was conducted using SAS. Discrepancies between the first and second data entries were reviewed and resolved by a senior investigator (WS).

After data cleaning and filing, bivariate analyses were conducted to relate frequently-reported foods to caries experience (D_1MFS scores). When the population was categorized into tertiles by D_1MFS scores, we selected the food items that were consumed at least once per day by at least 5% of the adult participants in the high-caries tertile, and again by at least 5% in the low-caries tertile. Some foods were reported in both tertiles, others were found only in one tertile and not the other. There were 29 such food items selected this way, and subsequent analysis was focused on them.

Factor analysis was conducted to identify the patterns of liquid and solid food consumption, and the resulting factor scores were then used as covariates in multivariable linear regression to relate dietary patterns to caries levels. Separate factor analyses were carried out for eight liquid food items (soft drinks were not included due to the high frequency of their consumption and strong bivariate association with caries) and 20 solid food items, using PROC FACTOR from the SAS package. The number of factors was determined by scree plots and by what made sense from the nutritional viewpoint. The resulting factor scores, along with the weekly servings of soft drinks, were then used as continuous independent variables in the multivariable linear regression model with D_1MFS values as the continuous dependent variable. Square root transformation was performed to improve normality. The multivariable linear regression was carried out by using PROC REGRESS available in SUDAAN, incorporating sampling design and weights. The other independent variables included were: (a) oral health variables: the number of times teeth were brushed during the last week; and PHP plaque scores; (b) demographic variables: age; education; income; employment status.

We carried out backward elimination model selection to determine the most parsimonious predictive model for caries. We began with the full model that included all independent variables, proceeding at each step by dropping from the model the variable with the highest p-value. (Demographic variables were retained because of their fundamental importance, regardless of their statistical significance). This process was repeated until only food, drink, or oral hygiene variables with a $p\text{-value} < 0.10$ remained in the model.

Results

The demographic characteristics and Body Mass Index of all adult participants who had at least one tooth ($n=1,005$), related to dental caries experience as measured by the D_1MFS index, are

shown in Table 1. Socioeconomic status (SES) is not included because the study population, by design, was clustered at the lower end of the SES spectrum.

Nutrient Intake and Body Mass Index

The study began with 1,021 adults. There were 200 FFQ interviews classed as invalid because they included far too many (≥ 17) or far too few (≤ 3) food items per day relative to national norms. Block Dietary Systems identified these records which were then eliminated from further dietary analyses. All analyses on diet and dietary patterns are therefore based on $n = 821$, the completed FFQs considered valid. There was still some missing data in a few FFQs, which is why the totals in some analyses are less than 821.

The Body Mass Index (BMI) is a broad measure of weight related to height. From the data in Table 1, it can be seen that nearly three-quarters of the participants were either overweight or obese.

Average daily energy intake for adult females was 2,090 kilocalories, which again is above average and consistent with the profile of overweight and obesity. The proportions of participants who were overweight or obese increased with increasing age: those aged under 26 were 68% overweight or obese, for those aged 26 to <46 years it was 74%, and for those 46 or older it was 78%. Adults ingested 13.5% of total energy from protein, 39.4% from fats, and 47.1% from carbohydrates (15.0% from sugars).

Yet another indicator of poor overall dietary quality comes with the reported consumption of fruits and vegetables. Only 7.4% reported eating two or more servings of vegetables per day and 7.8% two or more servings of fruit.

The most frequently reported food on a daily basis was tapwater. The second most-reported food consumed daily was sugared soda (reported by 30.3%), and the third was potato chips (22.1%). Fried chicken was ranked ahead of green salad. Food sources for sugars for these adults were predominantly liquid. Soft drinks dominated, with 19% of energy from sugars coming from soft drinks alone.

Caries Status and Plaque Deposits

The overall weighted kappa coefficient for inter-examiner reliability in recording caries was 0.83 and the weighted kappa coefficient for intra-examiner reliability was 0.74. Caries was commonplace and severe in this population, with 839 of the 1,005 dentate participants having one or more untreated cavitated lesions. In those aged 14-<35 years only 0.2% had no carious lesions. In those aged 35-<47 ($n = 140$) it was 1.9%, and in those aged 47 or more ($n = 38$) it was 3.6%.

The only plaque measure significantly related to D₁MFS scores was gingival plaque, i.e., visible plaque around the gingival margin after a disclosing rinse ($p < 0.02$). Total PHP scores, proximal surfaces and middle surface scores were not significantly related to caries experience.

Diet and Caries

For the independent variable in bivariate analyses we used tertiles of the proportion of total energy (we called this energy%) from the food item. By this measure D₁MFS scores were not related to the intake of protein or fat, but they were significantly related to carbohydrate intake (Table 2), despite the absence of a trend. Because carbohydrates include both non- and low-cariogenic starches [Lingstrom et al., 2000] as well as cariogenic simple sugars, when the same analysis was restricted to sugars (all kinds) the sugars-caries relationship was even more

significant (Table 3). D1MFS scores were not related to total PHP scores, but they were related to gingival oral hygiene scores (Table 4).

We also expressed the independent variable as daily servings of food items. No relation was found between D₁MFS scores and intakes of vegetables, fruit, grain, meat, dairy products, tomato juice, orange juice, other real fruit juices, drinks with added vitamin C, and milk. In bivariate analysis there was at first only a weak relation between D₁MFS and soft drink consumption, but this relationship was age-confounded. The younger adults drank more soft drinks and less coffee than the older adults, and when age-adjusted the D₁MFS scores were significantly related to soft drink consumption ($p < 0.01$; data not tabulated).

Factor analysis resulted in four factors for 20 solid food items and four factors for eight liquid food items (soft drinks were not included). The four solid food factors represented fruits, chips/candy, breakfast foods (e.g., eggs, sausage), and cheese snacks, and the four liquid food factors represented real fruit juice, milk, fruit-flavored drinks (e.g., HiC, Sunny Delight), and coffee/tapwater. These factor scores, along with the weekly servings of soft drinks, were then used as continuous independent variables in the multivariable linear regression with D₁MFS scores as the outcome. The full model is shown in Table 5. Age, frequency of soft drink consumption and presence of gingival plaque were significantly related to caries prevalence ($p < 0.05$).

The final model, after the backward model selection procedure was carried out, is shown in Table 6. The independent variables retained in the final model (apart from the demographic variables) were frequency of soft drink consumption and presence of gingival plaque. Age was negatively associated with caries severity, and soft drink consumption and presence of gingival plaque were positively associated with caries severity. These results are similar to those from the full model. Possible two-way interactions were also explored based on the final model, but none of them were statistically significant.

Discussion

Dietary patterns have been defined as multiple dietary attributes operationalized as a single exposure [Kant et al., 2004]. To measure dietary patterns in this study we first considered using an existing index from the literature. However, while a number of indexes to measure dietary quality have been devised, none has emerged as more complete than the others [Kant, 1996]. We developed our own Recommended Food Index, based on US Department of Agriculture recommendations, but it resulted in very low scores with little variance between individual participants. This indicated a population with a homogenous diet and low consumption of healthy foods, so we did not use the Recommended Food Index in this analysis. The approach adopted was to use factor analysis, which in this instance detects dietary patterns that reduce the number of variables by finding factors that comprise correlated dietary items [Kant, 2004]. Factor analysis has been shown to be a valid measure of dietary patterns when the factor loadings are compared with the results of a seven-day food diary [Togo et al., 2003].

The data in Table 1 are illustrative of the poor social conditions in this area. The vast majority of families have no male head of household, which in itself raises more social problems. The data also show that this is a severely overweight population. Having nearly three-quarters of the predominantly-female participants either overweight or obese is well above the national average (Flegal et al., 2002). Dietary quality is poor by several measures. The proportion of energy from fats is above the level recommended by the US Department of Agriculture, while the carbohydrates and protein levels are near the bottom of the recommended range. Only 7.4% reported consuming two or more servings of vegetables per day, and 7.8% two or more servings of fruit. Current recommendations are to consume 2-2½ cups of fruit and vegetables per day [US Department of Agriculture, 2005]. An alternative set of dietary guidelines has been published from Harvard University. These differ from those put out by the US Department of

Agriculture mostly in recommending that a higher proportion of energy come from fats, but only unsaturated fats, and less from carbohydrates [Harvard University, 2005] The diet of the participants in our study is still poor under the Harvard guidelines.

Dietary patterns of African-American adults is a subject that has not received much research attention, but our results are similar to what little information has been published. One study among African-Americans in North Carolina, drawn from all socioeconomic levels, found that participants' consumption of fruits and vegetables was below the levels advised in national recommendations [Gary, et al., 2004]. This pattern is even more pronounced in our Detroit study, where the study population, by definition, came from low socioeconomic levels and we did not see the contrasting dietary patterns that would be expected in a population drawn from all socioeconomic levels. It seemed to us that at least part of the reason is the poor availability of good-quality fruits and vegetables in sections of the city, with cultural and economic issues compounding this complex relationship.

The literature on the relationship between plaque deposits and caries is mixed. There are studies which conclude that plaque deposits (or more vaguely “poor oral hygiene”) are a risk factor for caries [Kleemola-Kujala and Rasanen, 1982; Mascarenhas, 1998], while others find no such association [Payette and Brodeur, 1992; Arrow, 1998; Ety et al., 1994]. Meticulous oral hygiene has been shown to reverse the caries process in accessible root lesions [Nyvad and Fejerskov, 1986]. The mixed literature is probably a result of studies' using inappropriate plaque measures (e.g., scoring plaque from tooth surfaces where caries is unlikely to be found), measuring caries only in the cavitated stage, and fluoride effects. In our study, overall PHP index scores were not related to caries levels, but the overall scores included plaque measures from areas of the tooth surface where caries was unlikely, e.g, the middle of the labial surface. When PHP scores were confined to gingival areas only, we found a strong relationship between plaque deposits and caries.

There was a time when cariogenic foods were thought of as being mostly solid foods, e.g., candy, chocolate, baked goods. The influential Vipeholm study [Gustaffson et al., 1954] in the 1950s emphasized the cariogenicity of “sticky” foods and downplayed the importance of sugar in liquid form. More recently, however, it has been documented that sugared beverages are a major risk factor for caries development in children [Levy et al., 2003; Marshall et al., 2003]. Our data confirm these findings and demonstrate that in this population the same picture applies to adults. It is noted that 55% of the adults' energy from sugars comes from a few types of drinks, for children the equivalent figure is 54% [Kolker et al., 2005]. This suggests similarities between the diets of the adults and the children.

It was this clear evidence that sugar sources in this population were predominantly liquid that led us to lift soft drinks out of the factor analysis and leave it as a separate covariable. Soft drinks retained a highly stable level of statistical significance throughout the backward elimination model selection, this stability would seem to emphasize their importance in the carious process. Other drinks in the ever-growing menu of available beverages, real fruit juices included, are sugary and hence are potentially cariogenic, but it is notable that all combinations of both liquids and solids identified by the factor analysis had dropped out of the final model while soft drinks remained. The model also retained its R^2 value of 0.24. The conclusion would seem to be that the quality of the overall diet by itself does not much influence caries in this population, but soft drink consumption in the presence of plaque retention around the gingival margins certainly does.

Our findings are that gingival plaque deposits and soft drinks are the major individual-level determinants of caries in this population. This leaves the door open for several intervention areas, i.e., improving both oral hygiene and dietary quality, specifically cutting back on soft

drink consumption. However, achieving permanent beneficial change in individual behavior will remain a challenge in this population unless there are also structural changes in the community and in key aspects of its social functioning. We cannot expect people living in deep poverty, without prospects for improvement, and for whom violence is a part of everyday life, to successfully adopt the habits of healthy diets and careful oral hygiene unless their physical and social environment is able to provide a measure of hope and dignity.

Acknowledgements

This study was supported by grant number U-54 DE 14261 from the National Institute for Dental and Craniofacial Research (NIDCR).

REFERENCES

- Arrow P. Oral hygiene in the control of occlusal caries. *Community Dent Oral Epidemiol* 1998;26:324–330. [PubMed: 9792124]
- Burt BA, Pai S. Sugar consumption and caries risk: a systematic review. *J Dent Educ* 2001;65:1017–1023. [PubMed: 11699972]
- Dreeben O. Health status of African Americans. *J Health Social Policy* 2001;14:1–17.
- Etty EJ, Henneberke M, Gruythuysen RJ, Woltgens JH. Influence of oral hygiene on early enamel caries. *Caries Res* 1994;28:132–136. [PubMed: 8156564]
- Flegal KM, Carroll MD, Ogden CL, Johnson CL. Prevalence and trends in obesity among US adults, 1999–2000. *JAMA* 2002;288:1723–1727. [PubMed: 12365955]
- Gary TL, Baptiste-Roberts K, Gregg EW, Williams DE, Beckles GL, Miller EJ III, Engelau MM. Fruit, vegetable and fat intake in a population-based sample of African Americans. *J Nat Med Assoc* 2004;96:1599–1605.
- Gustaffson BE, Quensel CE, Lanke LS, Lundquist C, Grahnen H, Bonow BE, Krasse B. The Vipeholm dental caries study. The effect of different levels of carbohydrates intake on caries activity in 436 individuals observed for five years. *Acta Odont Scand* 1954;11:232–364. [PubMed: 13196991]
- The Nutrition Source, knowledge of healthy eating. http://www.hsph.harvard.edu/nutritionsource/EDBH_update.html
- Kant AK. Dietary patterns and health outcomes. *J Am Dietetic Assoc* 2004;104:615–635.
- Kant AK. Indexes of overall diet quality: a review. *J Am Dietetic Assoc* 1996;96:785–791.
- Kant AK, Graubard BI, Schatzkin A. Dietary patterns predict mortality in a national cohort: the National Health Interview Surveys, 1987 and 1992. *J Nutr* 2004;134:1793–1799. [PubMed: 15226471]
- Kayrooz K, Moy TF, Yanek LR, Becker DM. Dietary fat patterns in urban African American women. *J Community Health* 1998;23:453–469. [PubMed: 9824794]
- Kleemola-Kujala E, Rasanen L. Relationship of oral hygiene and sugar consumption to risk of caries in children. *Community Dent Oral Epidemiol* 1982;10:224–233. [PubMed: 6958405]
- Kolker, J.; Burt, BA.; Sohn, W.; Ismail, AI. Program for the Annual Session of the International Association for Dental Research. Baltimore, MD: 2005. Disparities in dental caries: are sugared beverages a factor?.
- Levy SM, Warren JJ, Broffitt B, Hillis SL, Kanellis MJ. Fluoride, beverages and dental caries in the primary dentition. *Caries Res* 2003;37:157–165. [PubMed: 12740537]
- Lingstrom P, van Houte J, Kashket S. Food starches and dental caries. *Crit Rev Oral Bio Med* 2000;11:366–80. [PubMed: 11021636]
- Marshall TA, Levy SM, Broffitt B, Warren JJ, Eichenberger-Gilmore JM, Burns TL, Stumbo PJ. Dental caries and beverage consumption in young children. *Pediatr* 2003;112(3 Pt 1):184–191.
- Mascarenhas AK. Oral hygiene as a risk indicator of enamel and dentin caries. *Community Dent Oral Epidemiol* 1998;26:331–339. [PubMed: 9792125]
- Guidelines on overweight and obesity: electronic textbook. http://www.nhlbi.nih.gov/guidelines/obesity/e_txtbook/apndx/apndx5.htm
- Nyvad B, Fejerskov O. Active root surface caries converted into inactive caries as a response to oral hygiene. *Scand J Dent Res* 1986;94:281–284. [PubMed: 3461550]

- Ogden CL, Flegal KM, Carroll MD, Johnson CL. Prevalence and trends in overweight among US children and adolescents, 1999-2000. *JAMA* 2002;288:1728-1732. [PubMed: 12365956]
- Payette M, Brodeur JM. Comparison of dental caries and oral hygiene indices for 13- to 14- year-old Quebec children between 1977 and 1989-1990. *J Can Dent Assoc* 1992;58:921, 926, 932-922. [PubMed: 1292862]
- Pitts N. ICDAS: An international system for caries detection and assessment being developed to facilitate caries epidemiology, research and appropriate clinical management [editorial]. *Community Dent Health* 2004;21:193-198. [PubMed: 15470828]
- Podshadley AG, Haley JV. A method for evaluating oral hygiene performance. *Public Health Rep* 1968;83:259-264. [PubMed: 4967036]
- Ramfjord SP. Indices for prevalence and incidence of periodontal disease. *J Periodontol* 1959;30:51-59.
- Research Triangle Institute. SUDAAN Users' Manual, release 9.01, SAS-callable version. Research Triangle Institute; Research Triangle Park, NC: 2005.
- Ruottinen S, Karjalainen S, Pienihakkinen K, Lagstrom H, Niinikoski H, Salminen M, Ronnema T, Simell O. Sucrose intake since infancy and dental health in 10-year-old children. *Caries Res* 2004;38:142-148. [PubMed: 14767171]
- SAS Institute. SAS Release 9.0. SAS Institute; Cary NC: 2005.
- Sheiham A. Dietary effects on dental diseases. *Public Health Nutr* 2001;4(2B):569-591. [PubMed: 11683551]
- Togo P, Heitmann BL, Sorensen TI, Osler M. Consistency of food intake factors by different dietary assessment methods and population groups. *Br J Nutr* 2003;90:667-678. [PubMed: 13129474]
- Dietary guidelines for Americans 2005; key recommendations. <http://www.health.gov/dietaryguidelines/dga2005/recommendations.htm>

Table 1

Demographic characteristics and Body Mass Index (BMI)* related to caries experience (D_1^{MFS})** among participating adults with at least one tooth (n = 1,005) in Phase 1 of data collection by the Detroit Center for Research on Oral Health Disparities.

Characteristics of Adult Caregivers	N	Weighted Percent and S.E.	Weighted mean D_1^{MFS} and S.E.
Sex			
Male	55	5.5 ± 0.8	41.2 ± 3.9
Female	950	94.5 ± 0.8	47.8 ± 1.3
Age			
14 - 24 yrs	342	34.0 ± 1.8	34.9 ± 1.5
25 - 34 yrs	464	46.2 ± 2.1	47.5 ± 1.7
35 - 44 yrs	143	14.2 ± 1.0	68.7 ± 3.1
45 - 54 yrs	42	4.2 ± 0.5	73.4 ± 6.0
55+ yrs	14	1.4 ± 0.5	98.6 ± 10.9
Education			
7 - 12 yrs (less than high school)	456	45.5 ± 1.7	46.9 ± 1.7
13 - 14 yrs (high school graduate)	327	32.2 ± 1.9	47.2 ± 1.8
15 yrs (some college, no degree)	194	19.5 ± 1.9	47.3 ± 3.2
16 - 19 yrs (associate degree or higher)	28	2.8 ± 0.7	60.6 ± 4.3
Employment			
Yes (full time or part time)	389	38.7 ± 2.3	45.3 ± 1.8
No	616	61.3 ± 2.3	48.9 ± 1.5
Family Income			
< \$10,000	440	43.8 ± 2.4	47.1 ± 1.9
\$10,000 - \$19,000	278	27.7 ± 1.6	48.3 ± 2.3
\$20,000 - \$29,000	164	16.3 ± 1.1	46.2 ± 2.5
\$30,000 - \$39,000	85	8.4 ± 1.0	45.3 ± 2.7
\$40,000 or more	38	3.8 ± 0.7	55.6 ± 4.1
CDC Body Mass Index (BMI) level*			
Less than healthy (BMI < 18.5)	23	2.1 ± 0.6	54.2 ± 36.4
Healthy (BMI 18.5 to <25)	263	26.3 ± 1.7	47.6 ± 36.7
Overweight (BMI 25 to <30)	242	24.8 ± 1.4	49.5 ± 46.8
Obese (BMI 30+)	477	46.8 ± 1.5	46.1 ± 29.3

* Centers for Disease Control and Prevention, <http://www.cdc.gov/nccdphp/dnpa/bmi/bmi-adult.htm> Accessed June 9, 2005.

** Decayed, Missing, and Filled tooth surfaces in Adults, "D₁" signifies noncavitated as well as cavitated carious lesions.

Table 2

Mean D₁MFS scores for adults (n = 821) by tertiles of the proportion of energy (energy%) from carbohydrates.

Energy% from all carbohydrates	n	Mean D ₁ MFS	Standard Deviation	p
Lowest tertile ($\leq 44.1\%$ energy from carbohydrate)	272	45.6	28.5	0.03
Middle tertile ($44.1\% - \leq 50.4\%$ energy from carbohydrate)	271	51.3	32.4	
Highest tertile ($\geq 50.4\%$ energy from carbohydrate)	266	49.7	29.4	

* There were 12 participants with some missing data not included.

Table 3

Mean D₁MFS scores for adults (n = 821) by tertiles of the proportion of energy (energy%) from sugars.

Energy% from sugars	n	Mean D ₁ MFS	Standard Deviation	p
Lowest tertile (\leq 9.6% energy from sugars)	268	44.4	26.9	0.001
Middle tertile (9.6% - \leq 17.8% energy from sugars)	267	49.7	31.3	
Highest tertile (\geq 17.8% energy from sugars)	274	52.4	31.7	

* There were 12 participants with some missing data not included.

Table 4Mean gingival plaque scores for adults (n = 821) by tertiles of D₁MFS scores.

Mean D ₁ MFS	N*	Mean Gingival Plaque Score	Standard Deviation	p
Lowest tertile (<31)	268	0.76	0.37	0.02
Middle tertile (31 - <53)	261	0.79	0.37	
Highest tertile (53+)	260	0.88	0.40	

* There were 32 participants with some missing data not included.

Table 5Full regression model for caries predictors ($R^2 = 0.24$)

Independent Variables and Effects	Beta Coefficient	SE Beta	p
Intercept	4.62	0.34	0.00
Liquid Factor 1	-0.17	0.22	0.46
Liquid Factor 2	0.33	0.21	0.13
Liquid Factor 3	-0.07	0.15	0.67
Liquid Factor 4	0.18	0.21	0.38
Solid Food Factor 1	-0.04	0.21	0.85
Solid Food Factor 2	0.09	0.24	0.71
Solid Food Factor 3	-0.27	0.21	0.22
Solid Food Factor 4	0.17	0.21	0.43
Soft Drinks	0.02	0.01	0.02
Education: Not High School Graduate	Reference	-	-
Education: High School Graduate	0.02	0.16	0.92
Education: Some beyond High School	-0.01	0.24	0.97
Annual Income below \$10,000	Reference	-	-
Annual Income \$10,000 - \$19,999	-0.06	0.21	0.78
Annual Income \$20,000+	-0.01	0.22	0.98
Employment Yes	Reference	-	-
Employment No	0.31	0.18	0.10
Age Group <26 years	Reference	-	-
Age Group 26 - 30 years	0.54	0.24	0.03
Age Group 31+ years	2.01	0.20	0.00
No. times teeth brushed last week	0.01	0.02	0.51
Gingival Plaque Score	1.02	0.25	0.00

Table 6Final regression model for caries predictors by backward elimination selection procedure ($R^2 = 0.24$).

Independent Variables and Effects	Beta Coefficient	SE Beta	p
Intercept	4.60	0.24	0.00
Education: Not High School Graduate	0.00	0.00	-
Education: High School Graduate	0.13	0.15	0.39
Education: Some beyond High School	0.10	0.22	0.66
Annual Income below \$10,000	0.00	0.00	-
Annual Income \$10,000 - \$19,999	-0.05	0.21	0.81
Annual Income \$20,000+	-0.04	0.22	0.87
Employment Yes	0.00	0.00	-
Employment No	0.29	0.18	0.12
Age Group <26 years	0.00	0.00	-
Age Group 26 - 30 years	0.57	0.22	0.02
Age Group 31+ years	2.10	0.20	0.00
Gingival Plaque Score	1.03	0.22	0.00
Soft Drinks	0.02	0.01	0.01