

# Pannexin-1 mediates large pore formation and interleukin-1 $\beta$ release by the ATP-gated P2X<sub>7</sub> receptor

Pablo Pelegrin and Anmarie Surprenant\*

Department of Biomedical Science, University of Sheffield, Sheffield, UK

**P2X<sub>7</sub> receptors are ATP-gated cation channels; their activation in macrophage also leads to rapid opening of a membrane pore permeable to dyes such as ethidium, and to release of the pro-inflammatory cytokine, interleukin-1 $\beta$  (IL-1 $\beta$ ). It has not been known what this dye-uptake path is, or whether it is involved in downstream signalling to IL-1 $\beta$  release. Here, we identify pannexin-1, a recently described mammalian protein that functions as a hemichannel when ectopically expressed, as this dye-uptake pathway and show that signalling through pannexin-1 is required for processing of caspase-1 and release of mature IL-1 $\beta$  induced by P2X<sub>7</sub> receptor activation.**

*The EMBO Journal* (2006) 25, 5071–5082. doi:10.1038/sj.emboj.7601378; Published online 12 October 2006

**Subject Categories:** signal transduction; immunology

**Keywords:** hemichannel; inflammasome; ion channel; pro-inflammatory cytokines; purine receptors

## Introduction

Interleukin-1 $\beta$  (IL-1 $\beta$ ) is considered the initiator of the acute inflammatory response (Dinarello, 2005). Bacterial endotoxins and other inflammatory stimuli initiate its synthesis as a 31 kDa, biologically inactive, precursor molecule (pro-IL-1 $\beta$ ), which can then be cleaved to the bioactive 17 kDa mature IL-1 $\beta$  by proteolytically active caspase-1 (Martinon and Tschopp, 2004; Dinarello, 2005; Ferrari *et al.*, 2006). However, in the absence of a secondary stimulus, the processing and release of bioactive IL-1 $\beta$  is extremely inefficient. For example, more than 95% of the pro-IL-1 $\beta$  induced by the endotoxin lipopolysaccharide (LPS) remains unprocessed and dispersed throughout the cytoplasm until a secondary stimulus rapidly (within 2–5 min) induces the processing and release of bioactive IL-1 $\beta$  (Perregaux and Gabel, 1994; Mackenzie *et al.*, 2001; Kahlenberg and Dubyak 2004; Ferrari *et al.*, 2006). While viral and bacterial elements can subserve this role (Martinon and Tschopp, 2004; Dinarello, 2005), the only known physiological stimuli for this rapid processing and release of bioactive IL-1 $\beta$  are allogenic cytotoxic T cells and activation of the ATP-gated P2X<sub>7</sub> receptor (P2X<sub>7</sub>R) (Perregaux and Gabel, 1994; Perregaux *et al.*, 1996;

Mackenzie *et al.*, 2001; Kahlenberg and Dubyak 2004; Ferrari *et al.*, 2006).

The P2X<sub>7</sub>R, expressed on macrophage and other immune cells, is an ion channel gated by high concentrations of extracellular ATP, concentrations that are known to be present at sites of injury and inflammation (North, 2002; Ferrari *et al.*, 2006). Recent studies using transgenic mice lacking the P2X<sub>7</sub>R have most clearly demonstrated that this plasma membrane protein is directly responsible for ATP-mediated processing and release of IL-1 $\beta$  from LPS-primed cells (Solle *et al.*, 2001). These mice also exhibit altered bone formation and decreased inflammatory responses (Solle *et al.*, 2001; Ke *et al.*, 2003), making the P2X<sub>7</sub>R an attractive new target for anti-inflammatory drug discovery. Molecular mechanisms by which P2X<sub>7</sub>R activation couples to IL-1 $\beta$  processing and release remain unclear.

The P2X<sub>7</sub>R is unique among ion channels in that its activation not only opens a 'typical' ion channel selective for small cations including calcium but also leads to the gradual opening over seconds to minutes of a larger pore that allows passage of molecules up to 900 Da (North, 2002; Ferrari *et al.*, 2006). The 'large pore' properties of this receptor are usually assayed by uptake of fluorescent dyes such as ethidium and YoPro1 (North, 2002). Conventionally, it had been thought that the ion channel itself dilated to allow dye uptake; however, several recent studies have concluded that a distinct protein, or proteins, activated by P2X<sub>7</sub>R, more likely form this dye-uptake path (Schilling *et al.*, 1999; North, 2002; Faria *et al.*, 2005; Jiang *et al.*, 2005). It is not known whether this dye-uptake path plays a role in immune cell function but identification of a distinct P2X<sub>7</sub>-activated large pore, if present, should lead to significant new insight into the mechanisms underlying ATP-gated IL-1 $\beta$  release.

We initially sought this putative P2X<sub>7</sub>-activated pore by asking whether we could find a common membrane protein present in macrophage and macrophage-like cell lines that also could form a protein–protein interaction with the P2X<sub>7</sub>R and show functional properties of a dye-uptake pore. We hypothesized that a possible candidate may be a non-selective hemichannel pore. Two unrelated protein families could fulfil such a role: connexins or pannexins. Connexins are a multigene superfamily of proteins well known to form gap junctions; gap junctions are formed between cells by docking of connexin hemichannels, which are thought to exist as undocked oligomers prior to junctional formation (Spray *et al.*, 2002; Dhein, 2004; Griffith *et al.*, 2004). However, several previous studies have ruled out involvement of connexins in P2X<sub>7</sub>R activation and dye uptake (Steinberg and Di Virgilio, 1991; Hickman *et al.*, 1996). Pannexins are a recently identified three-membered family (Panx1, Panx2, Panx3) of mammalian proteins that have low sequence homology, but general structural similarity, to a family of non-mammalian proteins (innexins) that form gap junctions in invertebrate

\*Corresponding author. Department of Biomedical Science, University of Sheffield, Western Bank, Sheffield, South Yorkshire S10 2TN, UK. Tel.: +44 114 222 2366; Fax: +44 114 222 2360; E-mail: a.surprenant@shef.ac.uk

Received: 24 July 2006; accepted: 8 September 2006; published online: 12 October 2006

tissue (Panchin, 2005; Barbe *et al.*, 2006). Although no functional role for these proteins has been demonstrated *in situ*, ectopic expression of Panx1 cDNA in oocytes results in hemichannel-like membrane currents that can act as a conduit for release of intracellular ATP in single oocytes and the formation of junctional coupling between panx1 overexpressing oocytes (Barbe *et al.*, 2006). Moreover, very recent data have provided indirect evidence that panx1 may be responsible for ATP release from erythrocytes in response to mechanical or osmotic stimulation (Locovei *et al.*, 2006a).

Here, we find that panx1 is highly expressed in human and mouse macrophage, is upregulated by LPS stimulation, and co-immunoprecipitates with the P2X<sub>7</sub>R protein. Selective inhibition of panx1 by small interference RNA (siRNA) targeting panx1 and by a panx1-mimetic inhibitory peptide inhibited P2X<sub>7</sub>R-mediated dye uptake without altering the associated membrane current or calcium influx, and without altering P2X<sub>7</sub>R protein expression. Overexpression of panx1 increased the rate of P2X<sub>7</sub>R-mediated dye uptake and induced constitutive dye-uptake in the absence of P2X<sub>7</sub>Rs. Inhibition of panx1 with siRNA or the panx1-mimetic inhibitory peptide blocked caspase-1 cleavage as well as IL-1 $\beta$  processing and release from a repertoire of LPS-primed mouse and human macrophage in response to P2X<sub>7</sub>R activation. Thus, we have identified panx1 as the large pore pathway activated by P2X<sub>7</sub>Rs and as an upstream molecule essential for activation of the caspase-1/IL-1 $\beta$  inflammasome from activated macrophage by this receptor.

## Results

### **Panx1 tissue expression and association with P2X<sub>7</sub>R**

We used RT-PCR and quantitative RT-PCR (qPCR) to examine panx1 expression in a variety of immune cells and cell lines; we found high levels of mRNA for panx1 in human and mouse monocyte, macrophage and astrocyte cells and cell lines (Figure 1A). It was also present in non-immune cell lines (HEK, HeLa), which are extensively used to study heterologously expressed P2X<sub>7</sub>Rs. Phorbol esters, which up-regulate expression of P2X<sub>7</sub>Rs in macrophage (Humphreys and Dubyak, 1998), increased panx1 mRNA by four- to six-fold with further significant increased expression after LPS stimulation (Figure 1B). Because of these findings, and because a non-junctionally coupled hemichannel might be able to function as a dye-permeable pore (Barbe *et al.*, 2006; Locovei *et al.*, 2006a), we considered panx1 a potential candidate for a role in P2X<sub>7</sub>R-mediated dye uptake. We therefore cloned human panx1 from activated THP-1 macrophage (Supplementary Figure 1A), transiently overexpressed an epitope-tagged Panx1 protein in P2X<sub>7</sub>-expressing HEK cells and found that panx1 protein co-immunoprecipitated with P2X<sub>7</sub>Rs (Figure 1C). When transfected into HEK or HeLa cells, the panx1 protein was localized to the plasma membrane (Figure 1D, left-hand panel).

Activation of P2X<sub>7</sub>Rs induces rapid cytoskeletal re-arrangement of actin and other adhesion molecules, which leads to extensive membrane blebbing within 30–120 s (Mackenzie *et al.*, 2001, 2005; Morelli *et al.*, 2003). P2X<sub>7</sub>R-induced membrane blebbing is readily reversible when receptor activation is relatively brief (<10–15 min) and the P2X<sub>7</sub>R protein can be seen to remain in the plasma membrane of the blebs


as they swell and then retract (Mackenzie *et al.*, 2001, 2005; Morelli *et al.*, 2003). Overexpressing a GFP-fused Panx1 protein, we observed identical localization during P2X<sub>7</sub>R mediated membrane blebbing; that is, panx1 tracked identically in time and space with P2X<sub>7</sub> protein during the formation and retraction of membrane blebs (Figure 1D, right-hand panel).

### **Panx1 inhibition selectively blocks P2X<sub>7</sub>R induced dye-uptake**

We generated a series of siRNAs targeting Panx1 and synthesized several peptides mimicking residues in the ectodomain of panx1 in an attempt to selectively inhibit the expression and/or function of this protein (Supplementary Figure 1A and B). One of the siRNA constructs (siRNA70) was highly effective in knocking down endogenous expression of panx1 mRNA in HEK cells and THP-1 macrophage (Figure 1E). It also completely blocked exogenous expression of panx1 protein without altering protein levels of P2X<sub>7</sub>R (Figure 1F).

Panx1 siRNA70, but not scrambled siRNA or other panx1 targeted siRNAs, markedly inhibited the rate and magnitude of ATP-induced YoPro1 or ethidium uptake without altering membrane currents (Figure 2A, C and D). The selective inhibition of dye-uptake was unequivocally evidenced by simultaneous recording of membrane currents and ethidium fluorescence from single cells during whole-cell patch-clamp experiments (Figure 2A). We also generated complete ATP concentration–response curves for membrane currents and measured maximum current density in cells treated with scrambled siRNA or siRNA70. In the same population of cells in which siRNA70 inhibited ATP-mediated dye-uptake by 60–100% ( $n=6$  experiments), no significant change in ATP EC<sub>50</sub> at membrane currents (EC<sub>50</sub> = 215 ± 22  $\mu$ M scrambled siRNA versus 201 ± 35  $\mu$ M siRNA 70,  $n=4$ ) or ATP-evoked maximum currents (145 ± 37 pA/pF,  $n=14$  scrambled siRNA versus 129 ± 42,  $n=9$  siRNA 70) were observed (Figure 2A and F).


We similarly examined the actions of several panx1-mimetic peptides and found one, <sup>10</sup>panx1, (Supplementary Figure 1B and C) that potently inhibited P2X<sub>7</sub>-mediated dye uptake without altering other aspects of P2X<sub>7</sub>R activation (Figure 2B, E and F). The generation of this panx1-mimetic inhibitory peptide was of particular significance as it allowed us to readily and reliably examine inhibition of panx1 in cells in which efficient and consistent delivery of siRNA was extremely difficult (e.g. mouse J774 macrophage and alveolar macrophage obtained from human lung resections; Figure 2E and F). ATP-evoked dye-uptake was inhibited within 5–10 min exposure to <sup>10</sup>panx1 in HEK cells transfected with rat or human P2X<sub>7</sub>Rs (Figure 2B and F), in mouse J774 macrophage and human lung alveolar macrophage (Figure 2E and F), as well as in human THP-1 macrophage and HeLa or 1321-N1 astrocytes transfected with a rat P2X<sub>7</sub> expression vector (data not shown). In none of these cells did <sup>10</sup>panx1 alter ATP-evoked membrane currents, ATP concentration–response curves (Figure 2B and F), or cytosolic calcium transients (data not shown). These results rule out the possibility that panx1 siRNA or <sup>10</sup>panx1 peptide inhibit ATP-evoked dye-uptake by inhibition of the P2X<sub>7</sub>R or ion channel *per se*. Moreover, inhibition of panx1 did not significantly alter the kinetics or magnitude of ATP-evoked



**Figure 1** Panx1 is associated with P2X<sub>7</sub>R protein and its signalling. (A) Panx1 mRNA detected by RT-PCR from human and mouse cells as indicated; THP-1 macrophages, lung alveolar macrophage, Jurkat lymphocytes and mouse J774 macrophage show highest expression. (B) RT-PCR example and qPCR summary showing panx1 mRNA upregulation by PMA (0.5  $\mu$ M for 30 min) and further increased by PMA plus LPS (100 ng/ml for 4 h). (C) HEK cells stably expressing P2X<sub>7</sub>-ee receptor were transfected with Panx1-myc and immunoprecipitation (IP) of cell lysates with anti-myc Ab carried out followed by Western blotting (WB) with anti-ee Ab. (D) Immunohistochemical localization of panx1 (red) to plasma membrane in HEK cell co-transfected with panx1-ee and eGFP (green, cytoplasmic localization, left-hand panel); photo on right is HEK P2X<sub>7</sub>-expressing cell transfected with panx1-eGFP construct showing panx1 protein in blebs induced by ATP stimulation. (E, F) siRNA knockdown of endogenous levels of panx1 mRNA in HEK cells (E) and exogenously expressed panx1 protein (F). mRNA levels show mean  $\pm$  s.e.m. from qPCR assays for four separate experiments,  $P < 0.001$  for panx1 mRNA levels of cells transfected with siRNA70 versus scrambled siRNA. For WB, lysates were immunoblotted with anti-ee (panx1), anti-P2X<sub>7</sub>R and anti- $\beta$ -actin as loading control; note panx1 siRNA did not alter levels of P2X<sub>7</sub>R.

membrane blebbing ( $n = 8$ ), or phosphatidylserine exposure as assayed by annexin-V-FITC binding ( $n = 3$ ) (Mackenzie *et al*, 2001; Suadicani *et al*, 2006). Neither panx1 siRNA nor the panx1 inhibitory peptide altered membrane properties of HEK, HeLa or J774 cells (resting conductance, membrane capacitance and current-voltage relations). All experiments described above were carried out on physically isolated, uncoupled cells, but we also examined the effects of panx1 siRNA and <sup>10</sup>panx1 peptide on electrical coupling in clusters

of HEK293 cells. No change in membrane capacitance occurred in the presence of panx1 inhibition by either siRNA ( $n = 12$ ) or <sup>10</sup>panx1 peptide (200  $\mu$ M,  $n = 4$ ), whereas the nonspecific gap junction blocker heptanol (200  $\mu$ M) reversibly abolished junctional coupling within 5 min (membrane capacitance before/after heptanol  $68 \pm 3$  pF/  $11 \pm 0.7$  pF,  $n = 5$ ). The specific connexin 32-mimetic inhibitory peptide, gp27 (300  $\mu$ M) (Gudipaty *et al*, 2003), partially reduced electrical coupling (capacitance before/after




**Figure 2** Selective inhibition of Panx1 blocks P2X<sub>7</sub>-R-induced dye uptake but not ionic currents. (A) Representative traces of simultaneous recording of membrane current and ethidium uptake from control or panx1 siRNA70 transfected HEK cell expressing P2X<sub>7</sub>R in response to 3 mM ATP stimulation. (B) Similar recordings obtained from control cell and cell exposed to <sup>10</sup>panx1 peptide (200  $\mu$ M for 10 min). (C) Representative photos (5 min in ATP) and kinetic traces of ethidium uptake recorded from populations of P2X<sub>7</sub>R expressing or nonexpressing HEK cells as indicated after transfection with scrambled or panx1 siRNA70. Digitonin (100  $\mu$ M) applied at end of experiment to induce maximum dye uptake; siRNA did not alter ethidium fluorescence after digitonin treatment ( $n=20$  cells for each treatment and representative of 7 independent experiments). (D) Summary of all dye uptake obtained from siRNA experiments illustrated in A and C. Histograms show slope of dye uptake in response to maximum concentration of ATP (5 mM) from cells transfected with scrambled (control) or panx1-targetted siRNA; siRNA70 reduced ATP-evoked dye uptake by >80%. \*\* $P<0.001$ . (E) Kinetic traces of ethidium uptake from mouse J774 or human alveolar macrophage in the absence or presence of <sup>10</sup>panx1-mimetic peptide (100  $\mu$ M for 20 min); each trace is mean  $\pm$  s.e.m. of 50 cells in field of view and representative of 2–6 independent experiments. (F) Plots EC<sub>50</sub> concentration of ATP, maximum current amplitude and dye uptake induced by maximum ATP concentration (expressed as % control response) from P2X<sub>7</sub>R-expressing HEK cells transfected with panx1 siRNA70, and from HEK, J774 and human alveolar macrophage incubated for 10–30 min with <sup>10</sup>panx1 peptide as indicated. In all cases dye-uptake was reduced by >80% without alteration of membrane currents.

59  $\pm$  4 pF/26  $\pm$  3 pF,  $n=4$ ) in similar clusters of HEK cells. These control experiments confirm the specificity of the panx1 siRNA and peptide mimetic, and also rule out involvement of panx1 in formation of gap junctions in these cells.

**Panx1 is a non-selective hemichannel when overexpressed in mammalian cells**

Ectopic expression of panx1 in oocytes results in the appearance of hemichannel-like currents (Bao *et al*, 2004; Bruzzone



**Figure 3** Overexpression of panx1 induces constitutive dye-uptake and hemichannel-like currents selectively blocked by <sup>10</sup>panx1 and CBX. (A) Ethidium uptake in P2X<sub>7</sub>-negative HEK cells transfected with empty or panx1 expression vector; superfusion of cells with ethidium resulted in immediate uptake in panx1 transfected cells; each trace is mean  $\pm$  s.e.m. of 50 cells in field of view. (B) Similar experiment in P2X<sub>7</sub>-positive HEK cells; ethidium application did not result in constitutive dye uptake but the ATP-evoked response showed faster kinetics and maximum fluorescence; traces are mean  $\pm$  s.e.m. of 50 cells. Western blots obtained from cells used in these experiments are shown; panx1 overexpression did not change P2X<sub>7</sub>R protein expression. (C) Currents in response to ramp voltages (-120 to 80 mV) from control (GFP-transfected) or panx1 transfected HEK cell in normal extracellular solution, in solution containing the larger cation NMDG in place of sodium, in the presence of lanthanum (100  $\mu$ M), gadolinium (100  $\mu$ M) or CBX (5  $\mu$ M) as indicated. (D) Currents from panx1 transfected HEK cell in 0 (control), 5 and 12 min in presence of 100  $\mu$ M <sup>10</sup>panx1 peptide and 15 min after washout; complete inhibition was observed within 10–15 min. (E) Summary of all similar experiments where current at 60 mV is shown as % of control response; only <sup>10</sup>panx1 peptide and CBX (20  $\mu$ M) inhibited the current while other connexin hemichannel inhibitors (heptanol, gp27 peptide and replacing sodium with calcium) or Trp channel inhibitors (NMDG, lanthanum, gadolinium) were without effect. Removal of extracellular calcium, which activates connexin hemichannel currents, also was without effect on panx1 currents ( $n = 24$  for CBX and lanthanum, nine for <sup>10</sup>panx1 peptide and six for others).

*et al*, 2005; Panchin, 2005; Barbe *et al*, 2006). The existence of similar endogenous hemichannel currents in nonjunctional cells has not been reported, and nor have we observed similar currents, or constitutive dye uptake, in human or mouse macrophage, or any cell line we have used for transfection of P2X<sub>7</sub>Rs (Figure 3A and C). Thus, if indeed, such a non-selective hemichannel is involved in P2X<sub>7</sub>R dye-uptake, it must be nonfunctional in the absence of receptor activation. Therefore, we asked whether overexpression of panx1 protein resulted in hemichannel-like function in mammalian cells. Overexpression of panx1 protein in P2X<sub>7</sub>-negative HEK or Hela cells resulted in a constitutive uptake of ethidium that

was approximately 5–20% of the ATP-evoked ethidium uptake observed in P2X<sub>7</sub>-expressing cells (Figure 3A). In contrast, when panx1 was overexpressed in P2X<sub>7</sub>-positive cells, there was no constitutive ethidium uptake but the rate of dye uptake in response to ATP was significantly increased (slope of fluorescence increase =  $40 \pm 1$  for panx1 overexpression versus  $29 \pm 2.5$  for vector transfected,  $n = 4$ ; Figure 3B). Panx1 overexpression in P2X<sub>7</sub> expressing cells did not alter ATP-induced membrane currents recorded at -60 mV (no significant difference in ATP concentration-response or maximum current density;  $n = 12$ ). These results suggest that unstimulated P2X<sub>7</sub>R may negatively regulate panx1 function.

Overexpression of panx1 in non-P2X<sub>7</sub> expressing cells resulted in the appearance of membrane currents similar in most respects to those observed in previous studies using the oocyte expression system (Bao *et al.*, 2004; Bruzzone *et al.*, 2005; Barbe *et al.*, 2006). These currents were activated by depolarization, and showed a nonselective ionic permeability profile (currents were unaltered by replacing sodium with the larger cation, NMDG, or with K<sup>+</sup>, Ca<sup>2+</sup> or choline and they were also unaltered by changes in intracellular or extracellular Cl<sup>-</sup>, *n*=6 for each condition; Figure 3C and E). In agreement with results from oocyte studies (Bruzzone *et al.*, 2005), panx1-mediated currents were effectively blocked by relatively low concentrations of the nonspecific gap-junction blocker, carbenoxolone (CBX; Figure 3C and E), but not by other generally used connexin channel/gap junction blockers (heptanol, mefloquine and connexin-mimetic inhibitory peptide gp27; Figure 3E). They were also unaffected by high concentrations of lanthanum or gadolinium (0.1–1 mM; Figure 3C and E). The panx1-currents were readily and reversibly blocked by the selective panx1-mimetic inhibitory peptide <sup>10</sup>panx1 (Figure 3D and E). Steady-state inhibition by <sup>10</sup>panx1 required 4–15 min. Concentration–inhibition curves were obtained for the actions of <sup>10</sup>panx1 in inhibiting P2X<sub>7</sub>-mediated dye uptake and in inhibiting panx-1 mediated currents in panx-1 overexpressing cells; in both assays a similar half-maximal inhibitory concentration was obtained (30–50  $\mu$ M; Supplementary Figure 2A). These concentrations are within the range (50–1000  $\mu$ M) typically seen to be effective in other systems where peptide-mimetic inhibitors are commonly used (Hickman *et al.*, 1996; Gudipaty *et al.*, 2003; Mariathasan *et al.*, 2004; Elliott *et al.*, 2005). Steady-state inhibition by an IC<sub>50</sub> concentration of <sup>10</sup>panx1 required 10–20 min with washout being accomplished within 15–25 min (*n* = 4).

We also examined the actions of the nonspecific gap junction/hemichannel blockers on P2X<sub>7</sub>R mediated responses in our repertoire of immune and nonimmune cells in the absence of panx1 overexpression. CBX, but not heptanol, mefloquine or gp27, selectively inhibited the ATP-evoked dye uptake in all these cells (Figure 4A, D and E). In contrast, CBX produced a small, but significant *increase* in the associated ATP-evoked membrane currents and cytosolic calcium transients (Figure 4A and B). The selective inhibition of P2X<sub>7</sub>-mediated dye-uptake by CBX occurred over the same concentration range as did the CBX inhibition of the panx1-mediated currents in panx1-overexpressing cells with half-maximal inhibition occurring at 2–4  $\mu$ M CBX (Figure 4D). These results not only conclusively rule out involvement of connexins in P2X<sub>7</sub>R-induced dye-uptake but also suggest the inhibition of ATP-evoked dye-uptake by CBX may be due to inhibition of panx1. It is important to emphasize that all our experiments were carried out only on physically isolated, electrically uncoupled cells, thus obviating any complications due to junctional coupling between cells. In this regard, a recent study using confluent, electrically coupled, 1321-N1 astrocytes expressing P2X<sub>7</sub>Rs found that CBX, heptanol and mefloquine all equally inhibited ATP-evoked dye-uptake and cytosolic calcium transients, and concluded that these nonselective connexin channel blockers acted as P2X<sub>7</sub>R antagonists (Suadicani *et al.*, 2006). However, recordings of membrane currents, calcium transients and dye-uptake in physically isolated, single, 1321-N1 astrocytes

transfected with P2X<sub>7</sub>R clearly show that none of these nonselective connexin channel blockers inhibit P2X<sub>7</sub>Rs (Figure 4B, C and E).

### **Panx1 is required for release of mature IL-1 $\beta$ in response to P2X<sub>7</sub>R activation**


We next asked whether panx1 contributes to the physiological response in immune cells by measuring IL-1 $\beta$  release and processing from LPS-primed mouse J774 macrophage, human THP-1 cells and acutely isolated alveolar macrophage from human lung in response to P2X<sub>7</sub>R stimulation. Panx1 protein knockdown using siRNA70 transfection in human THP-1 cells significantly inhibited ATP-mediated IL-1 $\beta$  release while similar treatment with scrambled siRNA did not (Figure 5A). Neither of these siRNAs stimulated IL-1 $\beta$  release in the absence of P2X<sub>7</sub>R stimulation (Figure 5A).

The panx1-mimetic inhibitory peptide, <sup>10</sup>panx1, blocked ATP-mediated IL-1 $\beta$  release in mouse and human macrophage (Figure 5B–D); we also confirmed that the actions of ATP to induce IL-1 $\beta$  release were due to P2X<sub>7</sub>R activation by the use of the anti-P2X<sub>7</sub>R monoclonal antibody (Figure 5B). CBX also significantly inhibited ATP-mediated release from these macrophages although maximal inhibition by CBX was not as effective as <sup>10</sup>panx1 (Figure 5B). We used the connexin-mimetic inhibitor, gp27 (Spray *et al.*, 2002; Dhein, 2004), to confirm the lack of involvement of connexins and also as a control peptide; this peptide resulted in a small, but significant, increase in amounts of IL-1 $\beta$  secreted into the medium (Figure 5B). As expected from much previous work (Perregaux and Gabel, 1994; Mackenzie *et al.*, 2001; Gudipaty *et al.*, 2003; Kahlenberg and Dubyak, 2004; Ferrari *et al.*, 2006), the IL-1 $\beta$  released into the medium by activation of P2X<sub>7</sub>Rs was primarily the fully processed, 17 kDa form, although the unprocessed 31 kDa form was also often present in the medium as well (Figure 6A–C); however, no 17 kDa band was present in the medium after ATP stimulation when cells were incubated with <sup>10</sup>panx1 (Figure 6A–C) or with CBX (Figure 6B).

Because intracellular potassium depletion has been hypothesized as the underlying mechanism responsible for P2X<sub>7</sub>R-mediated release of bioactive IL-1 $\beta$  (Perregaux and Gabel, 1994; Martinon and Tschoop, 2004; Dinarello, 2005; Ferrari *et al.*, 2006), we asked whether inhibition of pannexin altered K<sup>+</sup> levels by direct measurement of cellular and released potassium from J774 cells during our cytokine release protocols. In agreement with previous studies (Perregaux and Gabel, 1994; Perregaux *et al.*, 1996; Kahlenberg and Dubyak, 2004), we found that ATP reduced intracellular K<sup>+</sup> by approximately 50% in LPS-primed cells with concomitant increased potassium in the medium, but the panx1-mimetic inhibitory peptide did not alter the P2X<sub>7</sub>R-mediated intracellular K<sup>+</sup> depletion (Figure 6D). Thus, intracellular K<sup>+</sup> depletion *per se* is not sufficient to induce P2X<sub>7</sub>R-evoked processing and release of IL-1 $\beta$  from LPS-primed macrophage.

### **Panx1 is required for caspase-1 cleavage in response to P2X<sub>7</sub>R activation**

Finally, we asked whether panx1 may be involved further upstream in the caspase cascade using immunoblot assays of caspase-1. We compared actions of <sup>10</sup>panx1 inhibitory peptide with actions of the selective caspase-1 inhibitor,




**Figure 4** CBX but not other connexin channel blockers blocks ATP-mediated dye-uptake without inhibiting P2X<sub>7</sub>R activation. (A, B) Membrane currents, YOPRO-1 uptake and Fluo4 calcium transients (as indicated) recorded from HEK (A) or 1321-N1 (B) cells expressing P2X<sub>7</sub>R. CBX (20 μM) effectively blocked YOPRO-1 uptake without altering currents or calcium transients. All recordings obtained from physically isolated, single cells. (C) ATP concentration–response curves for membrane current (recorded at –60 mV holding potential) from all experiments as illustrated in (A) and (B), in control (closed symbols) and in 20 μM CBX (open symbols) for HEK cells expressing rat or human P2X<sub>7</sub>R, for 1321-N1 cells expressing human P2X<sub>7</sub>R and for mouse J774 macrophage, as indicated. CBX had no effect. (D) Concentration–response curve for CBX inhibition of dye uptake (circles) in P2X<sub>7</sub>-expressing cells and for inhibition of panx1 currents recorded from HEK cells (squares) transfected with panx1 expression vector. (E) Summary of actions of several nonspecific connexin channel blockers to inhibit P2X<sub>7</sub>R-induced dye uptake from cells ectopically (HEK and 1321-N1) or endogenously (mouse J774 and human alveolar macrophage) expressing P2X<sub>7</sub>R. Concentrations used were CBX (20 μM), heptanol (200 μM), mefloquine (100 μM) and gp27 (1 mM); each value is mean ± s.e.m. from 4 to 12 experiments.

Ac-YVAD-AOM, which blocks caspase-1 cleavage and subsequent caspase-1 dependent processing and release of IL-1 $\beta$  (Nemeth *et al.*, 1997). Both <sup>10</sup>panx1 peptide and Ac-YVAD-AOM abolished ATP-mediated intracellular cleavage of caspase-1 as evidenced by absence of the caspase-1 p10 fragment in cells incubated with either of these compounds (Figure 7A), as well as IL-1 $\beta$  release of mature IL-1 $\beta$  into the medium (Figure 7B). Lactate dehydrogenase (LDH) release was also measured in each caspase and cytokine

release experiment as an assay for cell death but under no condition did LDH levels change significantly (Supplementary Figure 2B).

## Discussion

While dye-uptake has been a defining feature of, and useful assay for, the P2X<sub>7</sub>R for decades (precloning designated P2Z receptor), molecular mechanisms underlying this process and




**Figure 5** Panx1 blockade inhibits ATP-mediated IL-1 $\beta$  release from activated macrophage. (A) IL-1 $\beta$  release from LPS-primed THP-1 macrophages after transfection with scrambled or panx1 siRNA70. Neither siRNA alone induced IL-1 $\beta$  release nor did the scrambled siRNA alter ATP-induced release relative to untransfected cells, but panx1 siRNA70 significantly inhibited ATP-induced release; results are representative of three similar experiments.  $**P < 0.005$ . (B) IL-1 $\beta$  release from LPS-primed THP-1 macrophages under conditions indicated. CBX (20  $\mu$ M), <sup>10</sup>pan1 peptide (200  $\mu$ M) and the monoclonal anti-P2X<sub>7</sub>R Ab, but not the connexin blocking peptide gp27 (1 mM), significantly inhibited ATP-evoked IL-1 $\beta$  release. (C, D) Similar experiments performed on mouse J774 macrophage and human alveolar macrophage; <sup>10</sup>panx1 inhibitory peptide completely blocked ATP-mediated IL-1 $\beta$  release from these cells. Results are representative of 3–6 similar experiments in each case; error bars are standard deviation of triplicate samples.  $**P < 0.005$ .

whether it relates to immune cell function have remained unknown (Perregaux and Gabel, 1994; Hickman *et al*, 1996; North, 2002; Ferrari *et al*, 2006). A functional link between P2X<sub>7</sub>R-activated large pore formation and cytokine release in macrophage has long been postulated (see Ferrari *et al*, 2006); here we show that pannexin-1 is this functional link. Selective blockade of panx1 protein with panx1 siRNA or the panx1-mimetic inhibitory peptide revealed the essential role of this membrane protein in P2X<sub>7</sub>R-activation of the caspase-1 cascade. Our results suggest that panx1 may act as a nonselective hemichannel activated by P2X<sub>7</sub>R stimulation and that it is this hemichannel activity, rather than ion flow

through the P2X<sub>7</sub>R channel itself, that is responsible for activation of the caspase-1 cascade through the recently identified cryopyrin inflammasome (Mariathasan *et al*, 2006) in macrophage.

Early studies of ATP-activated dye uptake in macrophage had suggested that connexin hemichannels, which are permeable to molecules up to about 1000 Da, were themselves the P2Z/P2X<sub>7</sub>R (Beyer and Steinberg, 1991). This idea was ruled out by the molecular identification of the P2X<sub>7</sub>R as a member of the P2X family of ion channels (North, 2002) and by results showing that ATP does not induce dye-uptake in P2X<sub>7</sub>-receptor negative cells where connexin proteins are






**Figure 6** Panx1 blockade inhibits IL-1 $\beta$  processing but not intracellular K<sup>+</sup> depletion from LPS-primed macrophage. Western blot analysis of cell lysate and medium from human THP-1 macrophages (A), mouse J774 macrophage (B) and human alveolar macrophage (C). No 17 kDa form was present intracellularly under any condition, while medium contained both pro-IL-1 $\beta$  and fully processed, 17 kDa, IL-1 $\beta$  only after LPS priming and ATP treatment. No processed 17 kDa IL-1 $\beta$  was observed in the medium after treatment with <sup>10</sup>panx1 inhibitory peptide. (D) Intracellular and released K<sup>+</sup> (plotted as fraction of control) from parallel wells of J774 cells treated as in (B), *n* = 4.

endogenously expressed or heterologously overexpressed (Steinberg and Di Virgilio, 1991). Subsequently, two hypotheses have emerged to account for the dye-uptake: intrinsic P2X<sub>7</sub>R channel dilatation from cation-selective small channel to nonselective big pore or P2X<sub>7</sub>-receptor activated recruitment of an accessory pore-forming mechanism (Schilling *et al*, 1999; North, 2002; Faria *et al*, 2005; Jiang *et al*, 2005). The present results decisively support the latter conclusion and identify panx1 as the critical accessory molecule. Selective inhibition of panx1 expression by siRNA knock-down blocked P2X<sub>7</sub>-mediated dye-uptake independently of P2X<sub>7</sub>R activation; similar results were obtained with the specific panx1-mimetic inhibitory peptide. Overexpression of panx1 in mammalian cells resulted in the appearance of depolarization-evoked membrane currents that exhibited biophysical properties previously associated with hemichannel function (non-selective anion/cation permeability to molecules < 1 kDa) (Spray *et al*, 2002; Dhein, 2004; Griffith *et al*, 2004; Bruzzone *et al*, 2005) as well as constitutive ethidium uptake in P2X<sub>7</sub>-negative cells. The biophysical properties (large cation permeability) of the overexpressed pannexin-1 currents matched the properties of the ATP-evoked dye

uptake pathway in P2X<sub>7</sub>R-expressing cells. However, these currents were observed only with overexpression of panx1 but not in cells in which only endogenous levels of panx1 were present. If panx1 does function directly as a hemichannel, these findings indicate that panx1 is nonfunctional in the absence of P2X<sub>7</sub>R stimulation. Our data are compatible with a negative regulation of panx1 activity by unstimulated P2X<sub>7</sub>Rs, in that constitutive dye uptake was observed when panx1 was overexpressed in P2X<sub>7</sub>-negative cells but not in P2X<sub>7</sub>-positive cells even though dye uptake was significantly increased by P2X<sub>7</sub>R activation in the same cells. Overall, our data suggest that the panx1 protein, acting as a plasma membrane hemichannel, is in fact the dye-uptake pore. However, we cannot rule out an alternative possibility that panx1 allows dye uptake by altering the function of plasma membrane lipids and membrane transporters in light of our recent study showing that P2X<sub>7</sub>R activation in T-lymphocytes leads to a reversal of activity of the multidrug efflux transporter, MDR/p-glycoprotein, via alterations in membrane lipids (Elliott *et al*, 2005).

While it is well known that gap junctions form by the docking of two connexin hemichannels from opposing faces



**Figure 7** Panx1 inhibition blocks intracellular caspase 1 processing. Western blot analysis of mouse J774 macrophage for caspase-1 (A) and IL-1 $\beta$  (B). (A) Same gel is shown, upper lanes with short exposure and lower lanes with longer exposure; we used short exposure to assess p45 pro-caspase-1 protein levels in each lane and longer exposure to reveal processing and active caspase-1 p10 fragment. After LPS priming, ATP induced strong caspase-1 processing, which was prevented by either <sup>10</sup>panx1 inhibitory peptide or the caspase-1 inhibitor, Ac-YVAD-AOM. (B) Western blot for IL-1 $\beta$  from same set of experiments confirms ATP-mediated IL-1 $\beta$  processing and release are also prevented by <sup>10</sup>panx1 inhibitory peptide or caspase-1 inhibitor.

of adjacent plasma membranes, it has been a matter of much debate as to whether open (unopposed) hemichannels have a functional role in nonjunctional cells (Dhein, 2004; Locovei *et al*, 2006a). Most recently, hemichannel-like activity has been reported in hippocampal neurones in response to oxygen/glucose deprivation, and this activity was suggested to be consistent with activation of panx1 hemichannels, a conclusion based on inhibition by carbenxolone and lanthanum (Thompson *et al*, 2006). It is unlikely that panx1 hemichannels are responsible for the current observed in hippocampal neurones because we found that lanthanum had no effect on panx1-mediated currents or dye-uptake and panx1 current-voltage curves obtained in our study, and in previous studies on panx1 expression in oocytes, differ from the observed hemichannel activity in hippocampal neurones. However, panx2 is also found in hippocampal neurones and ectopically expressed panx1 and panx2 can form heteromeric channels in oocytes with distinct biophysical properties (Barbe *et al*, 2006). Further studies, particularly making use of the selective panx1-mimetic inhibitory peptide we have identified, may reveal whether a heteromeric panx1/2 channel may underlie this neuronal hemichannel activity.

The mechanisms underlying the upregulation and synthesis of IL-1 $\beta$  in response to Toll like receptor (TLR) activation by LPS and the steps in its processing via the caspase cascade are fairly well understood (Martinon and Tschopp, 2004; Dinarello, 2005; Ferrari *et al*, 2006). LPS activation of TLR triggers synthesis of pro-IL-1 $\beta$  and its cytoplasmic accumulation. Further secondary stimuli, particularly P2X<sub>7</sub>R activation by extracellular ATP, leads to the aggregation and activation of the inflammasome, the multiprotein cytoplasmic complex required to activate caspase-1 and so initiate caspase-1 proteolysis of pro-IL-1 $\beta$  into its bioactive 17 kDa form (Martinon and Tschopp, 2004; Dinarello, 2005; Ferrari *et al*, 2006). In contrast, how the inflammasome is formed and activated in response to this secondary stimulus remains a puzzle. Most recently, at least two distinct inflammasome activation pathways subsequent to TLR signalling have been delineated (Mariathasan *et al*, 2004, 2006; Kanneganti *et al*,

2006; Martinon *et al*, 2006; Sutterwala *et al*, 2006): they are both associated with the adaptor protein ASC and involve caspase-1 cleavage but activation can be further distinguished by involvement of ICE-protease activating factor (IPAF), or by an IPAF-independent involvement of another adaptor protein, cryopyrin/NALP3/CIAS1. The latter, but not the former, pathway can be activated by P2X<sub>7</sub>R stimulation as well as Gram-positive but not Gram-negative bacteria (Mariathasan *et al*, 2004). While these recent studies have provided significant new insights into the composition of distinct inflammasomes and specificity of their activation, they have not addressed the underlying mechanisms of inflammasome activation. It has generally been accepted that P2X<sub>7</sub>R-mediated cleavage of caspase-1 is due to depletion of intracellular potassium resulting from ion flow through the small ion channel and/or the large dye-uptake pore (Gudipaty *et al*, 2003; Mariathasan *et al*, 2004; Dinarello, 2005; Ferrari *et al*, 2006). However, our results rule out intracellular K<sup>+</sup> depletion via P2X<sub>7</sub>R activation as the causative mechanism because panx1 inhibition abolished release of IL-1 $\beta$  without altering cellular K<sup>+</sup> depletion.

While detailed signalling mechanisms will require further elucidation, panx1 must now be considered the key upstream molecule in P2X<sub>7</sub>R-mediated inflammasome activation. It is tempting to speculate that panx-1 hemichannels, which can act as conduits for ATP release when overexpressed (Bao *et al*, 2004; Barbe *et al*, 2006; Locovei *et al*, 2006b), may subserve a similar role when natively activated by P2X<sub>7</sub>R in macrophage. Concentrations of extracellular ATP (> 100  $\mu$ M) required to activate P2X<sub>7</sub>R may well be higher than free intracellular ATP concentrations, particularly as > 90% of all intracellular ATP is produced in, and bound to, mitochondria (Tarasov *et al*, 2004). Because ATP can directly activate several caspases in cell-free assays (Han *et al*, 2003), and because elevations in intracellular ATP levels are associated with caspase activation and IL-1 $\beta$  release (Eguchi *et al*, 1997), the existence of a conduit for *extracellular* ATP entry into the cell would provide a novel mechanism for direct and immediate activation of caspase-1 processing by the inflammasome.

## Materials and methods

### Cells and reagents

CBX, phorbol 12-myristate 13-acetate (PMA), heptanol, mefloquine hydrochloride, ethidium bromide, ATP, gadolinium and *Escherichia coli* LPS were from Sigma, lanthanum from Fisher Bioscience, and YoPro1 from Invitrogen. THP-1 and J774 macrophage were cultured in RPMI, HeLa and 1321-N1 astrocytes in DMEM and HEK293 cells in F-12 media, all supplemented with 10% fetal calf serum (Gibco). Pure populations of human lung macrophage were isolated as described previously (Chong *et al*, 2003; Mackenzie *et al*, 2003).

### Cloning, RT-PCR and qPCR

The full-length sequence encoding for human panx1 was amplified from PMA differentiated THP1 cDNA using PCR (Supplementary Figure 1A). c-myc (EQKLISEEDL), Glu-Glu (EYMPME) or eGFP tags were introduced in frame at the 3'-end of the coding sequence using overlapping PCR and Accuzyme<sup>TM</sup> DNA polymerase (Bioline) before cloning into pcDNA3.1(-) vector (Invitrogen); hence, incorporating a c-myc, Glu-Glu (ee) or eGFP fusion tag at the C terminus of the expressed protein. Cloned products were confirmed by sequencing. Human brain RNA (Ambion) and total RNA from other cells were isolated using RNeasy Mini kit (Qiagen), followed by reverse transcription using SUPERSRIPT<sup>TM</sup> III (Invitrogen) RNase H-reverse transcriptase with oligo-dT. Specific primers for panx1 (Supplementary Figure 1A), rat P2X<sub>7</sub> (5'-TGCCAGTTCAGG GAGGAATC-3' and 5'-ACAGTTCGAAGAAGTCCG-3') or  $\beta$ -actin (5'-ATGGATGATATATCGCCG-3' and 5'-CTAGAAGCATTTCGCGTG GAC-3') were used in PCR. The obtained product sizes for panx1 (610 bp), rat P2X<sub>7</sub> (585 bp) and  $\beta$ -actin (1.1 kb) were as expected from their mRNA sequences. Relative expression levels of panx1 and P2X<sub>7</sub> mRNA were measured using an iCycler iQ real-time PCR detection system (BioRad) and iQ SYBR Green Supermix (BioRad). Specific primers for Panx-1 (Supplementary Figure 1A) and rat P2X<sub>7</sub> (5'-TGTGCTACAGGTGCTACGCC-3' and 5'-GCCCTTCACTTCC GAAACTC-3') were purchased from Sigma-Genosys, GAPDH specific primers from Qiagen (QuantiTech Primer Assays). For each primer set the efficiency was >95% and a single product was seen on melt curve analysis. Relative expression levels were calculated using the 2<sup>- $\Delta\Delta$ Ct</sup> method normalizing to GAPDH expression levels (Livak and Schmittgen, 2001).

### siRNA and peptides

Small interference RNAs against human panx1 were *Silencer*<sup>TM</sup> (Ambion) pre-designed (Supplementary Figure 1A). siRNA (200 pmol) transfection was with Lipofectamine 2000 (Invitrogen), experiments were carried out 2 days posttransfection. The panx1 mimetic blocking peptide<sup>10</sup> panx1 (WRQAAFVDSY), the connexin 32 blocking peptide gp27 (SRPTEKTVFT) and other peptides (Supplementary Figure 1B) were synthesized by Sigma-Genosys and Alta Biosciences.

### Electrophysiology and fluorescence assays

Details are as described previously (Mackenzie *et al*, 2001; Jiang *et al*, 2005). Simultaneous whole-cell patch clamp recordings, capacitance tracking and YoPro1 or ethidium fluorescence were carried out under Fluor  $\times$ 40 objective on a Zeiss Axiovert 100 using EPC9 amplifier (HEKA) and Photonics monochromator imaging system. Other dye uptake and Fluo4 calcium imaging experiments were carried out using a Nikon confocal microscope under  $\times$ 20 or  $\times$ 40 objective; 20–60 isolated (not touching) cells were imaged in each experiment, averaged, and the slope of the fluorescence signal was used as the most accurate and consistent measurement for comparisons. Measurements of fluorescence intensity (fluorescence units, f.u.) at 100 s after compound application gave similar results but showed greater inter-experiment variability. All protocols and extracellular solutions were the same under both systems: YoPro1 (2  $\mu$ M) or ethidium bromide (20  $\mu$ M) was present throughout unless otherwise stated; because YoPro1 and GFP fluorescence overlap, ethidium (red spectrum) was used in all experiments involving transient transfections as we used eGFP co-transfection to identify transfected cells. Digitonin (100  $\mu$ M) was used at the end of the experiments to induce maximum dye uptake. Except where stated otherwise,

external solution was (mM) NaCl 147, HEPES 10, glucose 13, CaCl<sub>2</sub> 2, MgCl<sub>2</sub> 1 and KCl 2; internal solution was NaCl 147 or Cs-gluconate, HEPES 10, EGTA 10 and MgCl<sub>2</sub> 3. All recordings were performed only on single, isolated cells in order to preclude effects due to gap junctions.

### Co-immunoprecipitation, Western blotting and immunohistochemistry

HEK cells stably expressing P2X<sub>7</sub>-ee tagged receptor were transfected with panx1 c-myc cDNA or empty pcDNA3.1 (used as negative controls for anti-c-myc co-immunoprecipitations) and lysed in RIPA buffer (20 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM MgCl<sub>2</sub>, 1 mM CaCl<sub>2</sub>, 1% Triton X-100) supplemented with Complete protease inhibitor cocktail (Roche). Lysates were centrifuged to remove particulate matter, resolved in 4–10% polyacrylamide gels and transferred onto PVDF membranes (Millipore) by electro-blotting. For immunoprecipitations, lysates were pre-cleaned with protein G sepharose beads (Amersham Biosciences) before immunoprecipitation with mouse anti-c-myc (Covance) or control mouse IgG (Vector) and immunoblotted with rabbit anti-ee (Bethyl) or rabbit anti-c-myc (Covance). Immunohistochemistry was carried out using rabbit anti-ee Ab and photographed under the Nikon confocal microscope.

### IL-1 $\beta$ release and K<sup>+</sup> release

J774 macrophages, human lung macrophages or PMA differentiated THP-1 macrophage were stimulated with LPS (0.1–1  $\mu$ g/ml) for 4 h (Mackenzie *et al*, 2001; Kahlenberg and Dubyak, 2004), washed and preincubated for 30 min with 1  $\mu$ g/ml of anti-P2X<sub>7</sub> antibody (Buell *et al*, 1998), 50  $\mu$ M CBX, 100  $\mu$ M of cell-permeable and irreversible inhibitor of caspase-1 (Ac-YVAD-2,6-dimethylbenzoyloxymethyl ketone (AOM), caspase-1 inhibitor IV, Calbiochem) or 500  $\mu$ M of <sup>10</sup>panx1 or 1 mM gp27 peptide and ATP applied for 30 min. Supernatants were assayed for IL-1 $\beta$  by ELISA (Mackenzie *et al*, 2001) using anti-human IL-1 $\beta$  (Endogen) as the coating antibody, anti-human IL-1 $\beta$ -biotin (Pierce) as the detection antibody for THP-1 and for human lung macrophage. J774 supernatants were assayed using Quantikine<sup>®</sup> mouse IL-1 $\beta$  kit (R&D Systems) following the manufacturer instructions. For Western blot analysis, macrophage cell extracts were prepared as detailed above and supernatants were concentrated using 10-kDa nominal molecular weight cutoff filters (Millipore) and immunoblotted with anti-IL-1 $\beta$  antibody (3ZD, provided by the Biological Resources Branch of the National Cancer Institute, Frederick Cancer Research and Development Centre, Frederick, MD) or with anti-caspase-1 p10 rabbit polyclonal antibody for mouse (Santa Cruz). Cellular and released K<sup>+</sup> were measured against standards using the Cirox Vision inductively coupled plasma-atomic emission spectrometer (Spectro Analytical UK Ltd). For these experiments, cells were lysed in 1 ml of 10% nitric acid with parallel wells used for IL-1 $\beta$  assays.

### Statistical analysis

Average results are expressed as the mean  $\pm$  s.e.m. from the number of assays indicated. Data were analyzed by an unpaired Student's *t*-test to determine difference between groups using InStat (Graph-Pad) and Excel (Microsoft) software.

### Supplementary data

Supplementary data are available at *The EMBO Journal* Online (<http://www.embojournal.org>).

## Acknowledgements

We thank L Collinson and E Martin for molecular biology and tissue culture support. We are indebted to Dr P Peachell, Royal Hallamshire Hospital, University of Sheffield for invaluable help in providing human lung preparations. The Centre for Chemical Instrumental Analysis and Services, Department of Chemistry, University of Sheffield, analysed K<sup>+</sup> content of our samples. This work supported by the Wellcome Trust and a postdoctoral fellowship to PP from AstraZeneca Charnwood.

### Competing interests statement

The authors state that they have no competing financial interests.

## References

- Bao L, Locovei S, Dahl G (2004) Pannexin membrane channels are mechanosensitive conduits for ATP. *FEBS Lett* **572**: 65–68
- Barbe MT, Monyer H, Bruzzone R (2006) Cell–cell communication beyond connexins: the pannexin channels. *Physiology* **21**: 103–114
- Beyer EC, Steinberg TH (1991) Evidence that the gap junction protein connexin-43 is the ATP-induced pore of mouse macrophages. *J Biol Chem* **266**: 7971–7974
- Bruzzone R, Barbe MT, Jakob NJ, Monyer H (2005) Pharmacological properties of homomeric and heteromeric pannexin hemichannels expressed in *Xenopus* oocytes. *J Neurochem* **92**: 1033–1043
- Buell G, Chessell IP, Michel AD, Collo G, Salazzo M, Herren S, Gretener D, Grahames C, Kaur R, Kosco-Vilbois MH, Humphrey PP (1998) Blockade of human P2X<sub>7</sub> receptor function with a monoclonal antibody. *Blood* **92**: 3521–3528
- Chong LK, Suvarna K, Chess-Williams R, Peachell PT (2003) Desensitization of  $\beta$ 2-adrenoceptor-mediated responses by short-acting  $\beta$ 2-adrenoceptor agonists in human lung mast cells. *Br J Pharmacol* **138**: 512–520
- Dhein S (2004) Pharmacology of gap junctions in the cardiovascular system. *Cardiovasc Res* **62**: 287–298
- Dinarello CA (2005) Blocking IL-1 in systemic inflammation. *J Exp Med* **201**: 1355–1399
- Eguchi Y, Shimizu S, Tsujimoto Y (1997) Intracellular ATP levels determine cell death fate by apoptosis or necrosis. *Cancer Res* **57**: 1835–1840
- Elliott JI, Surprenant A, Marelli-Berg FM, Cooper JC, Cassidy-Cain RL, Wooding C, Linton K, Alexander DR, Higgins CF (2005) Membrane phosphatidylserine distribution as a non-apoptotic signalling mechanism in lymphocytes. *Nat Cell Biol* **7**: 808–816
- Faria RX, Defarias FP, Alves LA (2005) Are second messengers crucial for opening the pore associated with P2X<sub>7</sub> receptor? *Am J Physiol* **288**: C260–C270
- Ferrari D, Pizzirani C, Adinolfi E, Lemoli RM, Curti A, Idzko M, Panther E, Di Virgilio F (2006) The P2X<sub>7</sub> receptor: a key player in IL-1 processing and release. *J Immunol* **176**: 3877–3883
- Griffith TM, Chaytor AT, Edwards DH (2004) The obligatory link: role of gap junctional communication in endothelial dependent smooth muscle hyperpolarization. *Pharmacol Res* **49**: 551–564
- Gudipaty L, Munetz J, Verhoef PA, Dubyak GR (2003) Essential role for Ca<sup>2+</sup> in regulation of IL-1 $\beta$  secretion by P2X<sub>7</sub> nucleotide receptor in monocytes, macrophages and HEK-293 cells. *Am J Physiol* **285**: C286–C299
- Han BS, Hong HS, Choi WS, Markelonis GJ, Oh TH, Oh YJ (2003) Caspase-dependent and independent cell death pathways in primary cultures of mesencephalic dopaminergic neurons after neurotoxin treatment. *J Neurosci* **23**: 5069–5078
- Hickman SE, Semrad CE, Silverstein SC (1996) P2Z purinoceptors. *Ciba Found Symp* **198**: 71–83
- Humphreys BD, Dubyak GR (1998) Modulation of P2X<sub>7</sub> nucleotide receptor expression by pro- and anti-inflammatory stimuli in THP-1 monocytes. *J Leukoc Biol* **64**: 265–273
- Jiang LH, Rassendren F, Mackenzie A, Zhang YH, Surprenant A, North RA (2005) *N*-methyl-D-glucamine and propidium dyes utilize different permeation pathways at rat P2X<sub>7</sub> receptors. *Am J Physiol* **289**: C1295–C1302
- Kahlenberg JM, Dubyak GR (2004) Mechanisms of caspase-1 activation by P2X<sub>7</sub> receptor-mediated K<sup>+</sup> release. *Am J Physiol* **286**: C1100–C1108
- Kanneganti TD, Ozoren N, Body-Malapel M, Amer A, Park JH, Franchi L, Whitfield J, Barchet W, Colonna M, Vandenabeele P, Bertin J, Coyle A, Grant EP, Akira S, Nunez G (2006) Bacterial RNA and small antiviral compounds activate caspase-1 through cryopyrin/Nalp3. *Nature* **440**: 233–236
- Ke HZ, Qi H, Weidema AF, Zhang Q, Panupinthu N, Crawford DT, Grasser WA, Paralkar VM, Li M, Audoly LP, Gabel CA, Jee WS, Dixon SJ, Sims SM, Thompson DD (2003) Deletion of the P2X<sub>7</sub> nucleotide receptor reveals its regulatory roles in bone formation and resorption. *Mol Endocrinol* **17**: 1356–1367
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>- $\Delta\Delta$ C(T)</sup> method. *Methods* **25**: 402–408
- Locovei S, Bao L, Dahl G (2006a) Pannexin 1 in erythrocytes: function without a gap. *Proc Natl Acad Sci* **103**: 7655–7659
- Locovei S, Wang J, Dahl G (2006b) Activation of pannexin 1 channels by ATP through P2Y receptors and by cytoplasmic calcium. *FEBS Lett* **580**: 239–244
- Mackenzie A, Wilson HL, Kiss-Toth E, Dower SK, North RA, Surprenant A (2001) Rapid secretion of interleukin 1 $\beta$  by microvesicle shedding. *Immunity* **8**: 825–835
- Mackenzie AB, Chirakkal H, North RA (2003) Kv1.3 potassium channels in human alveolar macrophage. *Am J Physiol* **285**: L862–L868
- Mackenzie AB, Young MT, Adinolfi E, Surprenant A (2005) Pseudoapoptosis induced by brief activation of ATP-gated P2X<sub>7</sub> receptors. *J Biol Chem* **280**: 33968–33976
- Mariathasan S, Newton K, Monack DM, Vucic D, French DM, Lee WP, Roose-Girma M, Erickson S, Dixit VM (2004) Differential activation of the inflammasome by caspase-1 adaptors ASC and Ipaf. *Nature* **430**: 213–218
- Mariathasan S, Weiss DS, Newton K, McBride J, O'Rourke K, Roose-Girma M, Lee WP, Weinrauch Y, Monack DM, Dixit VM (2006) Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature* **440**: 228–232
- Martinon F, Tschopp J (2004) Inflammatory caspases: linking an intracellular innate immune system to autoinflammatory diseases. *Cell* **117**: 561–574
- Martinon F, Petrilli V, Mayor A, Tardivel A, Tschopp J (2006) Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature* **440**: 237–241
- Morelli A, Chiozzi P, Chiesa A, Ferrari D, Sanz JM, Falzoni S, Pinton P, Rizzuto R, Olson MF, Di Virgilio F (2003) Extracellular ATP causes ROCK I-dependent bleb formation in P2X<sub>7</sub>-transfected HEK cells. *Mol Biol Cell* **14**: 2655–2664
- Nemeth K, Bugovics G, Skekely J (1997) Antiapoptotic effect of benzyloxycarbonyl-aspartyl-( $\beta$ -tert-butyl ester)-bromomethylketone (Z-D(OtBu)-BMK), an intermediate of interleukin-1 $\beta$  converting enzyme inhibitors. *Int J Immunopharmacol* **19**: 215–225
- North RA (2002) Molecular physiology of P2X receptors. *Physiol Rev* **82**: 1013–1067
- Panchin YV (2005) Evolution of gap junction proteins—the pannexin alternative. *J Exp Biol* **208**: 1415–1419
- Perregaux D, Gabel CA (1994) Interleukin-1 beta maturation and release in response to ATP and nigericin. Evidence that potassium depletion mediated by these agents is a necessary and common feature of their activity. *J Biol Chem* **269**: 15195–15203
- Perregaux DG, Svensson L, Gabel CA (1996) Tenidap and other anion transport inhibitors disrupt cytolytic T lymphocyte-mediated IL-1 beta post-translational processing. *J Immunol* **157**: 57–64
- Schilling WP, Wasylyna T, Dubyak GR, Humphreys BD, Sinkins WG (1999) Maitotoxin and P2Z/P2X<sub>7</sub> purinergic receptor stimulation activate a common cytolytic pore. *Am J Physiol* **277**: C766–C776
- Solle M, Labasi J, Perregaux DG, Stam E, Petrushova N, Koller BH, Griffiths RJ, Gabel CA (2001) Altered cytokine production in mice lacking P2X<sub>7</sub> receptors. *J Biol Chem* **276**: 125–132
- Spray DC, Rozental R, Srinivas M (2002) Prospects for rational development of pharmacological gap junction channel blockers. *Curr Drug Targets* **3**: 455–464
- Steinberg TH, Di Virgilio F (1991) Cell-mediated cytotoxicity: ATP as an effector and the role of target cells. *Curr Opin Immunol* **3**: 71–75
- Suadicani SO, Brosnan CF, Scemes E (2006) P2X<sub>7</sub> receptors mediate ATP release and amplification of astrocytic intracellular Ca<sup>2+</sup> signaling. *J Neurosci* **26**: 1378–1385
- Sutterwala FS, Ogura Y, Szczepanik M, Lara-Tejero M, Lichtenberger GS, Grant EP, Bertin J, Coyle AJ, Galan JE, Askenase PW, Flavell RA (2006) Critical role for NALP3/CIAS1/cryopyrin in innate and adaptive immunity through its regulation of caspase-1. *Immunity* **24**: 317–327
- Tarasov A, Dusonchet J, Ashcroft F (2004) Metabolic regulation of the pancreatic  $\beta$ -cell ATP-sensitive K<sup>+</sup> channel. *Diabetes* **53**: S113–S122
- Thompson RJ, Zhou N, MacVicar BA (2006) Ischemia opens neuronal gap junction hemichannels. *Science* **312**: 924–927