

Chromosome Aberrations in Plants as a Monitoring System

by William F. Grant*

The potential of higher plants as a first-tier assay system for detecting chemical mutagens is evaluated. The use of plant tissue (primarily root tips and pollen mother cells) for studying the induction of chromosomal aberrations is one of the oldest, simplest, most reliable, and inexpensive methods available. Specific types of abnormalities have been induced by different classes of pesticides. Chromosome clumping, contraction, stickiness, paling, fragmentation, dissolution, chromosome and chromatid bridges, C-mitosis, and endoploidy have been reported in the literature. Examples of cytogenetic studies with pesticides demonstrating the usefulness of higher plants as a monitoring system are reviewed. Pesticides which cause chromosome aberrations in plant cells also produce chromosome aberrations in cultured animal cells. Frequently, the aberrations are identical. For example, studies have shown that compounds which have a C-mitotic effect on plant cells have the same effect on animal cells. It is recommended that plant systems be accepted as a first-tier assay system for the detection of possible genetic damage by environmental chemicals.

Introduction

Chromosome aberrations have been used as a measure of reproductive success in plants for many years and have been correlated with morphological and taxonomical changes, fertility-sterility relationships, mutations, and other characteristics. The first observation of a correlation between reduction in fertility and cytological abnormalities as a result of pesticide treatment dates back to 1931, when Kostoff observed seed set of tobacco plants to be greatly reduced after the plants had been fumigated with nicotine sulfate. In an examination of meiosis, Kostoff (1) found many chromosome irregularities which he considered to be the cause of the partial sterility of the plants. Subsequent studies with many mutagenic chemicals have shown that plant chromosomes exhibit many different types of aberrations some of which are specific for different chemicals or classes of chemicals.

In the present paper, some aspects of the relevance and reliability of chromosome aberrations in plants as a method for the detection of possible genetic damage by environmental agents are discussed. Examples of the effects of treatment with pesticides (that is, herbicides, insecticides, and fun-

gicides) are presented. In general, chromosome aberrations can provide both qualitative and quantitative data on the effects of exposure to a mutagen, and examples are given. In addition, parallels between chromosome aberrations caused by the same pesticides in plant and mammalian systems were to demonstrate that plant systems would serve as an excellent first-tier bioassay system.

Monitoring for Chromosome Abnormalities

One of the principal objectives of using chromosomes as a monitoring system is to determine whether or not a particular chemical is a clastogen (that is, capable of breaking chromosomes). If the chemical is a clastogen, then this would permit exchanges with subsequent cytological or genetic damage. At the same time it has been recognized that turbagens [chemicals which cause mitotic disturbances; a term proposed by Brøgger (2)], while not necessarily affecting DNA directly, may result in chromosome segregation errors, and therefore, should not be considered genetically insignificant.

Cytological aberrations in plants serve as an excellent monitoring system for the detection of environmental chemicals that may pose a genetic hazard. The plant systems which have proven most

* Genetics Laboratory, Macdonald Campus of McGill University, Ste. Anne de Bellevue, Quebec, HOA 1CO, Canada.

useful for this purpose have been reviewed recently by Nilan and Vig (3).

Chromosome aberrations may be detected in both mitotic and meiotic divisions. Structural rearrangements, which are most evident at metaphase and anaphase, are identical in somatic and gametic cells. Analysis of somatic chromosome aberrations may be carried out by using actively dividing root tip, stem apex, or pollen tube cells. Meiotic chromosome studies are usually carried out using pollen mother cells. In contrast, micronuclei are best detected at the quartet stage. Micronuclei, which vary in number and size, generally result from fragments or lagging chromosomes.

Types of Chromosome Abnormalities Induced by Pesticides

Nearly all of the common types of known cytological aberrations have been reported in plants following treatment with pesticides. The most frequently reported types will be discussed in the following sections.

Colchicine Mitosis

Levan (4) described colchicine mitosis as an inactivation of the spindle followed by a random scattering of the chromosomes over the cell. Delayed centromere division may result in the chromosomes assuming the characteristic C-pairs configuration in which sister chromatids, while remaining attached at the centromere, no longer remain adjacent to one another. C-mitotic compounds which interfere with the division of the cell nucleus are also classified as spindle poisons, mitotic poisons or antimitotic compounds.

There are a number of pesticides which are typical C-mitotic agents (Table 1). The carbamates, including barban (5), benomyl (6), carbaryl (7), chlor-

propham (5, 8), prothion (5, 9, 10) and diallate (11, 12); also BHC (13, 14) and the mercurials (15, 16, 17) are extremely active C-mitotic chemicals. The carbamates have been so effective as C-mitotic chemicals that several have been recommended for the artificial induction of polyploidy (18). Polyploidy has been induced as high as 16-ploid with the carbamate propham (10).

That plant systems are sensitive indicators of cytological aberrations is clear from studies by Fernandez-Gomez (19) and Fernandez-Gomez et al. (14) on the C-mitotic effect of the four isomers (α , β , γ , δ) of hexachlorocyclohexane. They found that the β isomer had no C-mitotic effect, the α isomer produced partial C-mitosis, while the γ and δ isomers resulted in complete C-mitosis. C-mitotic behavior in plants is a function of chemical concentration. If the C-mitotic agent is applied in too high a concentration, as with other chemicals (Fig. 1), mitosis may be completely arrested. Dilute solutions will induce partial or incomplete C-mitosis resulting in multipolar spindles, aneuploid nuclei, and micronuclei in addition to cells exhibiting normal mitoses. Thus plants are reliable indicators of C-mitotic behavior; and, as will be mentioned later, plant cells exhibit the same C-mitotic behavior as animal cells.

Table 1. Pesticides with C-mitotic activity.

Pesticides	
Mercury compounds	<i>Allium</i> , <i>Vicia</i> , <i>Crepis</i> , human leukocytes
Carbamates	<i>Allium</i> , <i>Avena</i> , <i>Secale</i> , <i>Hordeum</i> , <i>Zea</i> , <i>Tradescantia</i> , <i>Pisum</i>
Dieldrin	<i>Crepis</i>
Nitralin	<i>Allium</i> , <i>Tradescantia</i>
Hexachlorocyclohexane	33 species in 28 references; <i>Allium</i> , <i>Vicia</i> , <i>Zea</i> , <i>Triticum</i> , <i>Hordeum</i> , <i>Secale</i>




FIGURE 1. Chromosome aberrations in root tips of *Vicia faba* from treatments with three pesticides showing effect of toxicity from concentration and duration of treatment (dichloran and endrin, mean percentage of three concentrations, 100, 200 and 300 ppm; linuron, 200, 400 and 600 ppm). Data from Wu and Grant (55).

Binucleate, Multinucleate, and Polyploid Cells

Binucleate cells arise as a consequence of the inhibition of cell plate formation. These form a distinct sub-population of easily detected cells. Failure of cell plate formation in already binucleate cells may give rise to the multinucleate condition. Mitotic irregularities, such as incompleting anaphases or unequal distribution of the chromosomes to the daughter cells can result in aneuploid or even euploid cells.

Several pesticides are known to induce the binucleate and multinucleate conditions including bromacil (20), carbaryl (7), dinoseb (8), hexachlorocyclohexane (21–23), nitalin (24), and propham (9, 25, 26). Tri- and tetrapolar anaphases have also been reported following treatments with certain of the preceding pesticides.

The induction of euploid cells has been reported after treatment with hexachlorocyclohexane (13, 27–29), and of aneuploid cells from chloranil treatment of root tips of *Vicia faba* (30) and atrazine treatment of *Sorghum* (31).

Endoreduplication, in which chromosome duplication occurs without nuclear division has been reported after 2,4-D treatment (32).

Chromosome Condensation and Contraction

Chromosome condensation or contraction is the shortening and thickening of the chromosomes brought about by changes in chromosome coiling following chemical treatment during mitosis and meiosis. Chromosome contraction has been observed following treatment of *Tradescantia* root tips with mercury compounds (17) and some carbamates (18).

Chromosome Stickiness and Clumping

Klasterska et al. (33) and McGiff et al. (34) suggested that chromosome stickiness arises from improper folding of the chromosome fiber into single chromatids and chromosomes. As a result there is an intermingling of the fibers, and the chromosomes become attached to each other by means of subchromatid bridges.

Chromosome stickiness and clumping have been reported following treatment with a number of pesticides including asulam (35), carbaryl (7, 36, 37), 2,4-D and 2,4,5-T (38–40); demeton (41), isodrin (42), mercurials (17), pentachlorophenol (43), and phosphorin (44).

Chromosome Haziness or Paling

Haziness or paling of chromosomes probably results from a partial despiralization of the chromosome. Haziness or despiralization has been reported after treatment with the carbamates, chlorpropham and propham (45) and nitalin (24).

Interchromatid Connections

Chromatin fibers which join two sister chromatids at metaphase and presumably hold the chromatids together until anaphase have been termed interchromatid connections (46). Such interchromatid connections have been observed after treatment of *Tradescantia* and *Vicia faba* root tip cells with a mercurial fungicide (17).

Chromosome Dissolution

Chromosome dissolution refers to a complete breakdown in chromosome structure resulting in the formation of long, thin chromatin threads which possibly arise from an almost complete despiralization of the chromosome. Chromosome dissolution has been observed in barley cells after seed treatment with monuron. The long chromatin threads form bridges between aggregations of chromosomal material (47).

Chromosome Fragmentation

Chromosome fragmentation results from multiple breaks of the chromosome in which there is a loss of chromosome integrity. Fragmentation can range from partial to total disintegration of the chromosome (the latter is termed chromosome pulverization). Chromosome fragmentation in plant cells has been reported only rarely after treatment with pesticides.

Amer and Ali (43) reported that pentachlorophenol induced fragmentation of both mitotic and meiotic chromosomes of *Vicia faba*. Other pesticides which have been reported to induce fragmentation include ferbam in *Allium cepa* (48), linuron in *Hordeum* (49, 50) and simazine in *Vicia cracca* (51).

Intensely stained interphase micronuclei, termed chromatin bodies, which result from chromosome fragmentation or aberrations in the previous mitotic division, have been observed in root tip cells of *Vicia faba* from treatment with amitrole and *Allium cepa* after 2,4-D treatment (52).

Chromosome Breakage and Exchange

The most common abnormalities recorded in these categories are (a) chromosome and chromatid breaks, (b) acentric fragments, (c) chromatid and subchromatid exchanges, chromatid gaps (achromatic lesions), heterochromatic regions and sister chromatid exchanges at metaphase, (d) chromatid and chromosome bridges and side-arm bridges and fragments at anaphase. A very detailed classification system has been proposed by Savage (53).

Chromosome breaks, fragments, chromatid exchanges, and dicentric chromosomes are generally considered unstable aberrations; deletions, inversions, duplications, and translocations are considered stable aberrations. Chromosome breakage is now generally considered to involve the DNA molecule responsible for the linear continuity of the chromosome. Such aberrations are the result of unfinished repair or misrepair of DNA (54).

The specific type of aberration induced is a function of the time at which the interphase nucleus is exposed to a clastogen. Exposure in the G_1 phase of the mitotic cycle results in damage to the entire chromosome while treatment in the S or G_2 phase results in damage to individual chromatids. Following treatment in the S phase, the typical aberrations encountered are chromatid breaks and chromatid interchanges. Exposure in the G_2 phase gives rise mainly to chromatid breaks and chromatid gaps. However, it should be noted that cells undergoing additional mitoses usually contain aberrations of the chromosomal type.

Many pesticides are clastogens, producing chromosome breaks which may give rise to anaphase bridges and fragments (15, 47, 50, 51, 55, 56). Since pesticides are not a homogeneous class of chemicals, their mode of action may be very different. For example, Ehrenberg (personal communication) has stated that the physiological action of phenoxy acids in higher plants should be considered since such compounds might secondarily lead to disturbances, including heritable changes, and therefore, the mechanism by which chromosomal aberrations are produced with such compounds should be clarified.

Some pesticides have been shown to consistently induce aberrations in specific regions of the chromosome in contrast to the random distribution observed after irradiation. For example, the growth retarding chemical maleic hydrazide induces chromosome breakage largely in heterochromatic regions (57). Similarly, Nicoloff and Gecheff (58) have shown that in barley seeds, following treatment with ethylenimine, the greatest portion of aberrations were located in the centromere regions.

As a result, bridges were not formed and a large number of fragments were observed. On the other hand, chemicals may consistently produce chromosomal aberrations, but at the same time be ineffectual as clastogens. For example, pesticides which interfere with the spindle mechanism and thus induce C-mitosis generally possess only a very mild clastogenic effect.

Sister Chromatid Exchange

Sister chromatid exchange (SCE) involves a symmetrical exchange at one locus between sister chromatids. To my knowledge, the herbicide maleic hydrazide is the only pesticide that has been tested for the induction of SCE and it failed to induce SCE (59). Maleic hydrazide is an anomalous chemical, since it is a potent inducer of chromosome aberrations in plant cells (60) but it has never been reported to cause chromosome damage in mammalian cells.

Sensitivity

For a given class of chromosome aberrations, it has been shown that species vary in their sensitivity to pesticide treatment. For example, *Tradescantia* is less susceptible to chromosome breakage following pesticide treatment than *Vicia faba* (44). Barley is also less sensitive than *Vicia faba* (61). The susceptibility of a species to chromosome breakage has been shown to be related to level of ploidy, life-form and nuclear volume (62).

Use of Plants as a First-Tier Bioassay System

The question has been raised as to the relevance to human populations of data on chemically induced chromosome aberrations in plants (63). Furthermore, among the various test systems which have been recommended by a committee of the Environmental Mutagen Society, Committee 17 (64), no plant testing system has been included. However, a number of studies which have been carried out on pesticides indicate that there is an excellent correlation between chromosome abnormalities found in root-tip systems and those found in mammalian cell systems (Table 2). There is also a good correlation with mutagenic activity. It is true that the type of chromosome aberration induced by a specific chemical may not be the same in plant cells as in animal cells (63), but if a particular chemical will induce chromosome aberrations in one group, generally it will do so in the other as well. Fur-

Table 2. Comparison between chromosome-breaking and mutagenic effects of pesticides in plant and animals materials.

Compound	Chromosome aberrations					
	Plant root tips	Ref.	Mammalian cells in culture	Ref.	Mutagenic effect	Ref.
Apholate	+	(65)	+	(66)	?	
Atrazine	+	(31)	+	(67)	+	(68)
2,4-D	+	(52)	+	(69)	±	(58)
DDT	+	(70)	+	(71)	+	(72)
Dichlorvos	+	(73)	+	(74)	+	(75)
Dieldrin	+	(76)	+	(77)	±	(78)
Ethylene dibromide	-	(79)	-	(79)	+	(80)
Griseofulvin	+	(81)	+	(82)	-	(83)
Hempa	-	(65)	-	(84)	+	(85)
Heptachlor	+	(86)	+	(87)	+	(87)
Maleic hydrazide	+	(60)	-		+	(60)
Mercury compounds	+	(15)	+	(16)	?	
Phosphamidon	+	(50)	+	(88)	?	
2,4,5-T	+	(89)	+	(90)	±	(78)
Tepa	+	(65)	+	(91)	+	(85)

thermore, it has been shown that exactly the same morphologic "C-mitotic" picture occurs in plant and in animal tissue (92). This has been shown to be true for several mercurial compounds (15, 16) and griseofulvin (81), and possibly others. Thus, it is justified to assume that compounds which have a C-mitotic effect in plant tissue will induce the same effect in animal tissue.

Several higher plants provide unique and valuable systems for detecting and analyzing the effects of chemical mutagens (3). Such plants include maize (*Zea mays*), barley (*Hordeum vulgare*), tomato (*Lycopersicon*), mouse-ear cress (*Arabidopsis thaliana*), soybean (*Glycine max*), broad bean (*Vicia faba*), spiderwort (*Tradescantia*), onion (*Allium cepa*), Hawk's beard (*Crepis capillaris*), lily (*Lilium*), pea (*Pisum sativum*), and tobacco (*Nicotiana tabacum*). As a group, these plants offer systems for the analysis of almost all known genic and chromosomal aberrations which have been induced in eukaryotes by chemical or physical mutagens.

Some of the advantages in utilizing plant systems have been reviewed by previous authors (3, 93): (1) the chromosome organization of plants is similar to that of humans; (2) many plants are easy to grow; (3) some have short generation time; (4) the cost, handling, and space requirements are relatively small; (5) the cost and time of training technicians to handle a variety of end points following mutagen treatment is relatively small; (6) mutagenic effects can be studied under a wide range of environmental conditions such as large differences in pH, water content, temperature, and metabolic rates; (7) most of the plant systems have been in use for many years and are reliable systems which have been

adapted for newer techniques such as chromosome banding and sister-chromatid exchange studies.

Perhaps the most serious disadvantage of a plant system for the detection of genetic risks to man is the lack of similarity between vegetative and mammalian metabolism. Nevertheless, the positive correlation which has been noted between aberrations induced by the same chemical in plant root-tip cells and in cultured mammalian cells indicates that a plant root-tip system must be recognized as an appropriate first-tier assay system.

Numerous studies have demonstrated that plant chromosomes are sensitive indicators to environmental pollutants. In this paper pesticides have been used to illustrate the potential of plant systems as monitors of chromosome aberrations. Pesticides are a diverse and extensively used group of chemicals and they are known to induce a wide range of chromosome aberrations. It is evident that plant systems are simple, reliable, and inexpensive. The application of the results obtained from mutagenesis in plants to humans is just as valid as those from the diploid organism *Neurospora*, an accepted test organism. Higher plant systems appear to be excellent indicators of the cytotoxic, cytogenetic, and mutagenic effects of environmental chemicals; and, therefore, it is recommended that plant systems be accepted as a first-tier assay system for the detection of possible genetic damage resulting from the use of environmental chemicals.

Financial assistance from the National Research Council of Canada is gratefully acknowledged. I thank Alina E. Russell for technical assistance.

REFERENCES

- Kostoff, D. Heteroploidy in *Nicotiana tabacum* and *Solanum melongena* caused by fumigation with nicotine sulphate. *Bull. Soc. Bot. Bulgar.* 4: 87 (1931); *Biol. Abstr.* 8: 10 (1934).
- Brøgger, A. Chromosome damage in human mitotic cells after *in vivo* and *in vitro* exposure to mutagens. In: Export Conference on Genetic Damage in Man Caused by Environmental Agents, Oslo, 1977, Academic Press, New York, 1978.
- Nilan, R. A., and Vig, B. K. Plant test systems for detection of chemical mutagens. In: *Chemical Mutagens; Principles and Methods for Their Detection*. Vol. 4, A. Hollaender, Ed., Plenum Press, New York, 1976, 143.
- Levan, A. The effect of colchicine on root mitoses in *Allium*. *Hereditas* 24: 471 (1938).
- Mann, J. D., and Storey, W. B. Rapid action of carbamate herbicides upon plant cell nuclei. *Cytologia* 31: 203 (1966).
- Spasojević, V. Effect of fungicide Benlate on the mitosis of maize. *Arch. Poljop. Nauke* 27(100): 13 (1974).
- Amer, S. M., and Farah, O. R. Cytological effects of pesticides. III. Meiotic effects of *N*-methyl-1-naphthyl carbamate "Sevin." *Cytologia* 33: 337 (1968).
- Sawamura, S. 1965. Cytological studies on the effect of herbicides on plant cells *in vivo* II. Non-hormonic herbicides. *Cytologia* 30: 325 (1965).
- Doxey, D. The effect of isopropyl phenyl carbamate on mitosis in rye (*Secale cereale*) and onion (*Allium cepa*). *Ann. Bot.* 13: 329 (1949).
- Derenne, P. Effets morphologiques, physiologiques et cytologiques dus à l'action de l'isopropylphénylcarbamate sur les genres *Allium*, *Vicia* et *Hordeum*. *Bull. Inst. Agron. Stn. Rech. Gembloux* 21: 37 (1953).
- Morrison, J. W. Cytological effects of the herbicide "Avadex." *Can. J. Plant Sci.* 42: 78 (1962).
- Spasojević, V. Cytogenetical effect of Tuberite on *Zea mays*. *Arhiv. Poljopr. Nanke* 28: 119 (1975).
- Datta, N. Cytological effects of gammexane on somatic chromosomes of *Urginea coromandeliana* Hook. *F. Current Sci.* 35: 75 (1966).
- Fernandez-Gomez, M. E., Gimenez-Martin, G., and Lopez-Saez, J. F. Alteraciones en el ciclo de division celular inducidas por los isomeros del HCCH. II. Isomero delta. Possible mecanismo de accion del HCCH. *Genét. Ibér.* 19: 123 (1967).
- Fiskesjö, G. Some results from *Allium* tests with organic mercury halogenides. *Hereditas* 62: 314 (1969).
- Fiskesjö, G. The effect of two organic mercury compounds on human leukocytes in vitro. *Hereditas* 64: 142 (1970).
- Ahmed, M., and Grant, W. F. Cytological effects of the mercurial fungicide Panogen 15 on *Tradescantia* and *Vicia faba* root tips. *Mutat. Res.* 14: 391 (1972).
- Storey, W. B., Jordan, L. S., and Mann, J. D. Carbamate herbicides—new tools for cytological studies. *Calif. Agric.* 22(8): 12 (1968).
- Fernandez-Gomez, M. E. Alteraciones en el ciclo de division celular inducidas por los isomeros del HCCH. I. Isomeros alfa, beta y gamma. *Genét. Ibér.* 19: 103 (1967).
- Ashton, F. M., Cutter, E. G., and Huffstutter, D. Growth and structural modifications of oats induced by bromacil. *Weed Res.* 9: 198 (1969).
- Gimenez-Martin, G., Gonzalez-Fernandez, A., Lopez-Saez, J. F., and Fernandez-Gomez, E. Polymitosis with unbalanced nuclei. *Phyton* 23: 11 (1966).
- Gonzalez, A. Formacion y desarrollo de celulas binucleadas: Bimiosis. *Genet. Iber.* 19: 1 (1967).
- Baquar, S. R., and Khan, N. R. Effect of γ -hexachlorocyclohexane (HCCH) on the mitotic cells of *Pisum sativum* L. *Rev. Biol.* 7: 195 (1971).
- Gentner, W. A., and Burk, L. G. Gross morphological and cytological effects of nitratin on corn roots. *Weed Sci.* 16: 259 (1968).
- Ennis, W. B., Jr. Some cytological effects of *o*-isopropyl-*N*-phenyl carbamate upon *Avena*. *Am. J. Bot.* 35: 15 (1948).
- Canvin, D. T., and Friesen, G. Cytological effects of CDAA and IPC on germinating barley and peas. *Weeds* 7: 153 (1959).
- Quidet, P., and Hitier, H. 1948. Obtention de plantes polyploides par traitement à l'hexachlorocyclohexane et au sulfure de polychlorocyclane. *C. R. Acad. Sci.* 226: 833 (1948).
- Landa, Z. γ -Hexachlorocyclohexane as a polyploidy-inducing agent. *Biol. Plant.* 1: 151 (1959).
- Sharma, A. K., and Gosh, S. A comparative study of the effects of certain chemical agents on chromosomes. *Acta Biol. Acad. Sci. Hung.* 20: 11 (1969).
- Yakar, N. Mitotic disturbances caused by chloranil. *Am. J. Bot.* 39: 540 (1952).
- Liang, G. H., and Liang, Y. T. S. Effects of atrazine on chromosomal behavior in sorghum. *Can. J. Genet. Cytol.* 14: 423 (1972).
- Dvořák, J. Endopolyploidy in the roots of rye, *Secale cereale* L. *Biol. Plant.* 10: 112 (1968).
- Klásterská, I., Natarajan, A. T., and Ramel, C. An interpretation of the origin of subchromatid aberrations and chromosome stickiness as a category of chromatid aberrations. *Hereditas* 83: 153 (1976).
- McGill, M., Pathak, S., and Hsu, T. C. Effects of ethidium bromide on mitosis and chromosomes: A possible material basis for chromosome stickiness. *Chromosoma* 47: 157 (1974).
- Sterrett, R. B., and Fretz, T. A. Asulam-induced mitotic irregularities in onion root-tips. *HortScience* 10: 161 (1975).
- Amer, S. M. Cytological effects of *N*-methyl-1-naphthyl carbamate. *Naturwiss.* 51: 494 (1964).
- Amer, S. Cytological effects of pesticides. I. Mitotic effects of *N*-methyl-1-naphthylcarbamate "Sevin." *Cytologia* 30: 175 (1965).
- Crokér, B. H. Effects of 2,4-dichlorophenoxyacetic acid and 2,4,5-trichlorophenoxyacetic acid on mitosis in *Allium cepa*. *Bota. Gaz.* 114: 274 (1953).
- Nygren, A. Cytological studies of the effects of 2,4-D, MCPA, and 2,4,5-T on *Allium cepa*. *Ann. Roy. Agric. Coll. Sweden* 16: 723 (1949).
- Amer, S. M., and Ali, E. M. Cytological effects of pesticides. V. Effects of some herbicides on *Vicia faba*. *Cytologia* 39: 633 (1974).
- Gibson, P. B., and Beinhart, G. Abnormal meiosis in clover plants treated with organic phosphate pesticides. *Bull. S. C. Acad. Sci.* 31: 38 (1969).
- Scholes, M. E. The effects of aldrin, dieldrin, isodrin, endrin and DDT on mitosis in roots of the onion (*Allium cepa* L.). *J. Hortic. Sci.* 30: 181 (1955).
- Amer, S. M., and Ali, E. M. Cytological effects of pesticides. IV. Mitotic effects of some phenols. *Cytologia* 34: 1 (1969).
- Ahmed, M., and Grant, W. F. Cytological effects of the pesticides phosdrin and bladex on *Tradescantia* and *Vicia faba*. *Can. J. Genet. Cytol.* 14: 157 (1972).
- Herichová, A. Study of the effect of isopropyl-*N*-phenylcarbamate and isopropyl-*N*-(3 chlorophenyl) carbamate on chromosome structure and cytokinesis. *Acta F.R.N. Univ. Comen.-Physiol. Plant.* 1: 147 (1970).
- DuPraw, E. J. *DNA and Chromosomes*. Holt, Rinehart and Winston, New York, 1970.
- Wuu, K. D., and Grant, W. F. Chromosomal aberrations induced by pesticides in meiotic cells of barley. *Cytologia* 32: 31 (1967).

48. Prasad, I., and Pramer, D. Genetics effects of ferbam on *Aspergillus niger* and *Allium cepa*. *Phytopathology* 58: 1188 (1968).
49. Wu, K. D., and Grant, W. F. Induced abnormal meiotic behavior in a barley plant (*Hordeum vulgare* L.) with the herbicide Lorox. *Phyton* 23: 63 (1966).
50. Wu, K. D., and Grant, W. F. Morphological and somatic chromosomal aberrations induced by pesticides in barley (*Hordeum vulgare*). *Can. J. Genet. Cytol.* 8: 481 (1966).
51. Tomkins, D. J., and Grant, W. F. Monitoring natural vegetation for herbicide-induced chromosomal aberrations. *Mutat. Res.* 36: 73 (1976).
52. Mohandas, T., and Grant, W. F. Cytogenetic effects of 2,4-D and amitrole in relation to nuclear volume and DNA content in some higher plants. *Can. J. Genet. Cytol.* 14: 773 (1972).
53. Savage, J. R. K. Classification and relationships of induced chromosomal structural changes. *J. Med. Genet.* 12: 103 (1975).
54. Evans, H. J. 1977. Molecular mechanisms in the induction of chromosome aberrations. In: *Progress in Genetic Toxicology*. D. Scott, B. A. Bridges, and F. H. Sobels, Eds., Elsevier/North-Holland Biomedical Press, Amsterdam, 1977, p. 57.
55. Wu, K. D., and Grant, W. F. Chromosomal aberrations induced in somatic cells of *Vicia faba* by pesticides. *Nucleus* 10: 37 (1967).
56. Georgian, L. The comparative cytogenetic effects of aldrin and phosphamidon. *Mutat. Res.* 31: 103 (1975).
57. Price, M., and Schank, S. C. 1973. Chromosomal damage and abnormal seedling development in barley induced by chemical treatment with TIBA, maleic hydrazide and foramide. *Proc. Soil. Crop Sci. Soc. Fla.* 32: 41 (1973).
58. Nicoloff, H., and Gecheff, K. Methods of scoring induced chromosome structural changes in barley. *Mutat. Res.* 34: 233 (1976).
59. Perry, P., and Evans, H. J. Cytological detection of mutagen-carcinogen exposure by sister chromatid exchange. *Nature* 258: 121 (1975).
60. Grant, W. F., and Harney, P. M. Cytogenetic effects of maleic hydrazide treatment of tomato seed. *Can. J. Genet. Cytol.* 2: 162 (1960).
61. Grant, W. F. Cytogenetic factors associated with the evolution of weeds. *Taxon* 16: 283 (1967).
62. Mohandas, T., and Grant, W. F. A relationship between nuclear volume and response to auxin herbicides for some weed species. *Can. J. Bot.* 51: 1133 (1973).
63. Kihlman, B. A. 1971. Root tips for studying the effects of chemicals on chromosomes. In: *Chemical Mutagens: Principles and Methods of Their Detection*. Vol. 2, A. Hollaender, Ed., Plenum Press, New York, 1971, p. 489.
64. Committee 17, Council of the Environmental Mutagen Society. Environmental mutagenic hazards. *Science* 187: 503 (1975).
65. Ninan, T., and Wilson, G. B. Chromosome breakage by ethylenimines and related compounds. *Genetica* 40: 103 (1969).
66. Klassen, W., Chang, T. H., and Eide, P. E. Effect of apholate on chromosomes of germ cells in the grasshopper testes. *Can. J. Genet. Cytol.* 11: 829 (1969).
67. Yoder, J., Watson, M., and Benson, W. W. Lymphocyte chromosome analysis of agricultural workers during extensive occupational exposure to pesticides. *Mutat. Res.* 21: 335 (1973).
68. Plewa, M. J., and Gentile, J. M. Mutagenicity of atrazine: a maize-microbe bioassay. *Mutat. Res.* 38: 287 (1976).
69. Styles, J. A. Cytotoxic effects of various pesticides *in vivo* and *in vitro*. *Mutat. Res.* 21: 50 (1973).
70. Vaarama, A. Experimental studies on the influence of DDT insecticide upon plant mitosis. *Hereditas* 33: 191 (1947).
71. Kelly-Garvert, F., and Legator, M. S. Cytogenetic and mutagenic effects of DDT and DDE in a Chinese hamster cell line. *Mutat. Res.* 17: 223 (1973).
72. Vogel, E. The relation between mutational pattern and concentration by chemical mutagens in *Drosophila*. In: *Screening Tests in Chemical Carcinogenesis*. R. Montesano, H. Bartsch and L. Tomatis, Eds., IARC Sci. Publ., Lyon; Vol. 12, 1976, p. 117.
73. Sax, K., and Sax, H. J. Possible mutagenic hazards of some food additives, beverages and insecticides. *Japan. J. Genet.* 43: 89 (1968).
74. Bridges, B. A., et al. Mutagenicity of dichlorvos and methyl methanesulphonate for *Escherichia coli* WP2 and some derivatives deficient in DNA repair. *Mutat. Res.* 19: 295 (1973).
75. Hanna, P. J., and Dyer, K. F. Mutagenicity of organophosphorus compounds in bacteria and *Drosophila*. *Mutat. Res.* 28: 405 (1975).
76. Markaryan, D. S. Effect of dieldrin on the mitosis in *Crepis capillaris* sprout. *Genetika* 3: 55 (1967).
77. Markaryan, D. S. Cytogenetic effect of some chlororganic insecticides on the nuclei of mouse bone-marrow cells. *Genetika* 1: 132 (1966).
78. Grant, W. F. Cytogenetic effects of chlorinated hydrocarbon pesticides. In: *A Rational Evaluation of Pesticidal vs. Mutagenic/Carcinogenic Action*. R. W. Hart, H. F. Kraybill and F. J. De Serres, eds., DHEW-PHS-NIH-NCI Publ., U. S. Govt. Printing Office, Washington, D. C. 1978.
79. Kristofferson, U. Genetic effects of some gasoline additives. *Hereditas* 78: 319 (1974).
80. Nauman, C. H., Sparrow, A. H., and Schairer, L. A. Comparative effects of ionizing radiation and two gaseous chemical mutagens on somatic mutation induction in one mutable and two non-mutable clones of *Tradescantia*. *Mutat. Res.* 38: 53 (1976).
81. Paget, G. E., and Walpole, A. L. 1958. Some cytological effects of griseofulvin. *Nature* 182: 1320 (1958).
82. De Carli, L., Larizza, L., Simoni, G., Di Lernia, R., and Tedrici, F. Effect of griseofulvin on chromosomal complements of human diploid and heteroploid cell cultures. *Mutat. Res.* 21: 27 (1973).
83. Epstein, S. S., and Shafner, H. Chemical mutagens in the human environment. *Nature* 219: 385 (1968).
84. Chang, T.-H., and Klassen, W. Comparative effects of tremamine, TEPA, apholate and their structural analogs on human chromosomes *in vitro*. *Chromosoma* 24: 314 (1968).
85. Palmquist, J., and LaChance, L. E. Comparative mutagenicity of two chemosterilants, Tera and Hempa, in sperm of *Bracon hebetor*. *Science* 154: 915 (1966).
86. Scholes, M. E. The effects of chlordane and toxaphene on mitosis in roots of the onion (*Allium cepa* L.). *J. Hort. Sci.* 30: 12 (1955).
87. Cerey, K., Izakovic, V., and Ruttkay-Nedecka, J. Effect of heptachlor on dominant lethality and bone marrow in rats. *Mutat. Res.* 21: 26 (1973).
88. Georgian, L. The comparative cytogenetic effects of aldrin and phosphamidon. *Mutat. Res.* 31: 103 (1974).
89. Sawamura, S. Cytological studies on the effect of herbicides on plant cells *in vivo* I. Hormonic herbicides. *Cytologia* 29: 86 (1964).
90. Majumdar, S. K., and Hall, R. C. Cytogenetic effects of 2,4,5-T on *in vivo* bone marrow cells of Mongolian gerbils. *J. Hered.* 64: 213 (1973).
91. Adler, I. D., Ramarao, G., and Epstein, S. S. *In vivo* cytogenetic effects of trimethylphosphate and of Tera on bone marrow cells of male rats. *Mutat. Res.* 13: 263 (1971).
92. Hall, T. C. Chemotherapy of cancer. *N. Engl. J. Med.* 266: 129 (1962).
93. Kihlman, B. A. Root tips of *Vicia faba* for the study of the induction of chromosomal aberrations. *Mutat. Res.* 31: 401 (1975).