

- pulse ultrasound: its reproducibility, validation and variability. *Br J Dermatol* 1982;106:657-67.
- 12 Edwards C. The use of high frequency ultrasound for the study of structure and dimensions of skin. Manchester: University of Manchester Faculty of Technology, 1985. (PhD thesis.)
 - 13 Boyle J, Briggs JD, Mackie RM, Junor BJR, Aitchison TC. Cancer, warts and sunshine in renal transplant patients. *Lancet* 1984;ii:702-5.
 - 14 Shuttleworth D, Marks R, Griffin PJA, Salaman JR. Dysplastic epidermal changes in immunosuppressed patients with renal transplants. *Q J Med* 1987;64:609-16.
 - 15 Miller RG. *Simultaneous statistical inferences*. 2nd ed. Berlin: Springer Verlag, 1981:165-7.
 - 16 Gardner MJ, Altman DG. *Statistics with confidence—confidence intervals and statistical guidelines*. London: British Medical Journal, 1989:72.
 - 17 Reid DM, Nicholl JJ, Smith MA, Higgins B, Tothill P, Nuki G. Corticosteroids and bone mass in asthma: comparisons with rheumatoid arthritis and polymyalgia rheumatica. *Br Med J* 1986;293:1463-6.
 - 18 Crompton GK. Corticosteroids and bone mass in asthma. *Br Med J* 1987;294:123.
 - 19 Tan CY, Marks R, Payne P. Comparison of xeroderma radiographic and ultrasound detection of corticosteroid induced dermal thinning. *J Invest Dermatol* 1981;76:126-8.
 - 20 Dykes PJ, Marks R. An appraisal of the methods used in the assessment of atrophy from topical corticosteroids. *Br J Dermatol* 1979;101:599-609.
 - 21 Rook A, Wilkinson DS, Ebling FJE, Champion RH, Burton JL, eds. Glucocorticosteroid induced atrophy. In: *Textbook of dermatology*. London: Blackwell Scientific, 1986:1814.
 - 22 Dykes PJ, Hills S, Marks R. Assessment of the atrophogenicity potential of corticosteroids by ultrasound and by epidermal biopsy under occlusive and non-occlusive conditions. In: Christopher AL, ed. *Topical corticosteroid therapy: a new approach to safer drug therapy*. New York: Raven Press, 1988:111-21.
 - 23 David MJ. Skin trauma in patients receiving systemic corticosteroid therapy. *Br Med J* 1972;ii:614-6.
 - 24 Gottlieb NL, Penneys NS. Spontaneous skin tearing during systemic corticosteroid treatment. *JAMA* 1980;243:1260-1.
 - 25 McLeod DT, Capewell S, Law J, McLaren W, Seaton A. Intramuscular triamcinolone acetonide in chronic severe asthma. *Thorax* 1985;40:840-5.
 - 26 Gaddie J, Reid IW, Skinner C, Petrie GR, Sinclair DJM, Palmer KNV. Aerosol beclomethasone dipropionate—a dose response study in chronic bronchial asthma. *Lancet* 1973;ii:280-1.
 - 27 Kruszynska YT, Greenstone M, Home PD, Cooke NJ. Effect of high dose inhaled beclomethasone dipropionate on carbohydrate and lipid metabolism in normal subjects. *Thorax* 1987;42:881-5.
 - 28 Ali NJ, Capewell S, Ward M. Bone turnover during high dose inhaled corticosteroid therapy. *Thorax* 1989;44:900.
 - 29 Littlewood JM, Johnson AW, Edwards PA, Littlewood AE. Growth retardation in asthmatic children treated with inhaled beclomethasone dipropionate. *Lancet* 1988;ii:115-6.
 - 30 Karim AKA, Thompson GM, Jacob TJC. Steroid aerosols and cataract formation. *Br Med J* 1989;299:918.
 - 31 Brincat M, Yuen AWT, Studd JW, et al. Response of skin thickness and metacarpal index to estradiol therapy in post-menopausal women. *Obstet Gynecol* 1987;70:538-41.

(Accepted 3 April 1990)

A randomised controlled trial of surgery for glue ear

N A Black, C F B Sanderson, A P Freeland, M P Vessey

Abstract

Objective—To assess the effect of five different surgical treatments for glue ear (secretory otitis media) on improvement in hearing and, assuming one or more treatments to be effective, to identify the appropriate indications for surgery.

Design—Randomised controlled trial of children receiving (a) adenoidectomy, bilateral myringotomy, and insertion of a unilateral grommet; (b) adenoidectomy, unilateral myringotomy, and insertion of a unilateral grommet; (c) bilateral myringotomy and insertion of a unilateral grommet; and (d) unilateral myringotomy and insertion of a grommet. Children were followed up at seven weeks, six months, 12 months, and 24 months by symptom history and clinical investigations.

Setting—Otolaryngology department in an urban hospital.

Patients—149 Children aged 4-9 years who were admitted for surgery for glue ear and who had no history of previous operations on tonsils, adenoids, or ears and no evidence of sensorineural deafness. Inadequate follow up information on levels of hearing and on middle ear function was obtained from 22.

Main outcome measures—Mean hearing loss (dB) of the three worst heard frequencies between 250 and 4000 Hz, results of impedance tympanometry, and parental views on their child's progress.

Results—In the 127 children for whom adequate information was available ears in which a grommet had been inserted performed better in the short term (for at least six months) than those in which no grommet had been inserted, irrespective of any accompanying procedure. Most of the benefit had disappeared by 12 months. Adenoidectomy produced a slight improvement that was not significant, though was sustained for at least two years. The ears of children who had had an adenoidectomy with myringotomy and grommet insertion, however, continued to improve so that two years after surgery about 50% had abnormal tympanometry compared with 83% of those who had had only myringotomy and grommet insertion, and 93% of the group that had had no treatment. Logistic regression analyses identified preoperative hearing

level as the single best predictor of good outcome from surgery. Other variables contributed little additional predictive power.

Conclusions—If the principal objective of surgery for glue ear is to restore hearing then our study shows that insertion of grommets is the treatment of choice. The addition of an adenoidectomy will increase the likelihood of restoration of normal function of the middle ear but will not improve hearing. When deciding appropriate indications for surgery, a balance has to be made between performing unnecessary operations and failing to treat patients who might benefit from surgical intervention. Preoperative audiometry scores might be the best predictor in helping to make this decision.

Introduction

Glue ear, or otitis media with effusion, is the commonest reason for elective surgery in childhood.¹ In England and Wales in 1986 about 73 000 operations were carried out in NHS hospitals (based on hospital activity analyses for Oxford and for East Anglian regional health authorities) and a further 18 000 are estimated to have been performed in independent hospitals (J P Nicholl, personal communication). Despite the popularity of these operations considerable uncertainty exists about their efficacy and the appropriate indications for their use. Although the results of 15 randomised controlled trials concerning a total of 1549 children have been published since 1967, few of the studies can easily be compared.²⁻¹⁶ Variations of case definition, exclusion criteria, case severity, outcome measures, duration of follow up, and method of analysis have all contributed to the difficulty in achieving consensus. A further complication is that a variety of operative procedures in different combinations have been studied: adenoidectomy, myringotomy, and grommet (tympanostomy tube) insertion (table I).

Despite the difficulties entailed in making detailed comparisons between the trials it is possible to identify some consistent findings. Firstly, myringotomy results in little or no benefit.^{14,16} Secondly, myringotomy plus grommet insertion is effective for up to 12 months,^{2,3,14} though two studies found that this procedure was not

Health Services Research Unit, Department of Public Health and Policy, London School of Hygiene and Tropical Medicine, London WC1E 7HT

N A Black, MD, senior lecturer in public health medicine
C F B Sanderson, PHD, senior lecturer in health services research

Department of Otolaryngology, Oxfordshire Health Authority
A P Freeland, FRCS, consultant otolaryngologist

Department of Public Health and Primary Care, University of Oxford
M P Vessey, MD, professor of social and community medicine

Correspondence to: Dr Black.

Br Med J 1990;300:1551-6

TABLE I—Comparisons considered in published randomised controlled trials of surgery for glue ear 1967-89

Treatment 2	Treatment 1				
	No treatment	Myringotomy	Myringotomy and grommet	Adenoidectomy	Adenoidectomy and myringotomy
Adenoidectomy and grommet	Maw and Herod ²	Gates <i>et al</i> ¹	Roydhouse ⁴ Widemar <i>et al</i> ⁶ Gates <i>et al</i> ¹	Lildholdt ⁸ Maw and Herod ²	Richards <i>et al</i> ⁷ Bonding <i>et al</i> ⁵ Gates <i>et al</i> ¹
Adenoidectomy and myringotomy		Fiellau-Nikolajsen <i>et al</i> ¹⁰ Gates <i>et al</i> ¹	Gates <i>et al</i> ¹		
Adenoidectomy	Rynnel-Dagoo <i>et al</i> ¹¹ Bulman <i>et al</i> ¹² Maw and Herod ²		Maw and Herod ²		
Myringotomy and grommet	Brown <i>et al</i> ¹³ Maw and Herod ² Mandel <i>et al</i> ¹⁴ Zielhuis <i>et al</i> ¹⁵	Mandel <i>et al</i> ¹⁴ Gates <i>et al</i> ¹			
Myringotomy	Archard ¹⁶ Mandel <i>et al</i> ¹⁴				




FIG 1—Treatment groups resulting from randomisation

effective.^{13 15} Thirdly, adenoidectomy is effective,^{2 3 12} though again two studies found that it had no effect.^{10 11} Fourthly, grommet insertion and adenoidectomy are equally effective,^{2 3} though repeat surgery is needed more often after grommet insertion than after adenoidectomy. Finally, adenoidectomy combined with grommet insertion is no better than adenoidectomy alone^{2 3 6 8 9} or grommet insertion alone.^{2 5 17}

We had two objectives: to compare the relative effectiveness of the five different treatment strategies identified in table I and, assuming one or more treatment strategies to be effective, to identify the appropriate indications for surgery in the management of glue ear.

Methods

The parents of all children aged 4-9 years who were admitted to the Radcliffe Infirmary, Oxford, for surgery for bilateral glue ear between 1981 and 1986 were invited to allow their child to take part in the trial. Children who had previously had operations on their tonsils, their adenoids, or their ears and those in whom there was evidence of cleft palate or any sensorineural deafness were excluded. Children were also excluded if surgery for conditions other than glue ear was to be performed, such as adenoidectomy for alleviating gross nasal obstruction. The need for surgery was based on the clinical judgment of the otolaryngologist responsible for the care of each child, regardless of any findings on investigation.

Having obtained parental consent for inclusion in the trial, we randomly divided the children into one of four treatment groups: (a) adenoidectomy and bilateral myringotomy plus insertion of a unilateral grommet (standard Shepherd tympanostomy tube); (b) adenoidectomy plus a unilateral myringotomy and insertion of a grommet; (c) bilateral myringotomy plus insertion of a unilateral grommet; and (d) a unilateral myringotomy and insertion of a grommet (fig 1). Randomisation between the right ear and the left ear for grommet insertion was also carried out. Instructions about the treatment allocated were contained in sealed numbered envelopes. The contents

of the envelopes were determined with a table of random numbers. The clinicians who had obtained parental consent selected the next available envelope according to numerical sequence.

The minimum number of children that would be needed in the study to allow paired analysis and unpaired analysis to be performed was calculated based on the following assumptions. Firstly, we assumed that there would be a mean preoperative variation in hearing loss between a child's ears of 2 (SD 14.25) dB,¹³ and, secondly, that there would be a mean preoperative hearing loss of 32.5 (SD 11.4) dB.¹⁸ Finally, we thought that 10 dB should be the minimum difference in levels of hearing between treatments that might be regarded as clinically important and that the trial should have a 95% chance of detecting such a difference between two treatments at the 5% level of significance. These assumptions implied that about 104 children would be needed for paired analysis—that is, studying the difference in levels of hearing between the two ears in each child—and that about 136 would be needed for unpaired analysis—that is, comparison between treatment groups. We envisaged that about 10% of the children would be lost to follow up before the end of the study, so 149 children were entered into the study.

Information about the child's age, sex, social class (based on the father's occupation, or the mother's when the child was living in a single parent family), and history of symptoms (deafness, otalgia, nasal obstruction, and speech development) was recorded on a preoperative form that was completed by a doctor. In addition, pure tone audiometry (from 250 to 4000 Hz) and impedance tympanometry were carried out. When a myringotomy was performed a record was made of whether the middle ear was dry, contained serous fluid, or contained "glue."

Each child was followed up for two years and was reviewed at seven weeks, six months, 12 months, and 24 months. At each visit the following information was obtained: parental views on their child's progress, results of a pure tone audiogram, and results of a tympanogram. The children were not assessed by otoscopy because of the considerable interobserver variation associated with the observations. The audiometricians were blind to the treatment that the children had received. Children who did not attend their follow up appointment were sent another invitation. Attempts to get them to attend were abandoned only when they did not appear at three consecutive appointments.

Parental opinions on their child's treatment were defined as favourable, uncertain, or unfavourable. Parents were also asked to report any adverse side effects of treatment. In line with other trials audiometric performance was based on the mean hearing loss of the three worst heard frequencies.^{5 6 16} Tympanometry results were classified according to both the shape of

the recording and the pressure in the middle ear, and two categories were established: normal (A and C1) or abnormal (B and C2).¹⁹ Tympanometry was not performed on ears with grommets because valid recordings cannot be made when grommets are in place and are patent.

The organisers of the study recognised that after surgery the clinical management of each child remained the responsibility of the otolaryngologist concerned, and therefore any decision to carry out further or repeat surgery was beyond their control. The otolaryngologists were, however, asked to avoid further surgical treatment when possible. Data on repeat surgery were collected and analysed, but the children concerned were no longer followed up.

The statistical analyses consisted of: (a) a comparison of the findings before operation and after operation in the four treatment groups using contingency tables; (b) the proportions of children in each group who had to have repeat surgery, and the findings at reoperation; (c) paired analysis of the audiometric findings for the left ear and the right ear in the same child using *t* tests on the mean changes in hearing level since surgery; (d) independent comparisons of audiometric findings for the ears of different children after different surgical interventions using *t* tests on the mean changes in hearing level since surgery; (e) comparison of the proportions of children who had abnormal results on tympanometry and unfavourable parental opinion at follow up; (f) multivariate analysis to link the outcome

of grommet insertion to a set of preoperative variables using a range of outcome criteria.

Results

COMPARABILITY OF TREATMENT GROUPS

The children in the four treatment groups were comparable with regard to the stratification criteria of age, social class, and history of glue ear (table II) and findings on investigation (table III). The sex ratios differed, but there is no evidence to suggest that this would cause problems with confounding.

FOLLOW UP

Overall 48 (32%) children underwent further surgery for glue ear during the two year follow up period, the proportion varying with the initial treatment group. Children who had undergone an adenoidectomy were less likely to have further surgery (19% *v* 45%, $p < 0.01$), but this was not surprising as it is usually possible to undergo an adenoidectomy only once. A further 10 (7%) children either did not attend follow up appointments or moved from the area. Most of the loss to follow up occurred more than 12 months after the initial operation: 85% were seen at 12 months but only 61% at 24 months.

OUTCOMES

Audiometry (paired analysis)—Audiometric data were not obtained on every occasion in 22 children

TABLE II—Preoperative characteristics of the children according to treatment group. Values are numbers (percentages)

Characteristic	Treatment group			
	Adenoidectomy and bilateral myringotomy plus unilateral grommet (1) (n=37)	Adenoidectomy plus unilateral myringotomy and grommet (2) (n=38)	Bilateral myringotomy plus unilateral grommet (3) (n=37)	Unilateral myringotomy and grommet (4) (n=37)
Social class:				
Non-manual	13 (35)	16 (42)	12 (32)	13 (36)
Manual	18 (49)	18 (47)	20 (55)	21 (56)
Other	6 (16)	4 (11)	5 (13)	3 (8)
Pattern of deafness:				
Never	1 (3)	0	2 (5)	2 (5)
Fluctuating	23 (62)	21 (55)	24 (66)	18 (49)
Constant	13 (35)	17 (45)	11 (29)	17 (46)
Duration of deafness (months):				
≤9	4 (11)	7 (19)	6 (16)	11 (30)
10-18	14 (37)	13 (34)	13 (35)	9 (24)
>18	19 (51)	18 (47)	18 (49)	17 (46)
No of episodes of otalgia:				
None	11 (30)	15 (40)	12 (32)	12 (33)
1-3	20 (54)	12 (32)	16 (42)	14 (39)
≥4	6 (16)	11 (29)	9 (26)	11 (28)
Duration of otalgia (months):				
<6	5 (14)	2 (5)	4 (11)	5 (14)
6-12	8 (21)	6 (17)	9 (23)	11 (30)
>12	24 (64)	30 (79)	24 (65)	21 (57)
Nasal symptoms:				
None or mild	11 (30)	13 (35)	18 (48)	16 (44)
Moderate or severe	26 (70)	25 (65)	19 (52)	21 (56)
Speech development:				
Normal	30 (81)	33 (87)	29 (79)	33 (90)
Abnormal	7 (19)	5 (13)	8 (21)	4 (10)
Mean (SE) age (years)	6.3 (0.23)	6.6 (0.23)	6.1 (0.21)	6.0 (0.21)
Sex (male:female)	1.06	1.11	1.92	1.79

TABLE III—Preoperative investigations and operative findings according to treatment group. Hearing level is the mean of the three worst heard frequencies

Investigation or finding	Treatment group			
	Adenoidectomy and bilateral myringotomy plus unilateral grommet (1) (n=37)	Adenoidectomy plus unilateral myringotomy and grommet (2) (n=38)	Bilateral myringotomy plus unilateral grommet (3) (n=37)	Unilateral myringotomy and grommet (4) (n=37)
Hearing level (dB):				
Left ear	28.1	26.9	27.6	27.8
Right ear	29.6	29.1	29.2	27.2
Impedance (No (%) abnormal):				
Left ear	35 (95)	30 (79)	31 (84)	27 (73)
Right ear	35 (95)	36 (95)	29 (78)	29 (78)
Middle ear contents (No (%)):				
Dry	11 (30)	15 (39)	9 (24)	16 (44)
Serous	7 (19)	3 (8)	4 (10)	2 (5)
Glue	19 (51)	20 (53)	24 (66)	19 (51)

and therefore these children were omitted from the analysis. The effect of grommet insertion was assessed in each child by comparing the change in the hearing level (mean of the levels of the three worst heard frequencies) between the ears with and without a grommet. The data were initially analysed without taking into account any loss to follow up. The results are shown in table IV (raw data). Overall, ears in which a grommet had been inserted performed better in the short term (up to 12 months after surgery) than those in which no grommet had been inserted, irrespective of any accompanying procedures. Losses to follow up occurred for two reasons: repeat surgery and non-attendance at the outpatient clinic. The mean level of hearing of those needing further surgery had deteriorated by about 2 dB during the 12 months since their initial operation compared with an improvement of about 8 dB in those not requiring further intervention. It was possible that those children who had not attended their outpatient appointments had experienced a favourable outcome from surgery. To allow for these potential biases at the 12 and 24 month reviews we modified the raw data by assuming that without repeat surgery the levels of hearing would not have altered from the last recorded level. To test this assumption the analysis was repeated twice, allowing first for a deterioration of 10% in the hearing levels since the last recorded level and then for an improvement of 10%. These variant assumptions made little difference to the results.

Audiometry (independent comparisons)—Independent (rather than within child) comparisons of changes in mean audiometry scores for the different surgical interventions are shown in table V. It was apparent that myringotomy had no discernible effect compared with no treatment. In contrast, levels of hearing improved with both myringotomy plus grommet insertion and adenoidectomy. The outcome after the combined operation (adenoidectomy plus myringotomy and grommet insertion) confirmed these findings. There was little difference initially between the outcome of the combined operation and that obtained with myringotomy and grommet insertion alone (fig 2). Sensitivity analysis with different modifications to the raw data (again allowing for a deterioration of 10% in the levels of hearing since the last recorded level and an improvement of 10%) made little impact on the results, and comparisons of absolute values of the levels of




FIG 2—Independent comparisons of changes in mean audiometry scores for several different treatments using modified data

hearing on follow up, rather than changes from the preoperative levels, produced similar findings.

Tympanometry—In addition to the difficulties caused by the fairly high drop out rate during the second year of follow up the results of tympanometry were also affected by the lack of data during the first year of follow up for those ears in which a grommet had been inserted (because tympanometry could not be performed as a satisfactory seal cannot be achieved after grommet insertion). Because myringotomy had no effect on the levels of hearing the findings on tympanometry were analysed according to four groups (fig 3). Because of the preoperative differences in the proportions of abnormal readings changes in proportions were used in the analysis. During the second year the ears of children who had had an adenoidectomy continued to improve so that two years after surgery about half of them had abnormal tympanograms compared with 83% of those who had had a myringotomy plus grommet insertion, and 93% of those who had had either a myringotomy or no treatment.

TABLE IV—Within child comparison of change in mean results of audiometry (dB) with time after surgery according to treatment group: raw and modified data. Values are numbers (95% confidence intervals)

Treatment group comparisons	Time after operation (raw data)				Time after operation (modified data)	
	7 Weeks	6 Months	12 Months	24 Months	12 Months	24 Months
Adenoidectomy, myringotomy and grommet v adenoidectomy and myringotomy	8.1* (3.0 to 13.3)	2.8 (-1.9 to 7.4)	-1.0 (-6.1 to 4.0)	0.7 (-4.9 to 6.4)	-0.9 (-5.5 to 3.8)	0.2 (-4.9 to 5.3)
Adenoidectomy v adenoidectomy, myringotomy and grommet	3.3 (-0.5 to 7.1)	2.8 (-2.2 to 7.8)	1.9 (-3.6 to 7.4)	2.2 (-6.0 to 10.3)	2.3 (-2.8 to 7.4)	2.1 (-3.8 to 8.1)
Myringotomy and grommet v myringotomy	12.7* (7.9 to 17.5)	7.4* (1.4 to 13.4)	3.7 (-0.4 to 7.8)	0.9 (-2.7 to 4.6)	5.5* (0.9 to 10.1)	3.4 (-1.1 to 8.0)
Myringotomy and grommet v no treatment	3.4 (-0.9 to 7.6)	3.5* (0.1 to 6.9)	1.0 (-2.1 to 4.2)	-2.4 (-8.7 to 3.9)	2.0 (-1.0 to 5.1)	0.5 (-3.7 to 4.6)

*Significant *t* value ($p < 0.05$).

TABLE V—Independent comparisons of changes in mean audiometry scores (dB) with time after surgery: raw and modified data. Values are numbers (95% confidence intervals)

Treatment group comparisons	Time after operation (raw data)				Time after operation (modified data)	
	7 Weeks	6 Months	12 Months	24 Months	12 Months	24 Months
Myringotomy v no surgery	1.0 (-4.7 to 6.6)	-0.6 (-7.0 to 5.9)	-1.1 (-8.1 to 5.8)	-2.3 (-9.1 to 4.5)	1.2 (-5.3 to 7.8)	0.7 (-5.5 to 7.0)
Myringotomy and grommet v no surgery	11.7* (5.8 to 17.6)	8.0* (1.5 to 14.5)	4.8 (-2.4 to 11.9)	3.2 (-4.1 to 10.5)	4.3 (-2.2 to 10.8)	2.7 (-3.2 to 8.6)
Adenoidectomy v no surgery	4.5 (-1.3 to 10.4)	4.3 (-1.4 to 9.9)	4.3 (-3.1 to 11.6)	2.4 (-5.7 to 10.5)	3.2 (-3.5 to 10.0)	3.5 (-3.2 to 10.3)
Myringotomy and grommet v adenoidectomy	3.0 (-2.1 to 8.1)	1.2 (-4.1 to 6.6)	-1.4 (-7.5 to 4.8)	-3.5 (-11.4 to 4.6)	-0.2 (-5.9 to 5.5)	-2.7 (-8.7 to 3.3)
Adenoidectomy, myringotomy and grommet v no surgery	9.6* (4.3 to 14.8)	7.6* (2.1 to 13.0)	5.3 (-1.3 to 11.9)	5.9 (-1.9 to 13.3)	4.6 (-1.3 to 10.4)	5.9 (-0.2 to 12.0)
Adenoidectomy, myringotomy and grommet v adenoidectomy	6.9* (0.8 to 13.0)	3.8 (-2.6 to 10.2)	0.0 (-4.0 to 4.0)	4.3 (-4.4 to 13.0)	0.3 (-6.8 to 7.4)	2.6 (-4.7 to 9.8)
Adenoidectomy, myringotomy and grommet v myringotomy and grommet	2.0 (-2.3 to 6.4)	2.1 (-2.6 to 6.8)	2.4 (-2.7 to 7.6)	6.9* (0.3 to 13.7)	1.5 (-3.3 to 6.4)	5.1 (0.0 to 10.2)

*Significant *t* value ($p < 0.05$).

TABLE VI—Comparisons of power of different preoperative levels of hearing to predict relative improvements in mean audiometric score of at least 5 dB and at least 10 dB six and 12 months after surgery. Predictor refers to mean preoperative level of hearing taken to indicate operation

Predictor (dB)	Operation indicated		Operation not indicated		% Of whole group denied benefit from surgery (95% confidence interval)
	No	% With good outcome (95% confidence interval)	No	% With good outcome (95% confidence interval)	
<i>Improvement ≥ 10 dB six months postoperatively</i>					
30	53	45 (32 to 59)	74	14 (7 to 24)	8 (4 to 14)
25	79	38 (27 to 50)	48	8 (3 to 21)	3 (1 to 8)
20	96	33 (25 to 44)	31	6 (1 to 23)	2 (0 to 6)
15	117	28 (20 to 37)	10	10 (0 to 46)	1 (0 to 5)
<i>Improvement ≥ 10 dB 12 months postoperatively</i>					
30	53	36 (23 to 50)	76	11 (5 to 20)	6 (3 to 12)
25	80	29 (19 to 40)	49	8 (3 to 20)	3 (1 to 8)
20	100	25 (17 to 35)	29	7 (1 to 24)	2 (0 to 6)
15	121	22 (15 to 31)	8	0 (0 to 40)	0 (0 to 4)
<i>Improvement ≥ 5 dB six months postoperatively</i>					
30	53	57 (42 to 70)	74	34 (23 to 46)	20 (13 to 28)
25	79	53 (42 to 64)	48	27 (16 to 42)	10 (6 to 17)
20	96	50 (40 to 60)	31	23 (10 to 42)	6 (2 to 11)
15	117	44 (35 to 54)	10	30 (8 to 65)	2 (0 to 7)
<i>Improvement ≥ 5 dB 12 months postoperatively</i>					
30	53	51 (37 to 65)	76	28 (18 to 39)	17 (11 to 24)
25	80	46 (35 to 58)	49	22 (12 to 37)	9 (5 to 15)
20	100	40 (30 to 50)	29	28 (13 to 47)	6 (3 to 12)
15	121	36 (28 to 46)	8	50 (17 to 82)	3 (1 to 8)




FIG 3—Proportion of ears with abnormal impedance (B and C2) preoperatively that remained abnormal postoperatively for different treatment groups using modified data. No data were available for ears in which a grommet was in place

Parental opinion—It is difficult to assess the state of each of their child's ears separately. Parental opinion could therefore be used as an outcome measure only in relation to the four treatment groups. The parents of children who had had an adenoidectomy were more satisfied than those whose children had not (fig 4). This difference persisted throughout the two years of follow up so that by the end of the second year about half of the children who had not had an adenoidectomy were thought to be satisfactory compared with around 60-70% of those who had.

INDICATIONS FOR SURGERY

Logistic regression analyses were carried out to establish the appropriate indications for inserting grommets with or without adenoidectomy. The predictive power of a wide range of variables was considered: patient characteristics (age, sex, social class); symptoms (deafness, otalgia, nasal obstruction, speech); findings on investigation (hearing level, impedance); and findings at operation (middle ear contents). With the outcome criterion being defined as a relative improvement in hearing level of 10 dB after 12 months in the ear that had a grommet compared with the ear that did not, the data were examined for a

subset of variables that had some predictive power. The most useful were the preoperative hearing level and the contents of the middle ear. Other variables contributed little additional predictive power. As the purpose of this analysis was to provide a basis for decisions about whether to operate, further analyses omitted the contents of the middle ear as this information may be reliably obtained only during surgery.

The accuracy of using preoperative audiometry scores as the sole predictor of outcome was tested using various different mean (for left and right ears together) preoperative hearing levels as indicative of surgery and two levels of improvement (5 dB and 10 dB), at six and 12 months after the operation, as indicative of a satisfactory outcome (table VI). At six months the proportion of children who had an improvement of 10 dB or more was 38% among those whose preoperative hearing loss was 25 dB or more (95% confidence interval 27% to 50%). At 12 months this had dropped to 29% (95% confidence interval 19% to 40%). The corresponding figures among those whose preoperative hearing loss was less than 25 dB was 8% at both six months and 12 months (95% confidence interval 3% to 20%).

Discussion

This trial was designed to assess the effectiveness of surgery for glue ear rather than its efficacy. As such, no attempt was made to alter existing clinical practice—for example, by insisting that highly experienced senior surgeons assessed the children and performed the operations. Most of the surgery was performed by senior house officers and, with a steady turnover of medical staff, around 15 doctors of different grades were involved in the preoperative and postoperative care and assessment of patients. Recruitment of children took considerably longer than expected. This was due to failure by junior medical staff to attempt to recruit patients rather than a poor response rate. A comparison of the characteristics of the children included in the trial with those of the population of children undergoing surgery²⁰ suggested that those included were representative and that no selection bias had operated. We believe that the clinical management the children experienced was fairly typical of otolaryngological practice in England and Wales in the 1980s. The results obtained are therefore likely to reflect the effectiveness of current practice.

The only important methodological problem experienced was the higher than predicted number of children whom we were unable to follow up for two years. The principal reason for this was the clinicians'




FIG 4—Proportion of children whose parents thought that their child's condition was unfavourable or uncertain at various times postoperatively according to treatment group

(and occasionally the parents') dissatisfaction with a child's progress, which they believed warranted further surgical intervention. To cope with this problem the data were modified in the way we described. The results obtained with sensitivity analysis were robust to the various assumptions we made about those lost to follow up. Nevertheless, it is necessary to bear this adjustment in mind, particularly when considering data that related to the two year follow up.

It was clear that myringotomy plus grommet insertion produced a significant improvement in hearing which lasted for six to 12 months. Adenoidectomy resulted in only a modest improvement in hearing, though there was some evidence to suggest this was more long lasting than that obtained from the insertion of grommets. This view was supported by the finding that normal function of the middle ear (measured by impedance tympanometry) was restored in about half the children who underwent an adenoidectomy compared with only about 20% of children after myringotomy plus grommet insertion. If, however, the primary objective of surgery for glue ear is to restore hearing then this apparent advantage of adenoidectomy is irrelevant. To achieve a rapid and significant improvement in hearing myringotomy plus grommet insertion is the treatment of choice. The addition of an adenoidectomy produces little additional benefit. In this respect the results of this trial are consistent with those of several other studies.^{2,6,8,9} Considering the operative risks and the greater economic and social costs of adenoidectomy compared with myringotomy plus grommet insertion, our results offer little justification for continuing to use adenoidectomy in the routine treatment of glue ear. The finding that the proportion of parents who expressed satisfaction with the treatment that their child had received was higher among those whose children had had an adenoidectomy than in those whose children had not might be explained by the first group's knowledge that everything that might have been done had been done.

The need for clinicians to identify those children who would benefit from surgery is clear. Unfortunately, none of the 15 published randomised, controlled trials has considered the issue quantitatively. Our study has, however, investigated the sensitivity and specificity of preoperative findings in predicting the outcome of surgery. Despite the uncertainties surrounding the level of objectivity of audiometry this single measure appears to be a useful predictor of outcome. The use of preoperative hearing level both for ears that had grommets inserted and those that had not should have inhibited the effects of regression towards the mean.

Interpretation of the preoperative audiometry score as a predictor of outcome of surgery depends on the definition of a satisfactory outcome in terms of improvement in hearing and on attitudes to unnecessary operations on the one hand and to missed cases (children who might have benefited from surgery but who were not treated) on the other. The confidence intervals from this study were wide, but the implications for current practice are potentially dramatic. For example, if satisfactory outcome is defined as an improvement of 10 dB six months after surgery, and if a strategy of operating only on children with a hearing loss of 25 dB and above is adopted, then only 79 of the 127 children with complete data in this trial would have been operated on, of whom it might be expected that 30 would have benefited and 49 would not. Four children, however, who might have benefited would have been missed. Alternatively, setting the operation threshold at 20 dB would have resulted in 96 operations being performed, with 32 children expected to benefit, and two potential beneficiaries being missed. If the

children in this trial were representative of children operated on for glue ear in England and Wales in 1986 then the adoption of a policy of only operating when the preoperative hearing loss is at least 25 dB would have had the following implications. Firstly, the total number of operations would have been reduced from 91 000 to about 57 000, of which 21 000 would have achieved a satisfactory improvement of 10 dB or more. Secondly, however, nearly 3000 of the 34 000 children who would have been regarded as inappropriate for surgery under this policy would have been denied such an improvement.

These figures give only an indication of the scale of the problem. As in any trial, the sample used might not have been representative of the population of children undergoing surgery for glue ear and the effectiveness of the surgeons concerned might not have been typical. Also predictors that have been derived from one set of patients will generally not perform as well when used with another set, and greater precision is required. It will be necessary to test the predictors on other samples of children to confirm our results.

Finally, it is important to recognise that, as with most trials of surgery for glue ear, the effectiveness of the operations was assessed in terms of improvement in hearing. No attempt was made to determine any possible longer term effects—namely, improvements in language skills or in educational achievements.

We thank Rosemary Heddon for her help, members of the medical, nursing, and clerical staff of the department of otolaryngology, Radcliffe Infirmary, Oxford, Richard Gledhill and other audiometricians for taking on additional work, and Klim McPherson of the university department of community medicine and general practice for advice on design of the study. We are grateful for the financial support received from Oxfordshire Health Authority, Oxford Regional Health Authority, and the Department of Health and Social Security.

- Black NA. Surgery for glue ear—a modern epidemic. *Lancet* 1984;i:835-7.
- Maw AR, Herod F. Otoscopy, impedance, and audiometric findings in glue ear treated by adenoidectomy and tonsillectomy. A prospective randomised study. *Lancet* 1986;i:1399-402.
- Gates GA, Avery CA, Priboda TJ, et al. Effectiveness of adenoidectomy and tympanostomy tubes in the treatment of chronic otitis media with effusion. *N Engl J Med* 1987;317:1444-51.
- Roydhouse N. Adenoidectomy for otitis media with mucoid effusion. *Ann Otol Rhinol Laryngol* 1980;89(suppl 68):312-5.
- Widemar L, Svensson C, Rynnel-Dagoo B, et al. The effect of adenoidectomy on secretory otitis media: a two year controlled prospective study. *Clin Otolaryngol* 1985;10:345-50.
- Lildholdt T. Unilateral grommet insertion and adenoidectomy in bilateral secretory otitis media: preliminary report of the results in 91 children. *Clin Otolaryngol* 1979;4:87-93.
- Richards SH, Kilby D, Shaw JD, et al. Grommets and glue ears: a clinical trial. *J Laryngol Otol* 1971;85:17-22.
- Bonding P, Tos M, Poulsen G. Unilateral insertion of grommets in bilateral secretory otitis media. *Acta Otolaryngol* 1982;Suppl 386:161-2.
- To SS, Pahor AL, Robin PE. A prospective trial of unilateral grommets for bilateral secretory otitis media in children. *Clin Otolaryngol* 1984;9:115-7.
- Fiellau-Nikolajsen M, Hojstet PE, Felding JU. Adenoidectomy for eustachian tube dysfunction: long term results from a randomised controlled trial. *Acta Otolaryngol* 1982;Suppl 386:129-31.
- Rynnel-Dagoo B, Ahlborn A, Schiratzki H. Effects of adenoidectomy. A controlled two year follow up. *Ann Otol Rhinol Laryngol* 1978;87:272-8.
- Bulman CH, Brook SJ, Berry MG. A prospective randomised trial of adenoidectomy versus grommet insertion in the treatment of glue ear. *Clin Otolaryngol* 1984;9:67-75.
- Brown MJM, Richards SH, Ambegaokar AG. Grommets and glue ear: a five year follow up of a controlled trial. *J R Soc Med* 1978;71:353-6.
- Mandel EM, Bluestone CD, Paradise JL, et al. Efficacy of myringotomy with and without tympanostomy tube insertion in the treatment of chronic otitis media with effusion in infants and children: results for the first year of a randomised clinical trial. In: Lim DJ, Bluestone CD, Klein JO, et al, eds. *Recent advances in otitis media with effusion*. Philadelphia: Decker, 1984:308-12.
- Zielhuis GA, Rach GH, Van Den Broek P. Screening for otitis media with effusion in preschool children. *Lancet* 1989;i:311-4.
- Archard JC. The place of myringotomy in the management of secretory otitis media in children. *J Laryngol Otol* 1967;81:309-15.
- Black NA, Crowther J, Freeland A. The effectiveness of adenoidectomy in the treatment of glue ear: a randomised controlled trial. *Clin Otolaryngol* 1986;11:149-55.
- Tos M, Poulsen G. Secretory otitis media. Late results of treatment with grommets. *Archives of Otolaryngology* 1976;102:672-5.
- Paradise JL, Smith CG, Bluestone CD. Tympanometric detection of middle ear effusion in infants and young children. *Pediatrics* 1976;58:198-210.
- Black NA. The aetiology of glue ear—a case control study. *Int J of Pediatr Otorhinolaryngol* 1985;9:121-33.

(Accepted 30 March 1990)