

NOTES

A *Burkholderia pseudomallei* Type III Secreted Protein, BopE, Facilitates Bacterial Invasion of Epithelial Cells and Exhibits Guanine Nucleotide Exchange Factor Activity

Mark P. Stevens,¹ Andrea Friebe,² Lowrie A. Taylor,¹ Michael W. Wood,¹
Philip J. Brown,^{1†} Wolf-Dietrich Hardt,²
and Edouard E. Galyov^{1*}

Division of Environmental Microbiology, Institute for Animal Health, Compton Laboratory, Berkshire RG20 7NN, United Kingdom,¹ and Institute of Microbiology, Swiss Federal Institute of Technology Zurich, ETH Zentrum, CH-8092 Zurich, Switzerland²

Received 24 March 2003/Accepted 12 May 2003

We report the characterization of BopE, a type III secreted protein that is encoded adjacent to the *Burkholderia pseudomallei* bsa locus and is homologous to *Salmonella enterica* SopE/SopE2. Inactivation of *bopE* impaired bacterial entry into HeLa cells, indicating that BopE facilitates invasion. Consistent with this notion, BopE expressed in eukaryotic cells induced rearrangements in the subcortical actin cytoskeleton, and purified BopE exhibited guanine nucleotide exchange factor activity for Cdc42 and Rac1 in vitro.

Burkholderia pseudomallei is the etiological agent of melioidosis, a severe invasive infection of humans and animals that is endemic in subtropical areas (3, 6). Melioidosis has a remarkable capacity for latency. Development of disease 26 years after geographical exposure has been reported (21), and relapse is common even in patients treated with antibiotics (4). This is believed to result from the ability of *B. pseudomallei* to invade nonphagocytic host cells and to survive and replicate within phagocytes, where antibiotics may be less effective (14, 16, 25). The mechanism by which *B. pseudomallei* enters epithelial cells is poorly understood.

We and others have identified a putative type III protein secretion apparatus in *B. pseudomallei* (Bsa) similar to the *Salmonella enterica* Inv/Spa/Prg and *Shigella flexneri* Ipa/Mxi/Spa systems (1, 26, 31). Type III secretion systems are key virulence determinants of *Salmonella*, *Shigella*, and other gram-negative facultative intracellular pathogens and serve to inject bacterial proteins into target cells (reviewed in references 5, 10, 13, and 29). A subset of type III secretion system secreted proteins (translocators) is believed to interact with the eukaryotic cell membrane and mediate the delivery of secreted effector proteins. Once inside host cells the effector proteins subvert host cell processes to the benefit of the bacteria (reviewed in references 5 and 13).

Research in our laboratory and elsewhere has identified a number of *Salmonella* Inv/Spa/Prg secreted effector proteins

(Sops) and shown that several of these are delivered into eukaryotic cells by mechanisms dependent on secreted translocator proteins (Sips) (11, 34, 35). Mutations that disrupt the Inv/Spa/Prg apparatus and selected *sip* and *sop* genes inhibit bacterial invasion of epithelial cells and *Salmonella*-induced enteritis (reviewed in references 10, 33, and 36). Some Sop effector proteins possess eukaryote-like enzymatic activities. In particular, it has been shown that the SopE and SopE2 proteins promote bacterial invasion (2, 35) by acting as guanine nucleotide exchange factors (GEFs) for RhoGTPases that regulate the actin network (12, 27, 30). SopE acts as a GEF for Cdc42, Rac1, and Rab5 (8, 9, 12, 23, 27); however, SopE2 efficiently activates Cdc42 but not Rac1 (8, 30), indicating that SopE and SopE2 may activate different signaling cascades during *Salmonella* infection. Mutation of *Salmonella sopE* and *sopE2* reduces the induction of intestinal inflammatory and secretory responses in calves, suggesting that they play a role in *Salmonella*-induced enteritis (36, 37).

Recently we reported that mutations affecting putative components of the *B. pseudomallei* Bsa secretion and translocation apparatus impair intracellular survival of *B. pseudomallei* in murine macrophage-like cells and prevent escape of the bacteria from endocytic vesicles (31). Here we have investigated the role of a putative Bsa-secreted protein (BopE) that shares homology with the *Salmonella* SopE/SopE2 proteins. BopE is 27% identical over 168 amino acids to SopE and 28% identical over 139 amino acids to SopE2.

BopE is secreted by the Bsa type III secretion apparatus. To study expression and secretion of BopE, a BopE-glutathione-S-transferase fusion protein was generated and polyclonal antiserum was raised against BopE in rabbits. A DNA fragment encoding the domain of BopE proposed to be required for GEF activity (amino acid residues 78 to 261) was amplified

* Corresponding author. Mailing address: Division of Environmental Microbiology, Institute for Animal Health, Compton Laboratory, Berkshire RG20 7NN, United Kingdom. Phone: 44 1635 577291. Fax: 44 1635 577243. E-mail: edouard.galyov@bbsrc.ac.uk.

† Present address: Nuffield Department of Clinical Laboratory Sciences, University of Oxford, John Radcliffe Hospital, Oxford OX3 9DU, United Kingdom.




FIG. 1. Western blot analysis of BopE expression and secretion by *B. pseudomallei* 10276 wild type and *bsaZ*, *bipD*, and *bopE* mutant strains. Approximately 25 µg of total protein (A) or secreted protein (B) was probed with rabbit polyclonal antiserum to BopE₇₈₋₂₆₁ and detected with an anti-rabbit alkaline phosphatase conjugate. Molecular mass markers are shown on the left.

using the primers BopEGexBam (5'-CGGCAGCTATGGATCCACGGGCGACGCGAAAC-3') and BopEGexE1 (5'-CCACGCTGAATTCTCACGCGCCGTCC-3') and the product cloned into pGEX-2T (Amersham Biosciences, Little Chalfont, Buckinghamshire, England) via *EcoRI* and *BamHI* sites (underlined) in the primers. Following expression in *Escherichia coli* BL21(DE3) under isopropyl-β-D-thiogalactoside induction, the fusion protein was purified using glutathione Sepharose 4B resin and BopE₇₈₋₂₆₁ released from glutathione-S-transferase by thrombin digestion. A 12-week-old New Zealand White rabbit was immunized subcutaneously four times at 2-week intervals with ca. 100 µg of purified BopE₇₈₋₂₆₁ in Freund's incomplete adjuvant and serum collected 12 days after the final booster.

The BopE-specific antiserum was used to detect BopE in whole-cell and secreted protein fractions of *B. pseudomallei* strain 10276 and defined *bsaZ*, *bipD*, and *bopE* mutant strains described previously (31). *BsaZ* and *BipD* are homologous to the *Salmonella* SpaS and SipD proteins involved in secretion and translocation of Sop proteins, respectively. Bacteria were grown to stationary phase in Luria-Bertani broth, and culture supernatants were passed through 0.22-µm-pore-size filters prior to precipitation of secreted proteins with trichloroacetic acid (10% [vol/vol]). Approximately 25 µg of total protein (Fig. 1A) or secreted protein (Fig. 1B) was resolved by 4-to-15% sodium dodecyl sulfate-polyacrylamide gel electrophoresis and transferred to Immobilon-P membrane (Millipore, Bedford, Mass.). A 1:100 dilution of rabbit polyclonal antiserum to BopE₇₈₋₂₆₁ was used, and bound antibody was detected with an anti-rabbit alkaline phosphatase conjugate. As expected, BopE

was detected in whole-cell extracts of all strains except the 10276 *bopE* mutant (Fig. 1A). BopE secretion was dependent on the Bsa type III secretion apparatus, as no secretion was observed in a *bsaZ* mutant (Fig. 1B). In contrast, BopE secretion was elevated in *B. pseudomallei* lacking the putative translocator BipD (Fig. 1B). These data are consistent with the observation that *Salmonella sip* mutants secrete elevated levels of selected Sops (15, 35). Thus, our data suggest that the *B. pseudomallei* Bsa type III secretion apparatus is functional and that BopE is type III secreted.

BopE facilitates invasion of nonphagocytic cells. *B. pseudomallei* can invade and survive within nonphagocytic cells (14, 16). To assess the role of BopE in bacterial invasion, we quantified intracellular *B. pseudomallei* following infection of HeLa cells by strains 10276, 10276 *bipD*, and 10276 *bopE* by using a kanamycin protection assay. Previously we have been unable to detect significant invasion of HeLa cells by *B. pseudomallei* strain 10276 (31); however, we have found that invasion efficiency can be improved by centrifugation of the bacteria onto cell monolayers at 300 × g at the onset of infection. HeLa cells maintained in RPMI 1640 containing 10% (vol/vol) fetal calf serum were infected at a multiplicity of 10 with *B. pseudomallei* strains grown to stationary phase in Luria-Bertani broth at 37°C in a humidified 5% CO₂ atmosphere. One hour after bacterial inoculation, monolayers were washed three times and overlaid with medium containing kanamycin (250 µg/ml) to kill extracellular bacteria. After 6 h viable intracellular bacteria were released by gentle lysis using 0.1% Triton X-100 and enumerated by plating of serial dilutions. We detected a statistically significant reduction in invasion of HeLa cells by the 10276 *bopE* mutant compared to that of the wild type (*P* = 0.0464) (Fig. 2), indicating that BopE, like *Salmonella* SopE/SopE2, facilitates bacterial invasion of nonphagocytic cells. SopE acts in concert with other type III secreted proteins to promote *Salmonella* invasion (38). Recently it was reported




FIG. 2. Invasion of HeLa cells by *B. pseudomallei* 10276 wild type and *bipD* and *bopE* mutant strains. HeLa cells (5 × 10⁵) were infected at a multiplicity of infection 10 in triplicate for each assay and the results represent the arithmetic means (error bars show standard errors of the means) of results of four independent assays.




FIG. 3. Confocal micrographs showing rearrangement of the subcortical actin cytoskeleton following expression of BopE and SopE in transiently transfected HeLa cells. HeLa cells were transfected with pRK5myc-SopE or pRK5myc-BopE. F-actin was stained red with tetramethyl rhodamine isothiocyanate-conjugated phalloidin, and *myc*-tagged protein was stained green with a *myc*-specific monoclonal antibody detected with anti-mouse Alexa⁴⁸⁸ conjugate. Nontransfected cells are present in each field for comparison. Magnification, $\times 500$.

that the effector protein SopB, which possesses phosphatidylinositol phosphatase activity, influences *Salmonella* invasion (24, 39). It is likely that other type III secreted proteins influence invasion of nonphagocytic cells by *B. pseudomallei*. Consistent with this hypothesis we detected a highly significant reduction in invasion of HeLa cells by the 10276 *bipD* mutant ($P = 0.0058$). *B. pseudomallei* is a Centers for Disease Control and Prevention category B critical biological agent, and we were unable to *trans*-complement the *bopE* mutation owing to restrictions on genetic modification of the organism.

BopE expressed in eukaryotic cells induces rearrangements in the subcortical actin cytoskeleton. To assess the activity of BopE in eukaryotic cells, we amplified the entire coding sequence of *bopE* using the primers BopEpRKBam (5'-CTCCGATCCATGACTTACAACCCGAGAATCGGCGG-3') and BopEpRKE1 (5'-CTCGAATTCTCACGCGCCGTCCGCCGCTTCGTCGC-3') and cloned the product into pRK5myc (17) via *EcoRI* and *BamHI* (underlined) sites. This created a BopE fusion protein with a *myc* tag at the N terminus. As a control the *Salmonella enterica* serovar Typhimurium *sopE* gene was cloned into pRK5myc in the same way using the primers SopEpRKBam (5'-CTCGGATCCGTGACAAAAAT AACTTTATTTC-3') and SopEpRKE1 (5'-CTCGAATTCT CAGGGAGTGTGTTGGATATATT-3'). HeLa cells were transfected with pRK5myc-BopE or pRK5myc-SopE by using Lipofectamine (Invitrogen Life Technologies, Paisley, United

Kingdom). Twenty-four hours after transfection, cells were stained for the presence of the *myc*-tagged protein with a mouse monoclonal *myc*-specific antibody (Invitrogen) detected with anti-mouse Alexa⁴⁸⁸ conjugate (Molecular Probes, Leiden, The Netherlands). Filamentous actin was stained using tetramethyl rhodamine isothiocyanate-conjugated phalloidin and the cells were viewed using a Leica TCS NT confocal laser scanning microscope. In cells expressing the *myc*-tagged BopE and SopE proteins filamentous actin was abundant under the membrane and was associated with areas of "ruffling" (Fig. 3), suggesting that the proteins interfere with actin dynamics in eukaryotic cells. Such rearrangements were not detected in nontransfected cells present in the same field. Some SopE and BopE appeared to colocalize with regions of intense F-actin staining in membrane ruffles (Fig. 3).

BopE is a GEF. To determine if BopE possesses GEF activity, we employed fluorescence spectrometry using mGDP-loaded Rac1 (Fig. 4A) or Cdc42 (Fig. 4B) as a substrate (8). mGDP is a GDP derivative which is popular for kinetic studies because the fluorescence intensity of the mant moiety changes dramatically upon binding to GTPases. mGDP bound to Cdc42 has a fourfold-higher fluorescence intensity than unbound mGDP (20, 27, 28). Generally, the presence of the fluorophore has little effect on the kinetic parameters of G nucleotide release or GTP hydrolysis (7, 18, 19, 20, 32).

BopE₇₈₋₂₆₁ was expressed and purified as described above




FIG. 4. BopE acts as a GEF for the RhoGTPases Rac1 and Cdc42. The multiple turnover kinetics of guanine nucleotide exchange by BopE was analyzed by measuring the release of mantGDP from 10 μ M Rac₁₋₁₉₁·mantGDP (A) or Cdc42 Hs₁₋₁₉₂·mantGDP (B) in the presence of 1 mM GDP and 25 nM or 250 nM BopE using fluorescence spectrometry (excitation wavelength, 366 nm; emission wavelength, 440 nm; step size, 1; band-pass, 4) at 20°C in a buffer containing 40 mM HEPES–NaOH (pH 7.4), 100 mM NaCl, and 5 mM MgCl₂. Spontaneous dissociation of the RhoGTPase·mantGDP complex in the assay buffer or in assay buffer supplemented with 10 mM EDTA was measured as the control.

and mGDP-Rac1 or mGDP-Cdc42 was prepared as described previously (8, 9). In the assay buffer (no EDTA or BopE) mGDP dissociation from Rac1 was very slow. In contrast, fast dissociation of the mGDP-Rac1 complex was observed in the presence of 25 nM BopE and even faster in the presence of 250 nM BopE (Fig. 4A) ($k_{\text{obs}} = 0.48 \text{ s}^{-1}$). Similar observations were made using mGDP-Cdc42 as a substrate (Fig. 4B). These data demonstrate that BopE is an efficient GEF for Cdc42 and Rac1. The observed G-nucleotide exchange rates are lower than those observed with SopE from *Salmonella* serovar Typhimurium (8) but range in the same order of magnitude.

Taken together our observations suggest that *B. pseudomallei* enters epithelial cells by a mechanism dependent at least in part upon the Bsa type III protein secretion apparatus and one of its secreted proteins, BopE. It is likely that BopE is translocated into the host cell cytosol, where it may promote membrane ruffling by acting as a GEF for Cdc42 and Rac1. In *Salmonella* several Inv/Spa/Prg-secreted proteins (SopE, SopE2, and SopB) act in concert to promote bacterial uptake by non-phagocytic cells (38, 39), and *Salmonella* invasion probably evolved through the acquisition of new sequence elements (22). Given that a *B. pseudomallei* *bipD* mutant was impaired in invasion of HeLa cells to a greater extent than a *bopE* mutant, it is likely that other Bsa-secreted proteins may be involved in bacterial uptake. We are investigating the role of other putative type III secreted proteins in the host-cell interactions of *B. pseudomallei*.

REFERENCES

1. Attree, O., and I. Attree. 2001. A second type III secretion system in *Burkholderia pseudomallei*: who is the real culprit? *Microbiology* **147**:3197–3199.
2. Bakshi, C. S., V. P. Singh, M. W. Wood, P. W. Jones, T. S. Wallis, and E. E. Galyov. 2000. Identification of SopE2, a *Salmonella* secreted protein which is highly homologous to SopE and involved in bacterial invasion of epithelial cells. *J. Bacteriol.* **82**:2341–2344.
3. Brett, P. J., and D. E. Woods. 2000. Pathogenesis of and immunity to melioidosis. *Acta Trop.* **74**:201–210.
4. Chaowagul, W., Y. Suputtamongkol, D. A. B. Dance, A. Rajchanuvong, J. Pattaraarechachai, and N. J. White. 1993. Relapse in melioidosis: incidence and risk factors. *J. Infect. Dis.* **168**:1181–1185.
5. Cornelis, G. R., and F. van Gijsegem. 2000. Assembly and function of type III secretion systems. *Annu. Rev. Microbiol.* **54**:735–774.
6. Dance, D. A. 2000. Melioidosis is an emerging global problem. *Acta Trop.* **74**:115–119.
7. Franken, S. M., A. J. Scheidig, U. Krengel, H. Rensland, A. Lautwein, M. Geyer, K. Scheffzek, R. S. Goody, H. R. Kalbitzer, E. F. Pai, et al. 1993. Three-dimensional structures and properties of a transforming and a non-transforming glycine-12 mutant of p21H-ras. *Biochemistry* **32**:8411–8420.
8. Friebe, A., H. Ilchmann, M. Aepfelbacher, K. Ehrbar, W. Machleidt, and W.-D. Hardt. 2001. SopE and SopE2 from *Salmonella typhimurium* activate different sets of RhoGTPases of the host cell. *J. Biol. Chem.* **276**:34035–34040.
9. Friebe, A., and W.-D. Hardt. 2000. Purification and biochemical activity of *Salmonella* exchange factor SopE. *Methods Enzymol.* **325**:82–91.
10. Galán, J. E. 2001. *Salmonella* interactions with host cells: type III secretion at work. *Annu. Rev. Cell. Dev. Biol.* **17**:53–86.
11. Galyov, E. E., M. W. Wood, R. Rosqvist, P. B. Mullan, P. R. Watson, S. Hedges, and T. S. Wallis. 1997. A secreted effector protein of *Salmonella dublin* is translocated into eukaryotic cells and mediates inflammation and fluid secretion in infected ileal mucosa. *Mol. Microbiol.* **25**:903–912.
12. Hardt, W.-D., L. M. Chen, K. E. Schuebel, X. R. Bustelo, and J. E. Galán. 1998. *S. typhimurium* encodes an activator of Rho GTPases that induces membrane ruffling and nuclear responses in host cells. *Cell* **93**:815–826.
13. Hueck, C. J. 1998. Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiol. Mol. Biol. Rev.* **62**:379–433.
14. Jones, A. L., T. J. Beveridge, and D. E. Woods. 1996. Intracellular survival of *Burkholderia pseudomallei*. *Infect. Immun.* **64**:782–790.
15. Kaniga, K., D. Trollinger, and J. E. Galán. 1995. Identification of two targets of the type III protein secretion system encoded by the *inv* and *spa* loci of *Salmonella typhimurium* that have homology to the *Shigella* IpaD and IpaA proteins. *J. Bacteriol.* **177**:7078–7085.
16. Kespichayawattana, W., S. Rattanachetkul, T. Wanun, P. Utaincharoen, and S. Sirisinha. 2000. *Burkholderia pseudomallei* induces cell fusion and actin-associated membrane protrusion: a possible mechanism of cell-to-cell spreading. *Infect. Immun.* **68**:5377–5384.
17. Lamarche, N., N. Tapon, L. Stowers, P. D. Burbelo, P. Aspenstrom, T. Bridges, J. Chant, and A. Hall. 1996. Rac and Cdc42 induce actin polymerization and G1 cell cycle progression independently of p65PAK and the JNK/SAPK MAP kinase cascade. *Cell* **87**:519–529.
18. Lenzen, C., R. H. Cool, and A. Wittinghofer. 1995. Analysis of intrinsic and

- CDC25-stimulated guanine nucleotide exchange of p21ras-nucleotide complexes by fluorescence measurements. *Methods Enzymol.* **255**:95–109.
19. **Lenzen, C., R. H. Cool, H. Prinz, J. Kuhlmann, and A. Wittinghofer.** 1998. Kinetic analysis by fluorescence of the interaction between Ras and the catalytic domain of the guanine nucleotide exchange factor Cdc25Mm. *Biochemistry* **37**:7420–7430.
 20. **Leonard, D. A., T. Evans, M. Hart, R. A. Cerione, and D. Manor.** 1994. Investigation of the GTP-binding/GTPase cycle of Cdc42Hs using fluorescence spectroscopy. *Biochemistry* **33**:12323–12328.
 21. **Mays, E. E., and E. A. Ricketts.** 1975. Melioidosis: recrudescence associated with bronchogenic carcinoma twenty-six years following initial geographic exposure. *Chest* **68**:261–263.
 22. **Mirolid, S., K. Ehrbar, A. Weissmuller, R. Prager, H. Tschape, H. Russmann, and W.-D. Hardt.** 2001. *Salmonella* host cell invasion emerged by acquisition of a mosaic of separate genetic elements, including *Salmonella* pathogenicity island 1 (SPI1), SPI5, and sopE2. *J. Bacteriol.* **183**:2348–2358.
 23. **Mukherjee, K., S. Parashuraman, M. Raje, and A. Mukhopadhyay.** 2001. SopE acts as an Rab5-specific nucleotide exchange factor and recruits non-prenylated Rab5 on *Salmonella*-containing phagosomes to promote fusion with early endosomes. *J. Biol. Chem.* **276**:23607–23615.
 24. **Norris, F. A., M. P. Wilson, T. S. Wallis, E. E. Galyov, and P. W. Majerus.** 1998. SopB, a protein required for virulence of *Salmonella dublin*, is an inositol phosphatase. *Proc. Natl. Acad. Sci. USA* **95**:14057–14059.
 25. **Pruksachartvuthi, S., N. Aswapokee, and K. Thankerngpol.** 1990. Survival of *Pseudomonas pseudomallei* in human phagocytes. *J. Med. Microbiol.* **31**:109–114.
 26. **Rainbow, L., C. A. Hart, and C. Winstanley.** 2002. Distribution of type III secretion gene clusters in *Burkholderia pseudomallei*, *B. thailandensis* and *B. mallei*. *J. Med. Microbiol.* **51**:374–384.
 27. **Rudolph, M. G., C. Weise, S. Mirolid, B. Hillenbrand, B. Bader, A. Wittinghofer, and W.-D. Hardt.** 1999. Biochemical analysis of SopE from *Salmonella typhimurium*, a highly efficient guanosine nucleotide exchange factor for RhoGTPases. *J. Biol. Chem.* **274**:30501–30509.
 28. **Rudolph, M. G., P. Bayer, A. Abo, J. Kuhlmann, I. R. Vetter, and A. Wittinghofer.** 1998. The Cdc42/Rac interactive binding region motif of the Wiskott Aldrich syndrome protein (WASP) is necessary but not sufficient for tight binding to Cdc42 and structure formation. *J. Biol. Chem.* **273**:18067–18076.
 29. **Sansonetti, P. J.** 2001. Rupture, invasion and inflammatory destruction of the intestinal barrier by *Shigella*, making sense of prokaryote-eukaryote cross-talks. *FEMS Microbiol. Rev.* **25**:3–14.
 30. **Stender, S., A. Friebel, S. Linder, M. Rohde, S. Mirolid, and W.-D. Hardt.** 2000. Identification of SopE2 from *Salmonella typhimurium*, a conserved guanine nucleotide exchange factor for Cdc42 of the host cell. *Mol. Microbiol.* **36**:1206–1221.
 31. **Stevens, M. P., M. W. Wood, L. A. Taylor, P. Monaghan, P. Hawes, P. W. Jones, T. S. Wallis, and E. E. Galyov.** 2002. An Inv/Mxi-Spa-like type III protein secretion system in *Burkholderia pseudomallei* modulates intracellular behavior of the pathogen. *Mol. Microbiol.* **46**:649–659.
 32. **Tan, Y. C., H. Wu, W. N. Wang, Y. Zheng, and Z. X. Wang.** 2002. Characterization of the interactions between the small GTPase RhoA and its guanine nucleotide exchange factors. *Anal. Biochem.* **310**:156–162.
 33. **Wallis, T. S., and E. E. Galyov.** 2000. Molecular basis of *Salmonella*-induced enteritis. *Mol. Microbiol.* **36**:997–1005.
 34. **Wood, M. W., M. A. Jones, P. R. Watson, A. M. Siber, B. A. McCormick, S. Hedges, R. Rosqvist, T. S. Wallis, and E. E. Galyov.** 2000. The secreted effector protein of *Salmonella dublin*, SopA, is translocated into eukaryotic cells and influences the induction of enteritis. *Cell. Microbiol.* **2**:293–303.
 35. **Wood, M. W., R. Rosqvist, P. B. Mullan, M. H. Edwards, and E. E. Galyov.** 1996. SopE, a secreted protein of *Salmonella dublin*, is translocated into the target eukaryotic cell via a Sip-dependent mechanism and promotes bacterial entry. *Mol. Microbiol.* **22**:327–338.
 36. **Zhang, S., R. A. Kingsley, R. L. Santos, H. Andrews-Polymeris, M. Raffatellu, J. Figueiredo, J. Nunes, R. M. Tsois, L. G. Adams, and A. J. Baumler.** 2003. Molecular pathogenesis of *Salmonella enterica* serotype Typhimurium-induced diarrhea. *Infect. Immun.* **71**:1–12.
 37. **Zhang, S., R. L. Santos, R. M. Tsois, S. Mirolid, W.-D. Hardt, L. G. Adams, and A. J. Baumler.** 2002. Phage mediated horizontal transfer of the *sopE1* gene increases enteropathogenicity of *Salmonella enterica* serotype Typhimurium for calves. *FEMS Microbiol. Lett.* **217**:243–247.
 38. **Zhou, D., and J. Galán.** 2001. *Salmonella* entry into host cells: the work in concert of type III secreted effector proteins. *Microbes Infect.* **3**:1293–1298.
 39. **Zhou, D., L. M. Chen, L. Hernandez, S. B. Shears, and J. E. Galán.** 2001. A *Salmonella* inositol polyphosphatase acts in conjunction with other bacterial effectors to promote host cell actin cytoskeleton rearrangements and bacterial internalization. *Mol. Microbiol.* **39**:248–259.