

# Characterization of a Strong Dominant *phytochrome A* Mutation Unique to Phytochrome A Signal Propagation<sup>1</sup>

Rebecca C. Fry<sup>2</sup>, Jessica Habashi, Haruko Okamoto<sup>3</sup>, and Xing Wang Deng\*

Department of Molecular, Cellular, and Developmental Biology, Yale University, New Haven, Connecticut 06511-8104

Here, we report the isolation and characterization of a strong dominant-negative *phytochrome A* (*phyA*) mutation (*phyA-300D*) in *Arabidopsis*. This mutation carries a single amino acid substitution at residue 631, from valine to methionine (V631M), in the core region within the C-terminal half of PHYA. This PHYA core region contains two protein-interactive motifs, PAS1 and PAS2. Val-631 is located within the PAS1 motif. The *phyA-V631M* mutant protein is photochemically active and accumulates to a level similar to wild type in dark-grown seedlings. Overexpression of *PHYA-V631M* in a wild-type background results in a dominant-negative interference with endogenous wild-type *phyA*, whereas *PHYA-V631M* in a *phyA* null mutant background is inactive. To investigate the specificity of this mutation within the phytochrome family, the corresponding amino acid substitution (V664M) was created in the PHYTOCHROME B (PHYB) polypeptide. We found that the *phyB-V664M* mutant protein is physiologically active in *phyB* mutant and causes no interfering effect in a wild-type background. Together, our results reveal a unique feature in *phyA* signal propagation through the C-terminal core region.

One of the most important environmental factors affecting plant growth and survival is light. Plants employ arrays of photoreceptors to detect and respond to a broad spectrum of light (Kendrick and Kronenberg, 1994; McNellis and Deng, 1995). One of the best-studied groups of photoreceptors is the red (R)/far red (FR) light-absorbing phytochromes (Quail, 1997). In *Arabidopsis*, there is a family of five phytochromes whose apoproteins are encoded by a multigene family (*PHYA-PHYE*; Sharrock and Quail, 1989; Clack et al., 1994). Among the five phytochromes, the best characterized is the FR-absorbing phytochrome A (*phyA*). Like all phytochromes, *phyA* is capable of a photoreversible conformational change between the Pr and the Pfr forms in response to light. Whereas the Pfr form is rapidly degraded, the Pr form of *phyA* is stable in the cytosol. Upon exposure to R light, the Pr form is converted to the Pfr form, whereas exposure to FR light converts the Pfr form back to the Pr form. This photoconversion correlates with the absorption spectra of these forms, where Pr absorbs maximally at 666 nm and Pfr absorbs maximally at 730 nm (Quail, 1997). It is currently hypothesized that both the Pfr and the photo-

cycled Pr forms of *phyA* are the active species (Shinomura et al., 2000) and possess kinase activity (Yeh and Lagarias, 1998). Upon light perception, *phyA* and phytochrome B (*phyB*) migrate into the nucleus, where they presumably trigger downstream events (Kircher et al., 1999).

Since the discovery of the phytochrome system, the search has been on to identify the signal transduction mechanism through which light perception is coupled to changes in gene expression and cell physiology that control growth and development. Use of both biochemical and genetic means have identified candidate downstream-signaling components of the phytochrome pathway. Microinjection into hypocotyl cells of the phytochrome-deficient *aurea* mutant of tomato (*Lycopersicon esculentum*) has been used to biochemically assay the activities of various pharmacological agents and putative signaling intermediates (Bowler et al., 1994). Such studies suggest that heterotrimeric G proteins Ca<sup>2+</sup>-calmodulin and cyclic-guano-5' monophosphate may mediate phytochrome-induced responses. A reverse genetic approach has recently implicated the involvement of the only G- $\alpha$  protein in *Arabidopsis* in light inhibition of hypocotyl elongation (Okamoto et al., 2001).

Both yeast (*Saccharomyces cerevisiae*) two-hybrid protein-protein interaction screens and mutational approaches have resulted in identification of over a dozen putative downstream components and interacting partners of phytochromes (for review, see Quail, 2002; Wang and Deng, 2002). Those putative signaling components include both cytoplasmic and nuclear factors, some of which are transcription factors directly acting on light-responsive promoters. For *phyA*, the majority of the signaling components identified so far are nuclear localized. This is consistent with the view that light-activated *phyA* rapidly

<sup>1</sup> This work was supported by the National Institutes of Health (grant no. GM47850 to X.W.D.). X.W.D. was a National Science Foundation Presidential Faculty Fellow, and J.H. is a National Institutes of Health predoctoral trainee.

<sup>2</sup> Present address: Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139.

<sup>3</sup> Present address: The Department of Plant Sciences, University of Oxford, Oxford OX1 3RB, UK.

\* Corresponding author; e-mail xingwang.deng@yale.edu; fax 203-432-5726.

Article, publication date, and citation information can be found at [www.plantphysiol.org/cgi/doi/10.1104/pp.005264](http://www.plantphysiol.org/cgi/doi/10.1104/pp.005264).

**Table I.** Summary of the mutant screen

Isolate Name	Allele	Nature of Mutation
128	<i>fhv3-128</i>	Recessive
6476	<i>fhv3-6476</i>	Recessive
A2GO5	<i>phyA-300D</i>	Dominant
135	<i>phyA-301</i>	Recessive
158	<i>hy5-158</i>	Recessive
CS19960	<i>fhv1-19960</i>	Recessive
149	<i>fhv1-149</i>	Recessive
A2GO6	<i>fhv4-1</i>	Recessive

migrates into nucleus (Kircher et al., 1999) and triggers most of the signaling events there.

A plethora of studies have revealed that phytochrome molecules contain multiple functional domains (for review, see Quail, 1997; Park et al., 2000). The N-terminal half contains the chromophore attachment site and is responsible for the observed spectrum specificity of phyA and phyB. The C-terminal half seems to be responsible for dimerization and downstream signal transmission (Jordan et al., 1996). A region in the C-terminal half, designated as the "core region" or "Quail box," was revealed to be critically important for phytochrome signal propagation (Quail et al., 1995; Park et al., 2000). This core region in phyA contains two PAS motifs, presumed to be involved in protein-protein interaction. This region was shown to be important for interaction with the reported phytochrome-interactive partners (Ni et al., 1998; Choi et al., 1999). Although it was reported that the C-terminal half of phyA and phyB could be interchangeable for some R and FR responses (Quail et al., 1995), some aspect of distinction in the phyA and phyB signal propagation through their C-terminal core regions to downstream targets might be expected because there are a number of unique downstream components specific to either phyA or phyB.

Here, we report that in a mutant screen for an etiolated phenotype under continuous FR light (FRC), we have recovered a strong dominant-negative mutation of *PHYA*. This dominant interfering phenotype observed in the heterozygous state is even stronger than those previously reported for transgenic seedlings overexpressing phyA dominant-negative mutant forms (Boylan et al., 1994; Emmler et al., 1995). Characterization of this new *phyA* mutant revealed an insight specific to phyA-signaling propagation through its C-terminal domain.

## RESULTS

### Isolation of *phyA-300D*, a Strong Dominant-Negative Allele of *PHYA*

In an effort to isolate mutants impaired specifically in phyA signaling, long hypocotyl mutant seedlings specific for FRC were screened from several available T-DNA insertion mutagenized collections. Seven elongated hypocotyl mutants were identified (Table

I). Genetic complementation tests revealed that most of the mutations defined new alleles of known genes; however, one new locus, designated *FHY4*, was defined by a single-mutant allele (*fhv4-1*).

One of the mutants, now designated *phyA-300D*, exhibited a 3:1 mutant:wild-type phenotype segregation in a backcross to wild type (Table II). This result suggests that the *phyA-300D* mutation behaves as a dominant trait. This was further supported by the fact that during our allelism test between *phyA-300D* and various known phyA-signaling mutants, the F<sub>1</sub> progenies always displayed long hypocotyl phenotypes, contrasting to those of a normal recessive *phyA* allele (Table II; data not shown). In the F<sub>2</sub> populations, it was revealed that *phyA-300D* complements all other mutations except *phyA*, indicating that this may be a new allele of *PHYA*. As shown in Figure 1A, seedlings heterozygous for the *phyA-300D* allele possess long hypocotyls under FRC, similar to that of the homozygous seedlings. However, *phyA-300D* seedlings grown under other light wavelengths displayed normal de-etiolated phenotypes, suggesting that the *phyA-300D* phenotype is FRC specific (Fig. 1B).

To further verify this dominant mutant as a *phyA* allele, mapping analysis of the *phyA-300D* mutation was carried out (see "Materials and Methods"). As expected, *phyA-300D* was mapped to chromosome 1 near the *PHYA* locus (data not shown).


### *phyA-300D* Is Caused by a Single Amino Acid Change at Residue 631

To determine the exact nature of the *phyA-300D* mutation, overlapping regions covering the *PHYA* gene from the mutant background were PCR amplified and sequenced. This sequence analysis revealed two nucleotide changes, both from guanine (G) to adenine (A), affecting two adjacent codons of the *PHYA* open reading frame (Fig. 2). The first G to A mutation is silent, altering the codon 630 from ACG to ACA, both encoding the same amino acid, Thr.

**Table II.** Complementation analysis of *phyA* mutants

F<sub>1</sub> and F<sub>2</sub> progeny of the complementation crosses were grown in continuous far red light for 6 d before scoring seedling phenotypes and their segregation.

Cross	Generation	Seedlings with Short Hypocotyl	Seedlings with Long Hypocotyl
<i>phyA-300D</i> × WT	F <sub>1</sub>	0	69
	F <sub>2</sub>	188	677
<i>phyA-300D</i> × <i>phyA-1</i>	F <sub>1</sub>	0	2
	F <sub>2</sub>	0	400
<i>phyA-205</i> × WT	F <sub>1</sub>	0	14
	F <sub>2</sub>	105	370
<i>phyA-205</i> × <i>phyA-300D</i>	F <sub>1</sub>	0	9
	F <sub>2</sub>	0	200
<i>phyA-301</i> × WT	F <sub>1</sub>	30	0




**Figure 1.** Phenotype characterization of the dominant *phyA-300D* mutant. A, Three-day-old FRc-grown *phyA-300D* homozygote (-/-), heterozygote (-/+), and wild-type (WT) seedlings. Average hypocotyl lengths (in millimeters) are shown above each seedling ( $n = 25$ ) with SD shown in parentheses. B, Hypocotyl length of wild-type (WT) and *phyA-300D* seedlings grown in continuous white (W) and blue (B) light and Rc (R) and FRc (FR), or in darkness (D) for 6 d. Each column represents a mean value of the hypocotyl length (in millimeters) taken from 25 seedlings with the SD indicated by an error bar.

The second G to A nucleotide change is a missense mutation, which alters the codon at position 631 from GTG (encoding for Val) to ATG (encoding for Met). Thus, the overall consequence in *phyA-300D* is a Val to Met change at amino acid 631 (V631M) in the PHYA polypeptide (Fig. 2C). This Val-631 is located within the first of the two PAS motifs, within the core region or Quail box of the phytochrome C-terminal half (Fig. 2). The core region of *phyA* has been demonstrated to be critical in phytochrome signal propagation to downstream targets (Quail et al., 1995). However, all those reported mutations had been described as recessive in nature (Xu et al., 1995).


#### *phyA-205* Is an Independent Allele of *PHYA* with the Same Amino Acid Alteration as *phyA-300D*

It is intriguing that a V631M mutation of *PHYA* was previously reported for another allele, *phyA-205* (Reed et al., 1994). However, at that time *phyA-205* was described as a recessive mutation with an intermediate phenotype. We further examined the segregation property of the *phyA-205* backcross with wild type and our results clearly indicated that it is a strong dominant mutation (Table II). Furthermore, *phyA-205* failed to complement *phyA-300D* in both the F<sub>1</sub> and F<sub>2</sub> populations (Table II), as expected from their allelism. As shown in Figure 3A, seedlings heterozygous for the *phyA-205* mutation, showed long hypocotyls under FRc similar to that of homozygous mutant seedlings (Fig. 3A). Further sequence analysis of the *phyA-205* mutant confirmed that it contains the reported single G-to-A mutation in codon 631 (Reed et al., 1994; Fig. 2C). Thus, there are two independent mutations, *phyA-300D* and *phyA-205*, both resulting in the same V631M mutation in the PHYA protein (Fig. 2C). This result is consistent with the conclusion that the V631M mutation caused a dominant-negative phenotype for *phyA*.

Because the original characterization of *phyA-205* used growth media (GM) supplemented with 2% (w/v) Suc (Reed et al., 1994) and our studies used 0.3% (w/v) Suc, we examined the effect of Suc concentration on hypocotyl length in the *phyA-205* mutant. As shown in Figure 3, the high concentration of Suc indeed inhibited hypocotyl elongation under



**Figure 2.** The structural features of *phyA* and the molecular lesion of the *phyA-300D* mutation. A, Diagram of the Arabidopsis PHYA protein. The chromophore-binding site (C-323), core region, and its two PAS motifs are indicated. B, The conserved Val and its surrounding amino acid residues among four Arabidopsis phytochromes. Note that the Val is invariant among all known *phyA* species from higher plants, although it can be substituted by Ala in all other phytochromes among different higher plant species. The single-letter codes of amino acids are shown in bold. C, Nucleotide sequence comparison of the *PHYA* gene in *phyA-205*, *phyA-300D*, and wild type (WT) around the Val-631 region. Amino acids (in bold single-letter code) are shown above their codons. In both mutant alleles, there is a single identical amino acid residue mutation (V631M) in the translated proteins.




**Figure 3.** The *phyA-205* has a dominant-negative phenotype and is not dependent on Suc concentration. Average hypocotyl lengths (in millimeters) are shown above each seedling ( $n = 25$ ) with the SD shown in parentheses. A, Phenotypic comparison of 3-d-old FR light-grown *phyA-205* homozygous (-/-), heterozygous (-/+), and wild-type (WT) seedlings on 0.3% (w/v) Suc medium. B, Phenotypic comparison of 3-d-old FR light-grown *phyA-205* homozygous (-/-), heterozygous (-/+), and wild-type (WT) seedlings on 2% (w/v) Suc medium.

FRc. However, this inhibition was observed in both *phyA-205* homozygous and heterozygous seedlings. At either concentration of Suc, the heterozygous and homozygous mutant seedlings have similar hypocotyl lengths under the same FRc growth condition. *phyA-300D* homozygous and heterozygous mutant seedlings similarly also displayed this Suc inhibition of hypocotyl elongation (data not shown). This effect of Suc was apparently not unique to the dominant mutations, because both *phyA-1* and *phyA-301* (recessive mutations) displayed a similar decreased hypocotyl length under the higher Suc concentration.

### The PHYA-V631M Is Sufficient to Confer a Dominant-Negative Interference to *phyA*-Mediated FRc Inhibition of Hypocotyl Elongation

To test whether a V631M mutation in PHYA is sufficient to confer a FR-specific dominant-negative phenotype, we introduced a 35S promoter-driven *PHYA-V631M* and wild-type *PHYA* into wild-type Arabidopsis and a null mutant (*phyA-101*; Fig. 4A). When grown in FRc, transgenic seedlings expressing *PHYA-V631M* (from the *phyA-300D* allele) exhibited an etiolated phenotype (Fig. 4B). When grown complete darkness, there is no observable difference between transgenic and wild-type seedlings (Fig. 4C). However, overexpression of *PHYA-V631M* in a *phyA* null mutant did not rescue its mutant phenotype (Fig. 4D), further confirming the loss-of-function nature of the PHYA-V631M protein. Thus, our result implies that not only did *phyA-V631M* lose its ability to respond to FR light, but that the presence of




**Figure 4.** Expression of PHYA-V631M mutant protein is sufficient to cause a dominant interfering effect on endogenous *phyA* signaling. A, Diagrams of the transformation cassettes. The wild-type *PHYA* (WT) and *PHYA-V631M* full-length coding regions were driven by the cauliflower mosaic virus 35S promoter. B, Morphological comparison of 3-d-old FR light-grown seedlings of wild-type *PHYA* overexpressor (*PHYA* OE), wild-type (WT), and two mutant *PHYA-V631M* overexpressor (*PHYA-300D* OE) lines. *PHYA* protein immunoblots are shown in the bottom panel. C, Morphological comparison of 3-d-old dark-grown seedlings with *PHYA-V631M* overexpressor transgene in wild-type background (left) and the wild-type control (right). D, Morphology of 3-d-old FR light-grown seedling with *PHYA-V631M* overexpressor transgene in a *phyA* null mutant background. Scale bars in B through D = 1 mm.

PHYA-V631M is sufficient to interfere with endogenous wild-type *phyA* function in a dominant-negative fashion.

Interestingly, the degree of dominant-negative interference of the transgenic plants depends on mutant PHYA-V631M protein dose. As shown in Figure 4B, a high-expression level of mutant PHYA-V631M resulted in seedlings with an extremely elongated hypocotyl and closed cotyledons, resembling that of the *phyA* null mutants, whereas a lower expression level resulted in an intermediate phenotype, a long hypocotyl, and open but small cotyledons (Fig. 4B). As expected, expression of wild-type PHYA resulted in seedlings with short hypocotyls and open cotyledons (Fig. 4B), similar to the reported *phyA* overexpressor phenotype (Boylan et al., 1994).

#### The *phyA-V631M* Abundance Is Not Reduced in Mutant Seedlings Grown in Darkness But Is Slightly Higher under FR Light

As a first step to determine a possible molecular basis of the *phyA-V631M* mutation effect, the *phyA* protein levels of the *phyA-300D* and wild-type seedlings grown in darkness and in FRc were examined using immunoblot analysis. As shown in Figure 5, the level of *phyA* protein in dark-grown *phyA-300D* seedlings is as abundant as in wild-type seedlings. Interestingly, *phyA-300D* grown under FRc accumulated to about 2-fold higher levels of *phyA* protein than that of wild type. This result rules out the possibility that the *phyA-300D* phenotype is a result of lower PHYA levels under FR light. It should be noted that the dominant mutant phenotype of *phyA-300D* could be a result of increased levels of the mutant



**Figure 5.** Characterization of wild-type and PHYA-V631M stability under R and FR light. A, *phyA* levels in 3-d-old dark (D) and FR light-grown wild-type Columbia (WT) and *phyA-300D*, and in 3-d-old dark-grown seedlings exposed to Rc for 1 or 4 h (hr; top). The immunoblots were probed with a *phyA*-specific polyclonal antibody. An equal amount of total protein was loaded in each lane, as verified by CSN5 immunoblot (bottom). B, *phyA* levels in 3-d-old dark-grown (D) wild-type Columbia (WT), and *phyA-300D* seedlings, and in 3-d-old dark-grown seedlings exposed to FRc light for 1, 4, or 24 h (hr). An equal amount of total protein was loaded in each lane, as verified by immunoblot using anti-tubulin antibodies (bottom).

PHYA-V631M protein. However, this hypothesis would be argued against by the observation that the *phyA-105* mutant, which has a single amino acid change, accumulated to similar high levels of the PHYA-V631M mutant protein in FRc, and still exhibited completely recessive properties (Xu et al., 1995).

#### *phyA-V631M* Possesses Normal Light Triggered Degradation Kinetics and Photochemical Properties

It is known that *phyA* accumulates in darkness as the Pr form and is rapidly degraded upon exposure to light, because of the susceptible nature of the *phyA* Pfr form to proteasome-mediated protein degradation. Therefore, we examined the degradation kinetics of the *phyA-V631M* mutant protein compared with wild-type *phyA* protein upon exposure to continuous R light (Rc) or FRc. Dark-grown mutant and wild-type seedlings were exposed to Rc for 0, 1, and 4 h and harvested for immunoblot analysis. The degradation kinetics shown in Figure 5A indicates that there is no significant difference between the *phyA-V631M* mutant and wild-type *phyA* proteins. As expected, FRc treatment of dark-grown mutant or wild-type seedlings does not cause observable degradation of the *phyA* proteins (Fig. 5B). This result suggests that the mutant *phyA-V631M* protein was able to photoconvert from Pr to Pfr upon exposure to Rc, and only the Pfr form of *phyA-V631M* was labile and subjected to rapid degradation.


To test whether *phyA-V631M* protein retains typical *phyA* R-FR reversibility, the differential absorbance spectra for *phyA-300D* and wild-type seedlings were examined. To this end, *phyA-300D*, *phyA-101* (a *phyA* null allele), and wild-type seedlings were grown in darkness for 3 d and extracts were prepared under green safelight as described in "Materials and Methods." This assay revealed that *phyA-V631M* has an essentially identical spectral property as wild type (data not shown).

#### The PHYA-V631M Protein Retains Dimerization Capability


The Val residue at position 631 is located in the first PAS motif of PHYA that overlaps with the *phyA* dimerization region. Thus, we examined the ability of PHYA-V631M to form dimers *in vivo*. To this end, crude extracts from homozygous mutant and wild-type seedlings grown in darkness or FRc were subjected to native gel electrophoresis followed by immunoblot analysis (Fig. 6). The migration and amount of PHYA-V631M dimer from *phyA-300D* seedlings under native gel conditions are comparable with that of wild-type plants, indicating that the PHYA-V631M can effectively dimerize *in vivo*.

### Overexpression of *PHYB-V664M* in Wild-Type Plants Failed to Interfere with *phyB* Signaling

The *PHYA* and *PHYB* polypeptides are highly homologous (Sharrock and Quail, 1989). The region surrounding the Val-631 of *PHYA* and the corresponding region in *PHYB* is highly conserved (Fig. 2B). For *PHYB*, its Val residue at position 664 is equivalent to the Val at position 631 of *PHYA*. Thus, it is of interest to determine whether a corresponding *PHYB-V664M* mutant protein would confer a similar dominant-negative effect on *phyB* function. For this reason, we specifically created a corresponding *PHYB-V664M* mutant transgene under 35S promoter control. The transgene was transformed into wild-type *Arabidopsis* and a *phyB* null mutant (*phyB-101*; Fig. 7A). Homozygous transgenic lines were obtained from each construct, and their hypocotyl elongation under Rc was examined. Contrasting to *PHYA-V631M*, the *PHYB-V664M* mutant protein is clearly functional and active in both the *phyB* null mutant and wild-type seedlings in the lines tested. As shown in Figure 7B, *PHYB-V664M* can rescue the *phyB* null phenotype when overexpressed in the *phyB-101* mutant background. Furthermore, transgenic seedlings overexpressing the *PHYB-V664M* in a wild-type background exhibited a hyper-photomorphogenic phenotype with shorter hypocotyls than those of wild type (Fig. 7C), similar to seedlings overexpressing wild-type *PHYB* (Fig. 7A). Several *PHYB-V664M* transgenic lines that showed longer hypocotyl under Rc turned out to be cosuppression lines and had an undetectable level of *phyB* protein. This result indicates that *PHYB-V664M* behaves differently from *PHYA-V631M*. Our results imply that the mutant *PHYB-V664M* protein neither reduces its ability to respond to Rc, nor is capable of interfering with wild-type *phyB* signaling. Thus, the role of Val-631 in *phyA* defined in this work is specific to *phyA* and not applicable to *phyB*.



**Figure 6.** Native gel analysis of *phyA* dimerization in FR and dark-grown wild-type and mutant seedlings. Extracts were separated on a 4% to 20% (w/v) gradient non-denaturing gel and probed with a polyclonal *PHYA* antibody. The *phyA* dimer is marked by a triangle on the right. Equal loading was ensured by visualizing a protein band staining intensity (marked by control). The positions of two mass markers are indicated on the left side. The *phyA-101* is a null mutant control. A, FR light-grown seedlings (5 d); B, dark-grown seedlings (5 d).



**Figure 7.** The *PHYB-V664M* protein is physiologically active and does not confer an interfering effect on *phyB* signaling. The 3-d-old seedlings are shown on top, and their corresponding *PHYB* protein immunoblots are shown on the bottom. A, Three-day-old Rc light-grown wild type (WT), *PHYB* overexpressor (*PHYB* OE) in wild-type background, and the *phyB-101* null mutant. B, Three-day-old Rc light-grown seedlings of three independent *PHYB-V664M* overexpressor lines (L1, L2, and L3 of *PHYB-V664M* OE) in the *phyB-101* null mutant background. C, Three-day-old Rc light-grown seedlings of two different *PHYB V664M* overexpressor (*PHYB-V664M* OE) lines in a wild-type background. L'1 is an overexpressor line for *PHYB-V664M*, and L'2 is a cosuppression line.

## DISCUSSION

### *PHYA-V631M* Has a Strong Dominant-Negative Interfering Activity

Several lines of evidence presented in this report definitively show that *PHYA-V631M* is capable of conferring a strong dominant-negative interfering effect on normal *phyA* signaling. First, two independent alleles with the same V631M amino acid change exhibited a strong long hypocotyl phenotype under FR in the heterozygous state. The hypocotyl length of heterozygous mutant seedlings is nearly identical to that of their homozygous mutant seedlings. The only distinction between the heterozygous and homozygous seedlings is the slightly more advanced cotyledon development in the heterozygous mutants. The heterozygous seedlings exhibit opening and slight enlargement of the cotyledons after extended growth under FRc. Second, introduction of the *PHYA-V631M* transgene into wild-type *Arabidopsis* confirmed that *PHYA-V631M* is sufficient to confer the dominant-negative interfering effect. Our results imply that not only does the mutant *PHYA-V631M* protein reduce its ability to respond to FRc, it is also able to interfere with the ability of endogenous *phyA* to mediate FRc responses in a dominant-negative fashion. The *PHYA-V631M* is clearly photochemically active but not functional in mediating FRc responses, because its homozygous seedlings exhibited a phenotype essentially indistinguishable from that of the *phyA* null mutations. Furthermore, overexpression of *PHYA-V631M* in a *phyA* null mutant background also failed to show any phenotypic rescue (Fig. 4D).

Although a large number of *phyA* mutations within the two PAS motifs have been described, all of

them were reported as recessive (Reed et al., 1994; Xu et al., 1995). In fact, only one report described a slightly weak semidominant effect of *phyA* mutations under low-FR photon fluence conditions and with no sugar in the GM (Whitelam et al., 1993). In this case, heterozygous *PHYA/phyA* seedlings for two alleles, at least one is a null allele, exhibited slightly longer hypocotyls (3 versus 1 mm in wild type) under FRC but were still significantly shorter than those of *phyA* homozygous mutants (12–13 mm) under the same growth condition. However, this weak semidominant effect of *phyA* mutations is largely growth-condition dependent and was not observed by other researchers using different growth conditions (Reed et al., 1994; Xu et al., 1995). We also failed to observe a semidominant effect of the same null *phyA-1* allele (Whitelam et al., 1993) under our growth conditions, whereas the dominant effect observed in *PHYA-V631M* is very strong and consistent under our experimental conditions.

On the other hand, a dominant interfering effect has been reported for transgenic seedlings overexpressing several heterologous *phyA* species. For example, overexpression of three distinct oat (*Avena sativa*) *phyA* mutant forms (major deletions in very N- or C-terminal half) in *Arabidopsis* resulted in a strong dominant-negative interfering effect over endogenous *Arabidopsis phyA* function (Boylan et al., 1994). Overexpression of an amino-terminal deletion of rice (*Oryza sativa*) *phyA* in tobacco seedlings also resulted in a dominant interfering effect (Emmler et al., 1995). Last, overexpression of an intact fern *phyA* in *Arabidopsis* also caused a weak dominant interfering effect (Okamoto et al., 1997). A general explanation for those dominant interfering effects is that those mutant *phyA* molecules (oat or rice) or the too-diverged fern *phyA* may not be able to mediate a productive interaction with *phyA*-signaling partners, and thus, titrate out the available *phyA*-signaling partners. Nevertheless, comparing the degrees of the dominant-negative interfering phenotypes in those overexpressing lines, the *phyA-V631M* heterozygous mutant seedlings exhibited possibly the strongest interfering phenotype. Furthermore, this strong dominant-negative interfering phenotype in *phyA-300D* and *phyA-205* is caused by a single amino acid substitution.

### The Role of the PAS-Like Domain in *phyA*-Signaling Activity

Molecular and biochemical characterization of *phyA* indicates that the apoprotein folds into two major structural domains: a globular NH<sub>2</sub>-terminal domain cradling the covalently attached chromophore in a hydrophobic pocket, and a more extended C-terminal domain, with a short flexible hinge region connecting the two (Quail, 1997; Park et al., 2000). Evidence also indicates that the determi-

nants of the photosensory specificity of *phyA* and *phyB* to FRC and Rc reside in the NH<sub>2</sub>-terminal domains. In addition, the proximal end of the C-terminal domain, designated the core region or Quail box of the polypeptide, contains determinants necessary for regulatory activity or for the effective communication of perceived light signals to the cellular transduction circuitry. It is generally assumed that the phytochrome molecules, rather than directly binding to target gene promoters, relay information via signaling intermediates or directly to DNA-binding transcription factors (Quail, 2002; Wang and Deng, 2002).

The V631M mutation in *phyA-300D* and *phyA-205* falls within the “core region” of the PHYA protein (Fig. 2A). This structural domain has shown a propensity for light-signaling mutations (Quail, 1997). The severity of the homozygous *phyA-300D* mutant phenotype reveals a critical role of the Val residue within this core region for *phyA* function. This region of the PHYA protein has been shown to be a “hot spot” for missense mutations resulting in regulatory mutants that produce PHYA protein, but cannot transduce the light signal to elicit the photomorphogenic response. However, *phyA-V631M* is the first case where a strong dominant-negative interfering phenotype has been described, whereas other mutations in this region have been reported as recessive (Reed et al., 1994; Xu et al., 1995).

This core region of PHYA contains two PAS-like motifs and may be involved in phytochrome signal transduction by contacting downstream partners (Fig. 2A; Ni et al., 1998). Previous studies have shown that this core region of phytochrome interacts directly with phytochrome partners including NDPK2 (Choi et al., 1999) and PIF3 (Ni et al., 1998), supporting the conclusion that the PAS motifs or the core region defines protein-protein interactive surfaces. One possible explanation of the negative interference of the PHYA-V631M on *phyA* signaling is that the *phyA-V631M* mutant protein interacts too strongly with downstream positively acting components of the pathway. These stronger than wild-type interactions with downstream components for PHYA-V631M would compromise the capability of the activated target protein to initiate downstream events. In heterozygous plants, the PHYA-V631M protein would titrate out the downstream *phyA* target because of the higher affinity between PHYA-V631M and the target, thus, effectively preventing wild-type *phyA* from interacting with the target and blocking the proper *phyA*-mediated event. However, this is certainly not the only explanation, and there are other alternative possibilities. One alternative is that PHYA-V631M could have a stronger affinity to form a functionally inactive heterodimer with the wild-type PHYA. In heterozygous plants, this higher affinity to form nonfunctional PHYA-V631M:PHYA heterodimers could prevent the formation of active

wild-type *phyA* homodimers. As a result, the heterozygous plants would fail to respond to FRc. A third possibility is that the PHYA-V631M homodimer and its heterodimer with wild-type PHYA might have an elevated Pfr to Pf form dark reversion rate, which could potentially account for the interfering phenotype. An important task of future work will be to determine which mode of action is used in plants.

### The *phyA*-V631M Dominant-Negative Effect Is Not Applicable to *phyB*

Although there is a high-sequence homology between the Arabidopsis PHYA and PHYB apoproteins surrounding PHYA amino acid Val-631 (Fig. 2A), we demonstrated that the corresponding Val to Met mutation (V664M) in Arabidopsis *phyB* does not trigger a similar dominant-negative effect on *phyB* signaling. The PHYB-V664M mutant protein was active in both *phyB* null mutant and wild-type plants when introduced into the proper backgrounds. These results indicate that Val-631 is critical for PHYA-mediated FR signaling but not applicable to PHYB-mediated signaling. This specificity to *phyA* is also consistent with the observation that all known *phyA* proteins, including those from both monocots and dicots, have the invariant Val at the corresponding position (Xu et al., 1995). However, the Val residue is not absolutely conserved in other higher plant phytochromes, and in some cases, the corresponding position has changed to Ala (Xu et al., 1995). This observation is consistent with the conclusion that the role of Val-631 in *phyA* as defined by the V631M mutation is unique to *phyA*, and not applicable to *phyB*. Thus, it defines a key distinction in the manner through which *phyA* and *phyB* transduce light signal via their C-terminal core regions.

## MATERIALS AND METHODS

### Plant Materials, Mutant Screen, and Mapping

Arabidopsis mutants *phyA-1*, *phyA-101*, *phyA-205*, and *phyB-101* (Landsberg *erecta* ecotype) were described previously (Whitelam et al., 1993; Reed et al., 1994). Several available T-DNA insertion mutagenized collections were used, including those from the Arabidopsis Stock Center and Dr. Steven Dellaporta (Yale University; Galbiati et al., 2000). Either T<sub>2</sub> or T<sub>3</sub> seeds were screened for mutant seedlings exhibiting a complete (long hypocotyl and closed cotyledons) or partial etiolated (long hypocotyl and open cotyledons) phenotype after 3 d of growth under FRc. Seed sterilization and plating were essentially the same as previously described (Hsieh et al., 2000) except the GM was supplemented with 0.3% (w/v) Suc. To improve germination rate, the seeds on the plates were exposed to 24 h of white light after vernalization for 2 d at 4°C in the dark. The light conditions used were: Rc (100  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ), continuous blue light (50  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ), continuous white light (200  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ), and FRc (175  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ). Growth chambers (E-30 LED 2/3, Percival Scientific, Perry, IA) were maintained at 22°C constant temperature.

For mapping, mutant *phyA-300D* (Columbia ecotype) was crossed to wild-type Landsberg *erecta* ecotype. F<sub>2</sub> progenies were grown on 0.3% (w/v) GM plates and screened under FR for a short hypocotyl, whereas F<sub>3</sub> progeny were grown on 0.3% (w/v) GM plates and screened under FR light for a long hypocotyl. DNA from individual homozygous mutant seedlings was

prepared and used for PCR-based simple sequence length polymorphism or cleaved-amplified polymorphic sequence mapping (Bell and Ecker, 1994; Konieczny and Ausubel, 1993).

### Native Protein Gel and Immunoblot Analyses

Five-day-old FR or dark-grown seedlings were collected under green safelight for protein analysis. Seedlings were frozen in liquid nitrogen and ground with a mortar and pestle in 100  $\mu\text{L}$  of grinding buffer (10% [w/v] glycerol, 400 mM Suc, 50 mM Tris, pH 8.0, and 1 mM phenylmethylsulfonyl fluoride). For native gel analysis, 20  $\mu\text{g}$  of each sample was run on a 4% to 20% (w/v) non-denaturing gradient gel (Tris-Gly, pH 8.0). After electrophoresis, proteins were transferred to Immobilon-P membrane (Millipore, Bedford, MA) for immunoblot analysis. For the PHYA protein level, protein was extracted from 3-d-old dark-grown seedlings and from dark-grown seedlings subjected to Rc or FRc for the specified period of time. Protein concentration was determined by Bradford assay (Bio-Rad, Hercules, CA), loaded equally to SDS-PAGE, and transferred to nylon membrane. The protein-blot analysis was performed according to procedures described (Okamoto et al., 2001). Type-selected antibodies for *phyA* (Xu et al., 1995) and *phyB* (Ni et al., 1998) were used.

### Phytochrome Difference Spectrum

Arabidopsis protein extracts were prepared from 6-d-old dark-grown seedlings as described (Weller et al., 1996). The extract was dispensed into a spectrophotometer ultra-microcuvette and placed in a FR-LED chamber (200  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) for 360 s to induce photoconversion of phytochrome to the Pr form. After this exposure, the cuvette was placed into the chamber of a spectrophotometer (DU-70, Beckman Coulter, Fullerton, CA), and the baseline was taken. The cuvette was then exposed to R light for 180 s. R light was generated by using a light projector (Ektagraphics III AMT, Eastman Kodak, Rochester, NY) with a Kodak extra bright lamp module. The light passed through a Plexiglas filter (CBS Red 650, Westlake Plastics Co., Lenni, PA) and was reflected into the chamber of the spectrophotometer using a mirror. After R light exposure, an absorbance versus wavelength plot was generated, with readings taken every nanometer from 500 to 800 nm.

### DNA Sequencing of the *PHYA* Gene

Genomic DNA was isolated from *phyA-300D* and *phyA-205* mutants. A series of primers was synthesized based on the published *PHYA* sequence (Dehesh et al., 1994), and PCR was used to amplify the segments of the *PHYA* gene from the *phyA-300D* and *phyA-205* genomic DNA. PCR products were purified using a PCR purification kit (Qiagen USA, Valencia, CA) and sequenced.

### Construction of Full-Length *PHYA* and *PHYB* Overexpressing Plasmids and Arabidopsis Transformation

Total RNA was extracted from 6-d-old dark-grown seedlings of either wild type or the *phyA-300D* mutant using Trizol reagent (Invitrogen, Carlsbad, CA). cDNA was synthesized using a reverse transcriptase kit (Advantage RT for PCR kit, CLONTECH Laboratories, Palo Alto, CA). The entire coding region of *PHYA* was divided into two fragments for PCR amplification. The first fragment used a primer, 5'-GCG TCG ACA TGT CAG GCT CTA GGC CGA C-3', which covers the translation start codon of *PHYA* with an incorporated *SalI* cleavage site, and the second primer was 5'-AAT TTT GAG ATC ATT TAG CTT CG-3'. The second fragment used a primer covering the stop codon of *PHYA* with an incorporated *SmaI* cleavage site 5'-CGC CCG GGC TAC TTG TTT GCT GCA GCG-3', and the second primer was 5'-GAA TAC CAC ATG GAT TCA ACG-3'. The two PCR fragments were cloned into the pCR 2.1 TOPO cloning vector (Invitrogen). A unique internal enzyme site (Bst 1107) and a *XhoI* site within the TOPO vector were used to ligate the two PCR fragments to construct the full-length coding region for either the mutant or wild-type *PHYA* gene. Using the incorporated *SalI* site and a *SacI* site within the TOPO cloning vector, the full-length *PHYA* gene was ligated into the binary vector pZPY122 (Yamamoto et al., 1998), and the sequence was confirmed.



The wild-type full-coding region of *PHYB* was generated using a similar RT-PCR method as for *PHYA* above except that it used one primer set. The forward primer (5'-GCC CCG GGA TGG TTT CCG GAG TCG GG-3') was designed over the translation start codon of *PHYB* with an incorporated *SmaI* cleavage site. The reverse primer (5'-CGT CTA GAA CTA ATA TGG CAT CAT CAG CAT C-3') was designed over the stop codon of *PHYB* with an incorporated *XbaI* site. The generated PCR product was cloned into the pCR 2.1 TOPO cloning vector (Invitrogen) and sequenced. This wild-type *PHYB* cDNA clone was used as the template for generating the mutant *PHYB* construct via PCR that specifically incorporates an A-to-G nucleotide on codon 664 to convert Val to Met. As a result, a full-length *PHYB-V664M* mutant coding region was produced. Using the incorporated *SmaI* site and a *KpnI* site within the TOPO cloning vector, the full-length *phyB* genes were ligated into the binary vector pZPY122 (using the filled-in *XbaI* and *KpnI* sites within the vector; Yamamoto et al., 1998).

All *PHYA* and *PHYB* binary constructs were electroporated into the *Agrobacterium tumefaciens* strain GV3101 (MP90) and transformed into the proper *Arabidopsis* strains using the floral dip method (Clough and Bent, 1998).

## ACKNOWLEDGMENTS

We thank Haiyang Wang for reading and commenting on the manuscript and Dr. Peter Quail for *PHYA* and *PHYB* antibodies. We thank Steve Dellaporta for making his T-DNA tagged *Arabidopsis* collection available for the mutant screen. We are extremely grateful to Haiyang Wang for his help in *PHYA* cloning and to Matthew J. Terry for his guidance involving the difference spectra experiments.

Received March 6, 2002; returned for revision April 18, 2002; accepted May 9, 2002.

## LITERATURE CITED

- Bell CJ, Ecker JR (1994) Assignment of 30 microsatellite loci to the linkage map of *Arabidopsis*. *Genomics* **19**: 137–144
- Bowler C, Neuhaus G, Yamagata H, Chua NH (1994) Cyclic CMP and calcium mediate phytochrome phototransduction. *Cell* **77**: 73–81
- Boylan MT, Douglas N, Quail PH (1994) Dominant negative suppression of *Arabidopsis* photoresponses by mutant phytochrome A sequences identify spatially discrete regulatory domains in the photoreceptor. *Plant Cell* **6**: 449–460
- Choi G, Yi H, Lee J, Kwon Y, Soh M, Shin B, Luka Z, Hahn T, Song PS (1999) Phytochrome signaling is mediated through nucleoside diphosphate kinase 2. *Nature* **401**: 610–613
- Clack T, Mathews S, Sharrock RA (1994) The phytochrome apoprotein family in *Arabidopsis* is encoded by five genes: the sequences and expression of phyD and phyE. *Plant Mol Biol* **25**: 413–427
- Clough S, Bent A (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**: 735–743
- Dehesh K, Franci C, Sharrock RA, Somers DE, Welsch JA, Quail PH (1994) The *Arabidopsis* phytochrome A gene has multiple transcription start sites and a promoter sequence motif homologous to the repressor element of monocot phytochrome A genes. *Photochem Photobiol* **59**: 379–384
- Emmler K, Stockhaus J, Chua NH, Schafer E (1995) An amino-terminal deletion of rice phytochrome A results in a dominant negative suppression of tobacco phytochrome A activity in transgenic tobacco seedlings. *Planta* **197**: 103–110
- Galbiati M, Moreno MA, Nadzan G, Zourelidou M, Dellaporta SL (2000) Large-scale T-DNA mutagenesis in *Arabidopsis* for functional genomic analysis. *Funct Integr Genomics* **1**: 25–34
- Hsieh HL, Okamoto H, Wang M, Ang LH, Matsui M, Goodman H, Deng XW (2000) FIN219, an auxin-regulated gene, defines a link between phytochrome A and the downstream regulator COP1 in light control of *Arabidopsis* development. *Genes Dev* **14**: 1958–1970
- Jordan ET, Cherry JR, Walker JM, Vierstra RD (1996) The amino-terminus of phytochrome A contains two distinct functional domains. *Plant J* **9**: 243–257
- Kendrick RE, Kronenberg GHM, eds (1994) *Photomorphogenesis in Plants*, Ed 2. Kluwer Academic Publishers, Dordrecht, The Netherlands
- Kircher S, Kozma-Bognar L, Kim L, Adam E, Harter K, Schafer E, Nagy F (1999) Light quality-dependent nuclear import of the plant photoreceptors phytochrome A and B. *Plant Cell* **11**: 1445–1456
- Konieczny A, Ausubel FM (1993) A procedure for mapping *Arabidopsis* mutations using co-dominant ecotype-specific PCR-based markers. *Plant J* **4**: 403–410
- McNellis TW, Deng XW (1995) Light control of seedling morphogenetic pattern. *Plant Cell* **7**: 1749–1761
- Ni M, Tepperman JM, Quail PH (1998) PIF3, a phytochrome-interacting factor necessary for normal photoinduced signal transduction, is a novel basic helix-loop-helix protein. *Cell* **95**: 657–667
- Okamoto H, Matsui M, Deng XW (2001) Overexpression of the heterotrimeric G-protein alpha-subunit enhances phytochrome-mediated inhibition of hypocotyl elongation in *Arabidopsis*. *Plant Cell* **13**: 1639–1652
- Okamoto H, Sakamoto K, Tomizawa KI, Nagatani A, Wada M (1997) Photoresponses of transgenic *Arabidopsis* overexpressing the fern *Adiantum capillus-veneris* PHY1. *Plant Physiol* **115**: 79–85
- Park CM, Bhoo SH, Song PS (2000) Inter-domain crosstalk in the phytochrome molecules. *Semin Cell Dev Biol* **11**: 449–456
- Quail PH (1997) An emerging molecular map of the phytochromes. *Plant Cell Environ* **20**: 657–665
- Quail PH (2002) Phytochrome photosensory signalling networks. *Nat Rev Mol Cell Biol* **3**: 85–93
- Quail PH, Boylan MT, Parks BM, Short TW, Xu Y, Wagner D (1995) Phytochromes: photosensory perception and signal transduction. *Science* **268**: 675–680
- Reed JW, Nagatani A, Elich TD, Fagan M, Chory J (1994) Phytochrome A and phytochrome B have overlapping but distinct functions in *Arabidopsis* development. *Plant Physiol* **104**: 1139–1149
- Sharrock RA, Quail PH (1989) Novel phytochrome sequences in *Arabidopsis thaliana*: structure, evolution, and differential expression of a plant regulatory photoreceptor family. *Genes Dev* **3**: 1745–1757
- Shinomura T, Uchida K, Furuya M (2000) Elementary processes of photo-perception by phytochrome A for high-irradiance response of hypocotyl elongation in *Arabidopsis*. *Plant Physiol* **122**: 147–156
- Wang H, Deng XW (2002) Phytochrome signaling. In CR Somerville and EM Meyerowitz, eds, *The Arabidopsis Book*. American Society of Plant Biologists, Rockville, MD. doi/10.1199/tab.0074. <http://www.aspb.org/publications/arabidopsis>
- Weller JL, Terry MJ, Rameau C, Reid JB, Kendrick RE (1996) The phytochrome-deficient *pcd1* mutant of pea is unable to convert heme to biliverdin IXa. *Plant Cell* **8**: 55–67
- Whitelam GC, Johnson E, Peng J, Carol P, Anderson ML, Cowl JS, Harberd NP (1993) Phytochrome A null mutants of *Arabidopsis* display a wild type phenotype in white light. *Plant Cell* **5**: 757–768
- Xu Y, Parks BM, Short TW, Quail PH (1995) Missense mutations define a restricted segment in the C-terminal domain of phytochrome A critical to its regulatory activity. *Plant Cell* **7**: 1433–1443
- Yamamoto Y, Matsui M, Ang L, Deng XW (1998) Role of a COP1 interactive protein in mediating light-regulated gene expression in *Arabidopsis*. *Plant Cell* **10**: 1083–1094
- Yeh KC, Lagarias JC (1998) Eukaryotic phytochromes: light regulated serine/threonine protein kinases with histidine kinase ancestry. *Proc Natl Acad Sci* **95**: 13976–13981