

LeCTR1, a Tomato *CTR1*-Like Gene, Demonstrates Ethylene Signaling Ability in Arabidopsis and Novel Expression Patterns in Tomato¹

Julie Leclercq, Lori C. Adams-Phillips, Hicham Zegzouti, Brian Jones, Alain Latché, James J. Giovannoni, Jean-Claude Pech, and Mondher Bouzayen*

Unité Mixte de Recherche 990, Institut National de la Recherche Agronomique/Institut National Polytechnique-Ecole Nationale Supérieure Agronomique, Boite Postale 107 Auzeville, 31326 Castanet Tolosan cedex, France (J.L., H.Z., B.J., A.L., J.-C.P., M.B.); and United States Department of Agriculture-Agricultural Research Service and Boyce Thompson Institute for Plant Research, Tower Road, Ithaca, New York 14853 (L.C.A.-P., J.J.G.)

LeCTR1 was initially isolated by both differential display reverse transcriptase-polymerase chain reaction screening for tomato (*Lycopersicon esculentum*) fruit ethylene-inducible genes and through homology with the Arabidopsis *CTR1* cDNA. *LeCTR1* shares strong nucleotide sequence homology with Arabidopsis *CTR1*, a gene acting downstream of the ethylene receptor and showing similarity to the Raf family of serine/threonine protein kinases. The length of the *LeCTR1* transcribed region from ATG to stop codon (12,000 bp) is more than twice that of Arabidopsis *CTR1* (4,700 bp). Structural analysis reveals perfect conservation of both the number and position of introns and exons in *LeCTR1* and Arabidopsis *CTR1*. The introns in *LeCTR1* are much longer, however. To address whether this structural conservation is indicative of functional conservation of the corresponding proteins, we expressed *LeCTR1* in the Arabidopsis *ctr1-1* (*constitutive triple response 1*) mutant under the direction of the 35S promoter. Our data clearly show that ectopic expression of *LeCTR1* in the Arabidopsis *ctr1-1* mutant can restore normal ethylene signaling. The recovery of normal ethylene sensitivity upon heterologous expression of *LeCTR1* was also confirmed by restored glucose sensitivity absent in the Arabidopsis *ctr1-1* mutant. Expression studies confirm ethylene responsiveness of *LeCTR1* in various tissues, including ripening fruit, and may suggest the evolution of alternate regulatory mechanisms in tomato versus Arabidopsis.

The plant hormone ethylene is involved in a variety of developmental and physiological processes in plants, including senescence, fruit ripening, and abscission (Abeles et al., 1992; Lelièvre et al., 1997; Giovannoni, 2001). It also plays an important role in physiological responses to environmental stresses such as water deficit, mechanical wounding, and pathogen attack (Abeles et al., 1992). The unraveling of the molecular basis of the ethylene perception and signal transduction pathway has been enhanced by the use of Arabidopsis mutants altered in the seedling triple response (Guzmán and Ecker, 1990). The triple response is exhibited by seedlings treated with ethylene and results in: (a) inhibition of root elonga-

tion, (b) shortening and radial swelling of the hypocotyl, and (c) exaggerated curvature of the apical hook. Numerous loci have been identified and many corresponding genes cloned, representing various steps in ethylene signaling from receptors through transcription factors (Ecker, 1995; Johnson and Ecker, 1998; Stepanova and Ecker, 2000). The *ETR1* gene was the first to be cloned (Chang et al., 1993) and was shown to encode a functionally active ethylene receptor (Schaller and Bleecker, 1995). Subsequently, it has been demonstrated that *ETR1* belongs to a multigene family whose five members are differentially regulated (Schaller and Bleecker, 1995; Hua and Meyerowitz, 1998). Despite significant divergence at the structural and primary sequence level, all members of the ethylene receptor family are functionally active (Hua and Meyerowitz, 1998).

In contrast to the ethylene insensitivity phenotype conferred by the *ethylene response* mutation (*etr*), disruption of the *ctr1* locus confers constitutive ethylene response in the absence of the hormone. Epistatic studies revealed that the *CTR1* gene product acts downstream of the ethylene receptors (Kieber et al., 1993). Furthermore, the regulatory domain of *CTR1* was found to associate with *ETR1* and ethylene response sensor (*ERS1*) in yeast two-hybrid and in vitro

¹ This work was supported in part by the European Union (grant no. FAIR CT-95 0225), by the Institut National de la Recherche Agronomique (Action Transversale Structurante Tomate, grant no. 2001-2003), by the "Midi-Pyrénées" Regional Council (grant nos. 99009080 and 01002710 to M.B.), and by the National Science Foundation (grant nos. IBN-9604115 and DBI-9872617 to J.J.G.). This research forms part of the requirement for the degree of PhD for J.L.

* Corresponding author; e-mail bouzayen@ensat.fr; fax 33-5-62-19-35-73.

Article, publication date, and citation information can be found at www.plantphysiol.org/cgi/doi/10.1104/pp.009415.

protein association assays (Clark et al., 1998), raising the possibility that ethylene receptors directly regulate CTR1 activity.

In the tomato (*Lycopersicon esculentum*), initial inroads into ethylene perception were made via cloning of the *Never-ripe* gene, which proved to be a tomato ethylene receptor most like the ERS receptor of Arabidopsis (Hua et al., 1995; Wilkinson et al., 1995). Subsequent studies regarding ethylene perception and signal transduction have focused on the isolation and characterization of the receptor gene family and a family of putative transcription factors related to the Arabidopsis *EIN3* gene. In recent years, five *ETR1* homologs have been identified in tomato (Payton et al., 1996; Zhou et al., 1996; Tieman and Klee, 1999) and heterologous expression and complementation studies have been employed on a subset of *ETR* gene family members to demonstrate ethylene receptor activity (Wilkinson et al., 1995; Tieman et al., 2000).

Although two tomato sequences showing significant sequence homology with Arabidopsis *CTR1* have been reported, data addressing their functional significance are lacking. The Arabidopsis *CTR1* gene is reported to be constitutively expressed (Kieber et al., 1993). Constitutive expression was not the case for the tomato *LeCTR1* gene that we have shown previously to be regulated by ethylene and during fruit ripening (Giovannoni et al., 1998; Zegzouti et al., 1999). Isolation of *LeCTR1* and its regulation by ethylene and induction during ripening suggested the possibility that this step in the ethylene-signaling network may be a target for differential regulation in species displaying aspects of development critically dependent upon ethylene. Before addressing this question, however, it was first necessary to establish *LeCTR1* function.

Here, we show that the genomic structure of *LeCTR1* is highly conserved with the Arabidopsis *CTR1* gene and is capable of furnishing CTR1 function when expressed in the *ctr1-1* mutant of Arabidopsis. We also demonstrate that *LeCTR1* mRNA accumulates during fruit ripening and upon ethylene treatment not only in fruit but also in additional non-fruit tissues. Together, these results may suggest regulatory modification of a necessary component of ethylene signal transduction in tomato as compared with Arabidopsis.

RESULTS AND DISCUSSION

LeCTR1 Encodes a Putative Raf Kinase Protein

The ethylene-inducible *LeCTR1* differential display-derived fragment, initially called *ER50* (Zegzouti et al., 1999), was extended by primer extension to obtain a full coding sequence of the cDNA clone. Accurate *LeCTR1* sequence was confirmed by RACE-PCR of a partial cDNA isolated by screening a ripe fruit cDNA library with the Arabidopsis *CTR1* cDNA as probe, and followed by reverse transcriptase (RT)-PCR and sequencing of the resulting full-length cDNA (Giovannoni et al., 1998). Translation of the longest open


reading frame of the resulting cDNA sequence predicts a protein with a molecular mass of 92 kD and no obvious membrane-spanning domains.

The predicted *LeCTR1* protein shares significant homology with different members of the Raf family of Ser/Thr protein kinases of the class mitogen-activated protein kinase kinase kinase from both animals (e.g. *Rattus norvegicus* C-Raf-1) and plants. Database searches revealed that *LeCTR1* shows the strongest homology with *AtCTR1*, a negative regulator of ethylene signaling in Arabidopsis (Kieber et al., 1993). A similar level of homology was also found with other plant Raf kinases, like the *AtEDR1* (*Enhanced Disease Resistance 1*), an Arabidopsis gene shown to act as a negative regulator of defense responses (Frye et al., 2001) and *LeCTR2*, another tomato CTR-like protein (*LeCTR2/TCTR2*, GenBank accession no. AJ005077). Within the kinase domain of *LeCTR1*, *AtCTR1*, *AtEDR1*, and *LeCTR2*, there is perfect conservation of 11 subdomains typical of the catalytic site of Ser/Thr protein kinases (Hanks et al., 1988; Hanks and Quinn, 1991). These domains include two signature patterns. The first, spanning amino acids 562 to 569, corresponds to a typical ATP-binding site motif (GxGxxGxV; Schenk and Snaar-Jagalska, 1999) present in all protein kinases (Fig. 1). The second sequence signature (HRDLKxxN) located at amino acid 678 to 685 represents the consensus sequence for Ser/Thr protein kinases (Schenk and Snaar-Jagalska, 1999) and is well conserved in *AtCTR1* and both tomato homologs. The overall comparison of the full sequences indicate that *LeCTR1* shares between 58% and 62% identity at the protein sequence level with *AtCTR1*, *AtEDR1*, and *LeCTR2*. The kinase domain of *LeCTR1* exhibits more identity with *AtCTR1* (82%) than with *LeCTR2* (59%) and *AtEDR1* (62%), whereas the kinase domain of *LeCTR2* and *AtEDR1* exhibits 85% identity (Frye et al., 2001). In the N-terminal region, *LeCTR1* also shows higher identity with *AtCTR1* (55%) than with *LeCTR2* and *AtEDR1* (36%). However, the amino-terminal region lacks significant homology to the amino-terminal portion of Raf, suggesting that *LeCTR1* and *AtCTR1* may be regulated by different factors and/or by a distinct mechanism (Kieber, 1997). In summary, the *LeCTR1* sequence displays greater similarity to *AtCTR1* than *LeCTR2*, suggesting *LeCTR1* is more likely to bear *AtCTR1* function as a negative regulator of the ethylene transduction pathway in tomato.

Isolation and Structural Analysis of the Tomato *LeCTR1* Genomic Clone

The *LeCTR1* genomic clone has been obtained by PCR amplification of tomato genomic DNA. Comparison of genomic and cDNA sequences allowed the delineation of intron and exon positions. Both *AtCTR1* and *LeCTR1* have 15 exons and 14 introns

Figure 1. Sequence comparison between tomato LeCTR1 (AAI87456) and LeCTR2 (AJ005077), AtCTR1 (CAB82938), AtEDR1 (AAG31143), and *R. norvegicus* C-Raf (P11345) proteins. Identities between proteins are indicated by shaded squares. The kinase catalytic domain is located in the C-terminal side, including the 11 subdomains (roman numerals). The sequence consensus for the ATP-binding site and Ser/Thr protein kinase are also indicated (. and - - - - - , respectively). Arabic numbers indicate the starting of each exon for *LeCTR1* and *AtCTR1*. (Figure continues on facing page.)






Figure 2. Comparison of the genomic structure of the tomato *LeCTR1* (AY079048) and the Arabidopsis *AtCTR1* gene (L08790). Black portions represent the introns, white portions represent the exons, and gray portions represent the untranslated region. Arrows indicate that each exon of *LeCTR1* gene correspond to its homolog in the *AtCTR1* gene.

with the position and size of the exons perfectly conserved between the two species (Fig. 2). Despite this structural conservation, the *LeCTR1* introns are typically larger than those of *AtCTR1*. As a consequence, the overall size of the *LeCTR1* transcribed region from ATG to stop codon is more than twice that of *AtCTR1* (12,006 bp versus 4,701 bp, respectively). Such high conservation of genomic structure might be indicative of conserved function.

Reversion of the Arabidopsis *ctr1-1* Mutant Phenotype by Complementation with *LeCTR1*

Expression of LeCTR1 in the Arabidopsis ctr1-1 Mutant Restores the Wild-Type Phenotype

To assess the functional significance of *LeCTR1*, we attempted complementation of the *ctr1-1* mutant of Arabidopsis (Columbia ecotype) using a sense construct of the *LeCTR1* cDNA driven by the 35S promoter. Figure 3 shows the typical seedling phenotype of the *ctr1-1* mutant. Light-grown *ctr1-1* seedlings display a considerable delay in the opening of the apical hook and in cotyledon expansion, a greater darkening of the cotyledons, and significant reduction of root elongation. Dark-grown seedlings displayed a constitutive triple response (Kieber et al., 1993). Seventeen transgenic lines corresponding to independent transformation events harboring the 35S:*LeCTR1* sense construct were generated. Three transgenic lines presenting different levels of recovery of the wild-type phenotype were selected for detailed molecular and physiological analysis.

Light-grown transgenic lines displayed variable degrees of complementation of different aspects of the mutant phenotype (Fig. 3). For instance, adult plants from all three transformed lines displayed a wild-type phenotype in terms of rosette size and inflorescence development (Fig. 3, A and B). The cotyledon shape, color, and timing of development were completely identical between the complemented lines and wild-type control, whereas root length of transgenic plants was highly variable, ranging from the wild-type to *ctr1-1* mutant phenotypes (Fig. 3C). Specifically, line 27 grown in the light developed normal roots similar in size to those of

wild-type plants, whereas under identical growth conditions, line 104 has short roots only slightly longer than those of *ctr1-1*. Line 17 has roots with intermediate elongation. Moreover, etiolated seedlings of all *LeCTR1*-overexpressing lines displayed a gradual recovery of the hypocotyl elongation rate compared with the *ctr1-1* mutant (Fig. 3D).

Recovery of Normal Ethylene Response of the ctr1-1 Mutant Complemented with LeCTR1

To more fully assess the degree to which *LeCTR1* complementation restores the capacity of the *ctr1-1* mutant to respond to ethylene, we established a dose response curve of hypocotyl length in response to exogenous ethylene treatment (Fig. 4). After 3 d of growth in the absence of ethylene, hypocotyl length reached 80%, 72%, and 54% of that of wild-type size in lines 27, 17, and 104, respectively, compared with 33% in the non-complemented *ctr1-1* controls. When complemented seedlings were treated with ethylene, hypocotyl elongation decreased in correlation with increasing ethylene concentration. The minimal threshold ethylene concentration yielding scorable alteration of hypocotyl length was $0.1 \mu\text{L L}^{-1}$, whereas $1 \mu\text{L L}^{-1}$ resulted in identical inhibition of hypocotyl elongation in both wild-type and transgenic *ctr1-1* lines expressing *LeCTR1* (Fig. 4).

Analysis of apical hook curvature revealed that all three complemented lines responded to ethylene, though, consistent with the hypocotyl phenotype, the degree of response varied among the three lines. Specifically, lines 17 and 104 displayed a 90° (index 2) and 180° (index 3) apical curvature, respectively, even in the absence of ethylene treatment (Fig. 5), suggesting partial complementation for these two lines. Line 27 displayed a fully open hook in the absence of ethylene but exhibited hook formation at $0.1 \mu\text{L L}^{-1}$ ethylene as compared with a requirement of $1 \mu\text{L L}^{-1}$ for the wild-type control (Fig. 5). These observations show that expression of *LeCTR1* in the *ctr1-1* mutant was capable of restoring seedling responsiveness to ethylene both in the hypocotyl and in the hook. Although the hypocotyl response was triggered by the same ethylene concentration in wild-type and complemented mutant lines, hook curvature required less ethylene in the *LeCTR1*-expressing lines, suggesting differential sensitivity to the hormone in various complemented tissues.

LeCTR1 Expression Restores Glc Sensitivity to the ctr1-1 Mutant

Wild-type seedlings of Arabidopsis undergo growth arrest when cultivated in light in the presence of 6% (w/v) Glc. Exogenous ethylene allows seedlings to overcome Glc-induced inhibition of growth (Zhou et al., 1998). The *ctr1-1* mutant is capable of normal seedling growth in the presence of 6% (w/v)




Figure 3. Phenotypes of the transgenic *LeCTR1*-overexpressing lines (27, 17, and 104) compared with that of *Arabidopsis* wild type and the *ctr1-1* mutant. A, Adult plants at the rosette stage grown in the greenhouse. B, Adult plants at the flowering stage grown in the greenhouse. C, Four-day-old seedlings grown in the light. D, Four-day-old etiolated seedlings.

Glc, even in the absence of ethylene (Fig. 6A), presumably due to constitutive activation of ethylene signaling. Complementation of *ctr1-1* with *LeCTR1* resulted in recovery of Glc-induced growth inhibition similar to that shown by the wild type (Fig. 6A). When seedlings were supplemented with $10 \mu\text{L L}^{-1}$ of ethylene, all lines including wild-type control, overcame Glc-induced growth inhibition (Fig. 6B). The reversion of the *ctr1-1* phenotype relative to Glc tolerance was total and included all growth arrest-associated symptoms that had been previously described (Zhou et al., 1998). Specifically, *ctr1-1* seedlings expressing *LeCTR1* demonstrated ethylene reversible inhibition of: (a) expansion and greening of cotyledons, (b) abnormal development, and (c) root elongation (Fig. 6). These data further demonstrate the ability of *LeCTR1* to complement for the loss of CTR1 function in the *Arabidopsis ctr1-1* mutant.

Molecular Analysis of the Transformed Lines

The incorporation of the transgene in the three transformed lines selected for this study was confirmed by Southern-blot analysis. Figure 7A shows that line 27 contained two copies of the transgene, whereas lines 17 and 104 contained only one. As a consequence, homozygous progenies were easily se-

lected for lines 17 and 104, whereas for line 27, because of the presence of multiple insertions, it was difficult to ascertain whether the progenies were homozygous for all the insertions. Therefore, it was assumed that a mixed population of plants was used in the case of line 27. Northern-blot analysis clearly indicated the presence of transgene-derived transcripts in the transformed lines, but not in the wild type nor in the *ctr1-1* mutant. Moreover, *LeCTR1* transcripts accumulated to higher levels in line 27 as compared with the two other transgenic lines. The level of transgene expression correlated positively with the number of T-DNA insertions consistent with the apparent gene dose effect described for hypocotyl elongation. Line 27, which displayed the highest level of *LeCTR1* transcript accumulation, also showed the greatest degree of complementation (Fig. 7B).

To investigate the effect of *LeCTR1* at the level of ethylene-responsive gene expression, we analyzed the accumulation of an ethylene-inducible basic chitinase (Samac et al., 1990). Chitinase gene expression is ethylene dependent and has been shown to be constitutively expressed in the *ctr1-1* mutant (Kieber et al., 1993). As expected, in the absence of ethylene, the chitinase transcripts were undetectable in wild-type lines, while accumulating to substantial levels in the *ctr1-1* mutant (Fig. 7B). *LeCTR1*-complemented *ctr1-1* lines exhibited a dramatic reduction of chitinase ex-




Figure 4. Ethylene response of the transgenic *LeCTR1*-overexpressing lines (27, 17, 104) compared with that of wild type and the *ctr1-1* mutant. Etiolated seedlings untreated or treated with increasing ethylene concentration (0.01, 0.1, 1, and 10 $\mu\text{L L}^{-1}$) were grown for 3 d before monitoring hypocotyl length. Each histogram represents the mean of 30 measurements and the vertical bars indicate the confidence interval.

pression as compared with the untransformed mutant, indicating reversion toward the wild-type phenotype. However, there was no clear correlation between transgene copy number and the level of chitinase gene expression.

***LeCTR1* Gene Expression Is Regulated by Ethylene in Tomato**

A variety of tissues were harvested for RNA extraction and analyzed for levels of *LeCTR1* mRNA expression using real-time quantitative PCR (Fig. 8A). The *LeCTR1* message was detected at varying

levels in all tissues examined. The *LeCTR1* message increased, coincident with the onset of fruit ripening. *LeCTR1* transcript levels were relatively low in mature green fruit and increased during the breaker and 3 d post-breaker stages, followed by a decline during later fruit development.




Figure 5. Effect of ethylene on the apical hook curvature of the transgenic *LeCTR1*-complemented lines (27, 17, and 104) compared with that of wild type and the *ctr1-1* mutant. ○, Wild type; +, *ctr1-1* mutant; Δ, line 27; ◇, line 17; □, line 104. The level of apical curvature was estimated visually for 30 seedlings using a scale ranging from 0 to 4 (0, no apical hook; 1, 90° curvature; 2, 180° curvature; 3, beginning of hook formation; and 4, full hook). The experiment was repeated twice.




Figure 6. Effect of Glc on the development of the transgenic *LeCTR1*-complemented lines (27, 17, and 104) compared with wild type and the *ctr1-1* mutant. For each line, 50 seedlings are grown on Murashige and Skoog medium containing 6% (w/v) Glc during 10 d in the light. Wild type, *ctr1-1* mutant, or *LeCTR1*-complemented lines grown in a sealed box with air (A) or in the presence of 10 $\mu\text{L L}^{-1}$ ethylene (B). The experiment was repeated three times.




Figure 7. Molecular analysis of the transgenic *LeCTR1*-complemented lines (27, 17, and 104) compared with wild type and the *ctr1-1* mutant. A, Southern analysis of the transgenic lines with a *NPTII* probe. Numbers indicate the fragment size in kilobase pairs. Line 27 contained two insertions, whereas line 17 and 104 contained only one copy of the transgene. B, Northern-blot analysis of *LeCTR1* and basic chitinase transcript accumulation. Equal loading of the gel with the RNA samples is checked by ethidium bromide staining (bottom).

It has been shown previously that *LeCTR1* expression is inducible in mature green fruit treated with ethylene (Zegzouti et al., 1999). To more fully characterize the dynamics of ethylene responsiveness during fruit development and particularly at the onset of ripening, *LeCTR1* gene responsiveness to ethylene was examined in mature green fruit (Fig. 8B). *LeCTR1* responded relatively rapidly to ethylene treatment, demonstrating a 4-fold induction within 3 h of treatment. Modest levels of induction were maintained throughout the 24-h experimental time course and analysis of expression in ripening fruit indicated this induction is likely to persist throughout the later stages of ripening and senescence (Fig. 8A).

The highest levels of *LeCTR1* observed during flower development occur in association with senescence, which is also marked by considerable ethylene biosynthesis. Figure 8C shows that 4-week-old tomato plants demonstrated increased *LeCTR1* expression in response to ethylene in leaves (6-fold) and roots (2-fold). These results suggest that *LeCTR1* is ethylene inducible in a range of tomato tissues, in

contrast to the relative constitutive expression reported for its Arabidopsis counterpart, *AtCTR1* (Kieber et al., 1993). The fact that substantial induction of *LeCTR1* occurs during stages of significant ethylene-mediated developmental modification (e.g. ripening, senescence, and abscission) may reflect evolutionary modification of ethylene signal transduction pathway regulation to meet the signaling needs of tissues greatly influenced by ethylene action.


CONCLUSION

Although a number of CTR1-like sequences from tomato and other plant species are available in the gene databases, there is no experimental evidence regarding their putative involvement in the ethylene signal transduction pathway. The isolation of the *LeCTR1* genomic clone showed that its structural organization was very well conserved when compared with the Arabidopsis *CTR1* gene. The availability of a well-characterized Arabidopsis *ctr1-1* mutant offered a unique opportunity to investigate the predicted function of *LeCTR1* through heterologous expression. By converting the constitutive triple-response phenotype of the Arabidopsis *ctr1-1* mutant to a largely normal ethylene-responsive phenotype, we demonstrated that *LeCTR1* encodes a functional ethylene signal transduction component capable of interacting with upstream and downstream partners of the Arabidopsis CTR1 protein. These data strongly support that the ethylene response pathways in tomato and Arabidopsis are composed of conserved components, yet there are potential differences.

In contrast with what has been previously reported in Arabidopsis for *CTR1* (Kieber et al., 1993), we have shown that *LeCTR1* is ethylene inducible. In accordance, *LeCTR1* mRNA accumulation was up-regulated during tomato fruit ripening and in additional tissues synthesizing large amounts of ethylene. However, even though the data reported so far indicated that the expression of *AtCTR1* was not affected significantly by ethylene (Kieber et al., 1993), it must be stressed here that there is only limited information on ethylene inducibility of this gene in Arabidopsis. The discrepancy observed between the two species for the expression of the CTR gene might also arise from the fact that the Arabidopsis CTR protein is encoded by a single gene, whereas in the tomato, at least two CTR isoforms exist that are encoded by a small multigene family (L. Adams and J. Giovannoni, unpublished data).

In tomato, it has been shown that the five gene members encoding the ethylene receptors are differentially regulated at the transcriptional level (Lashbrook et al., 1998). Interestingly, the tomato ethylene receptor, NR (*LeETR3*), shows similar induction to *LeCTR1* during the course of fruit ripening in concert with increased ethylene evolution, which could be an

Figure 8. Ethylene-dependent and tissue-specific expression of *LeCTR1* in tomato. The levels of *LeCTR1* transcripts were assessed by real-time quantitative PCR. The experiments were carried out in triplicate. A, *LeCTR1* mRNA accumulation was monitored in the root (R), hypocotyl (H), cotyledon (C), unopened buds (B), flower at anthesis (FA), senescent flowers (FS), young fruit 7 DPA (IG), mature green fruit (MG), breaker fruit (Br), breaker + 3 (Br+3), breaker + 7 (Br+7), abscission zone (Ab), callus (Cal), and leaf (L). $\Delta\Delta Ct$ on the y axis refers to the fold difference in *LeCTR1* expression relative to the leaf. B, Ethylene responsiveness of *LeCTR1* in mature green fruit treated with 20 $\mu\text{L L}^{-1}$ ethylene. $\Delta\Delta Ct$ on the y axis refers to the fold difference in *LeCTR1* expression relative to the control. C, *LeCTR1* ethylene regulation in root and leaves. Ethylene treatment was performed as in B. $\Delta\Delta Ct$ on the y axis refers to the fold difference in *LeCTR1* expression relative to air-treated root and leaf, respectively.



effective method of regulating ethylene responsiveness of various plant tissues (Tieman et al., 2000). The observation that a second component of ethylene signaling in tomato, *LeCTR1*, demonstrates similar regulation suggests that multiple targets may exist for modulation of ethylene responsiveness in at least some plant species. It is striking that a negative regulator of ethylene response is induced during fruit ripening when just the opposite might be logically anticipated. This expression pattern may simply represent the ethylene-inducible nature of *LeCTR1* in a tissue producing large amounts of ethylene, though said induction may have little physiological relevance. The fact that tomato fruit produce ethylene in concentrations greatly exceeding those required to induce ripening would support this possibility (Oeller et al., 1991). Alternatively, induction of *LeCTR1* during ripening may represent a damping mechanism to prevent ripening and subsequent senescence from proceeding too rapidly. Analysis of *LeCTR1* expression in tomato cultivars demonstrating variable fruit ethylene evolution rates and/or

ripening times could clarify the role of *LeCTR1* induction during fruit ripening.

MATERIALS AND METHODS

Plant Material

Arabidopsis (ecotype Columbia) plants were grown under standard greenhouse conditions. *Agrobacterium tumefaciens*-mediated transformation was carried out using the pGA643 binary vector according to Bird et al. (1988). The sense construct was generated by cloning the full *LeCTR1* open reading frame under the transcriptional control of the cauliflower mosaic virus 35S promoter and the nopaline synthase terminator. The transformation protocol for in planta transformation of Arabidopsis was as described by Clough and Bent (1998). *A. tumefaciens* strain C58 carrying the binary plasmid pGA643 was grown to stationary phase in LennoxL-Broth medium at 28°C, 250 rpm. Cells were harvested by centrifugation for 20 min at room temperature at 5,500g and then resuspended in inoculation medium containing 5% (w/v) Suc and 0.05% (v/v) Silwet L-77 (OSI Specialties, Inc., Danbury, CT). Plants were inverted into this suspension such that all aboveground tissues were submerged, and plants were then removed after 3 to 5 s of gentle agitation. Plants were left in a low-light or dark location overnight and returned to the greenhouse the next day. Plants were grown for further 3 to 5 weeks until siliques were brown and dry. The selection of putative transformants was done on a 70 mg L⁻¹ kanamycin-containing agar medium.

Isolation of the Genomic Clone

Genomic DNA was extracted from 1 g of ground tomato (*Lycopersicon esculentum*) leaves. The powder was mixed with 5 mL of extraction buffer (2% [w/v] hexadecyl-trimethyl-ammonium bromide, 1.4 M NaCl, 20 mM EDTA, and 100 mM Tris-HCl, pH 8) and warmed at 65°C for 10 min. After a phenol/chloroform/isoamylalcohol and chloroform extraction, DNA was precipitated with 1 volume of isopropanol for 20 min on ice. After centrifugation (5 min for 2,000g), the pellet was resuspended in 10 mL of wash buffer (76% [v/v] ethanol and 10 mM ammonium acetate). After centrifugation (10 min for 2,000g), the DNA was resuspended in 200 μ L of sterile water. An RNase treatment was done at 37°C for 10 min. Several primers were chosen based on the cDNA sequence and polymerization chain reactions were performed on the genomic DNA (primer 1, GAGCTGAAAATTGCTA-ATGGCAG, with primer 2, CCATCCCATAAATCCAATAAAATCC; primer 3, GGATTTTATGGATTATGGGATGG, with primer 4, CTTCTCCAGCAG-GAGCTGCACCCC; primer 5, GGGGTGCAGCTCCTGCTGGAGAAG, with primer 6, CGCTTAAACTCCAGGCTTACC; primer 7, GGTAAGCCTG-GAGTTTTAAGCG, with primer 8, CTGTATCTGGTCTGTCATCGGTG; primer 9, CACCGATGACACGACCAGATACAG, with primer 10, TGAGAG-CAACTGCATGCTGTGTG; and primer 11, CTCCTACCTCCACCAGGT, with primer 12, CATAAGTTATACAAGAATCCTGGGC). The fragments were cloned and sequenced. Comparative analysis between the genomic clone and cDNA sequences allowed the delimitation of introns and exons.

Southern Analysis

The genomic DNA was extracted as described above. DNA (5 μ g) was digested with *EcoRV* enzyme. Digested DNA was separated in an agarose gel and blotted on a nylon membrane as described by Sambrook et al. (1989). A probe corresponding to the *NPTII* gene was labeled with [³²P]dCTP using a random primer kit (Ready-to-Go, Amersham Biosciences, Piscataway, NJ). Blots were hybridized with a fragment of *NPTII* gene in a buffer containing 0.3 volumes of 1 M sodium phosphate buffer (pH 7.2), 0.7 volumes of 10% (w/v) SDS, and 1:500 (v/v) volumes of 0.5 M EGTA, pH 8. Washes were carried out as described by Sambrook et al. (1989).

Northern Analysis

Leaves from Arabidopsis plants were collected at the rosette stage and total RNA was obtained from 0.5 g of leaf tissue ground in liquid nitrogen and extracted with phenol as previously described (Verwoerd et al., 1989), except that the extraction buffer was 100 mM Tris-HCl (pH 8.0), 100 mM LiCl, 10 mM EDTA, and 1% (w/v) SDS. Total RNA (8 μ g) was fractionated on a 1.2% (w/v) agarose gel containing formaldehyde in MOPS buffer (Sambrook et al., 1989) and then transferred onto GeneScreen membranes (PerkinElmer Life Sciences, Boston) following the manufacturer's instructions. Probes were labeled with [³²P]dCTP using a random primer kit (Ready-to-Go, Amersham Biosciences). Blots were hybridized, as described for the Southern analysis, with a fragment of the basic chitinase cDNA (Samac et al., 1990) or with a fragment of the *LeCTR1*. To check for equal loading, a reverse picture of the ethidium bromide-stained gel was used. Washes were carried out under high stringency (Sambrook et al., 1989).

Ethylene Treatment

Sterilized seeds were put on Murashige and Skoog agar medium plates and placed at 4°C for 4 d. Ethylene treatment was carried out in sealed boxes. For air control, contaminating ethylene was removed using KMnO₄. The different concentrations of ethylene applied were checked by gas chromatography and adjusted to 0.01, 0.1, 1, and 10 μ L L⁻¹. Hypocotyl measurements were made on 30 seedlings grown in darkness during 3 d with or without ethylene. The experiment was repeated three times and the level of apical curvature was estimated visually using a scale ranging from 0 to 4 (0, no apical hook; 1, 90° curvature; 2, 180° curvature; 3, beginning of hook formation; and 4, full hook). At a given ethylene concentration, the level of apical curvature was homogenous, i.e. >90% of the seedlings exhibited the same phenotype.

Mature green fruit were placed in a sealed chamber and gassed with 20 μ L L⁻¹ ethylene for 0 to 24 h. Four-week-old tomato plants were placed in a sealed chamber and gassed with or without 20 μ L L⁻¹ ethylene for 8 h.

Glc Sensitivity

Sterilized seeds were put on Murashige and Skoog agar medium containing 6% (w/v) filter-sterilized Glc. After 4 d at 4°C, plates were placed in light in a sealed box containing either air or 10 μ L L⁻¹ ethylene. Every 2 d, the boxes were opened to allow a renewal of the atmosphere and put back either with air or 10 μ L L⁻¹ ethylene. The experiment was stopped after 10 d of culture.

Quantitative RT-PCR

To obtain total RNA, the same protocol for northern analysis was used. The pellet was allowed to air dry and was resuspended in diethyl pyrocarbonate water. After quantification, 10 μ g of RNA was treated with DNase I (Promega, Madison, WI) and cleaned up with a phenol-chloroform extraction.

Real-time quantitative PCR was performed using 250 ng of total RNA for *LeCTR1* and 2.5 pg for 18S in a 20- μ L reaction volume using *Taq-Man* One-Step RT-PCR Master Mix reagents kit (PE-Applied Biosystems, Foster City, CA) on an ABI PRISM 7900HT sequence-detection system. PRIMER EXPRESS software (PE-Applied Biosystems) was used to design gene-specific primers and *Taq-Man* probes: *LeCTR1* forward primer, CAT-CCTCTTTCTACTGTGAGAAAATTTAGA; *LeCTR1* reverse primer, CATTCCCTGTATAAAAAACGTTTCAGTT; *LeCTR1* *Taq-Man* probe, VIC-CCAAGTCCATTAGCAATTTTCAGCTCAA-TAMRA; 18S forward primer, CGGAGAGGGAGCCTGAGAA; 18S reverse primer, CCCGTGT-TAGGATTTGGGTAATTT; and 18S *Taq-Man* probe, 6FAM-CGGCTACCA-CATCCAAGGAAGGCA-TAMRA. For *LeCTR1*, optimal primer concentration was 900 nM and optimal probe concentration was 250 nM. Optimal primer and probe concentrations for 18S were 300 and 125 nM, respectively. RT-PCR conditions were as follows: 48°C for 30 min, 95°C for 10 min, followed by 40 cycles of 95°C for 15 s and 60°C for 1 min. Samples were run in triplicate on each 384-well plate and were repeated on at least two plates for each experiment. For each sample, a Ct (threshold cycle) value was calculated from the amplification curves by selecting the optimal Δ Rn (emission of reporter dye over starting background fluorescence) in the exponential portion of the amplification plot.

Relative fold differences were calculated based on the comparative Ct method using the 18S as an internal standard. To demonstrate that the efficiencies of the *LeCTR1* and 18S primers and probes were approximately equal, the absolute value of the slope of log input amount versus delta Ct was calculated for both *LeCTR1* and 18S and was determined to be <0.1. To determine relative fold differences for each sample in each experiment, the Ct value for *LeCTR1* was normalized to the Ct value for 18S and was calculated relative to a calibrator (leaf for Fig. 8A and control tissues for Fig. 8, B and C) using the formula $2^{-\Delta\Delta Ct}$.

ACKNOWLEDGMENT

We are grateful to Simone Albert for excellent technical assistance in Arabidopsis genetic transformation and analysis of plant growth.

Received June 5, 2002; returned for revision July 14, 2002; accepted July 26, 2002.

LITERATURE CITED

- Abeles FB, Morgan PW, Salveit ME (1992) Ethylene in Plant Biology, Ed 2. Academic Press, New York
- Bird CR, Smith CJS, Ray JA, Moureau P, Bevan MW, Bird AS, Hughes S, Morris PC, Grierson D, Schuch W (1988) The tomato polygalacturonase gene and ripening-specific expression in transgenic plants. *Plant Mol Biol* 11: 651–662
- Chang C, Kwon SF, Bleecker AB, Meyerovitz EM (1993) *Arabidopsis* ethylene response gene *ETR1*: similarity of product two-component regulators. *Science* 262: 539–544
- Clark KL, Larsen PB, Wang X, Chang C (1998) Association of the *Arabidopsis* CTR1 Raf-like kinase with the ETR1 and ERS ethylene receptors. *Proc Natl Acad Sci USA* 95: 5401–5406

- Clough SJ, Bent AF** (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**: 735–743
- Ecker JR** (1995) The ethylene signal transduction pathway in plants. *Science* **268**: 667–675
- Frye CA, Tang D, Innes RW** (2001) Negative regulation of defense responses in plant by conserved MAPKK kinase. *Proc Natl Acad Sci USA* **98**: 373–378
- Giovannoni JJ** (2001) Molecular biology of fruit maturation and ripening. *Annu Rev Plant Physiol Plant Mol Biol* **52**: 725–749
- Giovannoni JJ, Kannan P, Lee S, Yen H-C** (1998) Genetic approaches to manipulation of fruit development and quality in tomato. In KE Cockshull, D Gray, GB Seymour, B Thomas, eds, *Genetic and Environmental Manipulation of Horticultural Crops*. CABI Publishing, New York, pp 1–15
- Guzmán P, Ecker JR** (1990) Exploiting the triple response of *Arabidopsis* to identify ethylene-related mutants. *Plant Cell* **2**: 513–523
- Hanks SK, Quinn AM** (1991) Protein kinase catalytic domain sequence database: identification of conserved features of primary structure and classification of family members. *Methods Enzymol* **200**: 38–62
- Hanks SK, Quinn AM, Hunter T** (1988) The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science* **241**: 42–52
- Hua J, Chang C, Sun Q, Meyerowitz EM** (1995) Ethylene insensitivity conferred by *Arabidopsis* *ERS* gene. *Science* **269**: 1712–1714
- Hua J, Meyerowitz EM** (1998) Ethylene responses are negatively regulated by a receptor gene family in *Arabidopsis thaliana*. *Cell* **94**: 261–271
- Johnson PR, Ecker JR** (1998) The ethylene gas signal transduction pathway: a molecular perspective. *Annu Rev Genet* **32**: 227–254
- Kieber JJ** (1997) The ethylene response pathway in *Arabidopsis*. *Annu Rev Plant Physiol Plant Mol Biol* **48**: 277–296
- Kieber JJ, Rothenberg M, Roman G, Feldman KA, Ecker JR** (1993) *CTR1*, a negative regulator of the ethylene response pathway in *Arabidopsis*, encodes a member of the Raf family of protein kinases. *Cell* **72**: 427–441
- Lashbrook CC, Tieman DM, Klee HJ** (1998) Differential regulation of the tomato *ETR* gene family throughout plant development. *Plant J* **15**: 243–252
- Lelièvre J-M, Latché A, Jones B, Bouzayan M, Pech J-C** (1997) Ethylene and fruit ripening. *Physiol Plant* **101**: 727–739
- Oeller PW, Wong LM, Taylor LP, Pike DA, Theologis A** (1991) Reversible inhibition of tomato fruit senescence by antisense 1-aminocyclopropane-1-carboxylate synthase. *Science* **254**: 427–439
- Payton S, Fray RG, Brown S, Grierson D** (1996) Ethylene receptor expression is regulated during fruit ripening, flower senescence and abscission. *Plant Mol Biol* **31**: 1227–1231
- Samac DA, Hironaka CM, Yallaly PE, Shah DM** (1990) Isolation and characterization of the genes encoding basic and acidic chitinase in *Arabidopsis thaliana*. *Plant Physiol* **93**: 907–914
- Sambrook J, Fritsch EF, Maniatis T** (1989) *Molecular Cloning: A Laboratory Manual*, Ed 2. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Shaller GE, Bleeker AB** (1995) Ethylene-binding site generated in yeast expressing the *Arabidopsis* *ETR1* gene. *Science* **270**: 1809–1811
- Schenk PW, Snaar-Jagalska BE** (1999) Signal perception and transduction: the role of protein kinase. *Biochem Biophys Acta* **1449**: 1–24
- Stepanova AN, Ecker JR** (2000) Ethylene signaling: from mutants to molecules. *Curr Opin Plant Biol* **3**: 353–360
- Tieman DM, Klee HJ** (1999) Differential expression of two novel members of the tomato ethylene-receptor family. *Plant Physiol* **120**: 165–172
- Tieman DM, Taylor MG, Ciardi JA, Klee HJ** (2000) The tomato ethylene receptors NR and LeETR4 are negative regulators of ethylene response and exhibit functional compensation within a multigene family. *Proc Natl Acad Sci USA* **97**: 5663–5668
- Verwoerd TC, Dekker BMM, Hoekema A** (1989) A small-scale procedure for the rapid isolation of plant RNAs. *Nucleic Acids Res* **17**: 2362
- Wilkinson JQ, Lanahan MB, Yen HC, Giovannoni JJ, Klee HJ** (1995) An ethylene inducible component of signal transduction encoded by *Never-ripe*. *Science* **257**: 967–971
- Zegzouti H, Jones B, Frasse P, Marty C, Maître B, Latché A, Pech J-C, Bouzayan M** (1999) Ethylene-regulated gene expression in tomato fruit: characterization of novel ethylene-response and ripening-related genes isolated by differential-display. *Plant J* **18**: 589–600
- Zhou D, Kalitatis P, Mattoo AK, Tucker M** (1996) The mRNA for an ETR1 homologue in tomato is constitutively expressed in vegetative and reproductive tissues. *Plant Mol Biol* **30**: 1331–1338
- Zhou L, Jank J-C, Jones TL, Sheen J** (1998) Glucose and ethylene signal transduction cross talk revealed by an *Arabidopsis* glucose-insensitive mutant. *Proc Natl Acad Sci USA* **95**: 10294–10299