

Differential prostaglandin production by microsomal fractions of rat pregnant uterus

I. DOWNING & K.I. WILLIAMS

Pharmacology Group, School of Pharmacy and Pharmacology, University of Bath, Bath BA2 7AY

Crude homogenates of decidual tissue from rat pregnant uteri synthesize prostaglandin (PG) F as estimated by parallel bioassay (Williams, Sneddon & Harney, 1974). As recent work has shown the increasing complexity of arachidonic acid metabolism we have developed a radiometric assay to allow more detailed analysis of products formed by the uterus.

Pregnant rats were killed (day 22 of pregnancy) and 20% homogenates of deciduum and myometrium prepared in Tris/acetate buffer (pH 8) containing hydroquinone 0.5 mM and reduced glutathione 2 mM. Microsomes were prepared by ultracentrifugation and samples incubated with 1 ml aliquots of buffer containing arachidonic acid (99 nmol) and [¹⁴C]-arachidonic acid (3.5 nmol) for 30 min at 37°C. Pilot experiments indicated these conditions were optimal for decidual tissue with regard to substrate and co-factor concentrations, pH and incubation time. Samples boiled before incubation served as controls. After solvent extraction residues were subjected to thin-layer chromatography (TLC) using the solvent chloroform 90: methanol 9: acetic acid 1: water 0.65 by volume. The radioactivity in consecutive 0.5 cm bands of each zone was then estimated by liquid scintillation counting. Authentic PGF_{2α}, PGE₂, [¹⁴C]-6-oxo PGF_{1α} (prepared as described by Cottee, Flower, Moncada, Salmon & Vane, 1977) and arachidonic acid were used as markers.

With decidual microsomes 23% of radioactivity on the TLC plate was converted to products other than monohydroxy acids (range 14–50%, 15 experiments). 50% of this radioactivity had an R_F value similar to PGE₂, 25% ran as PGD₂ while 20% had a mobility equivalent to PGF_{2α}. Myometrial substrate conversion was lower, 6% (range 2–15%, 15 experiments) and all the radioactivity was located in the PGE₂ zone. However, in this solvent system PGE₂ and 6-oxo PGF_{1α} have similar R_F values. Therefore the zones of radioactivity corresponding to PGE₂ were removed, eluted and rechromatographed using solvent systems devised to separate these two substances (Cottee *et al.*, 1977). Using these TLC systems we found that approximately 12% of radioactivity in the decidual PGE zone behaved as 6-oxo PGF_{1α} and 70% as PGE₂. However 84% of myometrial radioactivity appeared to be 6-oxo PGF_{1α} and only 6% PGE₂.

Thus decidual microsomes produce PGE₂ as the major product whereas myometrial preparations synthesize predominantly 6-oxo PGF_{1α}. Further investigation is needed to determine whether this myometrial PG production plays any part in regulating uterine function during pregnancy.

We thank the Wellcome Trust for a grant.

References

- COTTEE, F., FLOWER, R.J., MONCADA, S., SALMON, J.A. & VANE, J.R. (1977). Synthesis of 6-oxo PGF_{1α} by ram seminal vesicle microsomes. *Prostaglandins* (in press).
WILLIAMS, K.I., SNEDDON, J.M. & HARNEY, P.J. (1974). Prostaglandin production by the pregnant rat uterus *in vitro* and its relevance to parturition. *Pol. J. Pharmac. Pharm.*, 26, 207–215.

Accumulation of platelets at acute inflammatory sites

J.P. BOLAM & M.J.H. SMITH

Biochemical Pharmacology Research Unit, Department of Chemical Pathology, King's College Hospital Medical School, Denmark Hill, London SE5 8RX

The involvement of blood platelets in acute inflammatory reactions is a topic of some interest. One experimental approach has been the use of animals made thrombocytopenic by treatment with anti-platelet sera. Some workers have reported that such animals either show diminished responsiveness to reactions such as the active Arthus and Shwartzman (Margaretten & McKay, 1969, 1971) or a decreased appearance of platelet-like cells into developing inflammatory exudates (Smith, Walker, Ford-Hutchinson & Penington, 1976). In other animal

models of acute inflammation, such as hind paw oedemas, it has been reported that platelet-like bodies occur in the swollen paw (Rédei & Kelemen, 1969) but that the development of the oedema is not affected by pre-treatment of the animals with anti-platelet sera (Ubatuba, Harvey & Ferreira, 1975).

We have examined the distribution of platelets and erythrocytes in and around implanted inert sponges in the rat. Female albino Wistar rats (150–200 g) received either homologous platelets labelled with ⁵¹Cr (sodium chromate) according to the method of Radegran (1976) or homologous erythrocytes labelled with ⁵¹Cr according to the method of Gray & Sterling (1950). Eighteen hours after labelling, the animals were subjected to an inflammatory insult, either subdermal implantation of 3 to 6 saline soaked sponges or subdermal incisions without sponge implantation. The rats were killed after various time intervals and the sponges, the skin around the sponges or incision, skin from other areas and blood samples