

# Properties of Guaiacol Peroxidase Activities Isolated from Corn Root Plasma Membranes<sup>1</sup>

Angela Mika and Sabine Lüthje\*

Universität Hamburg, Institut für Allgemeine Botanik, Ohnhorststrasse 18, D-22609 Hamburg, Germany

Although several investigations have demonstrated a plasma membrane (PM)-bound peroxidase activity in plants, this study is the first, to our knowledge, to purify and characterize the enzymes responsible. Proteins were extracted from highly enriched and thoroughly washed PMs. Washing and solubilization procedures indicated that the enzymes were tightly bound to the membrane. At least two distinct peroxidase activities could be separated by cation exchange chromatography (pmPOX1 and pmPOX2). Prosthetic groups were identified in fractions with peroxidase activity by absorption spectra, and the corresponding protein bands were identified by *heme* staining. The activities of the peroxidase enzymes responded different to various substrates and effectors and had different thermal stabilities and pH and temperature optima. Because the enzymes were localized at the PM and were not effected by *p*-chloromercuribenzoate, they were probably class III peroxidases. Additional size exclusion chromatography of pmPOX1 revealed a single activity peak with a molecular mass of 70 kD for the native enzyme, whereas pmPOX2 had two activity peaks (155 and 40 kD). Further analysis of these fractions by a modified sodium dodecyl sulfate-polyacrylamide gel electrophoresis in combination with *heme* staining confirmed the estimated molecular masses of the size exclusion chromatography.

Peroxidases (EC 1.11.1.7, etc.) belong to a large family of enzymes that are ubiquitous in fungi, plants, and vertebrates. These proteins usually contain a ferriprotoporphylin IX prosthetic group and oxidize several substrates in the presence of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>; Penel et al., 1992; Vianello et al., 1997). In higher plants, the number of isoenzymes may be extremely high, up to 40 genes corresponding to isoperoxidases for each plant, and several other isoforms can be generated by posttranscriptional and posttranslational modifications (Welinder et al., 1996; De Marco et al., 1999).

Although many soluble intracellular and extracellular peroxidases have been characterized in detail (for refs., see Gaspar et al., 1982; Hiraga et al., 2001; Shigeoka et al., 2002), less is known about membrane-bound enzymes, in particular the peroxidases of plant plasma membranes (PMs). Evidence for a PM-bound peroxidase activity in higher plants has been demonstrated frequently. Lin (1982) reported an increased oxygen consumption by intact corn (*Zea mays*) root protoplasts in the presence of extracellular NADH. Pantoja and Willmer (1988) obtained similar results using guard cell protoplasts from *Commelina communis* in the presence of NAD(P) H. PMs isolated from several species and plant parts showed NAD(P) H oxidase activities, which were comparable with a peroxidase (Møller and Bérczi, 1986;

Askerlund et al., 1987; Vianello et al., 1990, 1997; De Marco et al., 1995; Zancani et al., 1995; Sagi and Fluhr, 2001). Because the application of detergents did not significantly affect the activity observed and because activity could be detected with intact protoplasts, peroxidase activity has been suggested to be located at the apoplastic surface of the PM.

The NADH oxidation by PM from cauliflower (*Brassica oleracea*) could be stimulated by phenolic substances or inhibited by typical effectors of peroxidases like catalase, superoxide dismutase, cyanide, or azide (Askerlund et al., 1987). In PM-enriched fractions of *Arabidopsis* and Chinese cabbage (*Brassica campestris* L. subsp. *pekinensis*) seedlings, oxidation of Trp was reported in the presence of H<sub>2</sub>O<sub>2</sub> (Ludwig-Müller et al., 1990; Ludwig-Müller and Hilgenberg, 1992). PM isolated from soybean (*Glycine max*) roots showed a peroxidase activity in the presence of substrates like *o*-dianisidine, guaiacol, and ascorbate (Vianello et al., 1997). The oxidation of ascorbate could be strongly stimulated by phenolic acids, like caffeic and ferulic acid. Guaiacol or *o*-dianisidine oxidation rates were increased by CaCl<sub>2</sub> and inhibited by potassium cyanide and azide. When proteins solubilized by SDS from non-washed PM were separated by SDS-PAGE, two bands (38 and 45 kD) could be detected by *heme* staining. Peroxidase activity of these bands was not demonstrated, and only one, less intensive band remained after partial washing of the membrane vesicles (Vianello et al., 1997).

In addition to these experiments, antibodies specific for apoplastic peroxidases were used to detect PM-bound peroxidases by immunogold labeling and electron microscopy in situ (Hu et al., 1989; Penel and Castillo, 1991; Crevecoeur et al., 1997). However,

<sup>1</sup> This work was supported by the Deutsche Forschungsgemeinschaft (grant no. DFG Lu 668/1-2) and by the University of Hamburg (PhD student's grant no. HmbNFG to A.M.).

\* Corresponding author; e-mail s.luthje@botanik.uni-hamburg.de; fax 49-40-82282-254.

Article, publication date, and citation information can be found at [www.plantphysiol.org/cgi/doi/10.1104/pp.103.020396](http://www.plantphysiol.org/cgi/doi/10.1104/pp.103.020396).

Askerlund et al. (1987) demonstrated that the presence of peroxidases in PM preparations depends largely on the final PM washing procedure, which decreases the level of peroxidases significantly. A PM-bound peroxidase has not yet been isolated and characterized from highly purified and properly washed PM (Bérczi and Møller, 2000).

In the present work, we demonstrate the occurrence of at least two distinct peroxidase activities (pmPOX1 and 2) in corn root PM. A purification protocol for the isolation of these enzymes was developed, and the properties of the partially purified proteins were investigated by comparing them with soluble peroxidase activities.

## RESULTS AND DISCUSSION

### Binding to the PM

To check if peroxidase activities were loosely bound to the PM or entrapped inside the vesicles, different washing procedures were carried out. Independent of the salt concentrations used a maximum of 40% of the activity could be washed off in the presence of 1 mM EDTA and 0.01% (w/v) Triton X-100, i.e.  $79\% \pm 7.2\%$  ( $n = 2$ ) of the activity remained in the PM at 150 mM KCl and  $60\% \pm 1.9\%$  ( $n = 4$ ) at 500 mM KCl, respectively. Using 1 mM EGTA instead of EDTA did not change this result. A combination of 150 mM KCl, 1 mM EDTA, 0.01% (w/v) Triton X-100, and 0.1% (w/v) CHAPS (i.e. a detergent:protein ratio of 6:1 [w:w]) removed  $62\% \pm 0.4\%$  ( $n = 2$ ) of the peroxidase activity from the PM.

Due to the fact that neither physiological or high salt concentrations in the presence of detergent and EDTA or EGTA nor high detergent concentrations were able to remove the activity completely from the PM, we conclude that these enzymes are probably tightly bound to the PM. Salts should have minimal effects on the micellar size of Triton X-100, whereas effects on the zwitterionic detergent CHAPS cannot be excluded. Thus, the presence of higher salt concentrations could change the critical micellar concentration of CHAPS, thereby increasing the proportion of washed off peroxidase activity as a result of partial solubilization.

However, because peroxidase activity remains in the low detergent phase after Triton X-114 solubilization and temperature-induced phase separation (data not shown), the peroxidases were probably not strongly hydrophobic. Independently of the detergent to protein ratio used, none of the detergents tested (Triton X-114, Triton X-100, CHAPS, or octylglucopyranoside) could solubilize the activity completely from the PM. The mechanism of the binding to the PM is unknown, but sequence analysis of intracellular peroxidases indicated that transmembrane domains may exist in plant peroxidases (Bunkelmann and Trelease, 1996; Jespersen et al., 1997; Nito et al., 2001).


In Arabidopsis, three genes encoding membrane-bound ascorbate peroxidases were found (Jespersen et al., 1997). One of the corresponding proteins is probably bound to microbodies by a C-terminal transmembrane domain like membrane-bound intracellular peroxidases of other plant species (Bunkelmann and Trelease, 1996; Nito et al., 2001). However, sequence analysis of these peroxidases revealed that they are class I peroxidases, which implies they cannot occur in the PM.

### Purification


Solubilization by CHAPS yields about  $30\% \pm 1\%$  ( $n = 2$ ) of the activity and increased up to  $73\% \pm 4\%$  ( $n = 5$ ) in the presence of the dipole aminocaproic acid. Two activity peaks (pmPOX1 and pmPOX2) could be separated by cation exchange chromatography (Fig. 1). Peroxidase activities were eluted at 115 and 395 mM KCl. The total activity was divided into  $59\% \pm 3\%$  and  $41\% \pm 2\%$  ( $n = 4$ ) for pmPOX1 and pmPOX2, respectively. Starting from washed PM (specific activity  $401 \pm 52 \text{ nmol min}^{-1} \text{ mg protein}^{-1}$ ;  $n = 5$ ), a 24.0- and 8.8-fold purification for peak fractions of pmPOX1 and pmPOX2 with an overall yield of 31.4% was achieved. To compare the properties of pmPOX with soluble peroxidases, activities of the washing fluid of the PM (wPOX) were concentrated and separated by the same protocol (Fig. 1). The elution profile obtained was similar to that from the PM-bound POX. The total activity was divided into 30% and 70% for wPOX1 and wPOX2, respectively.

### Relative Molecular Mass

As shown in Figure 2, pmPOX1 displayed a single peak after size exclusion chromatography. By modi-



**Figure 1.** Elution profiles of POX after cation exchange chromatography. Enzyme activities isolated from corn root PM were separated on a Uno S1 column. Bound proteins were eluted by a KCl gradient from 0 to 1 M. The flow rate was  $1 \text{ mL min}^{-1}$ , and fractions of 1.0 and 0.5 mL were collected. A, Separation of POX activities from washing fluid of PM (wPOX; ○). B, Elution profile of PM-bound peroxidase activities (pmPOX; ●). Enzyme activities were measured in the presence of 8.26 mM guaiacol and 8.8 mM  $\text{H}_2\text{O}_2$ .



**Figure 2.** Elution profiles of PM-bound POX after size exclusion chromatography. Peak fractions collected from several Uno S runs were combined, concentrated, and applied onto a Superdex 200 column. Proteins were separated by a flow rate of 0.5 mL min<sup>-1</sup>. The fraction size was automatically adjusted between 0.75 and 0.5 mL depending on increase or decrease in  $A_{280}$  (dotted line). Enzyme activities were measured as described in Figure 1. PM-bound peroxidase activities could be separated into three peaks.

fied SDS-PAGE and *heme* staining, a protein band with an apparent molecular mass of 70 kD could be identified (Fig. 3). However, pmPOX2 was clearly separated into two peaks after size exclusion chromatography (pmPOX2a and pmPOX2b; Fig. 2). In comparison with peak fractions eluted during the cation exchange chromatography, analysis of pmPOX2b showed a significant increase in intensity of a 40-kD band after *heme* staining (Fig. 3). pmPOX2a exhibited a protein band between 100 and 170 kD. Due to the modifications of the SDS-PAGE, these are molecular masses of whole enzymes, i.e. oligomers were not separated into subunits.


Molecular masses were also calculated by elution volumes of the size exclusion purification step in comparison with marker proteins. The native enzymes revealed apparent molecular masses of 70, 155, and 38 kD for pmPOX1, pmPOX2a, and pmPOX2b, respectively, confirming results obtained by gel electrophoresis and suggesting the presence of three distinct peroxidases at the plant PM. On the other hand, the separation of pmPOX2 into two peroxidase peaks by size exclusion chromatography could be due to proteins that were not fully solubilized and remained as aggregates (i.e. protein detergent or protein aggregates). However, the data obtained by SDS-PAGE excluded this hypothesis. Known class III peroxidases revealed molecular masses in a range of 28 to 60 kD (Hiraga et al., 2001) and a 70- or a 155-kD protein have not been described for soluble peroxidases from higher plants.

In PM isolated from soybean roots, 38- and 45-kD bands were identified by SDS-PAGE and *heme* staining (Vianello et al., 1997), masses comparable with that found for pmPOX2b (Fig. 3). However, both bands decreased in intensity after partial washing of

the PM vesicles with NaCl, and several apoplastic peroxidases with these molecular masses were identified in different plant species (Hendriks et al., 1991; Melo et al., 1996; De Marco et al., 1999; Blee et al., 2001). The molecular masses of pmPOX1 and pmPOX2a were different compared with the protein bands identified in soybean PM. However, this could be due to the different material.


### Prosthetic Groups

UV/visible absorption spectra of pmPOX1 and pmPOX2 were almost identical and typical for *heme*-containing proteins (e.g. Converso and Fernandez, 1995; Kvaratskhelia et al., 1997). Both pmPOXs exhibited a Soret peak at 416 nm (Fig. 4). In addition to these, the oxidized enzymes showed  $\alpha$ - and  $\beta$ -absorption bands at 607 and 528 nm, respectively. The Soret peak and the  $\alpha$ -band shifted to 425 and 559 nm when the proteins were reduced by sodium dithionite. The  $A_{416}/A_{280}$  values, which are a criterion of purity and *heme* content, were 0.5 and 0.2 for pmPOX1 and pmPOX2, suggesting that the enzymes were not purified to homogeneity, which was also shown by SDS-PAGE. Thus, the  $\alpha$ -absorption band at 607 nm cannot be definitely ascribed to the *heme* group of the peroxidase. The spectra of both pmPOXs more closely resemble those of guaiacol rather than ascorbate peroxidases (Chen and Asada, 1989; Converso and Fernandez, 1995; Kvaratskhelia et al., 1997). Peroxidase staining of the isolated proteins suggests a relatively strong binding of the *heme*




**Figure 3.** *Heme* staining of pmPOX fractions after modified SDS-PAGE. Electrophoresis was performed using a low concentrated SDS-PAGE, i.e. 0.1% (w/v) SDS in all solutions and gels without dithiothreitol or mercaptoethanol. Thus, the oligomers were not separated into their subunits. *Heme*-containing protein bands were visualized by their reaction with the peroxidase substrates tetramethylbenzidine (TMB) and H<sub>2</sub>O<sub>2</sub> as described in "Materials and Methods." Left, pmPOX1 (a) and pmPOX2 (b) are shown after cation exchange chromatography. Further purification of these fractions by size exclusion is presented on the right: pmPOX1 (c), pmPOX2a (d), and pmPOX2b (e). In addition, f shows pmPOX2b treated with 25 mM dithiothreitol. Bars indicate the corresponding molecular masses. After size exclusion chromatography, the PM-bound enzymes had apparent molecular masses of 70 and 40 kD for pmPOX1 and pmPOX2b, whereas pmPOX2a exhibited a broad protein band between 100 and 170 kD.






**Figure 4.** Absorption spectra of partially purified pmPOX1. Samples ( $1.1 \text{ mg protein mL}^{-1}$ ) containing the native enzyme (dashed line) were measured in  $50 \text{ mM}$  sodium phosphate buffer (pH 7.0) with buffer as reference. Ferric enzymes were reduced by the addition of approximately  $1.5 \text{ mM}$  dithionite (straight line). The spectra were measured at  $50 \text{ nm min}^{-1}$ .  $n = 2$  independent preparations showing identical results. The spectra indicate the presence of *heme* groups as the prosthetic group.

groups to the enzymes. Only pmPOX2b could be detected by *heme* staining after treatment with dithiothreitol and revealed the same molecular mass as without reducing compounds (Fig. 3). Thus, pmPOX2b was identified as a monomer. pmPOX1 and pmPOX2a did not reveal any visible band after the same treatment (data not shown). Conformational changes due to the cleavage of disulfide bridges within the molecules possibly resulted in a release of the *heme* groups.




**Figure 5.** Dependence of the guaiacol peroxidase activity of partially purified pmPOX on pH. The rates of guaiacol oxidation were determined under the standard assay conditions except that  $25 \text{ mM}$  sodium acetate (pH 4.0–5.0), MES (pH 5.5–6.5), or HEPES (pH 7.0–8.0) buffers were used. Data presented are average values  $\pm$  SD of  $n = 3$  experiments. ■, pmPOX1; ○, pmPOX2.



**Figure 6.** Dependence of the guaiacol peroxidase activity of purified pmPOX on temperature (Arrhenius plot). The rates of guaiacol oxidation were determined under the standard assay conditions except for temperature. Data presented are average values  $\pm$  SD of  $n = 3$  experiments. ■, pmPOX1; ○, pmPOX2.

#### pH Optimum and Kinetic Studies

The properties of POX, which were separated by cation exchange chromatography, were further characterized. As shown in Figure 5, the highest activity with guaiacol as a substrate was observed between pH 4.5 and 5.5 for pmPOX1, whereas pmPOX2 exhibited a pH optimum in the range of 5.0 to 6.0. With guaiacol as substrate, acidic pH optima have often been reported for the apoplastic peroxidases of several plant species (Hendriks et al., 1991; Melo et al., 1996; Nair and Showalter, 1996). Variations in pH optima could represent efficient regulatory means in vivo to shift optimal conditions from one isoenzyme to another and thereby favor different processes (De Marco et al., 1999).



**Figure 7.** Thermal stability of guaiacol peroxidase activities. Soluble and PM-bound POX were incubated at  $50^\circ\text{C}$  at different time slices. Data presented are average values  $\pm$  SD of  $n = 2$  experiments. ■, pmPOX1; ●, pmPOX2; □, wPOX1; ○, wPOX2.

**Table I.** Guaiacol-dependent activity in the absence or presence of typical peroxidase effectors

Peroxidase activity was measured with the partially purified enzymes after cation exchange chromatography in the presence of 8.26 mM guaiacol and 8.8 mM H<sub>2</sub>O<sub>2</sub> at pH 7.0 as described in "Materials and Methods." Data are given as mean  $\pm$  SD (n).

Substance	Concentration	Peroxidase Activity			
		pmPOX1	pmPOX2	wPOX1	wPOX2
Control		5.2 $\pm$ 0.1 (3)	1.6 $\pm$ 0.1 (3)	12.0 $\pm$ 0.3 (3)	16.8 $\pm$ 0.1 (3)
Control		100.0 $\pm$ 1.5 (3)	100.0 $\pm$ 4.9 (3)	100.0 $\pm$ 2.3 (3)	100.0 $\pm$ 0.3 (3)
KCN	1 mM	n.d. <sup>a</sup> (3)	n.d. (3)	1.2 $\pm$ 1.6 (3)	0.6 $\pm$ 0.8 (3)
Azide	1 mM	10.6 $\pm$ 0.4 (3) <sup>b</sup>	2.9 $\pm$ 0.6 (3) <sup>b</sup>	0.2 $\pm$ 0.3 (2) <sup>b</sup>	7.4 $\pm$ 0.9 (2) <sup>b</sup>
<i>p</i> -Chloromercuribenzoate (pCMB)	50 $\mu$ M	112.5 $\pm$ 7.1 (3)	111.1 $\pm$ 5.7 (3)	99.7 $\pm$ 9.6 (3)	96.4 $\pm$ 4.6 (3)
	200 $\mu$ M	105.7 $\pm$ 2.3 (3)	102.2 $\pm$ 5.0 (3)	101.4 $\pm$ 2.0 (3)	102.1 $\pm$ 8.3 (3)
	1 mM	110.2 $\pm$ 4.3 (3)	106.9 $\pm$ 0.6 (3)	54.5 $\pm$ 10.9 (3)	77.2 $\pm$ 9.1 (3)

<sup>a</sup> n.d., Not detectable. <sup>b</sup> pH 5.0.

The  $K_m$ s of both PM-bound peroxidase activities for guaiacol were comparable (12.2 mM for pmPOX1 and 14.3 mM for pmPOX2, calculated by Eadie-Hofstee plots).  $K_m$  values in a millimolar range are typical for peroxidases with artificial substrates like guaiacol. For instance, soluble peroxidases from kiwi-fruit (*Actinidia deliciosa*) and tomato (*Lycopersicon esculentum*) fruits had  $K_m$  values of 7.4 and 10 mM, respectively (Soda et al., 1991).

#### Temperature Optima and Thermal Stability

At low temperatures the enzyme activity of pmPOX2 was about 2-fold lower compared with pmPOX1 (Fig. 6). The activity of both protein fractions increased with higher temperatures. Although the activity of pmPOX2 more or less continuously increased in the range of 2°C to 51°C, pmPOX1 showed a maximum of activity at 43°C and decreased dramatically afterward.

In a second set of experiments, the thermal stability of soluble and PM-bound peroxidases was investigated (Fig. 7). All enzymes lost between 40% and 50% of their activities within 5 min at 50°C. During an incubation time of 3 h, the guaiacol peroxidase activities decreased exponentially to values between 5.7% and 34.3%. After 3 h, pmPOX1 showed twice the activity of pmPOX2. Most peroxidases from plants and animals seemed to have high temperature optima and show high thermal stabilities (Bakardjieva et al., 1996; Madhavan and Naidu, 2000). Apoplastic, cytosolic, and soluble peroxidases of several plant tissues showed temperature optima between 30°C and 60°C, the most between 50°C and 60°C (Soda et al., 1991; Bakardjieva et al., 1996; Nair and Showalter, 1996; Bernards et al., 1999; Loukili et al., 1999). Due to the fact that pmPOX1 had a lower temperature optima than pmPOX2, the latter enzyme seemed to be more stable (Fig. 6). However, for longer treatments of higher temperatures, pmPOX1 revealed a higher thermal stability (Fig. 7).

#### Effector Studies

As shown in Table I, classical peroxidase inhibitors like potassium cyanide or sodium azide caused a complete loss of the peroxidase activities or decreased the rates more than 90%. These results were consistent with the presence of *heme* groups as prosthetic groups.

The localization of the enzymes at the plant PM suggests that they may be part of the secretory pathway. According to Welinder et al. (1996), pCMB, a sulfhydryl inhibitor, is often used to distinguish between class I and class III peroxidases. As shown in Table I, this inhibitor did not effect the PM-bound activities of pmPOX1 or pmPOX2, indicating that SH groups did not participate in the active center or maintenance of the conformation of the isoenzymes. Thus, the PM-bound peroxidases were probably class III peroxidases. In contrast to the pmPOX, wPOX1 and wPOX2 were slightly inhibited in the presence of 1 mM pCMB.

Both PM-bound peroxidase activities were decreased by distinct concentrations of the lectins concanavalin A (Con A) and wheat germ agglutinin (WGA; Table II), whereas the *Ulex europaeus* agglutinin (UEA1) was without significant effect (data not shown). Inhibition of wPOX1 and wPOX2 was weak and occurred only at higher concentrations of Con A and WGA (Table II). The effects of lectins indicate glycosylation of the enzymes. These results are consistent with the finding of Vianello et al. (1997) that treatment of soybean roots with tunicamycin, an inhibitor of glycoprotein synthesis, reduced the guaiacol peroxidase activity of unwashed PM vesicles by 40%. Due to the possible glycosylation, which was also indicated by diffuse protein bands in SDS gels (Fig. 3), the real molecular masses of all identified proteins may be different from the calculated values. However, the structures of the proteins have to be further elucidated.

**Table II.** The effects of lectins on guaiacol-dependent peroxidase activity of POX

Guaiacol-dependent peroxidase activity was measured at pH 5.0 after cation exchange chromatography. Fractions were measured in absence or presence of different lectins. Pre-incubation was for 3 min. Data are given as mean  $\pm$  SD (*n*). For control rates, see Table I.

Substance	Concentration $\mu\text{g mL}^{-1}$	pmPOX1	pmPOX2	wPOX1	wPOX2
		% of control			
Control		100.0 $\pm$ 1.5 (3)	100.0 $\pm$ 4.9 (3)	100.0 $\pm$ 2.3 (3)	100.0 $\pm$ 0.3 (3)
Con A	1	79.8 $\pm$ 3.3 (4)	79.8 $\pm$ 2.0 (4)	97.9 $\pm$ 5.2 (3)	98.4 $\pm$ 3.6 (3)
	5	n.m. <sup>a</sup>	78.1 $\pm$ 2.8 (3)	111.5 $\pm$ 1.9 (3)	90.5 $\pm$ 4.1 (3)
WGA	1	78.2 $\pm$ 2.4 (3)	85.6 $\pm$ 1.9 (4)	106.1 $\pm$ 9.0 (3)	90.4 $\pm$ 1.9 (3)
	5	81.9 $\pm$ 4.9 (3)	n.m.	88.8 $\pm$ 2.6 (3)	83.6 $\pm$ 3.2 (3)

<sup>a</sup> n.m., Not measured.

$\text{Ca}^{2+}$  reduced the activity of pmPOX2 and wPOX2.  $\text{Mn}^{2+}$  had no effect on pmPOX1 or pmPOX2 (Table III). In contrast to the PM-bound enzymes, many peroxidases exhibit increased activities after treatment with  $\text{Ca}^{2+}$  or  $\text{Mn}^{2+}$  (Gaspar et al., 1982; Van Huystee et al., 1996; Greppin et al., 1999). Calcium probably maintains the conformation of the proteins, whereas  $\text{Mn}^{2+}$  could be involved in regulatory processes (Van Huystee et al., 1996). However, Loukili et al. (1999) characterized plant peroxidases that were not influenced by these ions. Furthermore,  $\text{Mn}^{2+}$  was not detectable in PM from corn roots (Lüthje et al., 1995). On the other hand, unwashed PM from soybean roots showed a 42% increase of activity in the presence of  $\text{CaCl}_2$  (Vianello et al., 1997). Possibly, this increase was caused by soluble peroxidases that were loosely attached to the PM or due to the different plant material.

DMSO had no effect at 0.5% (v/v), the final concentration of DMSO used in experiments with phenolic compounds as effectors (Table III). PM showed 90.4%  $\pm$  0.3% (*n* = 3) peroxidase activity in the presence of 2% (w/v) DMSO. pmPOX1 was not affected by this concentration, whereas pmPOX2 and the washed off peroxidase activities were inhibited. Detergents like Triton X-100 and Triton X-114 induced a decrease or increase of the enzyme activities.

The activities of cell wall-bound and apoplastic peroxidases have often been reported to be stimulated by phenolic substances, like ferulic acid and coumaric acid (Mäder and Füssl, 1982; Lobarzewski et al., 1996; De Marco et al., 1999). As shown in

Table IV, activities of the partially purified peroxidases increased to 220% and 400% of the control in the presence of ferulic acid as a substrate, whereas coumaric acid had no effect on pmPOX2 and wPOX2, and pmPOX1 and wPOX1 were slightly decreased. The phenolic compound propyl gallate inhibited the guaiacol-dependent peroxidase activity of all fractions. In the presence of IAA, the activity of pmPOX2 decreased slightly, whereas all other activities were not affected. The inhibitory effects of propyl gallate and IAA suggest a competition between the substrates, which was further indicated by their substrate specificity.

#### Substrate Specificity

Artificial electron donors were used by pmPOX1 in the following order: *o*-dianisidine > guaiacol > TMB  $\gg$  2,2'-azino-bis(3-ethylbenz-thiazoline-6-sulfonic acid) (ABTS; Table V). In contrast to pmPOX1, pmPOX2 showed a higher affinity for TMB than for guaiacol. Both pmPOXs oxidized natural substrates like phenolic acids and alcohols in the following order: coniferyl alcohol > ferulic acid > coumaric acid. Hydroxycinnamyl alcohol species are used by apoplastic peroxidases to participate in lignin polymerization, whereas hydroxycinnamic acids could be incorporated into suberin (for refs., see Hiraga et al., 2001).

In vitro IAA oxidation by peroxidases has been reported several times (Converso and Fernandez, 1995; Gazaryan and Lagrimini, 1996). This plant hor-

**Table III.** Guaiacol-dependent activity in presence of salts, solvents, or detergents

Guaiacol-dependent peroxidase activity was measured at pH 5.0 after cation exchange chromatography. Data are given as mean  $\pm$  SD (*n*).

Substance	Concentration	pmPOX1	pmPOX2	wPOX1	wPOX2
		% of control			
Control		100.0 $\pm$ 1.5 (3)	100.0 $\pm$ 4.9 (3)	100.0 $\pm$ 2.3 (3)	100.0 $\pm$ 0.3 (3)
$\text{CaCl}_2$	500 $\mu\text{M}$	108.2 $\pm$ 7.4 (4)	85.1 $\pm$ 1.7 (3)	95.0 $\pm$ 0.4 (2)	88.5 $\pm$ 4.8 (2)
$\text{MnCl}_2$	100 $\mu\text{M}$	105.3 $\pm$ 1.2 (3)	107.4 $\pm$ 3.4 (3)	94.4 $\pm$ 5.4 (3)	89.1 $\pm$ 1.3 (3)
	500 $\mu\text{M}$	102.3 $\pm$ 3.6 (3)	101.6 $\pm$ 2.4 (3)	99.0 $\pm$ 3.5 (3)	86.7 $\pm$ 0.5 (3)
Dimethyl sulfoxide (DMSO)	2% (v/v)	104.9 $\pm$ 8.6 (3)	77.0 $\pm$ 2.3 (3)	75.6 $\pm$ 4.9 (2)	78.0 $\pm$ 1.2 (2)
Triton X-100	0.02% (w/v)	98.2 $\pm$ 5.1 (3)	81.3 $\pm$ 6.7 (2)	n.m.	n.m.
Triton X-114	0.02% (w/v)	119.5 $\pm$ 3.5 (2)	119.5 $\pm$ 5.0 (2)	n.m.	n.m.

**Table IV.** *Guaiacol-dependent peroxidase activity in presence of different substrates*

Guaiacol-dependent peroxidase activity of the partially purified enzymes was measured at pH 5.0. Substrates were added to the assay at concentrations as indicated. Data are given as mean  $\pm$  SD (*n*).

Substance	Concentration	pmPOX1	pmPOX2	wPOX1	wPOX2
	$\mu\text{M}$	% of control			
Control		100.0 $\pm$ 1.5 (3)	100.0 $\pm$ 4.9 (3)	100.0 $\pm$ 2.3 (3)	100.0 $\pm$ 0.3 (3)
Coumaric acid	100	86.7 $\pm$ 2.4 (2)	111.0 $\pm$ 0.1 (2)	86.9 $\pm$ 2.8 (2)	101.9 $\pm$ 2.1 (2)
Ferulic acid	100	221.5 $\pm$ 8.2 (3)	402.0 $\pm$ 7.1 (2)	372.7 $\pm$ 3.9 (2)	264.1 $\pm$ 5.8 (2)
Propyl gallate	500	2.4 $\pm$ 0.8 (3)	5.0 $\pm$ 0.1 (3)	7.2 $\pm$ 0.2 (2)	5.7 $\pm$ 0.1 (2)
Indole-3-acetic acid (IAA)	10	98.5 $\pm$ 0.5 (3)	83.2 $\pm$ 0.2 (3)	102.7 $\pm$ 4.8 (3)	99.1 $\pm$ 6.0 (3)

none was used by pmPOX1, pmPOX2, and wPOX1, whereas the auxin was not oxidized by wPOX2 (Table V).

The highest peroxidase activities were reached with coniferyl alcohol as substrate for both pmPOX. Because the accumulation of the enzymes was different, the specific activities of the soluble POX were apparently higher than the specific activities of the pmPOX.

The washed off peroxidase activities could not only be distinguished from the PM-bound POX by their different substrate specificities for phenolic compounds and IAA but also by their ability to oxidize ascorbate (Table V). Only wPOX2 revealed an ascorbate peroxidase activity, suggesting that intracellular or extracellular soluble peroxidases were attached to the PM during the isolation procedure and removed by washing of the membranes. Also, both pmPOXs did not show any ascorbate peroxidase activity in presence of twice the amounts of enzyme into the assay (data not shown). However, the ability to oxidize ascorbate may have been lost during the purification process, as has been described for several ascorbate peroxidases extracted in the absence of ascorbate (Chen and Asada, 1989; Amako et al., 1994). Other cytosolic ascorbate peroxidases are resistant to depletion of ascorbate (Mittler and Zilinskas, 1991; Koshiba, 1993). Vianello et al. (1997)

reported ascorbate peroxidase activities at non-washed plant PM isolated in the absence of ascorbate from soybean roots, confirming the presented results.

In general, pmPOX1 and pmPOX2 showed more properties corresponding to apoplastic than to cytosolic peroxidases. On the other hand, a localization on the outside or inside of the plant PM cannot be concluded by these properties.

#### CONCLUDING REMARKS

The results of the present work demonstrate the presence of at least two distinct PM-bound peroxidase activities in corn roots. Although peroxidases are usually difficult to distinguish due to their similar characteristics (De Marco et al., 1999), both pmPOXs showed definitely distinct properties in dependence on substrate concentration, pH optima, temperature, and effectors. The biochemical characteristics of both activities are typical for class III peroxidases. Thus, it is the first time, to our knowledge, that enzymes of this class have been found with such high molecular masses in plants.

Until now, the physiological function of PM-bound POX is not clear, and several distinct functions have been postulated (Møller and Bérczi, 1986; Askerlund et al., 1987; Pantoja and Willmer, 1988; Ludwig-Müller et al., 1990; Ludwig-Müller and Hilgenberg,

**Table V.** *Substrate specificity of soluble and pmPOX*

Enzyme activities were measured in presence of 8.8 mM H<sub>2</sub>O<sub>2</sub> and given concentrations of common peroxidase substrates at pH 5.0. Data are given as mean  $\pm$  SD (*n*). Guaiacol oxidation rates are shown in Table I.

Substrate	Concentration	Relative Activity			
		pmPOX1	pmPOX2	wPOX1	wPOX2
	<i>mM</i>	%			
Guaiacol	8.26	100.0 $\pm$ 1.5 (3)	100.0 $\pm$ 4.9 (3)	100.0 $\pm$ 2.3 (3)	100.0 $\pm$ 0.3 (3)
ABTS	0.36	1.8 $\pm$ 0.1 (3)	10.9 $\pm$ 0.3 (3)	1.0 $\pm$ 0.4 (3)	7.6 $\pm$ 0.4 (5)
<i>o</i> -Dianisidine	0.127	133.1 $\pm$ 6.1 (4)	206.9 $\pm$ 10.7 (3)	124.9 $\pm$ 9.9 (3)	231.0 $\pm$ 4.6 (3)
TMB	0.083	86.0 $\pm$ 4.4 (3)	179.6 $\pm$ 8.5 (3)	85.6 $\pm$ 3.5 (3)	194.6 $\pm$ 9.4 (4)
Ascorbate	0.5	n.d. (4) <sup>a</sup>	n.d. (4) <sup>a</sup>	n.d. (3) <sup>a</sup>	20.6 $\pm$ 0.9 (3) <sup>a</sup>
Coniferyl alcohol	0.1	225.9 $\pm$ 4.6 (3)	288.3 $\pm$ 9.2 (3)	235.3 $\pm$ 16.4 (3)	177.9 $\pm$ 2.3 (3)
Coumaric acid	0.1	9.5 $\pm$ 0.2 (3)	4.3 $\pm$ 0.1 (3)	n.d. (3)	38.8 $\pm$ 1.6 (3)
Ferulic acid	0.1	71.1 $\pm$ 2.4 (3)	30.1 $\pm$ 1.5 (3)	121.0 $\pm$ 16.4 (3)	59.2 $\pm$ 3.3 (3)
IAA	0.2	60.3 $\pm$ 3.4 (4)	56.1 $\pm$ 4.3 (4)	14.2 $\pm$ 0.8 (3)	n.d. (3)

<sup>a</sup> pH 7.0



1992; De Marco et al., 1995; Zancani et al., 1995; Vianello et al., 1997). Due to the differences observed for pmPOX1 and pmPOX2, it is possible that these enzymes have distinct functions. Most of the possible functions like detoxifying or production of reactive oxygen species as signal mediators or antimicrobial agents at the interface cell wall/PM could be part of defense mechanisms against pathogen infection (Hiraga et al., 2001). A flavonoid-peroxidase reaction as a mechanism for H<sub>2</sub>O<sub>2</sub> scavenging was demonstrated by Yamasaki et al. (1997). Thus, protection and membrane repair mechanisms of the PM may also be possible.

However, the location (cytoplasmic or apoplasmic side of the PM), the binding properties to the PM, and the physiological function of PM-bound POX activities have to be further elucidated.

## MATERIALS AND METHODS

### PMs

PM have been prepared from 5-d-old corn (*Zea mays* L. cv Jet, Saatenunion, Hannover, Germany) roots by phase partitioning as described earlier (L uthje et al., 1998). The final pellet was stored at  $-80^{\circ}\text{C}$  until use.

### Solubilization of Membrane Proteins

Isolated PM were washed according to B erczi and M oller (1998) with minor modifications. PM were incubated in 25 mM sodium acetate-HCl (pH 4.0), 500 mM KCl, 1 mM EDTA, and 0.01% (w/v) Triton X-100 for 30 min at  $4^{\circ}\text{C}$  under continuous stirring to remove peripheral and adsorbed soluble proteins. Washed membranes were pelleted at 105,000g for 45 min at  $4^{\circ}\text{C}$ , resuspended in acetate buffer (25 mM sodium acetate-HCl [pH 4.0] and 1 mM EDTA), and solubilized with CHAPS at a detergent:protein ratio of 30:1 (w/v) in the presence of 0.5 mM aminocaproic acid. After incubation for 1 h at  $4^{\circ}\text{C}$ , solubilized proteins were separated by ultracentrifugation (1 h at 105,000g and  $4^{\circ}\text{C}$ ).

### Protein Purification

Proteins were purified by a combination of cation exchange chromatography and size exclusion using an HPLC-System (AKTA, Amersham Pharmacia Biotech, Freiburg, Germany) with a 10-mL super-loop. All of the following purification steps were performed at  $4^{\circ}\text{C}$ . Solubilized enzymes were applied on an Uno S1 column (HR 5/5, Bio-Rad, Munich) equilibrated with 25 mM sodium acetate-HCl (pH 4.0), 1 mM EDTA, 1% (w/v) glycerol, and 1 mM CHAPS. After loading, the matrix was washed with 10 column volumes of sodium acetate buffer, and bound proteins were eluted by a continuous KCl gradient (0–1 M KCl in sodium acetate buffer, flow rate, 1 mL min<sup>-1</sup>; total volume, 13 column volumes), followed by 2 column volumes of 1 M KCl. Fractions of 1 and 0.5 mL were collected for the flow through and gradient, respectively. Peak fractions of several Uno S runs were combined and concentrated using Centricon YM-10 concentrators (Millipore, Bedford, MA). Concentrated fractions (500  $\mu\text{L}$ ) or calibration proteins (thyroglobulin [669 kD], ferritin [440 kD], catalase [232 kD], aldolase [158 kD], bovine serum albumin [68 kD], horseradish peroxidase [44 kD], and ribonuclease A [13.7 kD], Amersham Pharmacia Biotech) were applied on a Superdex 200 column (HR 10/30, Amersham Pharmacia Biotech) equilibrated with 4 column volumes of phosphate buffer (50 mM Na<sub>3</sub>PO<sub>4</sub> [pH 7.0], 150 mM NaCl, 1 mM CHAPS, and 1 mM EDTA). Proteins were eluted by 1.5 column volumes of buffer. The flow rate was 0.5 mL min<sup>-1</sup>. The fraction size was automatically adjusted between 0.75 and 0.5 mL depending on the absorption ( $\lambda = 280\text{ nm}$ ). Estimates of the molecular masses of native pmPOX were calculated using a semilogarithmic plot of the molecular mass values for the calibration proteins against the elution volumes.

### SDS-PAGE

Successive steps of purification were monitored by SDS-PAGE, which were performed with 11% (w/v) polyacrylamide slab gels according to Laemmli (1970). Protein bands were visualized by the method of Merrill et al. (1984) using a silver staining kit (Bio-Rad).

PAGE for *heme* staining was performed at room temperature by modified SDS-PAGE. The final concentration of SDS was 0.1% (w/v) in all solutions and gels (Trost et al., 2000). Concentrated samples (0.9–5.0  $\mu\text{g}$  of protein) were diluted in loading buffer to final concentrations of 62.5 mM Tris-HCl, 0.1% (w/v) SDS, 10% (w/v) glycerol, and 0.002% (w/v) bromo-phenol blue without reducing compounds, and loaded onto the gels within 30 min without heating. Horseradish peroxidase as a positive control and each sample were loaded twice once on each one-half of the gel. The gels were cut in one-half after the run. One-half of the gel was used for silver staining, whereas the other one-half was stained in the presence of 6.3 mM TMB and 30 mM H<sub>2</sub>O<sub>2</sub> (Thomas et al., 1976). Because the running characteristics of monomers inside the gels were not effected by the lower SDS concentration, molecular mass standards (Broad Range, Bio-Rad) were used according to Laemmli (1970).

### Enzyme Assays

Peroxidase activities were measured as oxidation of guaiacol (8.26 mM,  $\epsilon = 26.6\text{ mm}^{-1}\text{ cm}^{-1}$ ) in the presence of 8.8 mM H<sub>2</sub>O<sub>2</sub> within 2 min. The assay (1 mL) contained 25 mM sodium acetate-HCl (pH 5.0) and 25  $\mu\text{L}$  of fraction. PM vesicles (50  $\mu\text{g}$  of protein) were measured in 25 mM sodium acetate (pH 5.0), 1 mM EDTA, and 0.01% (w/v) Triton X-100. The reaction was started by addition of guaiacol and followed spectrophotometrically (DU 7500, Beckmann, Munich) as the increase of absorption at 470 nm. Rates were corrected by chemical control experiments. The oxidation rates of other substrates were measured as increases or decreases in absorption using the same reaction mixture and assay conditions but with guaiacol replaced by ABTS ( $A_{405}$ ;  $\epsilon = 36.8\text{ mm}^{-1}\text{ cm}^{-1}$ ), coniferyl alcohol ( $A_{265}$ ;  $\epsilon = 7.5\text{ mm}^{-1}\text{ cm}^{-1}$ ), coumaric acid ( $A_{310}$ ;  $\epsilon = 16.6\text{ mm}^{-1}\text{ cm}^{-1}$ ), *o*-dianisidine ( $A_{460}$ ;  $\epsilon = 30.0\text{ mm}^{-1}\text{ cm}^{-1}$ ), ferulic acid ( $A_{310}$ ;  $\epsilon = 16.6\text{ mm}^{-1}\text{ cm}^{-1}$ ), IAA ( $A_{261}$ ;  $\epsilon = 3.2\text{ mm}^{-1}\text{ cm}^{-1}$ ), or tetramethyl-benzidine ( $A_{652}$ ;  $\epsilon = 39.0\text{ mm}^{-1}\text{ cm}^{-1}$ ). The peroxidase activity with ascorbate as the reducing substrate was determined in a reaction mixture containing 50 mM potassium phosphate (pH 7.0), 0.5 mM ascorbate, and 8.8 mM H<sub>2</sub>O<sub>2</sub>. Oxidation of ascorbate was followed by the decrease in  $A_{290}$  ( $\epsilon = 2.8\text{ mm}^{-1}\text{ cm}^{-1}$ ) within 1 min. The pH dependence of peroxidase activities was ascertained in 25 mM sodium acetate (pH 4.0–5.0), MES (pH 5.5–6.5), and HEPES (pH 7.0–8.0), respectively. Phenolic compounds were dissolved in 50% (v/v) DMSO, resulting in a final concentration of 0.5% (v/v) DMSO per assay. Absorption spectra were recorded in quartz cuvettes (1 cm) on a UV/Vis spectrophotometer (Uvikon 943, Kontron Instruments, Milano, Italy) with a scan speed of 50.0 min<sup>-1</sup>. Data presented were calculated with Microcal Origin (version 5.0, Additive GmbH, Friedrichsdorf/TS, Germany).

### ACKNOWLEDGMENTS

The authors appreciate support by Michael B ottger (University of Hamburg, Germany), helpful discussions with Alajos B erczi (Academy of Sciences, Szeged, Hungary), and critical reading of the manuscript by Richard Becket (University of Natal, Scottsville, South Africa).

Received January 13, 2003; returned for revision January 28, 2003; accepted March 3, 2003.

### LITERATURE CITED

- Amako K, Chen GX, Asada K (1994) Separate assays specific for ascorbate peroxidase and guaiacol peroxidase and for the chloroplastic and cytosolic isozymes of ascorbate peroxidase in plants. *Plant Cell Physiol* 35: 497–504
- Askerlund P, Larsson C, Widell S, M oller IM (1987) NAD(P)H oxidase and peroxidase activities in purified plasma membranes from cauliflower inflorescences. *Physiol Plant* 71: 9–19



- Bakardjieva NT, Cristova NV, Cristov K (1996) Reaction of peroxidase from different plant species to increased temperatures and the effect of calcium and zinc ions. In C Obinger, U Burner, R Ebermann, C Penel, H Greppin, eds, Proceedings of the IV International Symposium on Plant Peroxidases: Biochemistry and Physiology. University of Vienna, Austria, and University of Geneva, Switzerland, pp 345–351
- Bérczi A, Möller IM (1998) Characterization and solubilization of residual redox activity in salt-washed and detergent-treated plasma membrane vesicles from spinach leaves. *Protoplasma* **205**: 59–65
- Bérczi A, Möller IM (2000) Redox enzymes in the plant plasma membrane and their possible roles. *Plant Cell Environ* **23**: 1287–1302
- Bernards MA, Fleming WD, Llewellyn DB, Priefer R, Yang X, Sabatino A, Plourde GL (1999) Biochemical characterization of the suberization-associated anionic peroxidase of potato. *Plant Physiol* **121**: 135–146
- Blee KA, Juce SC, Richard G, Zimmerlin A, Davies DR, Bolwell GP (2001) Molecular identification and expression of the peroxidase responsible for the oxidative burst in French bean (*Phaseolus vulgaris* L.) and related members of the gene family. *Plant Mol Biol* **47**: 607–620
- Bunkelmann J, Trelease RN (1996) Ascorbate peroxidase: a prominent membrane protein in oilseed glyoxysomes. *Plant Physiol* **110**: 589–598
- Chen GX, Asada K (1989) Ascorbate peroxidase in tea leaves: occurrence of two isozymes and the differences in their enzymatic and molecular properties. *Plant Cell Physiol* **30**: 987–998
- Converso DA, Fernandez ME (1995) Peroxidase isozymes from wheat germ: purification and properties. *Phytochemistry* **40**: 1341–1345
- Crevecoeur M, Pinedo M, Greppin H, Penel C (1997) Peroxidase activity in shoot apical meristem from spinach. *Acta Histochem* **99**: 177–186
- De Marco A, Guzzardi P, Jamet É (1999) Isolation of tobacco isoperoxidases accumulated in cell-suspension culture medium and characterization of activities related to cell wall metabolism. *Plant Physiol* **120**: 371–382
- De Marco A, Pinton R, Fischer-Schliebs E, Varanini Z (1995) Possible interaction between peroxidase and NAD(P) H-dependent nitrate reductase activities of plasma membranes of corn roots. *J Exp Bot* **46**: 1677–1683
- Gaspar T, Penel C, Thorpe T, Greppin H, editors (1982) Peroxidases 1970–1980: A Survey of their Biochemical and Physiological Roles in Higher Plants. University of Geneva Press, Switzerland
- Gazaryan IG, Lagrimini LM (1996) Tobacco anionic peroxidase overexpressed in transgenic plants: aerobic oxidation of indole-3-acetic acid. *Phytochemistry* **42**: 1271–1278
- Greppin H, Wiater RG, Ginalska G, Lobarzewski J (1999) The cabbage peroxidase isoforms changes influenced by Ca<sup>2+</sup> and Mg<sup>2+</sup> ions. *Plant Peroxidase Newslett* **13**: 129–135
- Hendriks T, Wijnsman HJ, Van Loon LC (1991) Petunia peroxidase a: isolation, purification and characteristics. *Eur J Biochem* **199**: 139–146
- Hiraga S, Sasaki K, Ito H, Ohashi Y, Matsui H (2001) A large family of class III plant peroxidases. *Plant Cell Physiol* **42**: 462–468
- Hu C, Smith R, Van Huystee R (1989) Biosynthesis and localization of peanut peroxidases: a comparison of the cationic and the anionic isozymes. *Plant Physiol* **135**: 391–397
- Jespersen HM, Kjærsgård IVH, Østergaard L, Welinder KG (1997) From sequence analysis of three novel ascorbate peroxidases from *Arabidopsis thaliana* to structure, function and evolution of seven types of ascorbate peroxidase. *Biochem J* **326**: 305–310
- Koshiha T (1993) Cytosolic ascorbate peroxidase in seedlings and leaves of maize (*Zea mays*). *Plant Cell Physiol* **34**: 713–721
- Kvaratskhelia M, Winkler C, Thorneley RNF (1997) Purification and characterization of a novel class III peroxidase isoenzyme from tea leaves. *Plant Physiol* **114**: 1237–1245
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**: 680–685
- Lin W (1982) Responses of corn root protoplasts to exogenous reduced nicotinamide adenine dinucleotide: oxygen consumption, ion uptake and membrane potential. *Proc Natl Acad Sci USA* **79**: 3773–3776
- Lobarzewski J, Brzyska M, Greppin H (1996) The fungal peroxidase kinetics with some phenolics hydrogen donors in relation to lignin degradation. In C Obinger, U Burner, R Ebermann, C Penel, H Greppin, eds, Proceedings of the IV International Symposium on Plant Peroxidases: Biochemistry and Physiology. University of Vienna, Austria, and University of Geneva, Switzerland, pp 153–156
- Loukili A, Limam F, Ayadi A, Boyer N, Ouelhazi L (1999) Purification and characterization of a neutral peroxidase induced by rubbing tomato internodes. *Physiol Plant* **105**: 24–31
- Ludwig-Müller J, Hilgenberg W (1992) Tryptophan oxidizing enzyme and basic peroxidase isoenzymes in *Arabidopsis thaliana* (L.) Heynh.: are they identical? *Plant Cell Physiol* **33**: 1115–1125
- Ludwig-Müller J, Rausch T, Lang S, Hilgenberg W (1990) Plasma membrane bound high pI peroxidase isoenzymes convert tryptophan to indole-3-acetaldoxime. *Phytochemistry* **29**: 1397–1400
- Lüthje S, Niecke M, Böttger M (1995) Iron and copper in plasma membranes of maize (*Zea mays* L.) roots investigated by proton induced X-ray emission. *Protoplasma* **184**: 145–150
- Lüthje S, Van Gestelen P, Córdoba-Pedregosa MC, González-Reyes JA, Asard H, Villalba JM, Böttger M (1998) Quinones in plant plasma membranes: a missing link? *Protoplasma* **205**: 43–51
- Mäder M, Füssl R (1982) Role of peroxidase in lignification of tobacco cells: II. Regulation by phenolic compounds. *Plant Physiol* **70**: 1132–1134
- Madhavan ND, Naidu KA (2000) Purification and partial characterization of peroxidase from human term placenta of non-smokers: metabolism of benzo(a) pyrene-7,8-dihydrodiol. *Placenta* **21**: 501–509
- Melo NS, Calvete JJ, Thole HH, Welinder KG, Töpfer-Petersen E, Ferevereiro PS (1996) Cationic peroxidases from *Vaccinium myrtillus* cell suspension cultures. Preliminary sequence results. In C Obinger, U Burner, R Ebermann, C Penel, H Greppin, eds, Proceedings of the IV International Symposium on Plant Peroxidases: Biochemistry and Physiology. University of Vienna, Austria, and University of Geneva, Switzerland, pp 217–221
- Merrill CR, Goldman D, Van Keuren ML (1984) Gel protein stains: silver stain. *Methods Enzymol* **104**: 441–447
- Mittler R, Zilinskas A (1991) Purification and characterization of pea cytosolic ascorbate peroxidase. *Plant Physiol* **97**: 962–968
- Möller IM, Bérczi A (1986) Salicylhydroxamic acid-stimulated NADH oxidation by purified plasmalemma vesicles from wheat roots. *Physiol Plant* **68**: 67–74
- Nair AR, Showalter AM (1996) Purification and characterization of a wound-inducible cell wall cationic peroxidase from carrot roots. *Biochem Biophys Res Commun* **226**: 254–260
- Nito K, Yamaguchi K, Kondo M, Hayashi M, Nishimura M (2001) Pumpkin peroxisomal ascorbate peroxidase is localized on peroxisomal membranes and unknown membranous structures. *Plant Cell Physiol* **42**: 20–27
- Pantoja O, Willmer CM (1988) Redox activity and peroxidase activity associated with the plasma membrane of guard-cell protoplasts. *Planta* **174**: 44–50
- Penel C, Castillo FJ (1991) Peroxidases of plant plasma membranes, apoplastic ascorbate, and relation of redox activities to plant pathology. In FL Crane, DJ Morrè, H Loew, eds, Oxidoreduction at the Plasma Membrane, Vol II. CRC Press, Boca Raton, FL, pp 121–147
- Penel C, Gaspar T, Greppin H (1992) Plant Peroxidases: 1980–1990: Topics and Detailed Literature on Molecular, Biochemical, and Physiological Aspects. University of Geneva, Switzerland
- Sagi M, Fluhr R (2001) Superoxide production by plant homologues of the gp91(phox) NADPH oxidase: modulation of activity by calcium and by tobacco mosaic virus infection. *Plant Physiol* **126**: 1281–1290
- Shigeoka S, Ishikawa T, Tamoi M, Miyagawa Y, Takeda T, Yabuta Y, Yoshimura K (2002) Regulation and function of ascorbate peroxidase isoenzymes. *J Exp Bot* **53**: 1305–1319
- Soda I, Hasegawa T, Suzuki T, Ogura N (1991) Purification and some properties of peroxidase from kiwifruit. *Agric Biol Chem* **55**: 1677–1678
- Thomas PE, Ryan D, Levin W (1976) An improved staining procedure for the detection of the peroxidase activity of cytochrome P-450 on sodium dodecyl sulfate polyacrylamide gels. *Anal Biochem* **75**: 168–176
- Trost P, Berczi A, Sparda F, Sponza G, Marzadori B, Asard H, Pupillo P (2000) Purification of cytochrome b-561 from bean hypocotyls plasma membrane: evidence for the presence of two heme centers. *Biochim Biophys Acta* **1468**: 1–5
- Van Huystee RB, Rodriguez Marañón MJ, Wan L (1996) Peanut peroxidase, a trimetal glycoprotein. In C Obinger, U Burner, R Ebermann, C Penel, H Greppin, eds, Proceedings of the IV International Symposium on Plant Peroxidases: Biochemistry and Physiology. University of Vienna, Austria, and University of Geneva, Switzerland, pp 42–44

- Vianello A, Zancani M, Macri F** (1990) Hydrogen peroxide formation and iron ion oxidoreduction linked to NADH oxidation in radish plasmalemma vesicles. *Biochim Biophys Acta* **1023**: 19–24
- Vianello A, Zancani M, Nagy G, Macri F** (1997) Guaiacol peroxidase associated to soybean root plasma membranes oxidizes ascorbate. *J Plant Physiol* **150**: 573–577
- Welinder KG, Jespersen HM, Kjærsgård IVH, Ostergaard L, Abelskov AK, Hansen LN, Rasmussen SK** (1996) What can we learn from *Arabidopsis* peroxidases? *In* C Obinger, U Burner, R Ebermann, C Penel, H Greppin, eds, Proceedings of the IV International Symposium on Plant Peroxidases: Biochemistry and Physiology. University of Vienna, Austria, and University of Geneva, Switzerland, pp 173–178
- Yamasaki H, Sakihama Y, Ikehara N** (1997) Flavonoid-peroxidase reaction as a detoxification mechanism of plant cells against H<sub>2</sub>O<sub>2</sub>. *Plant Physiol* **115**: 1405–1412
- Zancani M, Nagy G, Vianello A, Macri F** (1995) Copper-inhibited NADH-dependent peroxidase activity of purified soya bean plasma membranes. *Phytochemistry* **40**: 367–371