

of 25 000 are expected.¹ New legislation may be needed.

This report was compiled on behalf of the staff at the Northern General Hospital, Sheffield, the staff at the Royal Hallamshire Hospital, Sheffield, and the many people who helped at the scene. The efforts of hundreds of professionals and volunteers were vital and greatly appreciated. We thank Neil Appleyard, David Edbrooke, David Dawson, Stuart Yates, Ian Winston, John Duncan, G A Baker, Charlie Cooper, Kath Sherry, Tim Shaw, and A Moss for access to the neuropsychological reports on some of the survivors.

- 1 Taylor P. *The Hillsborough stadium disaster. Interim report.* London: HMSO, 1989.
- 2 Shapiro HM. Neurosurgical anaesthesia and intracranial hypertension. In: Miller RD, ed. *Anaesthesia.* 2nd ed. Vol 2. Edinburgh: Churchill Livingstone, 1986:1573.
- 3 Landercasper J, Cogbill T. Long-term follow up after traumatic asphyxia. *J Trauma* 1985;25:838-41.
- 4 Ollivier d'A. Relation medicale des evenement survenus au Champs-de-Mars le 14 juin, 1837. *Annales d'Hygiene* 1837;18:486-9.
- 5 Sandiford JA, Sickler D. Traumatic asphyxia with severe neurological sequelae. *J Trauma* 1974;14:805-9.

- 6 Fred HL, Chandler FW. Traumatic asphyxia. *Am J Med* 1960;29:508-17.
- 7 Gorenstein L, Blair GK, Shandling B. The prognosis of traumatic asphyxia in childhood. *J Pediatr Surg* 1986;21:753-6.
- 8 Chaner KS, Edbrooke DL, Moores M, McHugh P, Michael S. Acute right heart strain after crushing injury at Hillsborough football stadium. *BMJ* 1989;299:1379-80.
- 9 Williams JS, Minken SL, Adams JT. Traumatic asphyxia reappraised. *Ann Surg* 1968;167:384-92.
- 10 Thompson A, Illescas FF, Chiu RC-J. Why is the lower torso protected in traumatic asphyxia? A new hypothesis. *Ann Thorac Surg* 1989;47:247-9.
- 11 Plum F, Posner JB. *The diagnosis of stupor and coma.* 3rd ed. Philadelphia: FA Davis, 1981.
- 12 Plum F, Posner JB, Hain RF. Delayed neurological deterioration after anoxia. *Arch Intern Med* 1962;110:18-25.
- 13 Lance J, Adams RD. The syndrome of intention or action myoclonus as a sequel to hypoxic encephalopathy. *Brain* 1963;86:111-35.
- 14 Staff of the Accident and Emergency Departments of Derbyshire Royal Infirmary, Leicester Royal Infirmary, Queen's Medical Centre, Nottingham. Coping with the early stages of the M1 disaster: at the scene and on arrival at hospital. *BMJ* 1989;298:651-4.
- 15 Sharpe DT, Roberts AHN, Barclay TL, Dickson WA, Settle JAD, Crockett DJ, et al. Treatment of burn casualties after the fire at Bradford City football ground. *BMJ* 1985;291:945-8.
- 16 NHS Management Executive. *Emergency planning in the NHS: health services arrangements for dealing with major incidents.* London: NHS Management Executive, 1990.

(Accepted 28 August 1991)

Doctor-patient communication: the Toronto consensus statement

Pretoria, South Africa
Michael Simpson,
MRCPsych, consultant
psychiatrist

Toronto-Bayview Regional
Cancer Centre, University
of Toronto, Toronto,
Ontario, Canada
Robert Buckman, FRCP,
medical oncologist

Centre for Studies in
Family Medicine,
University of Western
Ontario, London, Ontario,
Canada N6A 5C1
Moira Stewart, PhD,
professor of family medicine

Cancer Research
Campaign Psychological
Medicine Group, Christie
Hospital, Manchester M20
9BX
Peter Maguire, FRCPsych,
director

Department of Medicine,
New York University
School of Medicine, New
York, USA
Mack Lipkin, MD, director,
division of primary care

Brown University School of
Medicine, Providence,
Rhode Island, USA
Dennis Novack, MD,
associate professor of
community health

Ontario Cancer Institute,
Princess Margaret Hospital
Toronto, Ontario, Canada
James Till, PhD, senior
scientist

Correspondence to: Dr
Stewart.

BMJ 1991;303:1385-7

Michael Simpson, Robert Buckman, Moira Stewart, Peter Maguire, Mack Lipkin, Dennis Novack, James Till

Effective communication between doctor and patient is a central clinical function that cannot be delegated. Most of the essential diagnostic information arises from the interview, and the physician's interpersonal skills also largely determine the patient's satisfaction and compliance and positively influence health outcomes.¹⁻³ Such skills, including active listening to patients' concerns, are among the qualities of a physician most desired by patients.⁴ Increasing public dissatisfaction with the medical profession is, in good part, related to deficiencies in clinical communication. Studies in many countries have confirmed that serious communication problems are common in clinical practice.

This consensus statement addresses three issues: What are the most important facts we already know about doctor-patient communication? What are the most important things that could be done now to improve the situation? and, What are the most important unanswered questions?

What are the most important facts we already know about doctor-patient communication?

PROBLEMS IN PRACTICE

Communication problems in medical practice are both important and common. For example, 54% of patient complaints and 45% of patient concerns are not elicited by physicians.⁵ Psychosocial and psychiatric problems are common in general medical practice, but these diagnoses are missed in up to 50% of cases.^{6,7} In 50% of visits the patient and the doctor do not agree on the nature of the main presenting problem.^{8,9} In one study patients were interrupted by physicians so soon after they began describing their presenting problems (on average within 18 seconds) that they failed to disclose other significant concerns.¹⁰ Most complaints by the public about physicians deal not with clinical competency problems, but with communication problems,¹¹ and the majority of malpractice allegations arise from communication errors.¹² Residents or trainees¹³ and practising physicians¹⁴ have shown

substantial deficiencies when studied. Only a low proportion of visits with doctors include any patient education,¹⁵ and a surprisingly high proportion of patients do not understand or remember what their physicians tell them about diagnosis and treatment.¹⁶ Cultural differences also impede the work with patients.^{17,18}

Patient anxiety and dissatisfaction is related to uncertainty and lack of information, explanation, and feedback from the doctor. Yet doctors often misperceive the amount and type of information patients want. The language doctors use is often unclear, both as regards the use of jargon and in relation to a lack of the expected shared meanings of relatively common terms.²⁰⁻²³

COMMUNICATION PRACTICES IN RELATION TO OUTCOMES

The quality of clinical communication is related to positive health outcomes.^{24,25} Reduction in blood pressure was significantly greater in patients who, during visits to the doctor, had been allowed to express their health concerns without interruptions.²⁶ Concordance between physician and patient in identifying the nature and seriousness of the clinical problem is related to improving or resolving the problem.^{5,9,27}

Explaining and understanding patient concerns, even when they cannot be resolved, results in a significant fall in anxiety.²⁸ Greater participation by the patient in the encounter improves satisfaction and compliance¹ and outcome of treatment (for example, control of diabetes and hypertension).²⁴ The level of psychological distress in patients with serious illness is less when they perceive themselves to have received adequate information.^{29,30}

Beneficial clinical communication is feasible routinely in clinical practice and can be achieved during normal clinical encounters, without unduly prolonging them, provided that the clinician has learned the relevant techniques.^{31,32}

EDUCATIONAL ISSUES

To become effective communicators, physicians

must master a defined body of knowledge, skills, and attitudes.³³ Clinical communication skills do not reliably improve from mere experience. Examples of relevant areas of knowledge are psychiatry in relation to medicine (for example, diagnostic clues to depression, anxiety, somatisation problems) and the structure and functions of medical interviewing. Examples of skills needed within the interview are those of data gathering, forming and maintaining relationships, dealing with difficult issues (such as sexual history, breaking bad news, HIV), and imparting information; therapeutic skills and strategies are also necessary.³⁴⁻³⁸ These skills can be defined with behavioural criteria and can be reliably taught and assessed.³⁴⁻³⁹⁻⁴⁰ Helpful attitudes include a belief in the importance of a biopsychosocial perspective⁴¹ as well as an unconditional positive regard for patients.⁴² A physician's personal growth and self awareness are essential bases of effective communication.

Unfortunately, traditional medical education at all levels is generally ineffective in teaching clinical communication.⁴³⁻⁴⁵ Medical education is a stressful and sometimes abrasive experience that can produce cynicism and callousness.⁴⁶ Surveys of American and British medical schools showed that, although some important advances have occurred, there is extensive variability in the quality and intensity of courses offered.⁴⁷⁻⁴⁸

Much is known about the elements of an effective course on clinical communication. Highly structured programmes, in which specific skills are identified, demonstrated, practised, and evaluated, tend to be more effective than less structured programmes.⁴⁹⁻⁵⁰ Courses should include faculty development, quality control of teaching, and relatively low student to teacher ratios, providing students with multiple opportunities for practice and feedback. Use of videotape and audiotape reviews, role play, and standardised patients have been proved to be effective tools.³³⁻⁴⁰ Communication skills training should be taught at all levels of the curriculum: further development of skills is needed as students encounter more complex situations, and interpersonal skills, like other skills, benefit from reinforcement.⁵¹ Such training can be enhanced by partnership between clinicians and others—for example, behavioural and social scientists. The effects of such training on specific skills can persist over time⁵²⁻⁵³ and this effect is likely to be greater if there is coordination of teachers and courses at all levels. Coherent attention needs to be given to students' emotional issues, such as working with dying patients and anger at self inflicted ill health.⁵⁴⁻⁵⁵ Balint groups and other support and discussion groups have proved useful in this context.⁵⁶⁻⁵⁷

Few medical education programmes require demonstrations of competency in communication for graduation or certification. Though further development of valid and reliable assessments of communication competencies is anticipated⁵⁸ communication skills can already be assessed with useful accuracy with current evaluation techniques⁵⁹ to establish minimal standards of competence.⁶⁰

Each institution therefore requires faculty members capable of leading education for each clinical stage. The field of doctor-patient communication needs faculty development. The faculty development approach of the American Task Force on the Doctor and Patient combines didactic skills, learning skills, practice, task orientation, and a focus on personal response to the work and is an effective model. Another model with documented efficacy is the Manchester approach, which uses videotape feedback to prepare British general practice trainers to teach communication skills to their trainees.³⁷ Faculty members need access to training in research methods.

What are the most important things that could be done now to improve clinical communication by doctors?

Physicians should first encourage patients to discuss their main concerns without interruption or premature closure. This enhances satisfaction and efficacy of the consultation—yet, contrary to the expectation of many doctors, this need not take long: a maximum of 2½ minutes, or an average of 90 seconds.¹⁰ Doctors should also strive to elicit patients' preceptions of the illness and associated feelings and expectations.¹⁷⁻⁶¹⁻⁶⁵

Experience also supports the value of learning methods of active listening and empathy.³⁹⁻⁴¹⁻⁴² The appropriate use of open ended questions, frequent summaries, clarification, and negotiation are factors that positively affect the quality and quantity of information gathered; factors with a negative impact include inappropriate use of closed ended questions and premature advice or reassurance. Other important skills include giving clear explanations, checking the patient's understanding, negotiating a treatment plan,⁶⁶ and checking patients' attention to compliance.⁶⁷

To promote acquisition of these skills, facilities for videotaping and audiotaping and reviewing consultations should be provided. Institutions should consider requiring and facilitating such skills training with medical schools, hospitals, and private practices and within existing journal clubs, courses, and continuing medical education programmes. Licensing and specialty qualifying authorities should require proof of competence in clinical communications.

What are the most important unanswered questions and priorities?

QUESTIONS ABOUT CLINICAL ENCOUNTERS

What are the specific elements of communication that maximise patient satisfaction, collaborative autonomy, quality of life, enhancement of coping, and adaptation and recovery or rehabilitation while minimising conflict and litigation? Although a great deal is known about the early phases of the interview, less is known about information exchange and therapeutic strategies.

What physician, patient, family, and practice variables affect the encounter, and how do they interact? For example, what specific personal qualities of the physician are most influential? What patient attitudes, preferences, and beliefs are most important? How do health care and payment systems influence communication practices—and how should they be designed so as to optimise communication?

Relationships may vary according to the degree of control and involvement the patient prefers.⁶⁸ What are the influential dimensions and mechanisms of effective active participation by patients?

How do communication needs and practices vary in the course of continuing clinical care? Also, we need to explore the ethical aspects of communication and communication aspects of ethics.

QUESTIONS ABOUT EDUCATION ISSUES

A wide range of effective teaching methods and media have been developed—which are most effective and efficient for teaching which skills and strategies?

How does the stage of training or clinical experience of the learner affect optimal teaching methods and sequencing? How can teachers most effectively help learners identify their resistances to learning to encounter the full extent of their patients' problems, and how can such resistances be overcome?

Are all communication skills teachable? Which skills are the most and least teachable? How can students' learning styles be best assessed so that training can be most effectively tailored to individual needs?

How can educational, health care, and reimbursement systems be best adapted so as to motivate and reward faculty or trainers and practitioners to learn and teach these skills?

How can doctors best continue to develop their skills, apply them within their daily work, survive emotionally, and feel more satisfied? How can persistent behavioural, perceptual, and personal changes be produced? Self learning and self monitored feedback, distance learning, and serial workshops seem to be promising approaches for qualified doctors.

QUESTIONS ABOUT RESEARCH METHODS

Complementary development of alternative methods will allow the identification and description of aspects of clinical communication not already recognised or studied. Qualitative methods need to be encouraged, to complement the now standard quantitative and interaction analysis measures.

Conclusion

Sufficient data have now accumulated to prove that problems in doctor-patient communication are extremely common and adversely affect patient management. It has been repeatedly shown that the clinical skills needed to improve these problems can be taught and that the subsequent benefits to medical practice are demonstrable, feasible on a routine basis, and enduring. There is therefore a clear and urgent need for teaching of these clinical skills to be incorporated into medical school curriculums and continued into postgraduate training and courses in continuing medical education. If current knowledge is now implemented in clinical practice, and if the priorities for research are addressed, there may be material improvement in the relationship between patient and doctor.

- Roter DL. Patient participation in patient provider interaction: the effects of patient question-asking on the quality of interaction, satisfaction, and compliance. *Health Education Monographs* 1977;5:281-315.
- DiMatteo MR, DiNicola DD. *Achieving patient compliance—the psychology of the medical practitioner's role*. New York: Pergamon Press, 1982.
- Bartlett EE, Gravson M, Barker R. The effects of physician communication skills on patient satisfaction, recall and adherence. *J Chronic Dis* 1984;37:755-64.
- Matthews DA, Sledge WH, Lieberman PB. Evaluation of intern performance by medical inpatients. *Am J Med* 1987;35:936-44.
- Stewart MA, McWhinney IR, Buck CW. The doctor-patient relationship and its effect upon outcome. *J R Coll Gen Pract* 1979;29:77-82.
- Schulberg HC, Burns BJ. Mental disorders in primary care. *Gen Hosp Psychiatry* 1988;10:79-87.
- Freeling P, Rao BM, Paykel ES, Sireling LI, Burton RH. Unrecognised depression in general practice. *BMJ* 1985;290:1880-3.
- Starfield B, Steinwachs D, Morris I, Bause G, Siebert S, Westin C. Patient-doctor agreement about problems needing follow-up visit. *JAMA* 1979;242:344-6.
- Starfield B, Wray C, Hess K, Gross R, Birk PS, D'Lugoff BC. The influences of patient-practitioner agreement on outcome of care. *Am J Public Health* 1981;71:127-32.
- Beekman HB, Frankel RM. The effect of physician behaviour on the collection of data. *Ann Intern Med* 1984;101:692-6.
- Richards T. Chasms in communication. *BMJ* 1990;301:1407-8.
- Shapiro RS, Simpson DE, Lawrence SL, Talsky AM, Sobocinski KA, Schiedermayer DL. A survey of sued and nonsued physicians and suing patients. *Arch Intern Med* 1989;149:2190-6.
- Platt FW, McMath JC. Clinical incompetence: the interview. *Ann Intern Med* 1979;91:898-902.
- Byrne PS, Long BEL. *Doctors talking to patients*. London: Royal College of General Practitioners 1984.
- Waitzkin H. Doctor-patient communication: clinical implications of social scientific research. *JAMA* 1984;252:2441-6.
- Ley P. *Communication with patients: improving satisfaction and compliance*. London: Croom-Helm, 1988.
- Kleinman A, Eisenberg L, Good B. Culture, illness and care: clinical lessons from anthropologic and cross-cultural research. *Ann Intern Med* 1978;88:251-8.
- Waxler-Morrison N, Anderson JM, Richardson E, eds. *Cross-cultural caring: a handbook for health professionals in western Canada*. Vancouver: University of British Columbia Press, 1990.
- Faden RR, Becker C, Lewis C, Freeman J, Faden AI. Disclosure of information to patients in medical care. *Med Care* 1981;19:718-33.
- Mackillop WJ, Stewart WE, Ginsberg AD, Stewart SS. Cancer patients' perceptions of disease and its treatment. *Br J Cancer* 1988;58:355-8.
- Frances V, Korsch BM, Morris MJ. Gaps in doctor-patient communication: patient response to medical advice. *N Engl J Med* 1969;280:535-40.
- Simpson MA. *Clinical psycholinguistics: the languages of illness and healing*. New York: Irvington, 1980.
- Kaplan SH, Greenfield S, Ware JE. Assessing the effects of physician-patient

interactions on the outcomes of chronic disease. *Med Care* 1989;27 (suppl 3):S110-27.

- Headache Study Group of the University of Western Ontario. Predictors of outcome in headache patients presenting to family physicians—a one year prospective study. *Headache J* 1986;26:285-94.
- Orth JE, Stiles WB, Scherwitz L, Hennritus D, Vallbona C. Patient exposition and provider explanation in routine interviews and hypertensive patients' blood pressure control. *Health Psychol* 1987;6:29-42.
- Bass MJ, Buck C, Turner L, Dickie G, Pratt G, Robinson HC. The physician's actions and the outcome of illness in family practice. *J Fam Pract* 1986;23:43-7.
- MacLeod R. Patients with advanced breast cancer: the nature and disclosure of their concerns (dissertation). Manchester: University of Manchester, 1991.
- Fallowfield LJ, Baum M, Maguire GP. Effects of breast conservation on psychological morbidity associated with diagnostic and treatment of early breast cancer. *BMJ* 1986;293:1331-4.
- Fallowfield LJ, Hall A, Maguire GP, Baum M. Psychological outcomes of different treatment policies in women with early cancer outside a clinical trial. *BMJ* 1990;301:575-80.
- Greco RS, Pittenger RA. *One man's practice: effects of developing insight on doctor-patient transactions*. London: Tavistock, 1966.
- Stewart MA, Brown JB, Weston WW. Patient-centred interviewing part III: five provocative questions. *Canadian Family Physician* 1989;35:159-61.
- Lipkin M Jr, Quill TE, Napedano RJ. The medical interview: a core curriculum. *Ann Intern Med* 1984;100:277-84.
- Cohen-Cole SA. *The medical interview: a three function approach*. St Louis: Mosby, 1991.
- Lazare A, Lipkin M, Putnam SM. The functions of the medical interview. In: Lipkin M, Putnam SM, Lazare A, eds. *The medical interview*. New York: Springer-Verlag (in press).
- Novack DH. Therapeutic aspects of the clinical encounter. *J Gen Intern Med* 1987;2:346-55.
- Gask L, Goldberg D, Boardman A. Training general practitioners to teach psychiatric interviewing skills: an evaluation of group training. *Medical Education* 1991;25:444-51.
- Gask L. Evaluation of the impact of improving the psychiatric skills of general practitioners on patient satisfaction and outcome of care (thesis). Manchester: University of Manchester, 1991.
- Lipkin M. Medical interview and related skills. In: Branch WT, ed. *The office practice of medicine*. Philadelphia: WB Saunders, 1987:1287-306.
- Maguire P. Can communication skills be taught? *Br J Hosp Med* 1990;43:215-6.
- Engel GL. The need for a new medical model: a challenge for biomedicine. *Science* 1977;196:129-36.
- Rogers C. *Client-centered therapy—its current practice implications and theory*. Cambridge, Massachusetts: Riverside Press, 1951.
- Kahn GS, Cohen B, Jason H. The teaching of interpersonal skills in US medical schools. *J Med Educ* 1979;54:29-35.
- Kern DE, Grayson M, Barker LR, Roca RP, Cole KA, Roter D, Golden AS. Residency training in interviewing skills and the psychosocial domain of medical practice. *J Gen Intern Med* 1989;4:421-31.
- Maguire P. *Communication skills and patient care. Health care and human behavior*. New York: Academic Press, 1984:153-73.
- Simpson MA. *Medical education: a critical approach*. London: Butterworths, 1972.
- Novack DH, Volk G, Drossman DA, Putnam S, Lipkin M Jr. The teaching of medical interviewing and interpersonal skills: a survey of US medical schools. *Personal Communication*, 1991.
- Whitehouse CR. The teaching of communication skills in United Kingdom medical schools. *Medical Education* 1991;25:311-8.
- Hutter MJ, Dungy CI, Zakus GE, Moore DJ, Oh JE, Favret AC. Interviewing skills: a comprehensive approach to teaching and evaluation. *J Med Educ* 1977;52:328-33.
- Carroll JG, Monroe J. Teaching medical interviewing: a critique of educational research and practice. *J Med Educ* 1979;54:498-500.
- Engler CM, Saltzman GA, Walker ML, Wolf FM. Medical student acquisition and retention of communication and interviewing skills. *J Med Educ* 1981;56:572-9.
- Maguire P, Fairbairn S, Fletcher C. Consultation skills of young doctors: benefits of feedback training in interviewing as students persist [correction appears in *BMJ* 1986;293:26]. *BMJ* 1986;292:1573-6.
- Gask L, Goldberg D, Lesser AL, Millar T. Improving psychiatric skills of the general practice trainee: an evaluation of a group training course. *Medical Education* 1988;22:132-8.
- Wilkinson S. Factors which influence how nurses communicate with cancer patients. *J Advanced Nurs* 1991;16:677-68.
- Smith AC, Kleinman S. Managing emotions in medical school: students' contacts with the living and the dead. *Social Psychology Quarterly* 1989;52:56-69.
- Balint M. *The doctor, his patient, and the illness*. New York: International University Press, 1972.
- Reuben DB, Novack DH, Wachtel TJ, Wartman SA. A comprehensive support system for reducing house staff distress. *Psychosomatics* 1984;25:815-20.
- Stillman PL, Swanson DB, Smee S, Stillman AE, Ebert TH, Emmel VS, et al. Assessing clinical skills of residents with standardized patients. *Ann Intern Med* 1986;105:762-71.
- Overdia AB, Yager J, Heinrich RL. Assessment of medical interview skills in teaching psychiatry and behavior science. In: J Yager, ed. New York: Grunc and Stratton, 1982:485-500.
- Maguire P, Roe P, Goldberg D. The value of feedback in teaching interviewing skills to medical students. *Psychol Med* 1978;8:695-704.
- Pendleton D, Schofield T, Tate P, Havelock P. *The consultation: an approach to learning and teaching*. Oxford: Oxford University Press, 1984.
- Levenstein JH, Brown JB, Weston WW. The patient-centered clinical method: a model for the doctor-patient interaction in family medicine. In: Stewart M, Roter D, eds. *Communicating with medical patients*. Newbury Park, California: Sage, 1989:107-20.
- Weston WW, Brown JB, Stewart MA. Patient-centred interviewing part I: understanding patients' experiences. *Canadian Family Physician* 1989;35:147-51.
- Lassen LC. Connections between the quality of consultations and patients' compliance in general practice. *Fam Pract* 1991;8:154-60.
- Tuckett D, Boulton M, Olsen C, Williams A. *Meetings between experts*. London: Tavistock, 1986.
- Riccardi DM, Kurtz SM. *Communication and counselling in health care*. Springfield, Illinois: Charles C Thomas, 1987.
- Brown JB, Weston WW, Stewart M. Patient-centred interviewing part II: finding common ground. *Canadian Family Physician* 1989;35:151-8.
- Blanchard CG, Ruckdeschel JC, Fletcher BA, Blanchard EB. The impact of oncologists' behaviours on patient satisfaction with morning rounds. *Cancer* 1986;58:387-93.