

systemic hemodynamics in hypertension. *Am J Med* 1988;84(suppl 3B): 8-12.

11 Jespersen LT, Krusei LR, Sihm J, Pedersen OL. Differential effects of isradipine and atenolol on peripheral hemodynamics and arterial compliance. *Am J Med* 1989;86(suppl 4A):57-9.

12 Man in't Veld AJ, Schalekamp MADH. How intrinsic sympathomimetic activity modulates the haemodynamic responses to beta-adrenoceptor antagonists: clue to the nature of their antihypertensive mechanisms. *Br J Clin Pharmacol* 1982;13(suppl 2):2455-595.

13 SPSS Inc. *SPSS-X User's Guide*. 3rd ed. Chicago: SPSS, 1988.

14 National Center for Health Statistics. *Hypertension in adults 25-74 years of age*. Hyattsville, Maryland: NCHS, 1971-5.

15 Metropolitan Life Insurance Company. Metropolitan height and weight tables. *Stat Bull Metropol Insur Co* 1984;64:2-9.

16 Erne P, Bolli P, Bartel O, Kiowski W, Bühler FR. Factors influencing the hypotensive effects of calcium antagonists. *Hypertension* 1983;5(suppl 2):97-102.

17 Kusaka M, Ataraski K, Matsumoto K, Sumida Y, Matsuura H, Shingu T, et al. Plasma renin activity and norepinephrine as predictors for antihypertensive effects of nifedipine and captopril. *Am J Hypertens* 1991;4:735-9.

18 Nicholson JP, Rennick LM, Laragh JH. The antihypertensive effect of verapamil at extremes of dietary sodium intake. *Ann Intern Med* 1987;107: 329-34.

19 Freis ED, Reda DJ, Materson BJ. Volume (weight) loss and blood pressure response following thiazide diuretics. *Hypertension* 1988;12:244-80.

20 Frohlich ED. Obesity hypertension: converting enzyme inhibitors and calcium antagonists. *Hypertension* 1992;19(suppl 1):I-119-23.

21 Reisin E, Weed SG. Diuretics: a rational pharmacological approach for the treatment of obese hypertensives [abstract]. *Hypertension* 1991;17:405.

22 Messeri FH. *Cardiovascular drug therapy*. Philadelphia: Saunders, 1990.

23 Folkow B. Physiological aspects of primary hypertension. *Physiol Rev* 1982;62:347-504.

24 Koren MJ, Savage DD, Laragh JH, Devereux RB. Relation of hemodynamic parameters to prognosis in essential hypertension [abstract]. *Am J Hypertens* 1990;3:13a.

25 Tuck ML, Sowers JR, Dornfield L, Whitfield L, Maxwell M. Reductions in plasma catecholamines and blood pressure during weight loss in obese subjects. *Acta Endocrinol* 1983;102:252-7.

26 Sowers JR, Whitfield LA, Catania RA, Stern N, Tuck ML, Dornfield L, et al. Role of the sympathetic nervous system in blood pressure maintenance in obesity. *J Clin Endocrinol Metabol* 1982;54:1181-7.

27 Schulte W, Rüdell H, Schächinger H, Schmieder R. The efficacy of calcium antagonists and beta blockers on the hemodynamic response to stress. *Eur Heart J* (in press).

28 Schmieder R, Rüdell H, Neus H, Messeri FH, von Eiff AW. Disparate hemodynamic responses to mental challenge after antihypertensive therapy with betablockers and calcium entry blockers. *Am J Med* 1987;82:11-6.

(Accepted 22 June 1993)

Kidney granuloma in Crohn's disease

A J Archimandritis, M S Weetch

Gastrointestinal Centre,
Southern General
Hospital, Glasgow
G51 4TF
A J Archimandritis, honorary
registrar in gastroenterology
M S Weetch, registrar in
gastroenterology

Granulomas may be found in various tissues and organs of patients with Crohn's disease, such as lymph nodes, mesentery, peritoneum, liver, and lung and occasionally bones, joints, and skeletal muscle.^{1,2} We report an unusual case of Crohn's disease with granuloma formation in the kidneys.

Case report

A 22 year old man was admitted with episodic bloody diarrhoea, lassitude, anorexia, and weight loss over one year. BCG vaccination had been carried out in childhood; his grandmother had had Crohn's disease. Physical examination gave normal results, but his blood urea concentration was raised (10.0 mmol/l) and creatinine clearance reduced (44 ml/min). A chest x ray film, intravenous pyelogram, a barium meal and follow through examination, and a barium enema all gave negative results. On sigmoidoscopy patchy friability of the mucosa and discrete aphthoid ulcerations were found. A biopsy specimen showed an abnormal mucosa with severe inflammatory changes and fissuring; lymphocyte aggregations and sarcoid follicles were present in the submucosa. Crohn's disease was diagnosed and the patient treated with sulphasalazine and corticosteroids. Four weeks later he was symptom free; sigmoidoscopy showed an almost normal mucosa, but his blood urea concentration remained raised (12.0 mmol/l).

During the next six months the disease relapsed twice. On the second occasion a rectal biopsy speci-

men showed granulomas. Barium enema disclosed severe abnormalities in the caecum and ascending colon that were compatible with Crohn's disease.


Dramatic improvement followed treatment with sulphasalazine and oral corticosteroids. A year later there was a further relapse, but no improvement was observed despite administration of corticosteroids in high dosage. Barium enema showed that nearly all of the colon was affected. Serum alkaline phosphatase activity was 18.8 units (King-Armstrong method) and a liver biopsy specimen showed pericholangitis. His blood urea concentration ranged from 10.0 mmol/l to 16.6 mmol/l, and creatinine clearance was further reduced to 14-15 ml/min. A renal biopsy specimen showed chronic interstitial nephritis with local periglomerulofibrosis and granuloma formation (figure). A chest x ray film was normal; no tubercle bacilli were present in the urine. A Kveim test gave negative results, and the results of brucella serology were also negative. The patient's condition progressively deteriorated, with a rise in blood urea concentration to 19.9 mmol/l. Proctocolectomy was performed. He had an uneventful recovery and was discharged in good general condition without needing to be prescribed drugs.

Four years later he was still in remission with no specific treatment. His blood urea concentration had fallen to 9.1 mmol/l and his creatinine clearance increased to 29 ml/min. A second renal biopsy specimen showed mesangial prominence, loss and atrophy of renal tubules, and local interstitial fibrosis; no granulomas were found.

Comment

Liver granulomas occur in about 9.5% of patients with Crohn's disease,¹ although clinically overt liver involvement does not seem to be present in more than 2% of patients.³ To our knowledge, the presence of renal granulomas in patients with Crohn's disease has not been described before. It should be pointed out that granuloma formation in the kidneys is uncommon but may occur in tuberculosis, sarcoidosis, brucellosis, Wegener's granulomatosis, xanthogranulomatous pyelonephritis, malacoplakia, leprosy, and toxoplasmosis and as an adverse reaction to various drugs.^{4,5}

In our case all these causes of granuloma formation could reasonably be excluded from the patient's history; sulphasalazine, which may cause many untoward reactions such as haemolytic anaemia, temporal arteritis, and systemic lupus erythematosus, was used only after the renal impairment was discovered. In addition, the beneficial effect of proctocolectomy on renal function and the disappearance of



Renal biopsy specimen from patient with Crohn's disease showing interstitial nephritis with local periglomerulofibrosis and granuloma formation (arrow) (haematoxylin and eosin)

granulomas provide further evidence that the kidney granulomas in our patient were a manifestation of his Crohn's disease.

We thank Dr G P Crean for his advice and his permission to report this case.

- 1 Eade MN, Cooke WT, Williams JA. Liver disease in Crohn's disease. *Scand J Gastroenterol* 1971;6:199-204.
2 Whitehead R. Pathology of Crohn's disease. In: Kirsner JB, Shorter RG, eds.

- Inflammatory bowel disease*. 2nd ed. Philadelphia: Lean and Febiger, 1980:296-310.
3 Meyer JH, Sleisenger MH. Granulomatous disease of the colon. In: Sleisenger MH, Fordran JS, eds. *Gastrointestinal disease. Pathophysiology, diagnosis, management*. Philadelphia: WB Saunders, 1973:1350-68.
4 Cotran RS. Tubulointerstitial diseases. In: Brenner BM, Rector FC Jr, eds. *The kidney*. 2nd ed. Vol 2. Philadelphia: WB Saunders, 1981:1633-67.
5 Buchanan TM, Petersdorf RG. Brucellosis. In: Petersdorf RG, Adams RD, Braunwald E, Isselbacher KJ, Martin JB, Wilson JD, eds. *Harrison's principles of internal medicine*. 10th ed (international student ed). Tokyo: McGraw Hill, 1983:974-6.

(Accepted 14 June 1993)

Post-tropical screening: how useful is it?

Bernadette Carroll, Carol Dow, David Snashall, Thomas Marshall, Peter L Chiodini

Hospital for Tropical Diseases, London NW1 0PE

Bernadette Carroll, *research nurse*
Peter L Chiodini, *consultant parasitologist*

Foreign and Commonwealth Office, London SW1A 2AF
Carol Dow, *principal medical adviser*

United Medical and Dental Schools, St Thomas's Hospital, London SE1 7EH
David Snashall, *senior lecturer in occupational health*

London School of Hygiene and Tropical Medicine, London WC1E
Thomas Marshall, *lecturer*

Correspondence to:
Dr Carol Dow, FCO Medical and Welfare Department, Old Admiralty Building, The Mall, London SW1A 2AF.

BMJ 1993;307:541

Travellers to the tropics worry about catching "tropical" diseases. Those who live in the tropics for a while think they are at more risk than are short term visitors. Many believe, as do their doctors, that medical screening on return to the United Kingdom, even in asymptomatic individuals, will reveal conditions which could or should be treated. This study examines the usefulness of screening asymptomatic individuals who have returned from the tropics.

Methods and results

In 1991 at the Hospital for Tropical Diseases 1029 asymptomatic individuals presenting consecutively who had recently returned from the tropics were examined using a standard protocol comprising a questionnaire, a physical examination by a tropical medicine specialist, and a range (determined by the area visited) of laboratory tests. The subjects either worked for the Foreign and Commonwealth Office, the Overseas Development Administration, Voluntary Services Overseas, or the British Council or were trekkers. The subjects covered a wide range of ages (8 months-73 years), with 135 (13.1%) being under 14 years. Length of stay abroad ranged from three months to 45 years. Both sexes were represented equally. Most had been in sub-Saharan Africa, but all tropical areas of the world were represented.

Full blood count with white cell differential and stool microscopy for cysts, ova, and parasites were performed routinely, as was urine analysis by dipstick. Schistosomal and filarial serology was performed only if the questionnaire suggested the individual had been at risk. Additional tests were done where thought necessary.

Overall one in four of those screened had an abnormal result (see table). Urine analysis gave 116 abnormalities in 830 cases (14%). When followed up none of these abnormalities proved to be conditions resulting from life in the tropics. Stool microscopy for cysts, ova, and parasites showed abnormalities in 207 of 995 patients who were able to produce specimens (18.7%). The results were unaffected by area of travel or any other variable. The commonest abnormal finding was of cysts of *Entamoeba histolytica* or *Giardia lamblia*. Practically all these individuals were treated, despite being asymptomatic. Full blood count showed eosinophilia in 67 out of 852 samples (16.1%). In 26 people this was associated with parasitosis. Only one other blood count abnormality was of tropical origin, a case of post-diarrhoeal malabsorption. Schistosomal serology was positive in 72 out of 676 tests (10.7%). Positive results, more significant if associated with eosinophilia, were followed up and individuals treated when necessary. Thirteen positive results were

Abnormal findings on screening asymptomatic people returned from tropical areas

Investigations and findings	No
	<i>of pathogens</i>
Stool microscopy:	
Helminths	33
<i>Trichuris trichiura</i> ova	14
<i>Ascaris lumbricoides</i> ova	12
Hookworm ova and larvae	3
<i>Schistosoma mansoni</i> ova	2
Threadworm ova	1
Strongyloides larvae	1
Protozoa	174
<i>Entamoeba histolytica</i> cysts	115
<i>Giardia lamblia</i> cysts	39
<i>Blastocystis hominis</i> cysts	16
<i>Dientamoeba fragilis</i> trophozoites	4
	<i>of people</i>
Urine analysis	116
Albuminuria	58
Haematuria	33
Haematuria and albuminuria	10
Ketonuria	6
Glycosuria	4
Albuminuria and biliuria	2
Albuminuria and glycosuria	1
Ketonuria and haematuria	1
Albuminuria and ketonuria	1
Parasitosis in people with eosinophilia	26
Schistosomiasis	17
Schistosomiasis and filariasis	1
Filariasis	2
Hookworm infection	2
Strongyloidiasis	1
Hookworm infection and strongyloidiasis	1
Trichuriasis	2

obtained in enzyme linked immunosorbent assays for filaria, one of which was associated with detectable disease (onchocerciasis).

Physical examination by a tropical medicine specialist produced 387 abnormal findings, of which 108 were skin problems. Other findings reflected "non-tropical" diseases such as hypertension and obesity. The only specific tropical conditions diagnosed by specialist examination were one case of cutaneous larva migrans and two cases of splenomegaly, attributed to past malaria. The additional tests requested by the doctors produced few abnormal important results and did not prompt changes in management.

Comment

Screening of asymptomatic individuals returning from the tropics identified abnormalities in one in four. Most of these abnormalities were revealed by laboratory tests, some guided by answers to direct questions. They were performed in a specialist laboratory which would expect a high rate of positive findings.¹ Specialist physical examination of all patients added little to the practical management of these patients and thus can be limited to those considered to be at particular risk—for example, because of a history of malaria. Screening for tropical disease can be efficiently carried out by an informed health professional using structured history taking and relevant laboratory tests.

1 Mohr E, Mohr I. Statistical analysis of the incidence of positives in the examination of parasitological specimens. *J Clin Microbiol* 1992;30:1572-4.

(Accepted 2 July 1993)