

Coffea arabica L., a New Host Plant for *Acetobacter diazotrophicus*, and Isolation of Other Nitrogen-Fixing Acetobacteria

TERESITA JIMENEZ-SALGADO,¹ LUIS E. FUENTES-RAMIREZ,² ARMANDO TAPIA-HERNANDEZ,¹
MIGUEL A. MASCARUA-ESPARZA,¹ ESPERANZA MARTINEZ-ROMERO,²
AND JESUS CABALLERO-MELLADO^{2*}

Departamento de Genética Molecular, Centro de Investigación sobre Fijación de Nitrógeno, Universidad Nacional Autónoma de México, Cuernavaca, Morelos,² and Centro de Investigaciones Microbiológicas, Instituto de Ciencias, Universidad Autónoma de Puebla, Puebla, Puebla,¹ México

Received 23 December 1996/Accepted 12 June 1997

Acetobacter diazotrophicus was isolated from coffee plant tissues and from rhizosphere soils. Isolation frequencies ranged from 15 to 40% and were dependent on soil pH. Attempts to isolate this bacterial species from coffee fruit, from inside vesicular-arbuscular mycorrhizal fungi spores, or from mealybugs (*Planococcus citri*) associated with coffee plants were not successful. Other acid-producing diazotrophic bacteria were recovered with frequencies of 20% from the coffee rhizosphere. These N₂-fixing isolates had some features in common with the genus *Acetobacter* but should not be assigned to the species *Acetobacter diazotrophicus* because they differed from *A. diazotrophicus* in morphological and biochemical traits and were largely divergent in electrophoretic mobility patterns of metabolic enzymes at coefficients of genetic distance as high as 0.950. In addition, these N₂-fixing acetobacteria differed in the small-subunit rRNA restriction fragment length polymorphism patterns obtained with *Eco*RI, and they exhibited very low DNA-DNA homology levels, ranging from 11 to 15% with the *A. diazotrophicus* reference strain PAI 5^T. Thus, some of the diazotrophic acetobacteria recovered from the rhizosphere of coffee plants may be regarded as N₂-fixing species of the genus *Acetobacter* other than *A. diazotrophicus*. Endophytic diazotrophic bacteria may be more prevalent than previously thought, and perhaps there are many more potentially beneficial N₂-fixing bacteria which can be isolated from other agronomically important crops.

Almost 100 bacterial genera, of both the eubacteria and archaeobacteria, are capable of fixing N₂ (32). There may exist many more bacterial species or genera which can fix nitrogen since a majority of bacterial species are not presently culturable (31) and the search for diazotrophs in some environments has been relatively limited. Research on N₂-fixing bacteria endophytically associated with sugarcane led to the description of *Acetobacter diazotrophicus*, which is the only known nitrogen-fixing species of acetic acid-producing bacteria (13, 29). Similarly, in the last few years, the genus *Azoarcus* and its various species were described (16, 33), most of them recovered from the roots of Kallar grass (24). These findings suggest that many other endophytic N₂-fixing species may not yet have been described.

Looking for well-known N₂-fixing species and for new diazotrophs associated with previously untested plants or from new environments may provide a better picture not only of the distribution of N₂-fixation ability among bacterial taxa but also of the distribution and diversity of N₂-fixing bacterial populations.

In this work, we report the natural occurrence of diazotrophic acetic acid-producing bacteria in the rhizosphere and in tissues from different cultivars of seed-propagated coffee plants (*Coffea arabica* L.). Microbiological, biochemical, and genetic tests showed that a majority of these bacteria belong to the species *A. diazotrophicus*. We obtained evidence that strongly

supports the hypothesis that some of the strains could represent new N₂-fixing species of the genus *Acetobacter*.

MATERIALS AND METHODS

Locations and coffee cultivars. Coffee plant varieties grown in nurseries or under field conditions were collected from diverse geographic regions of Mexico up to 750 km apart. The origins of samples and the coffee varieties analyzed are summarized in Table 1.

Media and cultural conditions. N-free semisolid LGI medium supplemented with sugarcane juice at pH 4.5 (7) and cycloheximide (150 mg/liter) was used for enrichment culturing of N₂-fixing acetobacters. For isolation and culturing, acetic acid LGI agar plates supplemented with yeast extract (50 mg/liter) and cycloheximide (150 mg/liter) and potato agar plates with 10% cane sugar were used (7). N₂-fixing acetobacters were grown at 29°C in SYP medium (6) for all other assays.

Isolation. Care was taken to keep rhizosphere soil intact around the root. Later, the root samples were rinsed three times in sterile distilled water. Separately, coffee root and stem pieces were immersed in 1% chloramine T for 5 min and treated as described previously (11). Root and stem samples were macerated in a blender, and supernatant aliquots (100 µl) were placed in vials containing 5 ml of N-free semisolid LGI medium (7). Other vials were inoculated with 100-µl aliquots from a 1/10 (wt/vol) rhizosphere soil suspension. Also, five samples (10 g each) of ripening fruit from *Coffea arabica* cv. Garnica collected in the coffee-growing region of Huitzilán, Puebla State, Mexico, were surface sterilized and treated as mentioned above for root and stem samples. In attempts to recover *A. diazotrophicus* from inside vesicular-arbuscular mycorrhizal (VAM) fungal spores, 100 g of eight rhizosphere soil samples (four from Huitzilán and four from Tapachula, Chiapas, Mexico) was sieved and at least 60 VAM spores were isolated from each soil sample by the method described by Gerdemann and Nicolson (12). The VAM spores were surface sterilized with 1% chloramine T for 5 min and then washed four times with sterile distilled water. Spores without apparent damage were manually crushed and placed in vials containing N-free semisolid LGI medium as reported previously (23). In addition, 50 adult mealybugs identified as *Planococcus citri* were analyzed for N₂-fixing acetobacters. These were collected from aerial parts of coffee plants, cultivar Caturra, growing in fields at Atoyac, Guerrero State, Mexico. Groups of 10 insects were rinsed with 0.01% (vol/vol) Tween 20 in 10 mM MgSO₄ · 7H₂O until the liquid was clear. Insects were macerated in 1.0 ml of 10 mM MgSO₄ · 7H₂O, and 100-µl

* Corresponding author. Mailing address: Centro de Investigación sobre Fijación de Nitrógeno, UNAM, Apdo. Postal No. 565-A, Cuernavaca, Morelos, México. Phone: 73 13 16 97. Fax: 73 17 55 81. E-mail: mellado@cifn.unam.mx.

TABLE 1. Isolation frequencies of *A. diazotrophicus* recovered from coffee plant cultivars

Location	Cultivar	Plant age	pH of soil	Isolation (%)		
				Rhizp ^a	Root	Stem
Huitzilán, Puebla	Garnica	5 yr ^b	4.07	40.0	40.0	0.0
	Garnica	2 yr ^b	6.27	0.0	0.0	0.0
Xicotepec, Puebla	Catuai	2 mo ^c	4.74	ND ^d	0.0	20.0
	Catuai	6 mo ^b	4.00	ND	20.0	0.0
Atoyac, Guerrero	Caturra	1 yr ^b	3.64	15.0	20.0	0.0
	M. Novo	1 yr ^c	6.20	0.0	0.0	0.0
Tapachula, Chiapas	Caturra	5 mo ^c	5.40	30.0	0.0	0.0
	Caturra	5 mo ^c	5.80	20.0	0.0	0.0
	Caturra	3 mo ^c	5.30	40.0	0.0	0.0

^a Rhzp, rhizosphere (soil shaken off roots).

^b Coffee plants growing under field conditions.

^c Coffee plants growing in a nursery.

^d ND, not determined.

aliquots were inoculated into media for isolation of *A. diazotrophicus* as described previously (5).

Vials of inoculated N-free semisolid LGI medium were incubated at 30°C for 7 days. Thereafter, vials were replicated under the same conditions and assayed for acetylene reduction activity as described previously (21). Nitrogenase-positive vials with a yellow surface pellicle were streaked onto acetic LGI agar plates and incubated at 30°C. After 5 to 7 days, acid-producing dark-orange colonies suggested the presence of *A. diazotrophicus* (7). Colonies were streaked on potato agar plates to verify culture purity. In addition, atypical acid-producing isolates (referred to in the text as DOR and APL isolates) were also recovered from coffee rhizosphere samples from Tapachula. These isolates did not exhibit growth typical of *A. diazotrophicus* on LGI agar plates. DOR isolates were similar in their dark-orange color but formed very irregular smooth flat colonies. In addition, while *A. diazotrophicus* colonies are initially white and later become yellow-orange, DOR isolates are always orange. APL isolates showed a liquid-like appearance on the first days, but after 5 days, the isolates became dry and took on a yellowish color. One non-acid-producing mucoid strain (designated CFN-Cf 56) was also isolated due to its predominant growth on an LGI agar plate. This strain was selected based on its colony morphology, which was similar to that of a spontaneous, non-acid-producing mutant that was obtained from *A. diazotrophicus* SRT4 (1).

Identification. Isolate identification was based on colony morphology in culture media, on biochemical tests, and on genetic characteristics reported for *A. diazotrophicus* (5–7, 13). *A. diazotrophicus* PAI 5^T (ATCC 49037), kindly provided by J. Döbereiner, and UAP 5560, analyzed previously (6, 11), were used as controls.

MLEE. Each isolate was grown for 36 h in 40 ml of SYP medium and harvested by centrifugation; pellets were suspended and treated as described previously (6). Starch gel electrophoresis and selective staining of metabolic enzymes were done as described before (25). The analyzed enzymes were the same ones used in a previous study (5) and were assayed under the same conditions. Distinct combinations of alleles for 12 enzyme loci (multilocus genotypes) were designated as different electrophoretic types (ETs) (25). *A. diazotrophicus* strains (CFNE 501, CFNE 550, PAI 5^T, PAI 3, 1772, PSP 22, and PRC 1), corresponding to the reported seven ETs (5), were included as references in multilocus enzyme electrophoresis (MLEE) assays to determine the genetic relationships of coffee plant-associated isolates and *A. diazotrophicus*.

Total DNA isolation, DNA restriction, and filter blot hybridization. Total DNA was isolated as described previously by Ausubel et al. (3). DNA was digested with *EcoRI*, and restriction fragments were electrophoresed in vertical 1.0% agarose gels in Tris-acetate buffer (40 mM Tris-acetate, 2 mM EDTA [pH 8]) at 40V for 13 h at 4°C. Total DNA digests were transferred from gels to nylon filters by the Southern procedure as described before (6). The restriction fragment length polymorphism (RFLP) patterns of the *nifHDK* genes were determined by hybridization with a *HindIII-HindIII* 4.3-kb fragment containing the *nifHDK* genes from *A. diazotrophicus* UAP 5560 obtained from pUC19 derivative pNHAd4 (unpublished results). DNA-DNA homology was based on relative levels of hybridization to ³²P-labelled DNA from strain PAI 5^T. Amounts of DNA in gels were quantified as described before (5). Autoradiography was performed at –70°C for 24 h; filter lanes were cut and counted with a Beckman scintillation counter. The percentage of total hybridization was calculated for each strain tested. Hybridization patterns of small-subunit (SSU) ribosomal DNA (rDNA) genes were analyzed as described before (5), but in this study, total DNA also was digested with restriction enzymes *SphI* and *NcoI*. Genomic

DNA from coffee plant-associated isolates and from strains PAI 5^T and UAP 5560 of *A. diazotrophicus* were hybridized with an *Escherichia coli* SSU rRNA gene internal fragment from vector pKK3535 (4) corresponding to nucleotides 80 to 653. ³²P-labelled probes were prepared by nick translation.

SSU rDNA sequence alignment. To search discriminative restriction sites in the SSU rRNA genes for distinguishing *Acetobacter* from other bacteria, we aligned 11 reported SSU rDNA sequences of different strains of the family *Acetobacteraceae* and 29 sequences of strains from other members of the α subclass of the class *Proteobacteria* (α -*Proteobacteria*) with GCG software version 8.1-UNIX (Genetics Computer Group, Madison, Wis.). GenBank accession numbers for SSU rDNA sequences aligned are shown in Table 2.

RESULTS

Isolation. Typical yellow surface pellicles of nitrogen-fixing *Acetobacter* were observed in N-free LGI medium vials inoculated with rhizosphere soil, blended roots, and stems from different coffee plant varieties grown in various geographical areas of Mexico. On LGI agar plates, dark-orange colonies typical of *A. diazotrophicus* were observed (Fig. 1). Isolation frequencies from the rhizosphere, inside of roots, or stems ranged from 15 to 40% in plants grown in acid soils (Table 1). Additionally, from some rhizosphere samples, we recovered acid-producing DOR and APL isolates from LGI medium vials

TABLE 2. GenBank accession numbers used for the SSU rDNA sequence alignments

Species	Strain	Accession no.
<i>Acetobacter pasteurianus</i>	LMD 22.1	X71863
<i>Acetobacter aceti</i>	DSM 3508	X74066
<i>Acetobacter liquefaciens</i>	LMG 1382	X75617
<i>Acetobacter diazotrophicus</i>	PAI 5 ^T	X75618
<i>Acetobacter xylinum</i>	NCIB 11664	X75619
<i>Acetobacter hansenii</i>	NCIB 8746	X75620
<i>Acetobacter europaeus</i>	DSM 6160	Z21936
<i>Gluconobacter oxydans</i>	DSM 3503	X73820
<i>Gluconobacter asaii</i>	LMG 1390	X80165
<i>Gluconobacter cerinus</i>	LMG 1368	X80775
<i>Gluconobacter frateurii</i>	LMG 1365	X82290
<i>Acidomonas methanolica</i>	LMG 1668 ^a	X77468
<i>Acidiphilium</i> sp.	C-1	D30769
<i>Acidiphilium aminolytica</i>	101	D30771
<i>Acidiphilium angustum</i>	ATCC 35903	D30772
<i>Acidiphilium cryptum</i>	ATCC 33463	D30773
<i>Acidiphilium facilis</i>	ATCC 35904	D30774
<i>Acidiphilium organovorum</i>	ATCC 43141	D30775
<i>Acidiphilium rubrum</i>	ATCC 35905	D30776
<i>Acidiphilium</i> sp.	St 1-5	D86508
<i>Acidiphilium</i> sp.	St 1-7	D86509
<i>Rhodopila globiformis</i>	DSM 161	D86513
<i>Rhodopila globiformis</i>	ATCC 7950	M59066
<i>Rhodospirillum</i> sp.	MT-SP-3	D12703
<i>Rhizobium meliloti</i>	IAM12611	D14509
<i>Rhizobium leguminosarum</i>	IAM12609	D14513
<i>Rhodopseudomonas</i> sp.	IL-245	D15063
<i>Rhodobacter capsulatus</i>	ATCC 11166	D16428
<i>Rhodospirillum rubrum</i>	ATCC 11170	D30778
<i>Beijerinckia indica</i>	ATCC 9039	M59060
<i>Caulobacter</i> sp.	MCS 6	M83811
<i>Hyphomonas</i> sp.	MHS 3	M83812
<i>Hyphomicrobium vulgare</i>	MC-750	X53182
<i>Roseobacter litoralis</i>	ATCC 49556	X78312
<i>Azospirillum lipoferum</i>	ATCC 29708	X79729
<i>Azospirillum irakense</i>	103312	X79737
<i>Azospirillum brasilense</i>	Sp 7	X79739
<i>Azospirillum amazonense</i>	Y2	X79742
<i>Xanthobacter flavus</i>	JW/KR-E1	X94206
<i>Pedomicrobium manganicum</i>	ACM 3038	X97691

^a Substrain MB 58.




FIG. 1. Colony morphologies of N_2 -fixing acetobacters after 7 days at 29°C on LGI agar plates. (A and B) *A. diazotrophicus* PAI 5^T (A) and CFN-Cf 50 (B); (C) DOR isolate, strain CFN-Cf 55; (D) APL isolate, strain UAP-Cf 60; (E) mucoid strain CFN-Cf 56. The green color in LGI agar plates was turned to yellow by acid-producing isolates.

with yellow surface pellicles. These isolates reduced acetylene in pure culture but had clearly different morphologies from that of *A. diazotrophicus* on LGI agar plates (Fig. 1). These DOR and APL isolates were recovered from two rhizosphere samples (collected in Tapachula) with isolation frequencies of 20%. Strain CFN-Cf 56, which does not produce acid on LGI agar plates, was the only mucoid isolate recovered (Fig. 1).

No bacteria corresponding to the descriptions given above

were isolated from coffee plants growing at a pH higher than 6.0 nor from coffee fruit, VAM spores, or mealybugs (*P. citri*).

MLEE and genetic relationships. The origins of the coffee-associated N_2 -fixing isolates are shown in Table 3. The genetic relationships among the N_2 -fixing isolates associated with coffee plants and *A. diazotrophicus* strains recovered from known hosts are illustrated by the dendrogram shown in Fig. 2. Thirteen distinct ETs were identified among N_2 -fixing coffee iso-

TABLE 3. Origins of representative N_2 -fixing bacteria recovered from the coffee environment

MLEE division (ET) ^a	Type of isolate	Strain designation	Isolate recovered from:	Plant age	Cultivar	Location
I (1)	<i>A. diazotrophicus</i>	CFN-Cf13	Stem tissue	2 mo ^b	Catuai	Xicotepec, Puebla
I (11)	<i>A. diazotrophicus</i>	CFN-Cf50	Root tissue	6 mo ^c	Catuai	Xicotepec, Puebla
I (1)	<i>A. diazotrophicus</i>	UAP-Cf29	Rhizosphere	1 yr ^c	Caturra	Atoyac, Guerrero
I (9)	<i>A. diazotrophicus</i>	CFN-Cf52	Root tissue	1 yr ^c	Caturra	Atoyac, Guerrero
I (8)	<i>A. diazotrophicus</i>	UAP-Cf01	Rhizosphere	5 yr ^c	Garnica	Huitzilán, Puebla
I (8)	<i>A. diazotrophicus</i>	UAP-Cf05	Root tissue	5 yr ^c	Garnica	Huitzilán, Puebla
I (12)	<i>A. diazotrophicus</i>	UAP-Cf51	Rhizosphere	5 mo ^b	Caturra	Tapachula, Chiapas
I (14)	<i>A. diazotrophicus</i>	UAP-Cf53	Rhizosphere	3 mo ^b	Caturra	Tapachula, Chiapas
I (10)	<i>A. diazotrophicus</i>	UAP-Cf58	Rhizosphere	5 mo ^b	Caturra	Tapachula, Chiapas
I (13)	NAP ^d	CFN-Cf56	Rhizosphere	3 mo ^b	Caturra	Tapachula, Chiapas
III (15)	APL ^e	UAP-Cf59	Rhizosphere	5 mo ^b	Caturra	Tapachula, Chiapas
III (16)	APL	CFN-Cf60	Rhizosphere	3 mo ^b	Caturra	Tapachula, Chiapas
IV (17)	DOR ^f	CFN-Cf55	Rhizosphere	3 mo ^b	Caturra	Tapachula, Chiapas
IV (18)	DOR	UAP-Cf57	Rhizosphere	5 mo ^b	Caturra	Tapachula, Chiapas
V (19)	SAd ^g	CFN-Cf54	Rhizosphere	5 mo ^b	Caturra	Tapachula, Chiapas

^a Divisions and ETs were based on MLEE assays. More isolates included in divisions I and III to V were recovered, but only one of the many isolates recovered from each plant or rhizosphere sample was designated as a strain.

^b Coffee plants growing in a nursery.

^c Coffee plants growing under field conditions.

^d NAP, non-acid-producing isolate.

^e APL, acid-producing liquid isolate.

^f DOR, dark-orange isolate.

^g SAd, isolate with colonial features similar to those of *A. diazotrophicus*.




FIG. 2. Genetic relationships of ETs identified among *A. diazotrophicus* isolates recovered from well-known hosts and N₂-fixing acetobacters associated with coffee plants. A plus after the ET number indicates that the ET represents only coffee plant-associated nitrogen-fixing acetobacters, except for ET 1, which includes reported reference strains as well.

lates (multilocus genotype data are available upon request). Division I, with a genetic distance of 0.430, included six previously identified ETs (ET 1 to ET 6) (5) and six new closely related ETs (ET 8 to ET 12 and ET 14) from coffee-associated *A. diazotrophicus* isolates. In addition, division I included ET 13, which corresponds to an isolate (CFN-Cf 56) with no typical features of *A. diazotrophicus*. Moreover, isolates recovered from both the rhizosphere (e.g., strain UAP-Cf 29) and the inside of coffee plants (e.g., strain CFN-Cf 13) were identical to strains of *A. diazotrophicus* belonging to ET 1, previously identified (5, 6) as the predominant ET (e.g., UAP 5560 and CFNE 501) of the species. Division II contained only ET 7, a genetically distant group previously identified (5) among *A. diazotrophicus* strains isolated from sugarcane and *Pennisetum purpureum* in Brazil. Divisions III, IV, and V, which included ETs 15 to 19, diverged largely at a genetic distance of 0.950 from divisions I and II. Division III (ETs 15 and 16) contained only APL isolates, while division IV (ETs 17 and 18) included DOR isolates and division V (ET 19) grouped isolates with colonial features similar to those of *A. diazotrophicus* on acetic LGI agar plates.

Identification. Many isolates recovered from the inside of coffee plants and from the rhizosphere of these plants were identified as belonging to the species *A. diazotrophicus* on the basis of reported characteristics (5, 6, 7, 13) such as growth features on culture media, biochemical tests, and results of genetic approaches (Tables 4 and 5). Other isolates such as the mucoid strain CFN-Cf 56 and the DOR and APL strains differed from *A. diazotrophicus* in various phenotypic characteristics (Table 4 and carbon usage data not shown). Nevertheless, these isolates were able to grow at pH 5, oxidize ethanol to acetic acid in neutral and acid conditions, and oxidize acetate and lactate to CO₂ and H₂O (Table 4), phenotypic features which are considered (8, 29) fundamental for the identification of the genus *Acetobacter*.

Genetic characteristics. Total *Eco*RI DNA digests from coffee isolates, including those with different colony morpholo-

TABLE 4. Phenotypic characteristics of N₂-fixing acetic acid bacteria isolated from the coffee plant environment^a

Characteristic	I ^b		III		IV		V	
	PAI 5 ^{Tc}	UAP 5560 ^c	CFN-Cf 13	UAP-Cf 05	CFN-Cf 52	CFN-Cf 56	UAP-Cf 59	CFN-Cf 60
Gram stain	+	+	+	+	+	+	+	+
Oxidase	+	+	+	+	+	+	+	+
Catalase	+	+	+	+	+	+	+	+
Oxidation of ethanol to acetic acid	+	+	+	+	+	+	+	+
Oxidation of glucose to acetic acid	+	+	+	+	+	+	+	+
Oxidation of acetic acid to CO ₂ and H ₂ O	+	+	+	+	+	+	+	+
Oxidation of lactate to CO ₂ and H ₂ O	+	+	+	+	+	+	+	+
Water-soluble brown pigments on GYC ^f	+	+	+	+	+	+	+	+
Dark-orange colonies on LGI plates	+	+	+	+	+	+	+	+
Dark-brown colonies on potato agar with 10% sugar	+	+	+	+	+	+	+	+
Brownish colonies on potato agar with 10% sugar	+	+	+	+	+	+	+	+
Growth with 30% D-glucose	+	+	+	+	+	+	+	+
Growth with 30% sucrose	+	+	+	+	+	+	+	+
Yellow surface pellicle formation and pH below 3 in N-free semisolid LGI medium	+	+	+	+	+	+	+	+
C ₂ H ₂ reduction activity (N ₂ fixation)	+	+	+	+	+	+	+	+

^a Phenotypic characteristics were positive (+) or negative (-) for each strain.

^b Divisions based on MLEE assays.

^c *A. diazotrophicus* strains recovered from sugarcane used as controls.

^d Oxidation was observed up to day 4.

^e Cream-colored colonies, but producing a brownish liquid pigment.

^f GYC, 5% D-glucose-1% yeast extract-3% CaCO₃-2.5% agar (8).

TABLE 5. Genetic characteristics of some N₂-fixing acetobacters recovered from the coffee plant environment^a

MLEE division ^a (ET) ^a	Type of isolate ^b	Reference strain	Sizes (kb) ^c		DNA-DNA homology (%) ^d
			<i>nifHDK</i> genes	SSU rDNA genes	
I (3)	<i>A. diazotrophicus</i>	PAI 5 ^{Te}	9.0, 2.0, 1.25	9.3, 3.6, 2.3, 1.6	100.0
I (1)	<i>A. diazotrophicus</i>	UAP 5560 ^e	9.0, 2.0, 1.25	9.3, 3.6, 2.3, 1.6	104.0
I (8)	<i>A. diazotrophicus</i>	UAP-Cf 05	9.0, 3.1, 1.25	9.3, 3.6, 2.3, 1.6	ND ^f
I (9)	<i>A. diazotrophicus</i>	CFN-Cf 52	9.0, 2.0, 1.25	9.3, 3.6, 2.3, 1.6	83.0
I (10)	<i>A. diazotrophicus</i>	UAP-Cf 58	9.0, 2.0, 1.25	9.3, 3.6, 2.3, 1.6	78.0
I (11)	<i>A. diazotrophicus</i>	CFN-Cf 50	9.0, 2.0, 1.25	9.3, 3.6, 2.3, 1.6	96.0
I (12)	<i>A. diazotrophicus</i>	UAP-Cf 51	9.0, 2.0, 1.25	9.3, 3.6, 2.3, 1.6	72.0
I (14)	<i>A. diazotrophicus</i>	UAP-Cf 53	9.0, 2.0, 1.25	9.3, 3.6, 2.3, 1.6	77.0
I (13)	NAP	CFN-Cf 56	7.6, 3.5, 1.20, 1.0	9.3, 3.6, 2.3, 1.6	30.0
III (15)	APL	UAP-Cf 59	Not detected	8.6, 3.9, 3.6, 1.6	12.0
III (16)	APL	CFN-Cf 60	Not detected	8.6, 3.9, 3.6, 1.6	15.0
IV (17)	DOR	CFN-Cf 55	9.0, 2.0, 1.20	9.7, 3.6, 1.6	14.0
IV (18)	DOR	UAP-Cf 57	9.0, 2.0, 1.20	9.7, 3.6, 1.6	15.0
V (19)	SAd	CFN-Cf 54	6.6, 2.1, 1.15	9.7, 3.6, 2.8, 1.6	11.0

^a Divisions and ETs were based on MLEE assays.

^b Types described in Table 3, footnotes *d*, *e*, *f*, and *g*.

^c Bands from total *EcoRI* DNA fingerprints hybridized as described in Materials and Methods.

^d Homology to the control strain PAI 5^T.

^e *A. diazotrophicus* strains recovered from sugarcane used as controls.

^f ND, not determined.

gies, were hybridized to *A. diazotrophicus nifHDK* genes (Fig. 3). Three common hybridizing bands were observed for representative isolates of the 6 ETs from division I (Table 5), as reported previously (5, 6). In addition, isolates represented by strain UAP-Cf 05 (division I) and isolates grouped in division IV (e.g., CFN-Cf 55 and UAP-Cf 57) showed bands that differed from each other slightly in size (Table 5). Strain CFN-Cf 54 (division V) and the mucoid strain CFN-Cf 56 showed a more variable pattern of the *nifHDK* genes. No hybridizing bands were observed under stringent hybridization conditions with APL strains from division III (Fig. 3), even though pure cultures of these isolates were capable of reducing acetylene. This result seems to indicate that structural nitrogenase genes from APL isolates are largely divergent from *A. diazotrophicus nifHDK* genes.

RFLP analysis of *EcoRI* DNA digests from coffee plant-associated isolates showed four distinct hybridization patterns

to SSU rRNA genes (Fig. 4). Among the patterns obtained, two common hybridizing bands (3.6 and 1.6 kb) were observed. All isolates of division I showed the same pattern of hybridization (Table 5) as that observed previously in all *A. diazotrophicus* strains analyzed (5). N₂-fixing *Acetobacter* strains diverging at a large genetic distance from divisions I and II, according to the MLEE assays, presented different SSU rRNA hybridization patterns (Fig. 4 and Table 5). Isolates grouped in division IV did not have the 2.3-kb band which seems to correspond to the 3.9- and 2.8-kb bands observed in the strains from divisions III and V, respectively.

From the SSU rDNA sequence analysis, we inferred that Southern hybridization with a SSU rDNA probe of *SphI*-digested genomic DNA could be helpful in distinguishing members of the family *Acetobacteraceae* from other α -*Proteobacteria* (Fig. 5) and that *NcoI* digests could be used to distinguish the genera *Gluconobacter* and *Acetobacter* from *Acidiphilium* and *Rhodopila* (Fig. 5) (26). The majority of *Acetobacteraceae* spe-




FIG. 3. Autoradiogram of a Southern blot of total *EcoRI*-digested DNA hybridized with the *nifHDK* probe of *A. diazotrophicus* UAP 5560. Lanes: 1, strain UAP 5560 used as a control; 2 to 9, coffee plant-associated nitrogen-fixing strains CFN-Cf 54 (lane 2), CFN-Cf 56 (lane 3), UAP-Cf 05 (lane 4), CFN-Cf 57 (lane 5), UAP-Cf 59 (lane 6), UAP-Cf 13 (lane 7), CFN-Cf 55 (lane 8), and CFN-Cf 60 (lane 9).




FIG. 4. Autoradiogram of a Southern blot of total *EcoRI*-digested DNA hybridized with an internal 16S rDNA probe of *E. coli*. Lanes: 1, strain UAP 5560 used as a control; 2 to 9, coffee-associated nitrogen-fixing strains UAP-Cf 13 (lane 2), UAP-Cf 05 (lane 3), CFN-Cf 56 (lane 4), CFN-Cf 54 (lane 5), UAP-Cf 57 (lane 6), UAP-Cf 59 (lane 7), CFN-Cf 55 (lane 8), and CFN-Cf 60 (lane 9).




FIG. 5. Diagrammatic representation of distinctive restriction sites *SphI* and *NcoI* of SSU rRNA in *Acetobacteraceae* and phenotypically related bacteria. a, site not present in *G. oxydans* DSM 3503; b, site exclusively present in the *A. diazotrophicus* PAI 5^T sequence but not detected after Southern hybridization; c, site present in *Azospirillum lipoferum* ATCC 29708 and *Azospirillum amazonense* Y2; d, of 17 analyzed sequences, this site exclusively present in *Rhizobium meliloti* IAM 12611, *Rhizobium leguminosarum* IAM 12609, *Caulobacter* sp. strain MCS 6, *Hyphomonas* sp. strain MHS 3, and *Xanthobacter flavus* JW/KR-E1.

cies, including *Acidomonas methanolica*, have two internal *SphI* sites in their SSU rDNA, except for *A. diazotrophicus* PAI 5^T (accession number X75618), which supposedly has an extra *SphI* site at base 485 as deduced from the reported sequence (Fig. 5). *Rhodopila globiformis* (accession numbers D86513 and M59066) lacks one of the *SphI* sites. From the analysis of the *A. diazotrophicus* PAI 5^T SSU rDNA sequence (26), we expected to observe one hybridizing band of 450 bp with the probe used when the DNA was digested with *SphI*. However, only one SSU rRNA hybridizing band of 1.3 kb was observed in *A. diazotrophicus* PAI 5^T and UAP 5560. This band was conserved in all coffee plant-associated isolates. These conflicting results may be explained if the *A. diazotrophicus* sequence has an error at the *SphI* site. If such were the case, then the *Acetobacteraceae* and *Acidiphilium* spp. would have only two *SphI* conserved sites. *Gluconobacter* and *Acetobacter* SSU rDNA are characterized by two *NcoI* restriction sites. However, all *Acidiphilium* and *Rhodopila* species and *Gluconobacter oxydans* lack the *NcoI* restriction site at the base corresponding to nucleotide 110 of *A. diazotrophicus*. The rest of the α -*Proteobacteria* analyzed lack at least one site for each restriction enzyme. Genomic DNA from the strains recovered from the coffee plant environment, digested with *NcoI* and hybridized to the same SSU rRNA internal gene fragment, showed the expected 1.24-kb band (26) (data not shown).

The results of the DNA-DNA homology assays are shown in Table 5. The six strains of N₂-fixing acetobacters corresponding to division I (except strain CFN-Cf 56) analyzed were related to *A. diazotrophicus* PAI 5^T with DNA homology values of 72 to 96%, with a mean DNA homology of 81%. This value was consistent with the values of 86 and 84% reported previously (5, 13) for *A. diazotrophicus* strains recovered from sugarcane and other known hosts. The mucoid strain CFN-Cf 56 exhibited only 30% DNA homology to strain PAI 5^T. APL isolates (MLEE division III) and DOR acetobacters from division IV and strains from division V exhibited very low DNA homology levels, ranging from 11 to 15% with reference strain PAI 5^T.

DISCUSSION

It is considered that "the isolation of acetic acid bacteria and their assignment to either the genus *Acetobacter* or *Gluconobacter* generally pose few problems" (29). According to Swings (29), gram-negative or gram-variable aerobic bacteria that oxidize ethanol to acetic acid in neutral or acid media are can-

didates for the family *Acetobacteraceae*. This family is divided into the genera *Gluconobacter*, which includes three species, and *Acetobacter*, in which seven species have been identified (29). Only the species *A. diazotrophicus* is capable of fixing N₂ (13). On the basis of these and other phenotypic features used for a satisfactory identification (29), we considered that the diazotrophic isolates recovered from the coffee plant environment belong to the family *Acetobacteraceae*. Phenotypic identification was confirmed by the SSU rRNA genes obtained with total DNA digested with *NcoI* and *SphI* (data not shown). Moreover, we have considered it suitable to assign these N₂-fixing isolates to the genus *Acetobacter* because they were capable of oxidizing ethanol, first to acetic acid and then further to CO₂ and H₂O (overoxidation of ethanol), which is the main feature of the genus (8, 29). Other differential phenotypic characteristics analyzed (Table 4) were in agreement with descriptions for this genus (8, 29). By taking into account the differential phenotypic features at the species level (8, 29) and with support from the MLEE assays and the molecular characteristics reported previously, such as hybridization patterns of *nifHDK* genes and of SSU rDNA genes (5, 6) as well as DNA-DNA homology experiments, a majority of the N₂-fixing *Acetobacter* isolates (all strains from division I, excluding CFN-Cf 56) recovered from rhizosphere soil and from inside tissues of coffee plants were considered to belong to the species *A. diazotrophicus*. Although *A. diazotrophicus* strains were reported to form water-soluble brown pigments on GYC medium (7), some of the *A. diazotrophicus* strains (CFN-Cf 52, UAP-Cf 51, Cf 53, and Cf 58) recovered from the coffee plant environment did not produce them (Table 4). However, water-soluble brown pigment production is not a typical feature of the genus *Acetobacter* but rather of the genus *Frateuria* (30). Thus, the *A. diazotrophicus* isolates not producing water-soluble brown pigments could be considered more typical acetobacters.

A number of *Acetobacter* isolates recovered from the coffee plant rhizosphere, capable of fixing N₂ under microaerobic conditions, should not be assigned to the species *A. diazotrophicus* because remarkable differences were observed. We propose that the strains corresponding to ETs included in divisions III, IV, and V may be regarded as different N₂-fixing species of the genus *Acetobacter*. This is based on the fact that all of these isolates were easily differentiated from *A. diazotrophicus* by several morphological and biochemical traits, including the electrophoretic mobility patterns of metabolic enzymes, rendering coefficients of genetic distance as high as 0.950.

Furthermore, these *Acetobacter* isolates differed in SSU rRNA RFLP patterns, and they had a very low level of DNA homology with *A. diazotrophicus* PAI 5^T. These data are strong evidence to designate other diazotrophic species of the genus *Acetobacter*, but more N₂-fixing isolates from other coffee-producing areas of Mexico have to be isolated to provide an extended phenotypic and genetic analysis useful for taxonomic validation of a new species. This is specially true for strain CFN-Cf 56, which is a unique isolate with peculiar characteristics. For instance, on the basis of the MLEE data and SSU rRNA RFLP patterns, the strain CFN-Cf 56 should be regarded as belonging to the species *A. diazotrophicus*. However, on the basis of DNA-DNA homology values, this strain may be considered a new nitrogen-fixing species of the genus *Acetobacter*. Nevertheless, plasmid differences could account for the low DNA-DNA homology values between strain CFN-Cf 56 and strain PAI 5^T.

Natural habitats of acetic acid bacteria are sugar and alcohol solutions, with flowers and many fruits being excellent habitats (29). *A. diazotrophicus*, an endophytic bacterial species, occurs predominantly in vegetatively propagated plants (9). It has been recovered from inside tissues of sucrose-accumulating plants such as sugarcane (10, 11, 19), from a few samples of washed roots and aerial parts of *Pennisetum purpureum* cv. Cameroon, and from sweet potato stems and roots (9) as well as from different genera of mealybugs associated with sugarcane plants (2, 5). This species has not been recovered from other plants nor from nonrhizosphere soils collected from sugarcane fields or other sites (9, 19). However, *A. diazotrophicus* was detected in sugarcane rhizosphere soil by the indirect enzyme-linked immunosorbent assay method (20).

In this study, *A. diazotrophicus* was isolated mainly from coffee plant rhizosphere soils but also, in lower frequencies, from surface-sterilized stems and roots of coffee plants. Our results strongly contrast those of previous reports in which *A. diazotrophicus* isolation from the sugarcane rhizosphere was a rare event. The occurrence of VAM fungus species associated with coffee plants (28) could explain the frequent isolation of *A. diazotrophicus* from the rhizosphere since this bacterial species has been reported to occur inside VAM fungal spores (23), and these were not discarded from the soil inoculated into the culture medium. However, our results did not support the former possibility because we were unable to recover *A. diazotrophicus* from VAM spores. The recovery of N₂-fixing acetobacters from the rhizosphere, we suspect, could be in relation to the organic matter content present in the rhizosphere of coffee plants. While sugarcane is burned off before cutting, eliminating virtually all organic matter originating both from senescent and trash leaves, in coffee-producing areas, the falling fruit and leaves of these trees are largely accumulated in the soil. Perhaps this organic matter could protect bacteria against soil physicochemical factors. In addition, the organic matter degradation by microbial communities will enrich the rhizosphere with carbon (sugar) sources usable by acetobacters. Contrasting with previous results, our data demonstrated that *A. diazotrophicus* is capable of colonizing plants propagated through seeds in addition to plants propagated vegetatively.

Clearly, the distribution of *A. diazotrophicus* is wider than early reports indicated. Genotype ET 1 is extensively distributed, not only among the previously reported hosts (5, 6) but also among coffee plant isolates. Perhaps ET 1 strains have a large colonization capacity that could be related to the presence of a highly conserved plasmid (pAd170) that exists in most ET 1 *A. diazotrophicus* isolates (6). This plasmid has not been observed in isolates corresponding to other ETs (6; un-

published results). pAd170 was also observed in ET 1 isolates recovered both from the rhizosphere and inside coffee plants (data not shown).

Coffee-associated genotypes, except ET 1, were never identified among more than 70 *A. diazotrophicus* strains recovered from previously well-known hosts collected in diverse countries (5, 6). Because isolates of *A. diazotrophicus* recovered from the coffee plant environment are closely related genetically to isolates recovered from sugarcane, the existence of a common lineage is suggested.

It is worth noting that even though the isolation of *A. diazotrophicus* from internal tissues was infrequent, it was usually recovered from coffee plants grown in acid soils. The infrequency of recovery of *A. diazotrophicus* from coffee plant tissues may be related to the difficulties in homogenizing roots and stems, since these plants are highly lignified and very hard to blend. The presence of *A. diazotrophicus* in acid soils suggests that the transmission of this species into coffee plants could be through VAM fungi, as reported for sugarcane plants (22) and *Sorghum bicolor* (17). Also, we considered that transmission of *A. diazotrophicus* could be through mealybugs, as suggested previously (2), or directly into coffee plant fruit, as occurs in pineapple with other acetic acid bacteria (15). Nevertheless, we were not able to recover *A. diazotrophicus* nor any other N₂-fixing acetobacters from coffee plant fruit or mealybugs (*Planococcus citri*). From these results, we may speculate that *A. diazotrophicus* uses root tips and cracks at lateral root junctions to enter the coffee plants, as suggested for sugarcane plants (18).

Our results support the hypothesis that in nature there are many more N₂-fixing bacteria to be identified and also strongly suggest that endophytic diazotrophic bacteria are more prevalent than previously was thought.

Considering the great economic importance of coffee in the world, and the difficulties of obtaining nitrogen fertilizers (14), we consider that coffee-associated N₂-fixing acetobacters may be agronomically important because they could supply part of the nitrogen that the crop requires, as has been suggested in the case of sugarcane with its associated endophytic nitrogen-fixing bacteria.

ACKNOWLEDGMENTS

We are grateful to Michael Hynes and Michael Dunn for reviewing the paper and to R. Bustillos-Cristales, Minerva Rosas, L. Martínez-Aguilar, and M. E. Nava-Herrera for technical assistance. We thank C. Abarca-Ocampo (SAGAR-Guerrero) for his support in collecting coffee plants and the associated mealybugs.

This work was supported in part by grant UNAM-DGAPA-IN209496.

REFERENCES

- Arrieta, J., L. Hernández, A. Coego, V. Suárez, E. Balmori, C. Menéndez, M.-F. Petit-Glatron, R. Chambert, and G. Selman-Housein. 1996. Molecular characterization of the levansucrase gene from the endophytic sugarcane bacterium *Acetobacter diazotrophicus* SRT4. *Microbiology* **142**:1077-1085.
- Ashbolt, N. J., and P. E. Inkerman. 1990. Acetic acid bacterial biota of the pink sugarcane mealybug, *Saccharococcus sacchari*, and its environs. *Appl. Environ. Microbiol.* **56**:707-712.
- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl. 1987. Current protocols in molecular biology. John Wiley & Sons, Inc., New York, N.Y.
- Brosius, J., T. J. Dull, D. D. Sleeter, and H. F. Noller. 1981. Gene organization and primary structure of a ribosomal RNA operon from *Escherichia coli*. *J. Mol. Biol.* **148**:107-127.
- Caballero-Mellado, J., L. E. Fuentes-Ramírez, V. M. Reis, and E. Martínez-Romero. 1995. Genetic structure of *Acetobacter diazotrophicus* populations and identification of a new genetically distant group. *Appl. Environ. Microbiol.* **61**:3008-3013.
- Caballero-Mellado, J., and E. Martínez-Romero. 1994. Limited genetic diversity in the endophytic sugarcane bacterium *Acetobacter diazotrophicus*.

- Appl. Environ. Microbiol. **60**:1532–1537.
7. Cavalcante, V. A., and J. Döbereiner. 1988. A new acid-tolerant nitrogen fixing bacterium associated with sugarcane. *Plant Soil* **108**:23–31.
 8. De Ley, J., J. Swings, and F. Gosselé. 1984. Genus I. *Acetobacter* Beijerinck 1898, 215^{AL}, p. 268–274. In N. R. Krieg and J. G. Holt (ed.), *Bergey's manual of systematic bacteriology*, vol. 1. The Williams & Wilkins Co., Baltimore, Md.
 9. Döbereiner, J. 1993. History and new perspectives of diazotrophs in association with non-leguminous plants. *Symbiosis* **13**:1–13.
 10. Dong, Z., M. Heydrich, K. Bernard, and M. E. McCully. 1995. Further evidence that the N₂-fixing endophytic bacterium from the intercellular spaces of sugarcane stems is *Acetobacter diazotrophicus*. *Appl. Environ. Microbiol.* **61**:1843–1846.
 11. Fuentes-Ramírez, L. E., T. Jiménez-Salgado, I. R. Abarca-Ocampo, and J. Caballero-Mellado. 1993. *Acetobacter diazotrophicus*, an indoleacetic acid producing bacterium isolated from sugarcane cultivars of Mexico. *Plant Soil* **154**:145–150.
 12. Gerdemann, J. W., and T. H. Nicolson. 1963. Spores of mycorrhizal endogone species extracted from soil by wet sieving and decanting. *Trans. Br. Mycol. Soc.* **46**:235–244.
 13. Gillis, M., K. Kersters, B. Hoste, D. Janssens, R. M. Kroppenstedt, M. P. Stephan, K. R. S. Teixeira, J. Döbereiner, and J. De Ley. 1989. *Acetobacter diazotrophicus* sp. nov., a nitrogen-fixing acetic acid bacterium associated with sugarcane. *Int. J. Syst. Bacteriol.* **39**:361–364.
 14. Hall, C. A. S., and M. H. P. Hall. 1993. The efficiency of land and energy use in tropical economies and agriculture. *Agric. Ecosyst. Environ.* **46**:1–30.
 15. Hayward, A. C. 1974. Latent infections by bacteria. *Annu. Rev. Phytopathol.* **12**:87–97.
 16. Hurek, T., and B. Reinhold-Hurek. 1995. Identification of grass-associated and toluene-degrading diazotrophs, *Azoarcus* spp., by analyses of partial 16S ribosomal DNA sequences. *Appl. Environ. Microbiol.* **61**:2257–2261.
 17. Isopi, R., P. Fabri, M. Del Gallo, and G. Puppi. 1995. Dual inoculation of *Sorghum bicolor* (L.) Moench spp. *bicolor* with vesicular arbuscular and *Acetobacter diazotrophicus*. *Symbiosis* **18**:43–55.
 18. James, E. K., V. M. Reis, F. L. Olivares, J. I. Baldani, and J. Döbereiner. 1994. Infection of sugar cane by the nitrogen-fixing bacterium *Acetobacter diazotrophicus*. *J. Exp. Bot.* **45**:757–766.
 19. Li, R. P., and I. C. MacRae. 1991. Specific association of diazotrophic acetobacters with sugarcane. *Soil Biol. Biochem.* **23**:999–1002.
 20. Li, R. P., and I. C. MacRae. 1992. Specific identification and enumeration of *Acetobacter diazotrophicus* in sugarcane. *Soil Biol. Biochem.* **24**:413–419.
 21. Mascaraña-Esparza, M. A., R. Villa-González, and J. Caballero-Mellado. 1988. Acetylene reduction and indoleacetic acid production by *Azospirillum* isolates from cactaceous plants. *Plant Soil* **106**:91–95.
 22. Paula, M. A., S. Urquiaga, J. O. Siqueira, and J. Döbereiner. 1992. Synergistic effects of vesicular-arbuscular mycorrhizal fungi and diazotrophic bacteria on nutrition and growth of sweet potato (*Ipomoea batatas*). *Biol. Fertil. Soils* **14**:61–66.
 23. Paula, M. A., J. O. Siqueira, and J. Döbereiner. 1993. Ocorrência de fungos micorrízicos vesículoarbusculares e de bactérias diazotróficas na cultura da batata-doce. *Rev. Bras. Ci. Solo, Campinas* **17**:349–356.
 24. Reinhold-Hurek, B., T. Hurek, M. Gillis, B. Hoste, M. Vancanneyt, K. Kersters, and J. De Ley. 1993. *Azoarcus* gen. nov., nitrogen-fixing proteobacteria associated with roots of Kallar grass (*Leptochloa fusca* (L.) Kunth), and description of two species, *Azoarcus indigenus* sp. nov. and *Azoarcus communis* sp. nov. *Int. J. Syst. Bacteriol.* **43**:574–584.
 25. Selander, R. K., D. A. Caugant, H. Ochman, J. M. Musser, M. N. Gilmour, and T. S. Whittam. 1986. Methods of multilocus enzyme electrophoresis for bacterial population genetics and systematics. *Appl. Environ. Microbiol.* **51**:873–884.
 26. Sievers, M., W. Ludwig, and M. Teuber. 1994. Phylogenetic positioning of *Acetobacter*, *Gluconobacter*, *Rhodopila* and *Acidiphilium* species as a branch of acidophilic bacteria in the alpha-subclass of proteobacteria based on 16S ribosomal DNA sequences. *Syst. Appl. Microbiol.* **17**:189–196.
 27. Sievers, M., W. Ludwig, and M. Teuber. 1994. Revival of the species *Acetobacter methanolicus* (ex. Uhlig et al. 1986) nom. rev. *Syst. Appl. Microbiol.* **17**:352–354.
 28. Siqueira, J. O., O. Saggin, A. Colozzi-Filho, E. Oliveira, and P. T. G. Guimaraes. 1993. Ecology and application of VAM fungi in coffee crop in Brazil, p. 78. In *Abstracts of the 9th North American Conference on Mycorrhizae*. Guelph, Ontario, Canada.
 29. Swings, J. 1992. The genera *Acetobacter* and *Gluconobacter*, p. 2268–2286. In A. Balows, H. G. Trüper, M. Dworkin, W. Harder, and K.-H. Schleifer (ed.), *The prokaryotes. A handbook on the biology of bacteria: ecophysiology, isolation, identification, applications*, vol. III. Springer-Verlag, New York, N.Y.
 30. Swings, J., J. De Ley, and M. Gillis. 1984. Genus III. *Frateruia* Swings, Gillis, Kersters, De Vos, Gosselé and De Ley, 1980, 1547^{VP}, p. 210–213. In N. R. Krieg and J. G. Holt (ed.), *Bergey's manual of systematic bacteriology*, vol. 1. The Williams & Wilkins Co., Baltimore, Md.
 31. Torsvik, V., J. Goksoyr, and F. L. Daae. 1990. High diversity in DNA of soil bacteria. *Appl. Environ. Microbiol.* **56**:782–787.
 32. Young, J. P. W. 1992. Phylogenetic classification of nitrogen-fixing organisms, p. 43–86. In G. Stacey, R. H. Burris, and G. Evans (ed.), *Biological nitrogen fixation*. Chapman and Hall, New York, N.Y.
 33. Zhou, J., M. R. Fries, J. C. Chee-Sanford, and J. M. Tiedje. 1995. Phylogenetic analysis of a new group of denitrifiers capable of anaerobic growth on toluene and description of *Azoarcus toluolyticus* sp. nov. *Int. J. Syst. Bacteriol.* **45**:500–506.