

Signalling of abscisic acid to regulate plant growth

Axel Himmelbach, Monika Iten and Erwin Grill

Lehrstuhl für Botanik, Technische Universität München, Arcisstr. 16, D-80333 München, Germany

Abscisic acid (ABA) mediated growth control is a fundamental response of plants to adverse environmental cues. The linkage between ABA perception and growth control is currently being unravelled by using different experimental approaches such as mutant analysis and microinjection experiments. So far, two protein phosphatases, ABI1 and ABI2, cADPR, pH, and Ca^{2+} have been identified as main components of the ABA signalling pathway. Here, the ABA signal transduction pathway is compared to signalling cascades from yeast and mammalian cells. A model for a bifurcated ABA signal transduction pathway exerting a positive and negative control mechanism is proposed.

Keywords: abscisic acid; *Arabidopsis thaliana*; growth control; cell cycle; MAP kinase; signalling cascade

1. INTRODUCTION

ABA is involved in many physiological and developmental processes throughout the life cycle of plants. In the early phase of a plant's life, the phytohormone ABA regulates seed maturation and the maintenance of embryo dormancy. Later, at the onset of ontogenesis, it mediates several adaptational responses towards environmental cues such as desiccation, cold as well as salt stress, and acts as a negative growth regulator.

There are both fast and slow responses to ABA. The triggering of stomatal closure is the best-studied fast response to ABA and involves the orchestration of ion channel activities at the plasmalemma and tonoplast (MacRobbie 1995). The signal transduction pathway responsible for relaying the ABA signal to its targets (cation and anion channels (Blatt & Thiel 1993)) seems to be preformed and comprises an ABA-receptor(s), protein kinases and protein phosphatases (Pei *et al.* 1997).

Most ABA-mediated actions are characterized by a comparatively slow response and require differential gene regulation. A plethora of ABA-upregulated genes as well as a few ABA-downregulated genes such as *cab* and *cdc2a* (coding for the chlorophyll *ab* binding protein and a cell cycle-dependent kinase, respectively) have been characterized (Chandler & Robertson 1994). Several components involved in the signalling pathway responsible for modulating embryo dormancy have been described. This advance was made possible by isolating the affected genes from ABA-response mutants. Thus, the transcriptional regulators VPI from maize (McCarty *et al.* 1991) and its homologue from *Arabidopsis* ABI3 (Giraudat *et al.* 1992), two highly homologous members of the protein phosphatases 2C family ABI1 and ABI2 (Leung *et al.* 1994, 1997; Meyer *et al.* 1994; Rodriguez *et al.* 1998; Leube *et al.* 1998), and a farnesyl transferase ERA1 (Cutler *et al.* 1996) have been identified.

Of these signalling components only the two protein phosphatases seem to act also during the vegetative

growth phase of the plant. The analysis of the *abi1* and *abi2* loci supported a pleiotropic ABA-insensitive phenotype of the mutants (Koornneef *et al.* 1984; Finkelstein & Somerville 1990) and a dominant negative action of the mutant gene products associated with a strongly reduced enzymatic activity (Bertauche *et al.* 1996; Leung *et al.* 1997; Rodriguez *et al.* 1998; Leube *et al.* 1998). Both mutants are characterized by a wilted phenotype reflecting a lesion in stomatal regulation and by ABA-insensitive growth (Koornneef *et al.* 1984; Meyer *et al.* 1994).

To date, little is known about the molecular mechanism of ABA-mediated growth regulation. Here we will concentrate on this aspect of ABA responses.

2. TARGETS OF ABA ACTION MEDIATING GROWTH CONTROL

Approximately 30 years ago, it was observed that ABA inhibits DNA replication in root tips and embryos of *Fraxinus excelsior* (van Overbeek *et al.* 1967; Villiers 1968). Histochemical studies of root tissues supported the notion of an ABA-mediated inhibition of cell elongation and of an arrest in mitotic cell activity, presumably in the G1-phase (Nagl 1972; de la Torre *et al.* 1972).

Since these investigations it has been well established that ABA generally modifies plant growth by inhibiting shoot and root growth as an antagonist of auxin (Pilet & Chanson 1981; van Volkenburgh & Davies 1983; Kutschera & Schopfer 1986; Pilet & Saugy 1987). The role of ABA in modulating the growth rate of roots has been intensely studied and was found to be concentration-dependent. Whereas a negative correlation between ABA content in the elongation zone of maize roots and growth rate was observed (Saugy *et al.* 1989), low concentrations of ABA can even stimulate root growth, thus providing a basis for the explanation of the biphasic response of roots upon application of inhibitory ABA concentrations (Pilet & Barlow 1987). Exogenous ABA application inhibited root growth after an initial phase of increased growth lasting




Figure 1. Schematic signalling cascade of (a) the pheromone response in yeast and (b) the mitogenic response in mammalian cells. In both cases the signal is perceived by a receptor (REC) localized at the plasma membrane and relayed via an MAPK cascade into the nucleus. See the text for details.

several hours (Mulkey *et al.* 1983; Robertson *et al.* 1990). Inhibition of ABA biosynthesis by fluoridone or ABA-deficiency due to a mutational lesion, however, did not affect root growth of maize (Moore & Smith 1984) indicating that endogenous ABA is not a central growth regulator unless ABA levels are enhanced e.g. by cold or desiccation.

The root response towards inhibitory ABA concentrations is characterized by an inhibition of cell elongation and subsequent osmotic adjustment of the root apex. The reduction in root elongation was reciprocally correlated with an increase in both osmotic potential of the medium and the endogenous ABA level (Ribaut & Pilet 1991). The same adaptational response is observed under conditions of drought stress and coincides with both increased root hair formation and decrease in the size of the meristematic zone of the root tip (Robertson *et al.* 1990). The application of ABA mimics this response of roots (Schnall & Quatrano 1992). The low water potential response of wheat roots was observed at exogenous ABA concentrations as low as 0.1 μM ABA and was sufficient to increase the turgor pressure by approximately 50% (Jones *et al.* 1987).

In summary, these data imply that ABA imposes a bimodal growth control by regulating the potential of the cell to enlarge, possibly by turgor control, and by inducing mitotic growth arrest in plants in accordance with its role as a negative growth regulator.

3. PARADIGMS OF CELL CYCLE CONTROL

Mitotic growth arrest is well-known to be induced under adverse conditions in animal cells by preventing

the transition of G1 into the S-phase of the cell cycle (Ron 1994). The tumour suppressor p53 (Baker *et al.* 1990) and the retinoblastoma protein pRb (Weinberg 1995) were identified as central links in the pathway. Extensive phosphorylation of pRb at a checkpoint of the cell cycle (the restriction point) within the mid-G1 phase inactivates its growth-inhibitory action (Bartek *et al.* 1996). The multistep phosphorylation is initiated and controlled by the cell cycle machinery and provides a means to integrate internal and external signals. Central components of the mammalian cell cycle are the cyclins and their specific kinases (cdks), the proliferating cell nuclear antigen, and the cdk inhibitors such as p16, p21 and p27 (Sherr & Roberts 1995; Coats *et al.* 1996). Among the key regulators of S-phase entry are the cyclinD-cdk4/6 as well as cyclinE-cdk2 complexes that promote cell cycle progression by specific protein phosphorylation (Morgan 1995).

The pathway leading to mitotic growth arrest in the mating response of *Saccharomyces cerevisiae* has been unravelled through extensive mutant analyses (Levin & Errede 1995). A schematic presentation of the pathway is given in figure 1a. During mating of yeast cells, the release of a pheromone from a mating partner triggers a signalling cascade from the plasma membrane to the nucleus that finally activates a transcription factor and FAR1. The signalling pathway is activated by the pheromone acting on a membrane receptor that in turn activates a trimeric G-protein. The G-protein transmits the signal by initiating a phosphorylation cascade leading from STE20 and STE5 to the serine-threonine protein kinases STE11 and STE7, and subsequently to the kinase FUS3 that finally phosphorylates the transcription factor

STE12 and FAR1. FAR1 blocks DNA replication by inhibiting the cyclin-cdc2 kinase complex while STE12 regulates response-specific transcription.

The last steps of the signalling cascade are similar to the mitogenic response of mammalian cells that is induced by epidermal growth factor (figure 1b) and results in the opposite response, i.e. mitotic cell division (Robinson & Cobb 1997). Here, activation of a membrane-bound tyrosine receptor kinase activates, via the involvement of two receptor-associated proteins GRB2 and SOS, the small G-protein ras that activates a serine-threonine phosphorylation cascade by recruiting raf to the plasma membrane. The raf protein phosphorylates the mitogen activated protein kinase kinase (MEK) that regulates the activity of the MAP kinase (MAPK). The MAPK controls the heterodimeric transcription factor consisting of jun and fos but also regulates protein biosynthesis via the cell cycle-controlled S6 protein kinase (Blenis 1993). MEK and MAPK are structural homologues of STE7 and FUS3; hence both responses, cell cycle inhibition in yeast and cell cycle activation in mammals, rely on a conserved cascade of protein kinases.

In addition to the mitogenic response, mammalian MAPK pathways also regulate stress responses resulting in growth inhibition e.g. by the transforming growth factor TGF- β (Waskiewicz & Cooper 1995; Yamaguchi *et al.* 1995). Thus, distinct MAPK cascades emerge as general signalling pathways that regulate cell cycle progression.

4. SIGNALLING OF GROWTH INHIBITION

The question of how ABA could exert control of the cell cycle in plants is currently unresolved. One likely possibility is through a mechanism analogous to the mammalian system involving the regulation of the phosphorylation state of pRb and an MAPK pathway. In plants, indeed, the presence of the components of the cell cycle machinery (Doonan & Fobert 1997) as well as of MAPK cascades (Mizoguchi *et al.* 1996, 1997; Ligterink *et al.* 1997; Machida *et al.* 1997) have been unequivocally demonstrated. The observed structural conservation of the proteins is reflected by either functional complementation or protein-protein interactions among animal or yeast forms and plant components (Hirt *et al.* 1991; Dahl *et al.* 1995; Ball & Lane 1996; Ach *et al.* 1997a). This supports the view that cell cycle control mechanism are conserved throughout the eukaryotes.

Protein kinases homologous to Cdc2 were initially cloned from alfalfa (Hirt *et al.* 1991) and *Arabidopsis* (Ferreira *et al.* 1991; Hirayama *et al.* 1991) and identified as key components of the plant cell cycle machinery required for G1 to S-phase transition and for entry into mitosis (Martinez *et al.* 1992). By using a reporter gene under control of a cdc2 promoter, it was shown that Cdc2 expression is not always linked to cell proliferation but it always precedes cell division (Hemerly *et al.* 1993). Treatment of the transgenic plants with ABA resulted in the strong transcriptional down-regulation of the reporter gene. In contrast, auxin increased cdc2 gene expression within minutes after hormone application (John *et al.* 1993).

Several different cyclins including type A, B and D have been isolated (Hemerly *et al.* 1992; Day & Reddy

1994; Renaudin *et al.* 1996). Interestingly, several cyclins of the D-type revealed a pRb interaction motif (Fürst *et al.* 1996; Dahl *et al.* 1995; Soni *et al.* 1995). In fact, pRb proteins of maize have been identified and were capable of interacting with this cyclin type (Xie *et al.* 1996; Ach *et al.* 1997b) pointing to a similar role for these cyclins in cell cycle control as in the animal system. Recently, cdk inhibitors of plants were identified by their specific interaction with Cdc2 (Wang *et al.* 1997; de Veyler *et al.* 1997).

To understand the molecular mechanisms for the ABA-mediated control of plant growth and mitotic growth arrest, our group took a genetic approach to identify loci providing insensitivity to the growth inhibitory action of ABA in *Arabidopsis*. Mutants of *Arabidopsis* insensitive to the germination-inhibiting action of ABA have already been characterized as representatives of five independent loci, *abil* to *abi5* (Koornneef *et al.* 1984; Finkelstein 1994). In addition, response mutants hypersensitive towards ABA have been characterized (Cutler *et al.* 1996). However, of these mutants only *abil* and *abi2* seem to reveal an altered ABA sensitivity during vegetative growth. Additional ABA-insensitive *Arabidopsis* mutants were isolated by screening more than 1×10^5 seedlings of the chemically mutagenized M2 generation in the presence of growth-inhibiting concentrations of ABA. The intention was to identify all the signalling loci involved in the pathway leading to the growth inhibition by ABA (T. Ehrler, M. Iten and E. Grill, unpublished results). Genetic complementation analyses of the 22 identified mutant plants revealed lesions in eight new loci, *GCA1* to *GCA8* (growth control exerted by ABA). Physiological analyses, including stomatal regulation as well as dormancy of the mutant seeds, enabled the successful isolation of two additional pleiotropic mutants *gca1* and *gca2* that phenotypically resemble *abil* and *abi2*. All other loci of the mutants, *gca3* to *gca8*, affect, monotropically, the growth response.

Identification of the corresponding genes certainly will aid our understanding of how the hormonal signal controls cell growth and division. So far there is no evidence to support the involvement of an MAPK cascade in this response. However, the cascade has been implicated in the ABA-mediated antagonism of gibberellin action during barley seed germination (Knetsch *et al.* 1996)—a process in which a phospholipase D has recently been postulated to be involved (Ritchie & Gilroy 1998). In yeast and *Arabidopsis*, PP2Cs have been identified to counteract the activation of an MAPK cascade involved in osmoregulation and stress signalling, respectively (Maeda *et al.* 1994; Meskiene *et al.* 1998). Thus, the characterization of ABI1 and ABI2 as two PP2Cs (Leung *et al.* 1994, 1997; Meyer *et al.* 1994; Rodriguez *et al.* 1998; Leube *et al.* 1998) could suggest a similar role of these phosphatases as negative regulators of a MAPK cascade.

The unresolved question of ABA signal transduction has recently been tackled by the identification of cyclic ADP ribose as a second messenger of ABA action capable of inducing specifically ABA-dependent gene transcription (Wu *et al.* 1997). In detailed experiments Wu and co-workers demonstrated that NAD might serve as a precursor of cADP ribose and that an inhibitor of cADP ribose, 8-amino-cADP ribose, blocked both ABA signalling as well as cADP ribose action. cADP ribose




Figure 2. Model of a bifurcated ABA signal pathway exerting plant growth control. The binding of ABA to a specific receptor results in the generation of cADP ribose and a concomitant rise of the pH in the cytosol. cADP release triggers an increase in the concentration of cytosolic calcium. The pH signal activates the protein phosphatases ABI1 and ABI2 which dephosphorylate a postulated active repressor (R_a) of the signalling cascade thereby generating the inactive form R_{in} . The other branch of the pathway exerts a positive regulation and is activated by the released calcium ions. This leads to the presence of inactivated repressor and by several, so far unknown, steps to the control of cell enlargement and cell division. The gene products of the identified loci *GCA1* to *GCA8* represent candidates involved in this signalling pathway. *GCA1* and *GCA2* loci pleiotropically affect the ABA response while *GCA3* to *GCA8* seem to modulate specifically the ABA-mediated growth control.

seems to act by triggering the release of calcium ions into the cytosol (Allen *et al.* 1995; Wu *et al.* 1997) and by a second hitherto unknown mechanism, thereby providing specific regulation of ABA-modulated genes.

The specific signature of ABA and cADP ribose action observed via calcium release could be provided by the regulation of the proton concentration in the cytosol. ABA application to plant cells results within minutes in the alkalization of the cytosol by approximately 0.2–0.4 pH units, whereas an acidification concomitant with calcium release has been demonstrated after auxin administration (Gehring *et al.* 1990; Irving *et al.* 1992). The alkalization triggered by ABA would be sufficient to increase the PP2C activity of ABI1 and ABI2 by a factor of more than two (Leube *et al.* 1998, A. Himmelbach, unpublished results) and thereby could change the phosphorylation state of substrates critical for ABA signal relay. Guard cells of plants expressing the mutant *abil* phosphatase, although ABA-insensitive, do still respond

to ABA by a pH shift in the cytosol (Armstrong *et al.* 1995). Provided that these guard cells also show ABA-dependent calcium mobilization, then cADP ribose appears to act upstream of the protein phosphatases. Furthermore, because both ABI1 and ABI2 are involved in mediating ABA growth responses, cADP ribose must also be capable of regulating this response. Thus, cADP ribose emerges as a central player of ABA action.

In a speculative scenario, the signalling events leading to ABA-mediated growth control could be envisaged in the following way (see figure 2): ABA interacts with (a) receptor(s) generating by unknown steps a shift of the cytosolic pH into the alkaline and the formation of cADP ribose which results in an increase in the concentration of cytosolic calcium. The pH-shift activates ABI1 and ABI2 which in turn dephosphorylate a postulated negative regulator, a repressor, of ABA-signalling. This step inactivates the repressor and allows ABA-signal progression.

The behaviour of mutant forms of *abil* and *abi2* that act dominantly and provide ABA-insensitivity while revealing a strongly reduced enzymatic activity could be explained by sequestering the active repressor and not being able to inactivate it (Bertauche *et al.* 1996; Leung *et al.* 1997; Rodriguez *et al.* 1998; Leube *et al.* 1998; Sheen 1998). In accordance with the model, the rescue of the ABA-response by the application of serine–threonine kinase inhibitors to *abil*-expressing guard cells (Armstrong *et al.* 1995; Pei *et al.* 1997) could be explained by the generation of the dephosphorylated (inactivated) repressor. Interestingly, ABA signalling still required the presence of ABA in these experiments. Thus, ABA and possibly cADP ribose seem to act on a bifurcated signalling pathway by deactivating a negative control mechanism via ABI1 and ABI2, as well as by activating a positive control pathway in which calcium ions are involved as secondary messengers.

This work was funded by the Swiss Science Foundation and the EU biotech project BIO4-CT96-0062. The support of the Fonds der Chemischen Industrie is gratefully acknowledged.

REFERENCES

- Ach, R. A., Durfee, T., Miller, A. B., Taranto, P., Hanley-Bowdoin, L., Zambryski, P. C. & Gruissem, W. 1997a RRB1 and RRB2 encode maize retinoblastoma-related proteins that interact with a plant D-type cyclin and geminivirus replication protein. *Molec. Cell. Biol.* **17**, 5077–5086.
- Ach, R. A., Taranto, P. & Gruissem, W. 1997b A conserved family of WD-40 proteins binds to the retinoblastoma protein in both plants and animals. *Pl. Cell* **9**, 1595–1606.
- Allen, G. J., Muir, S. R. & Sanders, D. 1995 Release of Ca^{2+} from individual plant vacuoles by both InsP3 and cyclic ADP-ribose. *Science* **268**, 735–737.
- Armstrong, F., Leung, J., Grabov, A., Brearley, J., Giraudat, J. & Blatt, M. R. 1995 Sensitivity to abscisic acid of guard-cell K^+ channels is suppressed by *abil-1*, a mutant *Arabidopsis* gene encoding a putative protein phosphatase. *Proc. Natn. Acad. Sci. USA* **92**, 9520–9524.
- Baker, S. J., Markowitz, S., Fearson, E. R., Wilson, J. K. V. & Vogelstein, B. 1990 Suppression of human colorectal carcinoma cell growth by wild-type p53. *Science* **249**, 912–915.
- Ball, K. L. & Lane, D. P. 1996 Human and plant proliferating-cell nuclear antigen have a highly conserved binding site for the p53-inducible gene product p21WAF1. *Eur. J. Biochem.* **237**, 854–861.

- Bartek, J., Bartkova, J. & Lukas, J. 1996 The retinoblastoma protein pathway and the restriction point. *Curr. Opin. Cell Biol.* **8**, 805–814.
- Bertauche, N., Leung, J. & Giraudat, J. 1996 Protein phosphatase activity of abscisic acid insensitive 1 (ABI1) protein from *Arabidopsis thaliana*. *Eur. J. Biochem.* **241**, 193–200.
- Blatt, M. R. & Thiel, G. 1993 Hormonal control of ion channel gating. *A. Rev. Pl. Physiol. Pl. Molec. Biol.* **44**, 543–567.
- Blenis, J. 1993 Signal transduction via the MAP kinases: proceed at your own RSK. *Proc. Natn. Acad. Sci. USA* **90**, 5889–5892.
- Chandler, P. M. & Robertson, M. 1994 Gene expression regulated by abscisic acid and its relation to stress tolerance. *A. Rev. Pl. Physiol. Pl. Molec. Biol.* **45**, 113–141.
- Coats, S., Flanagan, W. M., Nourse, J. & Roberts, J. M. 1996 Requirement of p27Kipl for restriction point control of the fibroblast cell cycle. *Science* **272**, 877–880.
- Cutler, S., Ghassemian, M., Bonetta, D., Cooney, S. & McCourt, P. 1996 A protein farnesyl transferase involved in abscisic acid signal transduction in *Arabidopsis*. *Science* **273**, 1239–1241.
- Dahl, M., Meskiene, I., Bogre, L., Ha, D. T., Swoboda, I., Hubmann, R., Hirt, H. & Heberle-Bors, E. 1995 The D-type alfalfa cyclin gene *cycMs4* complements G1 cyclin-deficient yeast and is induced in the G1 phase of the cell cycle. *Pl. Cell* **7**, 1847–1857.
- Day, I. S. & Reddy, A. S. N. 1994 Cloning of a family of cyclins from *Arabidopsis thaliana*. *Biochem. Biophys. Acta* **1218**, 1796–1799.
- de la Torre, C., Diez, J. L., Lopez-Saez, J. F. & Gimenez-Martin, G. 1972 Effect of abscisic acid on the cytological components of the root growth. *Cytologia* **37**, 197–205.
- de Veylder, L., Segers, G., Glab, N., Casteels, P., Montagu, M. V. & Inze, D. 1997 The *Arabidopsis* Cks1At protein binds the cyclin-dependent kinases Cdc2aAt and Cdc2bAt. *FEBS Lett.* **412**, 446–452.
- Doonan, J. & Fobert, P. 1997 Conserved and novel regulators of the plant cell cycle. *Curr. Opin. Cell Biol.* **8**, 824–830.
- Ferreira, P. C. G., Hemerly, A. S., van Montagu, M. & Inze, D. 1991 The *Arabidopsis* functional homolog of the p34-cdc2 protein kinase. *Pl. Cell* **3**, 531–540.
- Finkelstein, R. R. 1994 Mutations at two new *Arabidopsis* ABA response loci are similar to the *abi3* mutations. *Pl. J.* **5**, 765–771.
- Finkelstein, R. R. & Somerville, C. R. 1990 Three classes of abscisic acid (ABA)-insensitive mutations in *Arabidopsis thaliana*. *Pl. Physiol.* **94**, 1172–1179.
- Fürst, R. A., Soni, R., Murray, J. A. & Lindsey, K. 1996 Modulation of cyclin transcript levels in cultured cells of *Arabidopsis thaliana*. *Pl. Physiol.* **112**, 1023–1033.
- Gehring, C. A., Irving, H. R. & Parish, R. W. 1990 Effects of auxin and abscisic acid on cytosolic calcium and pH in plant cells. *Proc. Natn. Acad. Sci. USA* **87**, 9645–9649.
- Giraudat, J., Hauge, B. M., Valon, C., Smalle, J., Parcy, F. & Goodman, H. M. 1992 Isolation of the *Arabidopsis* ABI3 gene by positional cloning. *Pl. Cell* **4**, 333–344.
- Hemerly, A., Bergounioux, C., van Montagu, M., Inze, D. & Ferreira, P. 1992 Genes regulating the plant cell cycle: isolation of a mitotic-like cyclin from *Arabidopsis thaliana*. *Proc. Natn. Acad. Sci. USA* **89**, 3295–3299.
- Hemerly, A. S., Ferreira, P., de Almeida-Engler, J., van Montagu, M., Engler, G. & Inze, D. 1993 CDC2a expression in *Arabidopsis* linked with competence for cell division. *Pl. Cell* **5**, 1711–1723.
- Hirayama, T., Imajuku, Y., Anai, T., Matsui, M. & Oka, A. 1991 Identification of two cell-cycle-controlling *cdc2* gene homologs in *Arabidopsis thaliana*. *Gene* **105**, 159–166.
- Hirt, H., Pay, A., Gyogyey, J., Bako, L., Nemeth, K., Bogre, L., Schweyen, R. J., Heberle-Bors, E. & Dudits, D. 1991 Complementation of a yeast cell cycle mutant by an alfalfa complementary DNA encoding a protein kinase homologous to p34-cdc2. *Proc. Natn. Acad. Sci. USA* **88**, 1636–1640.
- Irving, H. R., Gehring, C. A. & Parish, R. W. 1992 Changes in cytosolic pH and calcium of guard cells precede stomatal movements. *Proc. Natn. Acad. Sci. USA* **89**, 1790–1794.
- John, P. C. L., Zhang, K., Dong, C., Diederich, L. & Wightman, F. 1993 P34-cdc2 related proteins in control of cell cycle progression, the switch between division and differentiation in tissue development, and stimulation of division by auxin and cytokinin. *Austr. J. Pl. Physiol.* **20**, 503–520.
- Jones, H., Leigh, R. A., Tomos, A. D. & Jones, R. G. W. 1987 The effect of abscisic acid on cell turgor pressures, solute content and growth of wheat roots. *Planta* **170**, 257–262.
- Knetsch, M. L. W., Wang, M., Snaar-Jagalska, B. E. & Heimovaara-Dijkstra, S. 1996 Abscisic acid induces mitogen-activated protein kinase activation in barley aleurone protoplasts. *Pl. Cell* **8**, 1061–1067.
- Koornneef, M., Reuling, G. & Karszen, C. M. 1984 The isolation of abscisic acid-insensitive mutants of *Arabidopsis thaliana*. *Physiol. Pl.* **61**, 377–383.
- Kutschera, U. & Schopfer, P. 1986 In vivo measurement of cell-wall extensibility in maize coleoptiles: effects of auxin and abscisic acid. *Planta* **169**, 437–442.
- Leube, M. P., Grill, E. & Amrhein, N. 1998 ABI1 of *Arabidopsis* is a protein serine/threonine phosphatase highly regulated by the proton and magnesium ion concentration. *FEBS Lett.* **424**, 100–104.
- Leung, J., Bouvier-Durand, M., Morris, P.-C., Guerrier, D., Chefdor, F. & Giraudat, J. 1994 *Arabidopsis* ABA response gene ABI1: features of a calcium-modulated protein phosphatase. *Science* **264**, 1448–1452.
- Leung, J., Merlot, S. & Giraudat, J. 1997 The *Arabidopsis* ABSCISIC ACID-INSENSITIVE2 (ABI2) and ABI1 genes encode homologous protein phosphatases 2C involved in abscisic acid signal transduction. *Pl. Cell* **9**, 759–771.
- Levin, D. E. & Errede, B. 1995 The proliferation of MAP kinase signaling pathways in yeast. *Curr. Opin. Cell Biol.* **7**, 197–202.
- Ligterink, W., Kroj, T., Zurnieden, U., Hirt, H. & Scheel, D. 1997 Receptor-mediated activation of a MAP kinase in pathogen defense of plants. *Science* **276**, 2054–2057.
- McCarty, D. R., Hattori, T., Carson, C. B., Vasil, V. & Lazar, M. 1991 The viviparous-1 developmental gene of maize encodes a novel transcriptional activator. *Cell* **66**, 895–905.
- Machida, Y., Nishihama, R. & Kitakura, S. 1997 Progress in studies of plant homologs of mitogen-activated protein (MAP) kinase and potential upstream components in kinase cascades. *Crit. Rev. Pl. Sci.* **16**, 481–496.
- MacRobbie, E. A. 1995 ABA-induced ion efflux in stomatal guard cells: multiple actions of ABA inside and outside the cell. *Pl. J.* **7**, 565–576.
- Maeda, T., Wurgler-Murphy, S. M. & Saito, H. 1994 A two-component system that regulates an osmosensing MAP kinase cascade in yeast. *Nature* **369**, 242–245.
- Martinez, M. C., Jorgenson, J. E., Lawton, M. A., Lamb, C. J. & Doerner, P. W. 1992 Spatial pattern of *cdc2* expression in relation to meristem activity and cell proliferation during plant development. *Proc. Natn. Acad. Sci. USA* **89**, 7360–7364.
- Meskiene, I., Bogre, L., Glaser, W., Balog, J., Brandstötter, M., Zwerger, K., Ammerer, G. & Hirt, H. 1998 MP2C, a plant protein phosphatase 2C, functions as a negative regulator of mitogen-activated protein kinase pathways in yeast and plants. *Proc. Natn. Acad. Sci. USA* **95**, 1938–1943.
- Meyer, K., Leube, M. P. & Grill, E. 1994 A protein phosphatase 2C involved in ABA signal transduction in *Arabidopsis thaliana*. *Science* **264**, 1452–1455.

- Mizoguchi, T., Irie, K., Hirayama, T., Hayashida, N., Yamaguchi-Shinozaki, K., Matsumoto, K. & Shinozaki, K. 1996 A gene encoding a mitogen-activated protein kinase kinase is induced simultaneously with genes for a mitogen-activated protein kinase and an S6 ribosomal protein kinase by touch, cold, and water stress in *Arabidopsis thaliana*. *Proc. Natn. Acad. Sci. USA* **93**, 765–769.
- Mizoguchi, T., Ichimura, K. & Shinozaki, K. 1997 Environmental stress response in plants: the role of mitogen-activated protein kinases. *Trends Biotech.* **15**, 15–19.
- Moore, R. & Smith, J. D. 1984 Growth, graviresponsiveness and abscisic acid content of *Zea mays* seedlings treated with fluridone. *Planta* **162**, 342–344.
- Morgan, D. O. 1995 Principles of CDK regulation. *Nature* **374**, 131–134.
- Mulkey, T. J., Evans, M. L. & Kuzmanoff, K. M. 1983 The kinetics of abscisic acid action on root growth and gravitropism. *Planta* **157**, 150–157.
- Nagl, W. 1972 Selective inhibition of cell cycle stages in the *Allium* root meristem by colchicine and growth regulators. *Am. J. Bot.* **59**, 346–351.
- Pei, Z. M., Kuchitsu, K., Ward, J. M., Schwarz, M. & Schroeder, J. I. 1997 Differential abscisic acid regulation of guard cell slow anion channels in *Arabidopsis* wild-type and *abil* and *abi2* mutants. *Pl. Cell* **9**, 409–423.
- Pilet, P. E. & Barlow, P. W. 1987 The role of abscisic acid in root growth. A critical examination. *Pl. Sci. Lett.* **21**, 99–106.
- Pilet, P. E. & Chanson, A. 1981 Effect of abscisic acid on maize root growth. A critical examination. *Pl. Sci. Lett.* **31**, 117–122.
- Pilet, P. E. & Saugy, M. 1987 Effect on root growth of endogenous and applied IAA and ABA. A critical reexamination. *Pl. Physiol.* **83**, 33–38.
- Renaudin, J. P. (and 13 others) 1996 Plant cyclins: a unified nomenclature for plant A-, B- and D-type cyclins based on sequence organization. *Pl. Molec. Biol.* **32**, 1003–1018.
- Ribaut, J. M. & Pilet, P. E. 1991 Effects of water stress on growth, osmotic potential and abscisic acid content of maize roots. *Physiol. Pl.* **81**, 156–162.
- Ritchie, S. & Gilroy, S. 1998 Abscisic acid signal transduction in the barley aleurone is mediated by phospholipase D activity. *Proc. Natn. Acad. Sci. USA* **95**, 2697–2702.
- Robertson, J. M., Hubick, K. T., Yeung, E. C. & Reid, D. M. 1990 Developmental responses to drought and abscisic acid in sunflower roots. I. Root growth, apical anatomy, and osmotic adjustment. *J. Exp. Bot.* **41**, 325–338.
- Robinson, M. J. & Cobb, M. H. 1997 Mitogen-activated protein kinase pathways. *Curr. Opin. Cell Biol.* **9**, 180–186.
- Rodriguez, P. L., Benning, G. & Grill, E. 1998 ABI2, a second protein phosphatase 2C involved in abscisic acid signal transduction in *Arabidopsis*. *FEBS Lett.* **421**, 185–190.
- Ron, D. 1994 Inducible growth arrest: new mechanistic insights. *Proc. Natn. Acad. Sci. USA* **91**, 1985–1986.
- Saugy, M., Mayor, G. & Pilet, P. E. 1989 Endogenous ABA in growing maize roots: light effects. *Pl. Physiol.* **89**, 622–627.
- Schnall, J. A. & Quatrano, R. S. 1992 Abscisic acid elicits the water-stress response in root hairs of *Arabidopsis thaliana*. *Pl. Physiol.* **100**, 216–218.
- Sheen, J. 1998 Mutational analysis of protein phosphatase 2C involved in abscisic acid signal transduction in higher plants. *Proc. Natn. Acad. Sci. USA* **95**, 975–980.
- Sherr, C. J. & Roberts, J. M. 1995 Inhibitors of mammalian G1 cyclin-dependent kinases. *Genes Dev.* **9**, 1149–1163.
- Soni, R., Carmichael, J. P., Shah, Z. H. & Murray, J. A. 1995 A family of cyclin D homologs from plants differentially controlled by growth regulators and containing the conserved retinoblastoma protein interaction motif. *Pl. Cell* **7**, 85–103.
- van Overbeek, J., Loeffler, J. E. & Mason, M. I. R. 1967 Dormin (abscisin II), inhibitor of plant DNA synthesis. *Science* **156**, 1497–1499.
- van Volkenburgh, E. & Davies, W. J. 1983 Inhibition of light-stimulated leaf expansion by abscisic acid. *J. Exp. Bot.* **35**, 110–120.
- Villiers, T. A. 1968 An autoradiographic study of the effect of the plant hormone abscisic acid on nucleic acid and protein metabolism. *Planta* **82**, 342–354.
- Wang, H., Fowke, L. C. & Crosby, W. L. 1997 A plant cyclin-dependent kinase inhibitor gene. *Nature* **386**, 451–452.
- Waskiewicz, A. J. & Cooper, J. A. 1995 Mitogen and stress response pathways: MAP kinase cascades and phosphatase regulation in mammals and yeast. *Curr. Opin. Cell Biol.* **7**, 798–805.
- Weinberg, R. A. 1995 The retinoblastoma protein and cell cycle control. *Cell* **81**, 323–330.
- Wu, Y., Kuzma, J., Marechal, E., Graeff, R., Lee, H. C., Foster, R. & Chua, N.-H. 1997 Abscisic acid signaling through cyclic ADP-ribose in plants. *Science* **278**, 2126–2130.
- Xie, Q., Sanz-Burgos, P., Hannon, G. J. & Gutierrez, C. 1996 Plant cells contain a novel member of the retinoblastoma family of growth regulatory proteins. *EMBO J.* **15**, 4900–4908.
- Yamaguchi, K., Shirakabe, K., Shibuya, H., Irie, K., Oishi, I., Ueno, N., Taniguchi, T., Nishida, E. & Matsumoto, K. 1995 Identification of a member of the MAPKKK family as a potential mediator of TGF-beta signal transduction. *Science* **270**, 2008–2011.