



## What's new at the back of the eye?

### EDITORIAL



#### Ivo Kocur

VISION 2020 Global Coordinator, World Health Organization, Prevention of Blindness and Deafness, Chronic Diseases and Health Promotion, CH-1211, Geneva 27, Switzerland.

### The changing demographic pattern of visual impairment

VISION 2020 and the prevention of visual impairment are moving into the political and public spotlight. Recent WHO discussions on the growing incidence of chronic, age-related, non-communicable diseases indicate that visual health and its preservation are now receiving attention. Recent publications illustrate the changing demographic pattern of visual impairment, implications for public health and possible interventions to control sight-threatening conditions. Both *The State of the World's Sight*, *VISION 2020: the Right to Sight, 1999-2005*,<sup>1</sup> published jointly by WHO and the International Agency for the Prevention of Blindness (IAPB), and an article in the *World Health Bulletin*, November 2004,<sup>2</sup> illustrate these trends. The recent WHO data on blindness give clear evidence: among the causes of blindness, the share of chronic, age-related, non-communicable potentially blinding eye conditions is dramatically increasing (Figure 1, over page). *The State of the World's Sight* provides an insight into what has thus far been achieved in the prevention of visual impairment through international alliances and collaboration since the launch of VISION 2020 in 1999. Attempts to modernise the attitude of health care providers towards the preservation of visual health have been further cultivated by the representatives of many WHO member states when adopting, in 2003, the Resolution on the Elimination of Avoidable Blindness, and when preparing a new resolution for the World Health Assembly in 2006. What does all this mean for eye health care professionals – new challenges, new opportunities?

*Editorial continues over page* ➤



An ophthalmologist examining the posterior segment of the eye. UK

Richard Leung

### IN THIS ISSUE...

#### EDITORIAL

- 1 **What's new at the back of the eye?**  
Ivo Kocur

#### ARTICLES

- 4 **What's new in age-related macular degeneration?**  
David Yorston
- 6 **Can we grow new retina?**  
Geeta K Vemuganti

#### REPORT

- 8 **What will be new at the back of the eye in the year 2020?**  
Shaheen Shah reports from the World Ophthalmology Congress, including a glossary

#### EVIDENCE-BASED EYE CARE

- 11 **Cochrane Eyes and Vision Group (CEVG) systematic review activity on back of the eye treatments**  
Cochrane Eyes and Vision Group

#### HOW TO...

- 12 **How to prescribe spectacles for presbyopia**  
R nee du Toit

#### EXCHANGE

Including an inexpensive tool for examining the fundus, using eye shields, a free histopathology service and a KAP study from South India

#### NEWS AND NOTICES

Including findings from the Reader Survey





The journal is produced in collaboration with the World Health Organization



Volume 19 | Issue No. 57 | March 2006

#### Editor

Victoria Francis

#### Editorial Committee

Professor Allen Foster  
Dr Clare Gilbert  
Dr Murray McGavin  
Dr Ian Murdoch  
Dr GVS Murthy  
Dr Daksha Patel  
Dr Richard Wormald  
Dr David Yorston

#### Regional Consultants

Dr Grace Fobi (Cameroon)  
Professor Gordon Johnson (UK)  
Dr Susan Lewallen (Tanzania)  
Dr Wanjiku Mathenge (Kenya)  
Dr Babar Qureshi (Pakistan)  
Dr Yuliya Semenova (Kazakhstan)  
Dr B R Shamanna (India)  
Professor Hugh Taylor (Australia)  
Dr Andrea Zin (Brazil)

#### Advisors

Dr Liz Barnett (Teaching and Learning)  
Catherine Cross (Infrastructure and Technology)  
Sue Stevens (Ophthalmic Nursing and Teaching Resources)

#### Administration

Ann Naughton (Administrative Director)  
Anita Shah (Editorial/Administrative Assistant)

#### Editorial Office

Community Eye Health Journal  
International Centre for Eye Health  
London School of Hygiene and Tropical Medicine,  
Keppel Street, London WC1E 7HT, UK.  
Tel: +44 207 612 7964/72  
Fax: +44 207 958 8317  
Email: Anita.Shah@Lshtm.ac.uk

#### Information Service

Sue Stevens  
Email: Sue.Stevens@Lshtm.ac.uk  
Tel: +44 207 958 8168

#### On-line Edition ([www.jceh.co.uk](http://www.jceh.co.uk))

Sally Parsley  
Email: admin@jceh.co.uk

Community Eye Health Journal is published four times a year and **sent free to developing country applicants**. French, Chinese and Indian editions are also available. Please send details of your name, occupation and postal address to Community Eye Health Journal, at the address above. Subscription rates for applicants elsewhere: one year UK£28/US\$45; two years UK£50/US\$80. Send credit card details or an international cheque/banker's order made payable to London School of Hygiene and Tropical Medicine to the address above.

#### Website

Back issues are available at  
[www.jceh.co.uk](http://www.jceh.co.uk)

Content can be downloaded in both HTML and PDF formats.

© International Centre for Eye Health, London  
Articles may be photocopied, reproduced or translated provided these are not used for commercial or personal profit. Acknowledgements should be made to the author(s) and to Community Eye Health Journal. All graphics by Victoria Francis unless stated otherwise.

ISSN 0953-6833

The journal is produced in collaboration with the World Health Organization. Signed articles are the responsibility of the named authors alone and do not necessarily reflect the policies of the World Health Organization. The World Health Organization does not warrant that the information contained in this publication is complete and correct and shall not be liable for any damages incurred as a result of its use. The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned.

## EDITORIAL Continued

### Meeting the challenge through specialised eye care services

One challenging aspect of the latest WHO data on the changing pattern of blindness prevalence in the world is that the demand for high-quality eye care services is expected to increase. Predicted population growth, along with the increase in life expectancy, will substantially alter the picture of future recipients of eye care. The numbers of people affected by sight-threatening eye conditions requiring highly specialised health services are likely to increase.

One such condition is diabetes. Diabetic retinopathy accounts for about 5 per cent of the global pattern of blindness. WHO is expecting a significant increase in the incidence of diabetes in all geographic regions. For instance, in many highly populated countries in Asia, the number of individuals suffering from diabetes will double by 2030.

Glaucoma is another example which, from the public health perspective, may be even more challenging. Controlling blindness from glaucoma requires early detection, life-long treatment and the compliance of patients. It is estimated that glaucoma is responsible for 1.2 per cent of global blindness. However, numerous unresolved issues in glaucoma control deprive many populations from effectively planned and delivered specialised eye care services.

The WHO data also indicate the growing threat of age-related macular degeneration. The increasing share of this eye condition is progressively reshaping the ratio of avoidable sight-threatening eye conditions to those in which conventional eye care is thus far failing. Cataract is another age-related eye condition causing avoidable blindness. Given that it has been one of the major stimuli for technological research and development in eye care, the failure of modern society to adequately control



Richard Leung

Direct ophthalmoscopy provides a magnified view of the fundus.UK

## ‘The recent WHO data on blindness give clear evidence: among the causes of blindness, the share of chronic, age-related, non-communicable potentially blinding eye conditions is dramatically increasing.’

cataract blindness is particularly shameful. The last WHO data indicate that cataract is still responsible for almost half of global blindness. Its cure is known, and several options for its surgical treatment have been tested and broadly implemented to reasonably fit healthcare budgets, even in the poorest societies. Demographic changes and longevity, along with the lack, for the foreseeable future, of knowledge about cataract prevention, will further increase the demand for cataract surgical services. Age-related macular degeneration poses a different set of challenges, as the condition cannot be prevented, and only some forms are amenable to very expensive, time-consuming interventions which are of limited benefit. The mainstay will need to be rapid expansion of low vision services, along with raising awareness among the general public and eye care professionals.


The implications of demographic changes and the ageing of populations are a growing concern. However, preventable and treatable sight-threatening eye conditions occurring in children must also remain among the top priorities. The World Health Bulletin of November 2004<sup>2</sup> also sent a strong warning signal emphasising the need to address uncorrected refractive errors, an often overlooked public health issue relevant to all age groups. The importance of refractive errors in the pattern of visual impairment is likely to be substantially greater than generally assumed.

### The need for targeted action and an interdisciplinary approach

The current situation presents us with a new opportunity for targeted action. The preservation of visual health requires, more than ever, a coordinated approach by health care providers, who will develop comprehensive, integrated eye care services. An effectively designed eye care component in primary, secondary and tertiary health care is an essential prerequisite to respond adequately to the changing pattern of avoidable visual impairment. Over the long term, vertical interventions and projects oriented to a single eye disease may no longer bring the desired achievements.

The nature of the causes of blindness which WHO considers to be preventable and/or treatable, and therefore avoidable, requires an interdisciplinary approach for long-term control. For example, controlling diabetic retinopathy is complex, requiring: health education to prevent diabetes; early diagnosis of diabetes; comprehensive, cost-

Figure 1: Global causes of blindness as a percentage of total blindness in 2002



effective screening programmes to detect diabetics with treatable retinopathy; laser treatment and follow-up; and patient education and compliance. In many countries, attitudes about diabetes control are being reshaped by constituting national diabetes control programmes, rather than by pursuing the control of this disease through uncoordinated efforts within individual medical sub-specialties. In those countries where the number of diabetic patients is rapidly increasing, introducing control programmes on the scale needed will be extremely challenging. We need to educate more ophthalmologists to manage diabetic retinopathy, make appropriate technology available, develop new programmes, establish new partnerships, and develop a major public health education strategy.

### Integrating eye care into national health plans

Health care providers in the most affluent societies and in many other countries with modern health care systems are becoming increasingly interested in comprehensive, integrated care. In this context, eye care providers have three responsibilities: to be ready to explain the known and tested options in blindness control to national

health care planners and providers; to be proactive in health care policy development; and to take part in updating national health care strategies where eye care should be addressed. It would be disappointing to lose the opportunity for integrating eye care into national health care plans because of a failure to provide health care planners with an adequate briefing on the management of sight-threatening conditions. Several national health care areas should be explored, especially neonatal care, health care programmes for mothers and children, preventive health care programmes for working-age populations as well as for the elderly. As the world copes more adequately with the challenges of blindness control, eye care professionals will play a vital role in advocating for comprehensive health care which integrates and adequately addresses visual health.

#### References

- 1 World Health Organization and International Agency for the Prevention of Blindness. The state of the world's sight. VISION 2020: The Right to Sight 1999 - 2005. Geneva: WHO, 2005.
- 2 Resnikoff S, Pascolini D, Etya'ale D, Kocur I, Pararajasegaram R, Pokharel G, Mariotti S. Global data on visual impairment in the year 2002. Bulletin of the World Health Organization 2004; 82: 844-85. [www.who.int/bulletin/volumes/82/11/en/844.pdf](http://www.who.int/bulletin/volumes/82/11/en/844.pdf)