

Plasmid-Mediated Tet M in *Haemophilus ducreyi*

MARILYN C. ROBERTS

Department of Pathobiology, School of Public Health and Community Medicine, University of Washington,
Seattle, Washington 98195

Received 8 February 1989/Accepted 2 June 1989

A tetracycline-resistant *Haemophilus ducreyi* strain isolated in the United States was shown to carry a 34-megadalton plasmid which hybridized with the 1.8-kilobase *KpnI-HindIII* Tet M probe. The complete Tn916 transposon hybridized with five different bands from this plasmid, suggesting homology throughout the length of the transposon.

The Tet M determinant has been found in the chromosome of commensal *Neisseria* species (14, 28) and gram-negative anaerobic species (28), as well as in a variety of gram-positive species (5, 12, 28). This gene codes for a protein that binds to the ribosomes, protecting them from tetracycline (3), and shares amino acid sequence homology with various elongation factors (30).

Five tetracycline-resistant *Haemophilus ducreyi* strains isolated in Jacksonville, Fla., were obtained from the Centers for Disease Control. The *H. ducreyi* strains were grown and cleared lysates were prepared (19, 22). Duplicate dot blots were hybridized with a 1.8-kilobase (kb) *KpnI-HindIII* internal Tet M fragment from plasmid pUW-JKB1 (2, 31) or an internal 1.27-kb *HincII* Tet B fragment from plasmid pRT11 (17, 32). Four of the strains hybridized with the Tet B probe as described previously (17) and were not examined further, while the fifth strain (86.039418) hybridized only with the Tet M probe and was chosen for further study.

The DNA from this strain was prepared and electrophoresed through an agarose gel (19). A small plasmid which hybridized with the TEM β -lactamase gene from plasmid pJ13 was observed (4, 27) and thus assumed to be a β -lactamase plasmid (1, 7). A large plasmid of 34 megadaltons (MDa) was also seen as a faint band in some DNA preparations and was named pMR9418. The pMR9418 DNA was found to hybridize with the Tet M probe, while no hybridization was observed between the Tet M probe and the chromosome or the small plasmid.

DNAs from strain 86.039418, *Enterococcus faecalis* strain DS160 (5), and Tc^r *Neisseria gonorrhoeae* strain 83.022650 (15, 18) (Table 1) were treated with the restriction enzyme *HincII* or *HincII* and *EcoRI*. Southern blots were prepared and probed with the 1.8-kb Tet M determinant or the Tn916 probe. The Tn916 probe used was a 17.7-kb fragment from plasmid pAM120, which carries the complete Tn916 DNA sequence, originally from DS160 (5). When the 1.8-kb Tet M probe was used, all three DNAs had one hybridizing band. This is similar to what has been observed for other Tet M-containing species (2, 23, 27, 28). With the Tn916 probe, at least five hybridizing bands were seen with the *H. ducreyi* DNA, and multiple bands were seen with the *E. faecalis* DNA (5, 12) (Fig. 1). In contrast, the *N. gonorrhoeae* DNA hybridized only with a single band. A 4.9-kb *HincII* Tet M fragment from this *N. gonorrhoeae* strain has been cloned and shown to contain the structural Tet M determinant, which hybridizes with one band with either probe (my unpublished data). Therefore, I conclude that the *N. gonorrhoeae* does not contain the entire transposon, while the *H. ducreyi* strain appears to have homology along the whole

length of the Tn916 transposon. This is similar to the Tc^r *Clostridium difficile* strains, except in this species the Tet M determinant is located in the chromosome (12).

In most of the species with the Tet M determinant, including streptococci, the determinant is located in the chromosome (5, 12, 23, 24, 27, 28). The one exception is the 25.2-MDa plasmid (14, 15, 18). This plasmid is highly related to the indigenous gonococcal 24.5-MDa conjugative plasmid, and we have hypothesized that it was created by the transposition of the Tet M determinant into the 24.5-MDa plasmid (14, 18). We examined whether plasmid pMR9418 was related to the 25.2-MDa plasmid family by using a 24.5-MDa plasmid, pMR388, which had >60% of its DNA sequences in common with pMR650 but does not carry the Tet M determinant, as a probe (14). The large R plasmids described previously in *H. ducreyi* share 70 to 80% of their DNA sequences in common with R plasmids isolated from *H. influenzae* (1, 9, 22, 29). We used a well-characterized *H. influenzae* plasmid, RSF007 (8), to determine whether the *H. ducreyi* plasmid was related to this plasmid class as well. Southern blots were prepared and hybridized under stringent conditions (50% formamide). No homology was found between plasmids pMR9418 and pMR388, or with the RSF007 plasmid, suggesting that the pMR9418 plasmid is not highly related to either plasmid family.

We examined whether pMR9418 could be transferred to other *Haemophilus* spp. We chose, as recipients, the well-characterized *H. influenzae* strains Rd (29) and G32, carrying the 4.1-MDa β -lactamase plasmid (RSF0885) isolated from *H. parainfluenzae* HR-8 (8). The *H. influenzae* strains were grown as described previously (22), while the *H. ducreyi* strain was grown overnight on four agar plates supplemented with tetracycline. The *H. ducreyi* cells were removed and suspended in 1 ml of brain heart infusion to which broth-grown *H. influenzae* was added. The suspension was mixed, plated directly on four agar plates, incubated for 16 h, and then removed. A 1:10 dilution series was cultured on agar plates supplemented with rifampin, rifampin and tetracycline, rifampin and ampicillin, tetracycline and ampicillin, ampicillin, or tetracycline at the following concentrations (micrograms per milliliter): tetracycline, 10; streptomycin, 250; rifampin, 10; ampicillin, 10. Plates were incubated at 36.5°C in 5% CO₂ and were examined for growth at 24 and 48 h. Colonies that grew on the antibiotic-containing media were scored for streptomycin resistance if the recipient strain was Rd, and all transconjugants were screened for plasmids.

Both recipients, Rd and G32, acquired tetracycline resistance from the donor *H. ducreyi* strain at a frequency that

TABLE 1. Bacterial strains and their plasmids

| Strain | Plasmid (MDa) | Resistance ^a | Reference |
|---|-------------------|---|------------|
| <i>H. ducreyi</i> 86.039418 | 7.0, 34 (pMR9418) | Ap ^r Tc ^r | This study |
| <i>H. influenzae</i> Rd G32(RSF0885) ^b | None 4.1 | Ery ^r Rif ^r Str ^r Ap ^r | 29 8 |
| <i>N. gonorrhoeae</i> 83.022650 | 25.2 (pMR650) | Tc ^r | 18 |
| <i>E. faecalis</i> DS160 | None | Tet ^r | 5 |

^a Plasmid-mediated resistance: Ap^r, ampicillin; Tc^r, tetracycline. Chromosomal resistance: Ery^r, erythromycin; Rif^r, rifampin; Str^r, streptomycin; Tet^r, tetracycline (original Tn916).

^b RSF0885 is an *H. parainfluenzae* plasmid which has been transformed into *H. influenzae* G32 (8).

ranged from 10⁻⁶ to 10⁻⁷ per recipient (Table 2). When the transconjugants were lysed, a plasmid of 34 MDa was found in all strains and was shown to hybridize with the 1.8-kb Tet M probe.

Cross-mating experiments were done by using the filter-mating technique (29) (Table 2). Both strains Rd and G32 could transfer the 34-MDa plasmid at higher frequencies (10⁻¹ to 10⁻⁴) compared with the original *H. ducreyi* donor strain (Table 2). This is not unusual, since transfer frequencies are often higher when both donor and recipient are from the same species than when the two represent different species (11, 25).

H. ducreyi R plasmids, in general, do not mobilize the small β-lactamase plasmids to recipients by conjugation. Only one cryptic *H. ducreyi* plasmid has been described which does (7). There was apparent transfer of the ampicillin determinant into strain Rd from the *H. ducreyi* strain. However, we could not maintain the transconjugants for more than one passage on ampicillin media and were never




FIG. 1. Southern blot of 1.0% agarose gel probed with a ³²P-radiolabeled 17.7-kb fragment carrying the complete Tn916 transposon. Lane A, *Hinc*II-restricted DNA from *E. faecalis* DS160; lane B, *Hinc*II- and *Eco*RI-restricted DNA from *N. gonorrhoeae* 83.022650; lane C, *Hinc*II- and *Eco*RI-restricted DNA from *H. ducreyi* 86.039418. Numbers on the side represent sizes, in kilobases, of the *Hind*III lambda and ϕ X174 *Hae*II fragments.

TABLE 2. Transfer of tetracycline and ampicillin plasmids by conjugation^a

| Donor | Frequency with given recipient | | |
|-----------------------|--------------------------------|-------------------------|---------------------------------|
| | Rd | | G32(RSF0885) (Tc ^r) |
| | Tc ^r | Ap ^r | |
| 86.039418 | 3.2 × 10 ⁻⁷ | 4.5 × 10 ^{-6b} | 2.6 × 10 ⁻⁶ |
| 83.022650 | 2.6 × 10 ⁻⁶ | | 1.4 × 10 ⁻⁴ |
| G32(RSF0885, pMR9418) | 1.4 × 10 ⁻⁴ | 2.3 × 10 ^{-6b} | ND ^d |
| G32(RSF0885, pMR650) | 7.8 × 10 ⁻⁶ | 6.5 × 10 ^{-6c} | ND |
| Rd(pMR9418) | ND | | 3.0 × 10 ⁻¹ |
| Rd(pMR650) | ND | | 3.0 × 10 ⁻⁵ |

^a Average of three or more separate experiments.

^b Apparent transfer of β-lactamase plasmids, but not confirmed by agarose gel.

^c Confirmed transfer of β-lactamase plasmids, validated by agarose gel visualization.

^d ND, Not done.

able to demonstrate the β-lactamase plasmid in the transconjugants. This suggests that pMR9418 could not mobilize the small β-lactamase plasmids. Thus, plasmid pMR9418 is similar to other conjugative *Haemophilus* R plasmids in this characteristic and differs from the cryptic *H. ducreyi* conjugative plasmid described previously (7). This, along with the fact the 25.2-MDa plasmid retained its ability to mobilize β-lactamase plasmids (25, 26), suggests that the ancestor of pMR9418 may not be the cryptic *H. ducreyi* plasmid described previously, but more work is needed to confirm this hypothesis.

We transferred plasmid pMR650 from *N. gonorrhoeae* into the two *H. influenzae* recipients by conjugation (Table 2). These transconjugants could then act as donors in further matings (Table 2). When an *H. influenzae* strain carried both the 25.2-MDa plasmid and the 4.1-MDa β-lactamase plasmid, it could transfer both plasmids into a recipient and the β-lactamase plasmid was stable and could be visualized by agarose gels. The resulting transconjugants, which carried both the 25.2- and 4.1-MDa plasmids, could then act as donors and transfer both plasmids to other strains (data not shown).

We have shown that plasmid pMR650 from *N. gonorrhoeae* can transfer into the genus *Haemophilus*, suggesting that at some time we will see clinical isolates with the 25.2-MDa plasmid. This, together with the discovery of the *H. ducreyi* plasmid pMR9418, indicates that it is possible for strains of *Haemophilus* spp. to acquire the Tet M determinant. It also demonstrates that the antibiotic resistance gene pool in *H. ducreyi*, and probably in all bacteria, is not stagnant and indicates that, even in a bacterial population which has a high carriage of one tetracycline determinant (Tet B) (1), a second unrelated determinant can be introduced. How common the Tet M determinant is in *H. ducreyi* is not known and will require the screening of new Tc^r strains. However, as carriage of the Tet M determinant increases in the bacterial populations of humans, the probability that the Tet M determinant will be introduced into other strains within the genus *Haemophilus* would be expected to increase.

I am grateful to S. B. Levy for providing the Tet B probe, to J. S. Knapp for providing the *H. ducreyi* strains, and to J. Lansciardi for technical assistance.

This work was supported by Public Health Service grant AI24136 from the National Institute of Allergy and Infectious Diseases.

LITERATURE CITED

1. Albritton, W. L., I. W. MacClean, L. A. Slaney, A. R. Ronald, and H. G. Deneer. 1984. Plasmid-mediated tetracycline resistance in *Haemophilus ducreyi*. *Antimicrob. Agents Chemother.* **25**:187-190.
2. Brown, J. T., and M. C. Roberts. 1987. Cloning and characterization of *tetM* from a *Ureaplasma urealyticum* strain. *Antimicrob. Agents Chemother.* **31**:1852-1854.
3. Burdett, V. 1986. Streptococcal tetracycline resistance mediated at the level of protein synthesis. *J. Bacteriol.* **165**:564-569.
4. Burdett, V., J. Inamine, and S. Rajagopalan. 1982. Heterogeneity of tetracycline resistance determinants in *Streptococcus*. *J. Bacteriol.* **149**:995-1004.
5. Clewell, D. B., and C. Gawron-Burke. 1986. Conjugative transposons and the dissemination of antibiotic resistance in *Streptococci*. *Annu. Rev. Microbiol.* **40**:635-659.
6. Dang Van, A., G. Bieth, and D. H. Bouanchaud. 1975. Resistance plasmidique a la tetracycline chez *Haemophilus influenzae*. *C.R. Acad. Sci.* **280**:1321-1323.
7. Deneer, H. G., L. Slaney, I. W. MacClean, and W. L. Albritton. 1982. Mobilization of nonconjugative antibiotic resistance plasmids in *Haemophilus ducreyi*. *J. Bacteriol.* **149**:726-732.
8. Elwell, L. P., J. De Graaff, D. Seibert, and S. Falkow. 1975. Plasmid-linked ampicillin resistance in *Haemophilus influenzae* type b. *Infect. Immun.* **12**:404-410.
9. Elwell, L. P., J. R. Saunders, M. H. Richmond, and S. Falkow. 1977. Relationships among some R plasmids found in *Haemophilus influenzae*. *J. Bacteriol.* **131**:356-362.
10. Gaillard, J. L., P. Berche, and P. Sansonetti. 1986. Transposon mutagenesis as a tool to study the role of hemolysin in the virulence of *Listeria monocytogenes*. *Infect. Immun.* **52**:50-55.
11. Genco, C. A., J. S. Knapp, and V. L. Clark. 1984. Conjugation of plasmids of *Neisseria gonorrhoeae* to other *Neisseria* species: potential reservoirs for the β -lactamase plasmid. *J. Infect. Dis.* **150**:397-401.
12. Hachler, H., F. H. Kayser, and B. Berger-Bachi. 1987. Homology of a transferable tetracycline resistance determinant of *Clostridium difficile* with *Streptococcus (Enterococcus) faecalis* transposon Tn916. *Antimicrob. Agents Chemother.* **31**:1033-1038.
13. Jones, J. M., S. C. Yost, and P. A. Pattee. 1987. Transfer of the conjugal tetracycline resistance transposon Tn916 from *Streptococcus faecalis* to *Staphylococcus aureus* and identification of some insertion sites in the staphylococcal chromosome. *J. Bacteriol.* **169**:2121-2131.
14. Knapp, J. S., S. R. Johnson, J. M. Zenilman, M. C. Roberts, and S. A. Morse. 1988. High-level tetracycline resistance resulting from TetM in strains of *Neisseria* spp., *Kingella denitrificans*, and *Eikenella corrodens*. *Antimicrob. Agents Chemother.* **32**:765-767.
15. Knapp, J. S., J. M. Zenilman, J. W. Biddle, G. H. Perkins, W. E. DeWitt, M. L. Thomas, S. R. Johnson, and S. A. Morse. 1987. Frequency and distribution in the United States of strains of *Neisseria gonorrhoeae* with plasmid-mediated, high-level resistance to tetracycline. *J. Infect. Dis.* **155**:819-822.
16. Levy, S. B., A. Buu-Hoi, and B. Marshall. 1984. Transposon Tn10-like tetracycline resistance determinants in *Haemophilus parainfluenzae*. *J. Bacteriol.* **160**:87-94.
17. Marshall, B., M. Roberts, A. Smith, and S. B. Levy. 1984. Homogeneity of transferable tetracycline-resistance determinants in *Haemophilus* species. *J. Infect. Dis.* **149**:1028-1029.
18. Morse, S. A., S. R. Johnson, J. W. Biddle, and M. C. Roberts. 1986. High-level tetracycline resistance in *Neisseria gonorrhoeae* is result of acquisition of streptococcal *tetM* determinant. *Antimicrob. Agents Chemother.* **30**:664-670.
19. Roberts, M., L. P. Elwell, and S. Falkow. 1977. Molecular characterization of two beta-lactamase-specifying plasmids isolated from *Neisseria gonorrhoeae*. *J. Bacteriol.* **131**:557-563.
20. Roberts, M., and S. Falkow. 1977. Conjugal transfer of R plasmids in *Neisseria gonorrhoeae*. *Nature (London)* **266**:630-631.
21. Roberts, M., P. Piot, and S. Falkow. 1979. The ecology of gonococcal plasmids. *J. Gen. Microbiol.* **114**:491-494.
22. Roberts, M. C., L. A. Actis, and J. H. Crosa. 1985. Molecular characterization of chloramphenicol-resistant *Haemophilus parainfluenzae* and *Haemophilus ducreyi*. *Antimicrob. Agents Chemother.* **28**:176-180.
23. Roberts, M. C., S. L. Hillier, J. Hale, K. K. Holmes, and G. E. Kenny. 1986. Tetracycline resistance and *tetM* in pathogenic urogenital bacteria. *Antimicrob. Agents Chemother.* **30**:810-812.
24. Roberts, M. C., and G. E. Kenny. 1987. Conjugal transfer of transposon Tn916 from *Streptococcus faecalis* to *Mycoplasma hominis*. *J. Bacteriol.* **169**:3836-3839.
25. Roberts, M. C., and J. S. Knapp. 1988. Host range of the conjugative 25.2-megadalton tetracycline resistance plasmid from *Neisseria gonorrhoeae* and related species. *Antimicrob. Agents Chemother.* **32**:488-491.
26. Roberts, M. C., and J. S. Knapp. 1988. Transfer of β -lactamase plasmids from *Neisseria gonorrhoeae* to *Neisseria meningitidis* and commensal *Neisseria* species by the 25.2-megadalton conjugative plasmid. *Antimicrob. Agents Chemother.* **32**:1430-1432.
27. Roberts, M. C., L. A. Koutsky, D. LeBlanc, K. K. Holmes, and G. E. Kenny. 1985. Tetracycline-resistant *Mycoplasma hominis* strains containing streptococcal *tetM* sequences. *Antimicrob. Agents Chemother.* **28**:141-143.
28. Roberts, M. C., and B. J. Moncla. 1988. Tetracycline resistance and TetM in oral anaerobic bacteria and *Neisseria perflava-N. sicca*. *Antimicrob. Agents Chemother.* **32**:1271-1273.
29. Roberts, M. C., C. D. Swenson, L. M. Owens, and A. L. Smith. 1980. Characterization of chloramphenicol-resistant *Haemophilus influenzae*. *Antimicrob. Agents Chemother.* **18**:610-615.
30. Sanchez-Pescador, R., J. T. Brown, M. Roberts, and M. S. Urdea. 1988. Homology of the TetM with translational elongation factors: implications for potential modes of tetM conferred tetracycline resistance. *Nucleic Acids Res.* **16**:1218.
31. Sanchez-Pescador, R., J. T. Brown, M. Roberts, and M. S. Urdea. 1988. The nucleotide sequence of the tetracycline resistance determinant tetM from *Ureaplasma urealyticum*. *Nucleic Acids Res.* **16**:1216-1217.
32. Wray, L. V., Jr., R. A. Jorgensen, and W. S. Reznikoff. 1981. Identification of the tetracycline resistance promoter and repressor in transposon Tn10. *J. Bacteriol.* **147**:297-304.