

PostScript

LETTERS

If you have a burning desire to respond to a paper published in *Occupational and Environmental Medicine*, why not make use of our "rapid response" option?

Log on to our website (www.occenvmed.com), find the paper that interests you, and send your response via email by clicking on the "eLetters" option in the box at the top right hand corner.

Providing it isn't libellous or obscene, it will be posted within seven days. You can retrieve it by clicking on "read eLetters" on our homepage.

The editors will decide as before whether to also publish it in a future paper issue.

Musicians playing wind instruments and risk of lung cancer: is there an association?

Lung cancer is an important public health problem. Tobacco is its main risk factor. Occupation is also an important risk factor. Some jobs have shown higher risks than others, but few investigations have asked about activities or hobbies in leisure time¹ in relation to the risk of lung cancer.

A case-control study was performed between 1999 and 2000 in the Santiago de Compostela Health District (Galicia, northwest Spain). A total of 132 cases with confirmed diagnosis of lung cancer and 187 controls were enrolled. Controls underwent trivial surgery at the same hospital as did the cases. A personal interview about lifestyle and activities (past and present) was conducted by a trained researcher.

We found that, besides tobacco and occupational exposure to carcinogens, some leisure time activities were risk factors for lung cancer.¹ Among the cases there were two musicians who played wind instruments, whereas there were no wind instrument players among the controls. The two cases had been playing the clarinet and trombone for 35 and 30 years respectively. Both were ex-smokers (moderate smokers) and played music as a hobby. They had epidermoid lung cancer and were diagnosed at 57 and 76 years of age.

Since in our population the prevalence of persons playing musical instruments and specifically wind instruments is extremely low, we think that this activity might be a risk factor in development of lung cancer. The very low number of persons playing this type of musical instrument is probably a reason for the lack of studies focused on this activity, as many occupational studies of lung cancer and occupation are based on registries of workers. One study² found an increased mortality rate of lung cancer for a category that included painters, potters, musicians, and actors—an inhomogeneous category that did not allow us to extrapolate results. The results were not adjusted according to smoking history.

This hobby requires inspiration and breathing of large volumes of air, making the lung alveoli expand more than in other people. This fact could facilitate the penetrance of carcinogens in the cells of the lung epithelium, and this could be more harmful in smokers. We have found no other studies that have reported this possible association. It would therefore be necessary to explore this association in greater samples of professionally exposed persons in order to ascertain whether this finding is consistent or due to chance.

A Ruano-Ravina, A Figueiras, J M Barros-Dios

Department of Preventive Medicine and Public Health, University of Santiago de Compostela, Spain

Correspondence to: Dr J M Barros-Dios, Department of Preventive Medicine and Public Health, School of Medicine, C/ San Francisco s/n, Universidad de Santiago de Compostela, 15782 Santiago de Compostela, La Coruña, Spain; mrbarrros@usc.es

References

- 1 Ruano-Ravina A, Figueiras A, Barros-Dios JM. Noxious exposures in leisure time and risk of lung cancer: A neglected exposure? *Epidemiology* 2002;13:235-6.
- 2 Minder CE, Beer-Porizek V. Cancer mortality of Swiss men by occupation, 1979-1982. *Scand J Work Environ Health* 1992;18(suppl 3).

How important is personal exposure assessment in the epidemiology of air pollutants?

The paper by Harrison and colleagues¹ and the accompanying editorial by Cherrie² in the October 2002 issue of *Occupational and Environmental Medicine* address the important issue of personal exposure assessment (of air pollutants) in environmental epidemiology. After reading both papers we would like to make some comments with regard to the design, conduct and statistical analysis of the study by Harrison *et al* and at the same time answer the question raised by Cherrie in his editorial.

Coming from the occupational exposure assessment arena it is interesting to see that our environmental colleagues are still relying to a large extent on static (microenvironmental) sampling and even rely on shadowing to represent personal exposure. The latter brought back memories of old occupational hygiene textbooks with pictures of technicians standing with a sampling probe in the breathing zone of a worker (clearly hindered while carrying out his work task). It is interesting to note that Dr Cherrie's very relevant earlier work³ on whether wearing sampling pumps affects exposure (it hardly did) was not mentioned in both papers.

The paper by Harrison and colleagues¹ clearly states as one of its goals to answer the question "Does modelling through the use of microenvironment measurements and activity diaries produce reliable estimates of personal exposure to air pollutants?". However, in the only setting where personal exposures were actually measured (phase 1, volunteers; with regard to phase 2 we do not think that shadowing results can be seen as equivalent to personally measured exposure) it is hard to grasp from both fig 1 and table 2 which exposure was actually modelled (1 hour averages,

2-3 day averages) and how (a formula was only provided for measurements within the susceptible groups).

When comparing direct personal measurements for CO and PM₁₀ with the modelled results, the authors exclude all data which are not directly comparable—that is, when the volunteer spent most of their time out of house, and all the data for smokers. It is therefore not surprising that good correlations were found between personal and static measurement results. Why were smokers excluded? Was their measured CO exposure representing a different kind of CO leading to a different health effect? We know that excluding smokers or people with unventilated gas heaters is common practice in the statistical analyses of environmental exposures, but this would only make sense if we were expecting different risks from the same exposure originating from different sources.

In fig 1 the authors present 120 comparable data points for 11 individuals; given the repeated nature of the sampling these data points cannot be seen as statistically independent. Putting a simple regression line through these points is therefore not correct and application of a mixed effects model would have been more appropriate. Besides that, when estimating environmental exposure, for instance, for a panel study, we are interested in the full range of exposures both in the temporal and spatial sense (not only for the room with the static sampler). However, Harrison *et al* conclude, "... modelled personal exposure is unable to reflect the variability of measured personal exposures occasioned by the spread of concentrations within given microenvironments".

Both Cherrie and Harrison *et al* claim that microenvironmental sampling would be a good alternative for direct personal exposure measurements that supposedly are "costly and time consuming". However, the costs for sampling microenvironments in a general population study will be far greater if we want to measure all the microenvironments people end up in (for instance, in table 1 seven environments are indicated, and most of them will most likely be different for each study participant). In addition, it will be practically impossible to measure some of these environments as the authors point out. In their study, it was not possible to collect data for all appropriate microenvironments, even for a comparatively small number of subjects.

Recently, a very insightful paper was presented at the X2001 conference in Gothenburg. Seixas and colleagues⁴ showed that in a study to assess occupational noise exposure, a task based methodology (analogous to microenvironmental sampling in environmental exposure assessment) could only account for 30% of variability in daily exposures. They even considered this estimate somewhat optimistic since their estimated noise exposures were derived from the same data on which the daily average exposures were estimated. In addition they clearly pointed out that using simple task based averages that artificially compress exposure variability resulted in a very substantial negative bias in the estimated daily exposure.

In our opinion, we should aim to collect personal exposure measurements when estimating exposure for epidemiological studies.