

Dietary linoleic acid, immune inhibition and disease

Alastair M Sammon

Summary

Review of the evidence available in published literature supports a radical change in viewpoint with respect to disease in countries where maize is the predominant dietary component. In these countries, the pattern of disease is largely determined by a change in immune profile caused by metabolites of dietary linoleic acid. High intake of linoleic acid in a diet deficient in other polyunsaturated fatty acids and in riboflavin results in high tissue production of prostaglandin E2, which in turn causes inhibition of the proliferation and cytokine production of Th1 cells, mediators of cellular immunity. Tuberculosis, measles, hepatoma, secondary infection in HIV and kwashiorkor are all favoured by this reduction in cellular immunity. Diet-associated inhibition of the Th1 subset is a major contributor to the high prevalence of these diseases found in areas of sub-Saharan Africa where maize is the staple.

Keywords: maize; linoleic acid; prostaglandin E2; cellular immunity; kwashiorkor; diet

Tuberculosis, measles, hepatoma, secondary infection in HIV, and kwashiorkor are all favoured by a reduction in cellular immunity. For example, the prevalence of infection with the tubercle bacillus is reported as 34% in Africa and 28% in Western Europe. Yet expected active tuberculosis for 1990 was 220/100 000 for Africa but only 31/100 000 for Europe and other Western countries.¹ A probable 11 300/100 000 of the population of Transkei, South Africa, had active tuberculosis in 1977.² The incidence of active tuberculosis is less a function of prevalence of infection with the bacillus, and more a function of host resistance. High prevalences of tuberculosis have in the past been blamed on social deprivation and dietary deficiency but the variation in the rate of incidence of the disease in different countries is difficult to blame entirely on these factors. For example, the annual notification rate per 100 000 in South Africa is 170 where the GNP is \$2740; in Chad, where the GNP is \$177, the notification rate is 30 per 100 000.

HIV infection in sub-Saharan Africa has a higher rate of progression to AIDS or to death³ than in developed countries. It has also been suggested that there is a higher risk of seroconversion after exposure.⁴ Several reasons have been suggested to explain its rapid heterosexual spread, including poverty, ignorance, the high rate of sexually transmitted disease (STD), cultural practices,⁵ and higher immune activation,⁴ but there are several other possible factors. For example, a seasonal pattern was reported in Uganda,⁶ which suggests an environmental influence on either the contracting of the disease or on its progression. Similarly, STDs, repeated pregnancies, and immune activation⁷ have been blamed for the more rapid progression of HIV to AIDS and to death, with tuberculosis as the most common opportunistic infection.⁸ There is so far inadequate evidence to lay the blame for the very different course of the disease in Africa fully on any or all of these factors.

Kwashiorkor is a disease of children in areas where maize is the main dietary component, and it has been suggested that dietary deficiencies are the source of the problem. Sepsis is a frequent terminal event, and there is poor resistance to diseases such as tuberculosis and measles. Measles in Africa does not follow the relatively benign course that occurs in western countries. The disease and its complications are very severe, and are significant contributors to childhood mortality. It is frequently associated with malnutrition, particularly kwashiorkor.

In all these diseases, general factors with undefined mechanisms have been blamed. This paper puts forward the concept of a specific pathological mechanism related to a relative excess of dietary linoleic acid, as a major factor in the incidence and progression of these diseases.

Methods

The relevant literature was found by a review of appropriate journals and literature, Medline searches, and cross-references, and a hypothesis formulated on the basis of published evidence.

Results

Maize is a high yield crop which grows easily in relatively poor soils, and whose use has been spreading up through Africa for over a century, starting from South Africa where it has been the major dietary constituent for most of this century. It contains 5-6% unsaturated fatty acids, the predominant fat being linoleic acid. It is almost the sole provider of calories in some communities in East, Central and Southern Africa and is increasingly used now in West Africa. The main disease caused by vitamin deficiency in communities who are heavily dependent on maize, is pellagra, but ariboflavinosis also occurs.

Department of Surgery, Level 7,
University of Bristol, Bristol Royal
Infirmary, Bristol BS2 8HW, UK
A M Sammon

Accepted 2 November 1998




Figure 1 Diet and PGE₂ production

HIGH LINOLEIC ACID CAUSES HIGH PROSTAGLANDIN E₂ PRODUCTION

Linoleic acid is a precursor of prostaglandin E₂ (PGE₂). High dietary linoleic acid leads to high intragastric PGE₂,⁹ and, by implication, high PGE₂ levels in the portal vein (figure 1). Symptoms suggestive of a high level of gastric PGE₂ have been reported in a community whose main diet is maize.¹⁰ A volunteer study showed higher urinary PGE₂ levels in volunteers on a high linoleic acid diet than in controls.¹¹ Salivary gland and gingival PGE₂ production is high when linoleic acid intake is high.¹² Plasma PGE₂ in rats was higher in the group fed a diet rich in linoleic acid than in other groups.¹³

Dietary fat intake is reflected in the composition of plasma and body adipose tissue.¹⁴ Thus, diet has a long-term effect on the availability of linoleic acid, and on plasma levels of linoleate and its metabolites. High levels of production of PGE₂ is further enhanced by an absence of other polyunsaturated fatty acids (PUFAs) in the diet, such as fish oils, which contain eicosapentaenoic acid and docosahexaenoic acid, and which inhibit the conversion of linoleic acid to PGE₂.¹⁵ Isomeric PUFAs in hydrogenated foods such as margarine, may also competitively inhibit utilisation of natural PUFAs.

RIBOFLAVIN DEFICIENCY ENHANCES HIGH PGE₂ PRODUCTION

Riboflavin deficiency favours the conversion of linoleic acid to PGE₂. Riboflavin provides the sulphhydryl group for glutathione production, and riboflavin deficiency thus results in glutathione deficiency.¹⁶ Glutathione exerts an inhibitory regulatory influence on prostaglandin synthesis¹⁷ and, in its absence, PGE₂ production rises.¹⁸

A diet high in linoleic acid, low in other polyunsaturated fatty acids and low in vitamins, including riboflavin, is characteristic of the areas of Africa where tuberculosis, HIV, measles and kwashiorkor are rife.




Figure 2 PGE₂ and tuberculosis

PGE₂ DEPRESSES CELLULAR IMMUNITY

The Th1 subset of T-helper lymphocytes mediate delayed-type hypersensitivity reactions. They produce interferon-gamma (IFN-gamma), interleukin-2 (IL-2), and tumour necrosis factor-beta. They are quite distinct from the Th2 subset, which mediate humoral immune response.

PGE₂ depresses cellular immunity associated with the Th1 subset of T cells. IL-2 and IFN-gamma producing ability¹⁹ are reduced, this effect being exerted at the level of priming of the naive T cell.²⁰ Where there is uncontrolled production of PGE₂, there will be uncontrolled and inappropriate depression of cellular immunity, and increased susceptibility to infection. An illustration of this is the depression of cellular immunity which follows haemorrhage and is inhibited by ibuprofen,²¹ an inhibitor of prostaglandin production.

EXCESSIVE LINOLEIC ACID DEPRESSES CELLULAR IMMUNITY

Studies showing that excessive dietary linoleic acid is immunosuppressive, but that concurrent interference with prostaglandin production prevents or reduces depression of cellular immunity,²² provide some support for this theory. The immune depression of trauma and burns appears to be prostaglandin associated, and can be similarly blocked with *n*-3 PUFAs.²³

DEPRESSED CELLULAR IMMUNITY AND DISEASE

Tuberculosis

Protection against tubercle infection is primarily mediated by Th1 cells, and when these are inhibited or reduced in number, host defence is compromised (figure 2). Tuberculosis is strongly associated with the diminished type 1 response²⁴ and depressed CD4+ count characteristic of HIV infection,²⁵ and has been noted to occur at a higher CD4+ count than most other opportunistic infections.²⁶ This may be due to the early loss of Th1 immunity reported in HIV infection at a stage when CD4+ counts are near-normal.²⁷ Multi-drug-resistant tuberculosis is associated with a poor Th1 response.²⁸

This leads to the conclusion that dietary immune deficiency of Th1 function increases susceptibility to tuberculosis. Delayed hypersensitivity reactions to tuberculin have been shown to be reduced in malnourished children.²⁹

HIV infection

HIV infection is associated with a similar immune depression to that caused by PGE₂ (figure 3). There is reduced Th1 subset proliferation and cytokine production. This can take effect before a major drop occurs in total lymphocyte or CD4+ count.²⁷

A pre-existing dietary depression of Th1 subset responsiveness will allow a more rapid progression of HIV infection to AIDS, at least partly by encouraging




Figure 3 PGE₂, HIV, and AIDS




Figure 4 PGE₂ and kwashiorkor




Figure 5 PGE₂ and hepatoma

AIDS-defining opportunistic infections. There will be susceptibility to life-threatening infection early in HIV infection, before AIDS develops, due to the additive effect of dietary and HIV-related immune depression. A chronically reduced subset 1 response of dietary aetiology may also predispose to HIV sero-conversion on exposure, since natural defence against HIV is most probably vested in the Th1 response.³⁰

Kwashiorkor

In kwashiorkor, in addition to the main features of oedema, growth failure, diarrhoea and irritability,³¹ cellular immunity is grossly depressed,³² and affected children are highly susceptible to infectious disease (figure 4).³³ Children with kwashiorkor have decreased T cells in the peripheral blood,³⁴ with preferential loss of CD4+ cells.³⁵

In one study, 25 children who died with kwashiorkor were subjected to autopsy; 21 of them were found to have a 'nutritional thymectomy'.³³ Watts³⁶ confirmed that thymic atrophy is common in malnutrition; in children under one year old, it was most marked in those with kwashiorkor, in whom the atrophy was very gross. Williams³¹ noted that cod liver oil, an excellent source of eicosapentaenoic acid, and therefore an inhibitor of PGE₂ production, is an effective treatment for kwashiorkor.

Measles

In severely malnourished children, measles carries a high mortality rate. In the central hospital in Lesotho, five of 28 malnourished children who contracted measles died.³⁷ Measles virus control is mediated by the Th1 subset of T cells.³⁸ The reasons for the severe course of the disease may include a low level of virus clearing activity, as well as susceptibility to secondary infection.

Hepatoma

Hepatoma is very common in much of Africa. In rat studies, it has been shown that dietary linoleic acid increases hepatoma growth³⁹ (figure 5), while indomethacin inhibits it.⁴⁰ Similarly, fish oil has been found to inhibit hepatic carcinogenesis,⁴¹ and decrease growth.⁴² It is probable that this effect of linoleic acid is mediated through PGE₂. Portal PGE₂ has also been shown to suppress liver-associated immunity, and promote the formation of hepatic metastases.⁴³

Discussion

In the Gambia, evidence has been found of a lifelong change of immune competence brought about by foetal malnutrition.⁴⁴ While this may cause an irreversible change, arising from specific episodes of intervention or deficiency, in this article we are talking about a nutritional, and potentially reversible type of immune disturbance. Where population growth, poverty and social disorder coexist, maize with its high yield and ability to grow with little attention or rainfall, is a very valuable crop. However, a diet almost exclusively of maize is seriously flawed, and is a root cause of many of what have been termed 'the diseases of poverty'.

Other diseases, whose principal protection arises from cellular immunity (eg, hydatid disease, malaria, leishmaniasis, syphilis and leprosy), may also be favoured in this situation.

Immune profile shift due to a temporary dietary cause implies the possibility of readjustment of the immune profile by diet. An increase in dietary *n*-3 PUFAs is more efficient than a decrease in *n*-6 acids in reducing PGE₂ synthesis.^{13 15} This suggests an approach to the problem which may be both effective and achievable.

The author wishes to acknowledge Dr S Glover and Dr DJ Unsworth of Southmead Hospital, Bristol, for their help in the preparation of this paper.

1 Sudre P, ten Dam G, Kochi A. Tuberculosis: a global overview of the situation today. *Bull World Health Organ* 1992;70:149-59.
 2 Fourie PB, Gatter EMS, Glatthaar E, Kleeberg HH. Follow-up tuberculosis prevalence survey of Transkei. *Tubercle* 1980;61:71-9.
 3 Gilks CF. The clinical challenge of the HIV epidemic in the developing world. *Lancet* 1993; 342:1037-9.
 4 Weissman D, Barker TD, Fauci, AS. The efficiency of acute infection of CD4+ cells is markedly enhanced in the setting of antigen-specific immune activation. *J Exp Med* 1996; 183:687-92.

5 Mhalu FS, Lyamuya E. Human immunodeficiency virus infection and AIDS in east Africa: challenges and possibilities for prevention and control. *East Afr Med J* 1996;73:13-9.
 6 Smallman-Raynor MR, Cliff AD. Seasonality in tropical AIDS: a geographical analysis. *Int J Epidemiol* 1992;21:547-67.
 7 Georges AJ, Georges-Courbot MC. (Epidemiology of HIV infection in Africa). *Rev Prat* 1990;40:2131-5.
 8 Shafer RW, Edlin BR. Tuberculosis in patients infected with human immunodeficiency virus: perspective on the past decade. *Clin Infect Dis* 1996;22:683-704.

9 Grant HW, Palmer KR, Kelly RW, Wilson NH, Misiewicz JJ. Dietary linoleic acid, gastric acid, and prostaglandin secretion. *Gastroenterology* 1988;94:955-9.
 10 Sammon AM. Letter to the editor. *Dis Esophagus* 1994;7:220.
 11 Blair IA, Prakash C, Phillips MA, Dougherty RM, Iacono JM. Dietary modification of omega 6 fatty acid intake and its effect on urinary eicosanoid excretion. *Am J Clin Nutr* 1993;57: 154-60.
 12 Alam SQ, Bergens BM, Alam BS. Arachidonic acid, prostaglandin E2 and leukotriene C4 levels in gingiva and submandibular glands of rats fed

- diets containing n-3 fatty acids. *Lipids* 1991;26:895-900.
- 13 Narisawa T, Takahashi M, Kotanagi H, et al. Inhibitory effect of dietary perilla oil rich in the n-3 polyunsaturated fatty acid alpha-linolenic acid on colon carcinogenesis in rats. *Jpn J Cancer Res* 1991;82:1089-96.
 - 14 Marangoni F, Mosconi C, Galella G, Galli C. Increments of dietary linoleate raise liver arachidonate, but markedly reduce heart n-6 and n-3 fatty acids in the rat. *Lipids* 1992;27:624-8.
 - 15 Raedersdorf D, Moser U. Influence of an increased intake of linoleic acid on the incorporation of dietary (n-3) fatty acids in phospholipids and on prostanoid synthesis in rat tissues. *Biochim Biophys Acta* 1992;1165:194-200.
 - 16 Taniguchi M, Hara T. Effects of riboflavin and selenium deficiencies on glutathione and its relating enzyme activities with respect to lipid peroxide content of rat livers. *J Nutr Sci Vitaminol (Tokyo)* 1983;29:283-92.
 - 17 Capdevila JH, Morrow JD, Belosludtsev YY, Beauchamp DR, DuBois RN, Falck JR. The catalytic outcomes of the constitutive and the mitogen inducible isoforms of prostaglandin H2 synthase are markedly affected by glutathione and glutathione peroxidase(s). *Biochemistry* 1995;34:3325-37.
 - 18 Pelliccione NJ, Karmali R, Rivlin RS, Pinto J. Effects of riboflavin deficiency upon prostaglandin biosynthesis in rat kidney. *Prostaglandins Leukot Med* 1985;17:349-58.
 - 19 Gold KN. Modulation of helper T cell function by prostaglandins. *Arthritis Rheum* 1994;37:925-33.
 - 20 Katamura K, Shintaku N, Yamauchi Y, et al. Prostaglandin E2 at priming of naive CD4+ T cells inhibits acquisition of ability to produce IFN-gamma and IL-2, but not IL-4 and IL-5. *J Immunol* 1995;155:4604-12.
 - 21 Ertel W, Morrison MH, Meldrum DR, Ayala A, Chaudry IH. Ibuprofen restores cellular immunity and decreases susceptibility to sepsis following hemorrhage. *J Surg Res* 1992;53:55-61.
 - 22 Bennett M, Uauy R, Grundy SM. Dietary fatty acid effects on t-cell-mediated immunity in mice infected with mycoplasma pulmonis or given carcinogens by injection. *Am J Pathol* 1987;126:103-13.
 - 23 Ayala A, Chaudry IH. Dietary n-3 polyunsaturated fatty acid modulation of immune cell function before or after trauma. *Nutrition* 1995;11:1-11.
 - 24 Zhang M, Gong J, Iyer DV, Jones BE, Modlin RL, Barnes PF. T cell cytokine responses in persons with tuberculosis and human immunodeficiency virus infection. *J Clin Invest* 1994;94:2435-42.
 - 25 Sudre P, Hirschel B. Risk factors for tuberculosis among HIV-infected patients in Switzerland. Swiss HIV cohort study. *Eur Respir J* 1996;9:279-83.
 - 26 Crowe SM, Carlin JB, Stewart KI, Lucas CR, Hoy JF. Predictive value of CD4 lymphocyte numbers for the development of opportunistic infections and malignancies in HIV-infected persons. *J Acquir Immune Defic Syndr* 1991;4:770-6.
 - 27 Khoo SH, Wilkins EGL, Fraser IS, Hamour AA, Stanford JL. Lack of skin test reactivity to common mycobacterial antigens in human immunodeficiency virus infected individuals with high CD4 counts. *Thorax* 1996;51:932-5.
 - 28 McDyer JF, Hackley MN, Walsh TE, Cook JL, Seder RA. Patients with multidrug-resistant tuberculosis with low CD4+ T cell counts have impaired Th1 responses. *J Immunol* 1997;158:492-500.
 - 29 Ziegler HD, Ziegler PB. Depression of tuberculin reaction in mild and moderate protein-calorie malnourished children following BCG vaccination. *John Hopkins Med J* 1975;137:59-64.
 - 30 Barcellini W, Rizzardì GP, Velati C, et al. In vitro production of type 1 and type 2 cytokines by peripheral blood mononuclear cells from high-risk HIV-negative intravenous drug users. *AIDS* 1995;9:691-4.
 - 31 Williams C. A nutritional disease of childhood associated with a maize diet. *Lancet* 1935;Nov 16:1151.
 - 32 Chowdhury FA, Miah RA, Begum M, Rahman KM. The transformation in vitro of peripheral lymphocytes of malnourished children. *Bangladesh Med Res Counc Bull* 1993;19:67-70.
 - 33 Purtilo DT, Connor DH. Fatal infections in protein-calorie malnourished children with thymolymphatic atrophy. *Arch Dis Child* 1975;50:149-52.
 - 34 Fakhir S, Ahmad P, Faridi MA, Rattan A. Cell-mediated immune responses in malnourished host. *J Trop Pediatr* 1989;35:175-8.
 - 35 Chandra RK. Numerical and functional deficiency in T helper cells in protein energy malnutrition. *Clin Exp Immunol* 1983;51:126-32.
 - 36 Watts T. Thymus weights in malnourished children. *J Trop Pediatr* 1969;15:155-8.
 - 37 Tolboom JJ, Ralitapole-Maruping AP, Kabir H, Molatseli P, Anderson J. Severe protein-energy malnutrition in Lesotho, death and survival in hospital, clinical findings. *Trop Geogr Med* 1986;38:351-8.
 - 38 Finke D, Brinckmann UG, ter Muelen V, Liebert UG. Gamma interferon is a major mediator of antiviral defense in experimental measles virus-induced encephalitis. *J Virol* 1995;69:5469-74.
 - 39 Keler T, Barker CS, Sorof S. Specific growth stimulation by linoleic acid in hepatoma cell lines transfected with the target protein of a liver carcinogen. *Proc Natl Acad Sci USA* 1992;89:4830-4.
 - 40 Holmberg SB, Naredi PL, Lindner GO, et al. Influence of zymosan (a non-specific macrophage stimulator) and of indomethacin on liver tumours - an experimental study in rats. *J Cancer Res Clin Oncol* 1995;121:463-8.
 - 41 Ramesh G, Das UN. Effect of dietary fat on diethylnitrosamine induced hepatocarcinogenesis in Wistar rats. *Cancer Lett* 1995;1-2:237-45.
 - 42 Fox JC, Hay RV. Eicosapentaenoic acid inhibits cell growth and triacylglycerol secretion in McA-RH7777 rat hepatoma cultures. *Biochem J* 1992;286:305-12.
 - 43 Okuno K, Jinnai H, Lee YS, et al. A high level of prostaglandin E2 (PGE2) in the portal vein suppresses liver-associated immunity and promotes liver metastases. *Surg Today* 1995;25:954-8.
 - 44 Moore SE, Cole TJ, Poskitt EME, et al. Season of birth predicts mortality in rural Gambia. *Nature* 1997;388:434.