

ORIGINAL ARTICLE

Declining prevalence of STI in the London sex industry, 1985 to 2002

H Ward, S Day, A Green, K Cooper, J Weber

Sex Transm Infect 2004;**80**:374–378. doi: 10.1136/sti.2003.009068

See end of article for authors' affiliations

Correspondence to: Helen Ward, Division of Primary Care and Population Health Sciences, Imperial College London W2 1PG, UK; h.ward@imperial.ac.uk

Accepted for publication 3 March 2004

Objectives: To describe major changes in the London sex industry between 1985 and 2002 and assess the implications for sexually transmitted infection (STI) risk.

Method: A descriptive study comparing women who first attended a sex work clinic between 1996 and 2002 and those first attending from 1985 to 1992; a nested case-control study. 1050 female sex workers took part. The setting was a specialist clinical service for sex workers based in a London genitourinary medicine (GUM) clinic, and fieldwork in west London. The main outcome measures were reported condom use and prevalent STI.

Results: Over the period of the study there was a significant increase in the proportion of sex workers not born in the United Kingdom (from 25% to 63%, $p < 0.001$), and women entered sex work at an older age (median 24 years compared with 20 years, $p < 0.001$). Condom use increased (with the exception of oral sex). There was a significant decline in the proportion of participants reporting a previous STI (32% compared with 80%, $p < 0.001$) and the prevalence of acute STI declined from 25% to 8% ($p < 0.001$). Acute STI was associated with younger age, younger age at first sex work, being new to sex work, and inconsistent condom use. In a multivariate analysis unprotected sex with clients was the only significant risk.

Conclusion: Major restructuring of the sex industry, including the shift to a primarily migrant workforce, has been associated with a steep decline in acute STI, undermining popular assumptions that migrant sex workers are central to the ongoing STI epidemic. We attribute the decline in acute STI to an increase in safer sex.

Female sex workers in London have had a relatively low risk of sexually transmitted infections (STI), including HIV, and a high level of condom use.^{1–3} Peer norms and networks, together with specialist health services, have contributed to health and safety in the industry in the United Kingdom and many parts of Europe.^{4–8}

However, by the second half of the 1990s, specialist health services in Europe were reporting increasing numbers of non-European prostitutes in the local workforce (see www.europap.net). This demographic change, reflecting global population movements, posed a challenge to health promotion campaigns: how to sustain the high levels of safer sex characteristic of professional prostitution? It has been unclear whether this migration, often from regions of the world with higher risks of STI and HIV, would increase risks among sex workers in Europe. A few studies have found higher risks of HIV and STI in migrant sex workers, but others have not.^{9–12}

In the United Kingdom there has also been an increase in the rate of bacterial STI in the UK population generally, including outbreaks of syphilis, and a crisis in access to sexual health services in the United Kingdom.^{13, 14} We hypothesised that changes within the sex industry would weaken established norms of occupational health and safety. Given the concurrent STI epidemic in the United Kingdom, we were concerned that sex workers would be at greater risk of infection. We therefore established a study to describe changes in the sex industry workforce and to analyse the impact of these changes on sexual health.

METHODS

From 1985 to 2002, female sex workers attending the Department of Genitourinary Medicine at St Mary's Hospital were invited to attend a specialist service, the Praed Street Project Clinic, for screening and health care. In

addition to referrals from the walk-in sexual health clinic, we received referrals through linked outreach services to local streets and sex work venues, and a drop-in centre run by the Praed Street Project.² From 1985 to 1992 and again from 1996 to 2002, baseline data were collected on 1050 women at their first clinic visit using a standardised proforma. The data were linked to results of diagnostic and screening tests. Women were screened for one or more of gonorrhoea, chlamydia, trichomonas, syphilis, HIV, and hepatitis B. Screening was based on routine diagnostic methods (see Ward *et al*).³ Gonorrhoea was detected using Gram stain with culture confirmation. Direct immunofluorescence was used to detect chlamydia from cervical specimens until 1993; subsequently an enzyme immunoassay (EIA) test was used until 1998 when a nucleic acid amplification test was introduced; trichomoniasis was diagnosed through direct microscopy of a wet mount from vaginal secretions. A diagnosis of new genital herpes was based on positive culture from a clinical lesion in someone with no previous history. Serum samples were tested for antibodies to HIV-1 and HIV-2 using routine ELISA screening tests with confirmation; hepatitis B core antibody was used as evidence of past infection; syphilis was diagnosed using TPHA, VDRL, and fluorescent treponemal antibody (FTA) tests. Newly diagnosed syphilis was distinguished by clinical history and serological profile.

Baseline data on the first 402 women included in this article have previously been published in a separate analysis.³ Baseline data were not collected in a standard way from 1993 to 1995 and therefore these years have not been included in most analyses, with the exception of country of birth which

Abbreviations: EIA, enzyme immunoassay; FTA, fluorescent treponemal antibody; GUM, genitourinary medicine; STI, sexually transmitted infections

was asked of all patients registering with the service throughout, and is therefore included for completeness in the description of trends (see fig 1).

Members of the research team carried out fieldwork during two phases, 1988–91 and 1998–2002, visiting areas of street prostitution, magistrates' courts, sex work businesses (flats and agencies), prisons, and drugs projects.

From 1989 to 1991 we made 184 fieldwork visits to three street areas and three magistrates' courts, meeting 274 different women. In 1999–2001, field work included twice weekly attendance at the Praed Street Project drop-in clinic for 12 months, six visits to another local drop-in, visits to 12 saunas, visits to 48 flats in two parts of London, one agency, two street areas, and one magistrates' court. In both periods we collected phone numbers from adverts in local phone boxes and newspapers. Fieldwork involved one or more members of the team visiting the areas and talking to people associated with the sex industry. Most of this fieldwork was conducted alongside outreach services, including health promotion relating to safer sex and drug use. Notes were kept from observations and informal interviews. Where possible data were collated on the numbers of sex workers/sex businesses and their characteristics. We also enumerated the working population locally. In this paper, we use these fieldwork data to assess potential biases associated with the clinic sample.

Data were analysed by reading through notes made at the time, transcripts of interviews, and collating information on numbers and some demographic data about contacts made.

In the clinic and during fieldwork, we communicated primarily in English, but could also converse in Spanish, Portuguese, and French. Translators were sometimes available for other languages but, more commonly, non-English speakers communicated through friends and colleagues.

Early year of recruitment to the study was the strongest determinant of STI and so, to look at other factors associated with STI, we carried out a nested case-control study matching for year of recruitment. Cases were defined by having one or more STI (gonorrhoea, chlamydia, trichomoniasis, infectious or early latent syphilis, first episode genital herpes, or first episode genital warts) at recruitment. Two controls were randomly selected for each case from those sex workers who had registered in the same year and screened negative. The sample size of the case-control study was based on the total number of cases eligible.

Data management and analysis

Data from all studies were stored without individual identifying information. Numeric data were entered into a spreadsheet, checked, and exported to SPSS for analysis. Fieldwork notes were entered into a word processing package and then exported into a qualitative analysis program for analysis. We created four broad categories for country of birth in order to analyse demographic changes: United Kingdom, developed, developing, and transitional economies (former Soviet states and east/central Europe). Condom use was analysed using two categories: always or not always. We compared those recruited in the earlier period, from 1985 to 1992, with those recruited later from 1996 to 2002. Simple

Table 1 Comparison of sex workers registering with clinic 1985–92 with those in 1996–2002

	1985–92 (n = 498)	1996–2002 (n = 552)	p Value
Age			
Range	16–59	16–46	
Median (IQR)	25 (21, 29)	26 (23, 31)	<0.001
Missing	2	3	
Country of birth			
UK	346 (74.9%)	203 (36.9%)	<0.001
Developed	90 (19.5%)	89 (16.2%)	NS
developing	22 (4.8%)	149 (27.1%)	<0.001
Transitional	4 (0.9%)	109 (19.8%)	<0.001
Missing	36	2	
Months of sex work			
Range	1–324	1–216	
Median (IQR)	24 (10, 60)	12 (4, 36)	0.004
Missing	127	128	
Age at start of sex work			
16 or under	42 (11.3%)	21 (5.0%)	0.01
17–19	108 (29.1%)	51 (12.1%)	<0.001
20–29	189 (50.9%)	268 (63.8%)	<0.001
30 or older	32 (8.6%)	80 (19.0%)	<0.001
Range	11–50	12–52	
Median (IQR)	20 (18, 24)	24 (21, 28)	
Missing	127	132	
Sex work sector			
Club/sauna	52 (12.5%)	71 (14.3%)	NS
Escort/madam	161 (38.8%)	190 (38.4%)	NS
Private	46 (11.1%)	16 (3.2)	<0.001
Flat	81 (19.5%)	170 (34.3%)	<0.001
Street	74 (17.8%)	16 (3.2%)	<0.001
Other	1 (0.2%)	32 (6.5%)	<0.001
Missing	83	57	
Past STI*			
Gonorrhoea	177/352 (50.3%)	36/494 (7.3%)	<0.001
Chlamydia	111/352 (31.5%)	71/494 (14.4%)	<0.001
Genital herpes	49/352 (13.9%)	30/494 (6.1%)	<0.001
Syphilis	9/352 (2.6%)	13 (2.6%)	NS
Any†	281/352 (79.8%)	159/494 (32.2%)	<0.001

*Past STI based on self report.

†Gonorrhoea, chlamydia, trichomoniasis, genital herpes, genital warts, syphilis.




Figure 1 Type of country of birth of sex workers by year of first attending the Praed Street Project Clinic, 1985–2002*. (*In this figure data have been included for all years from 1985 to 2002 using routine clinic registration data for the years 1993 to 1995. See methods for description of the country categories.)

univariate analysis was carried out for most comparisons using χ^2 tests for categorical variables, and Mann-Whitney U test to compare continuous variables. In the case-control study we carried out univariate logistic regression to produce odds ratios and 95% confidence intervals. Multivariate analysis was carried out using forward conditional logistic regression, testing in the model the effect of those variables significantly associated on univariate analysis, or those variables of borderline significance that are known determinants of risk. Fieldwork notes and transcripts were analysed by content; workplaces and the local working population were mapped.

Consent and ethics

Approval was obtained from the local research ethics committee. In the clinic, baseline data were requested as part of the registration visit, and specific consent was obtained for interviews. Screening was conducted with appropriate counselling and consent; infections were managed according to standard clinic protocol. During fieldwork, consent was obtained for interviews and for access to workplaces and projects.

RESULTS

Clinic sample

In all, 1050 registered with the clinic during the two study periods, 1985–1992 and 1996–2002 (see table 1). Later recruits were older, less likely to have been born in the United Kingdom, and attended earlier in their sex work careers; 40% attended within the first 6 months. The age at

which women had entered prostitution increased significantly; fewer women reported first work before the age of 20 (17% compared with 40% in the earlier group), and more started in their 30s or older (19% compared with 9% in the earlier group). More women worked in flats or “other” ways, referring primarily to direct advertising on the internet and the adult entertainment industry. At first visit the proportion of women reporting past STI declined substantially, from 80% to 32%.

Figure 1 shows the change in geographical origin of sex workers in more detail, including figures for the intervening years, which were available from routine data. The proportion of workers of UK origin fell in the mid-1990s; the biggest increase was in sex workers from the transitional economies of eastern Europe and the former Soviet Union, and from developing countries in Asia, South America, and Africa.

Condom use for vaginal sex with clients increased to 98% by 1990, and has stayed at that level. In contrast, condom use for oral sex with clients has declined; consistent use was reported by 41% of women in 1985–6, rising to 91% by 1990; it has subsequently declined with only 66% of women reporting universal use in 2002 ($\chi^2 = 21.94$, $p \leq 0.001$). The proportion of women using condoms for anal sex with clients rose to 94%, and more women reported selling this type of sex, increasing from five in the earlier period (1% of total) to 34 in the later period (6% of total). Systematic condom use with non-commercial partners increased from 8% of women in 1985–6 up to 44% in 2003.

Table 2 shows the results of screening carried out at first visit. The prevalence of acute STI, including gonorrhoea and chlamydia, decreased significantly from the earlier to the later group. Newly diagnosed syphilis increased but not significantly along with HIV, past hepatitis B, and previous syphilis. HIV and hepatitis B were most strongly associated with injecting drug use in the earlier group³ and with origin from high prevalence areas in the later group.

Analysis of vulnerability to infection

In the nested case-control study, we included 128 cases of acute STI and 256 controls. Cases included 59 *Chlamydia trachomatis*, 34 newly diagnosed genital warts, 23 trichomoniasis, 22 gonorrhoea, nine first clinical episode genital herpes, and two early latent syphilis. Most women (112) had a single infection, but 12 had two infections and four had three. Table 3 shows factors associated with an acute STI. Univariate analysis suggested that younger age now, younger age at first sex work, and being new to sex work were all associated with increased risk. Having a boyfriend increased the risk, and using condoms with him, and using condoms with all clients for vaginal sex reduced the risk. Drug use, working outside London or the United Kingdom, being non-UK born, and workplace were not associated with increased risk. The results of a multivariate analysis are also shown in

Table 2 Baseline prevalence of infections comparing sex workers registering with clinic 1985–92 with those in 1996–2002*

	1985–92		1996–2002		Change
	Prevalence	%	Prevalence	%	
Any acute STI†	99/398	24.9	29/367	7.9	–68% (p<0.001)
Gonorrhoea	18/396	4.5	4/364	1.1	–75% (p<0.01)
Chlamydia	42/393	10.7	17/354	4.8	–55% (p<0.01)
HIV	4/357	1.1	3/200	1.5	+36% (NS)
Hepatitis B‡	27/417	6.5	28/307	9.1	+40% (NS)
Syphilis previous treatment	9/367	2.5	13/279	4.7	+88% (NS)
Syphilis newly diagnosed	3/367	0.8	6/279	2.2	+175% (NS)

*The denominators vary in this table since not all sex workers consented to screening for all STIs.

†One or more of gonorrhoea, chlamydia, trichomoniasis, first episode genital herpes, first episode genital warts, infectious syphilis.

‡Hepatitis B anti-core positive, indicating previous infection.