

Geographical distribution of tuberculosis notifications in national surveys of England and Wales in 1988 and 1993: report of the Public Health Laboratory Service/British Thoracic Society/Department of Health Collaborative Group

L P Ormerod, A Charlett, C Gilham, J H Darbyshire, J M Watson

Abstract

Background—The geographical distribution of tuberculosis in England and Wales and changes since 1983 were examined using data from the 1988 and 1993 national surveys of tuberculosis notifications.

Methods—Notification rates for England and Wales in 1988 and 1993 were calculated for geographical areas using Office for National Statistics (ONS) mid year population estimates. Those for the standard regions and the Greater London boroughs were calculated for the main ethnic groups. Those for the counties and local authorities were calculated for all ethnic groups combined. These were compared using data from the 1983 national survey as a baseline.

Results—Wide regional variations in notification rates persist with Greater London having the highest rates. Rates in the ethnic group from the Indian subcontinent (ISC) were high in all regions, whilst those of the white ethnic group varied fourfold. Twenty seven of the 33 London boroughs showed increased rates in 1993 compared with 1988. In general, those local authority areas with high rates had high proportions of notifications in individuals of ISC ethnic origin, emphasising the continuing important contribution of ethnic minority groups to local tuberculosis rates. The number of local authority areas with notification rates four times the national average increased, but the number of areas with low or zero rates increased even more.

Conclusions—The distribution of tuberculosis in England and Wales continues to vary markedly by geographical area. The distribution is becoming increasingly polarised with a growth in the number of areas with very high rates of notifications and a greater increase in the number of areas with very few notifications. Patients from ethnic minorities continued to contribute a substantial and increasing proportion of all reported tuberculosis cases in most regions in 1988 and 1993. These findings have important implications for the provision of tuberculosis services in England and Wales.

(Thorax 1998;53:176-181)

Keywords: tuberculosis; geographical distribution; England and Wales

The decline in tuberculosis notification rates in Britain¹ and many other industrialised countries ceased in the mid to late 1980s, and in some countries small increases then followed. Tuberculosis occurring in people born in or originating from parts of the world with high rates of tuberculosis has been identified as the major contributor to these changes. Within individual countries there are variations in tuberculosis rates in different geographical areas, such as those previously described for England and Wales.^{2,3} Considerable variation also exists in the geographical distribution of the population subgroups at high risk of tuberculosis. Consistent with the recent overall notification rates,¹ the number of cases in the 1988⁴ and 1993⁵ surveys increased. The proportion of tuberculosis notifications from ethnic minority groups is continuing to increase in England and Wales, being 45% in 1983,⁶ 47% in 1988,⁴ and 56% in 1993.⁵

In view of the recent changes in tuberculosis notification rates overall, and since the last reported geographical analysis was based on data from the 1983 National Tuberculosis Notification Survey,³ it is timely to examine the current geographical distribution and changes over time using the data from the more recent national surveys during 1988⁴ and 1993.⁵

The standardised rates of tuberculosis for both the 1988 and 1993 surveys have been calculated and, using 1983 as the baseline,⁶ the changes in the rates since 1983 in the geographical subdivisions of England and Wales have been examined.

Methods

Details of the survey methods have already been reported.^{4,5} During the six month survey periods the survey team was sent a copy of all notification forms received by the Medical Officers of Environmental Health (MOEH) in 1988, and from Consultants in Communicable Disease Control (CCDC) in 1993 for the 403 local authorities in England and Wales. These were checked against lists received from the Office of Population Censuses and Surveys (OPCS) which gave details of age, sex, and type

Chest Clinic,
Blackburn Royal
Infirmery, Blackburn,
Lancashire BB2 3LR,
UK (for the Joint
Tuberculosis
Committee of the
British Thoracic
Society)
L P Ormerod

Public Health
Laboratory Service
Statistics Unit, London
NW9 5EQ, UK
A Charlett
C Gilham

Medical Research
Council, HIV Clinical
Trials Centre,
University College
London Medical
School, London WC1E
6AU, UK
J H Darbyshire

Public Health
Laboratory Service
Communicable
Disease Surveillance
Centre, London NW9
5EQ, UK
J M Watson

Correspondence to:
Dr L P Ormerod.

Received 21 August 1997
Returned to authors
5 November 1997
Revised version received
3 December 1997
Accepted for publication
3 December 1997

Table 1 Notifications in the six month surveys in 1988 and 1993 (and estimated annual rate/100 000) in the standard regions of England and Wales (with Greater London shown separately) by the main ethnic groups

Region	All ethnic groups			White			ISC		
	Population* (1000s)	Tuberculosis notifications	Estimated annual rate	Population* (1000s)	Tuberculosis notifications	Estimated annual rate	Population* (1000s)	Tuberculosis notifications	Estimated annual rate
A. 1988									
North	3037	98	6.2	3002	83	5.3	21	12	109.0
Yorkshire/Humberside	4855	252	10.0	4624	149	6.2	158	92	113.0
East Midlands	3925	215	10.6	3769	104	5.3	104	103	192.0
East Anglia	2004	37	3.6	1977	31	3.0	10	3	59.2
Greater London	6616	680	19.9	5434	246	8.8	515	338	126.9
Rest of South East	10418	262	4.9	10105	171	3.3	144	76	102.2
South West	4540	92	3.9	4474	81	3.5	12	8	130.1
West Midlands	5151	335	12.6	4790	142	5.7	238	160	130.1
North West	6283	342	10.5	6041	201	6.4	148	124	162.2
England	46829	2313	9.6	44216	1208	5.3	1349	916	131.3
Wales	2823	95	6.5	2779	86	6.0	16	6	73.5
England and Wales	49652	2408	9.4	46995	1294	5.3	1365	922	130.7
B. 1993									
North	3047	111	6.9	3014	92	5.8	16	15	168.2
Yorkshire/Humberside	4928	247	9.5	4681	106	4.3	177	132	140.4
East Midlands	4034	210	9.8	3848	88	4.3	124	112	170.1
East Anglia	2103	35	3.1	2049	27	2.5	22	8	66.8
Greater London	6728	924	25.9	5273	290	10.4	557	370	125.2
Rest of South East	10697	244	4.3	10332	136	2.5	207	82	74.6
South West	4702	90	3.6	4630	73	3.0	17	9	95.7
West Midlands	5222	381	13.8	4764	144	5.7	313	204	122.7
North West	6311	383	11.4	6035	176	5.5	183	175	179.5
England	47772	2625	10.4	44631	1132	4.8	1621	1107	128.8
Wales	2874	81	5.3	2830	70	4.7	18	7	72.8
England and Wales	50646	2706	10.1	47462	1202	4.8	1639	1114	128.2

*Based on 1988 and 1993 Labour Force Surveys.

of disease of the notified cases to confirm that information on all notifications during the survey had been received. In addition, a special form was completed for each patient notified by the clinician in all of the surveys, which included results of age, sex, ethnic origin, country of birth, and bacteriological and histological results. The classification of ethnic origin was similar to that used in previous surveys.^{2,7} Patients were classified by the notifying physician as being of White, Indian,

Pakistani, Bangladeshi, Black Caribbean (West Indian), Black African, Arab, Chinese, other (including mixed), or unknown ethnic origin. In the analysis of this report the Indian, Pakistani and Bangladeshi ethnic groups have been combined as the Indian Subcontinent (ISC) ethnic group, and all other non-white ethnic groups have been combined as "other" ethnic groups. Due to seasonal variations in the number of reported notifications, the annual number of cases have been estimated by multiplying the numbers from the two six month surveys by an appropriate scaling factor (1.953 in 1983⁶; 1.9348 in 1988⁴; 1.8864 in 1993⁵). As with previous geographical analyses,^{2,3} all notified cases include those with a previous history of tuberculosis.




Figure 1 Relationship between the estimated number of patients of non-white non-ISC ethnic origin with infectious (smear and culture positive) respiratory tuberculosis per 100 000 of the white population and the estimated notification rate (for all forms of tuberculosis) in the white population in the 33 London boroughs in (A) 1988 ($R_s = 0.35$) and (B) 1993 ($R_s = 0.69$).

DENOMINATORS

For local government purposes in 1983–93 England was divided into eight standard regions which were further subdivided into 46 counties; seven of the latter were largely urban and were designated metropolitan counties. Wales was divided into eight counties. The 54 authority areas of which 33 were in Greater London. The population estimates for 1983, 1988, and 1993 were taken from the appropriate mid year estimates of the Office for National Statistics (ONS, formerly OPCS). In this report individuals for whom the place of birth or date of entry into the UK was not obtained have been distributed proportionally amongst those for whom it was known. Population data were available to calculate notification rates for the main ethnic groups for the regions of England and Wales, and for the 33 boroughs of Greater London. However, population data by ethnic group were not available for the counties and local authorities,

Table 2 Greater London Boroughs: estimated annual notification rates for 1983, 1988 and 1993 for all ethnic groups combined and details of 1993 population and notifications

Borough	Estimated annual rate/100 000			1993		
	1983	1988	1993	Population* (1000s)	% notified	
					ISC	% Non-ISC Non-white
Barking & Dagenham	17.2	9.2	11.7	146	44	0
Barnet	18.2	27.3	29.7	305	58	21
Bexley	12.7	3.5	6.0	220	43	0
Brent	92.3	43.5	47.2	248	60	24
Bromley	7.2	2.6	6.5	292	20	10
Camden	33.4	24.8	38.5	181	19	30
City of London	0.0	0.0	0.0	4	0	0
Croydon	19.3	13.5	18.7	323	50	25
Ealing	45.4	30.1	46.2	286	64	21
Enfield	16.6	9.0	20.2	262	46	14
Greenwich	15.8	11.8	22.8	215	27	42
Hackney	27.3	18.7	39.5	191	20	30
Hammersmith & Fulham	32.7	24.0	13.3	155	18	36
Haringey	30.2	24.8	32.9	212	14	51
Harrow	37.3	33.3	42.0	207	65	26
Havering	5.7	4.9	4.9	232	0	33
Hillingdon	17.8	9.9	18.9	240	46	42
Hounslow	36.2	21.6	27.4	207	70	10
Islington	33.2	18.6	22.6	175	14	38
Kensington & Chelsea	32.7	22.9	6.3	149	20	20
Kingston upon Thames	8.9	7.3	13.7	138	40	30
Lambeth	37.5	18.5	43.7	259	23	45
Lewisham	20.0	17.9	20.4	240	12	39
Merton	17.5	8.0	22.6	175	33	33
Newham	72.2	52.9	49.2	226	59	25
Redbridge	24.9	12.8	19.5	233	29	25
Richmond upon Thames	7.2	7.4	4.5	167	0	0
Southwark	20.8	28.3	22.2	229	0	48
Sutton	8.2	7.9	0.0	173	0	0
Tower Hamlets	56.4	38.7	60.1	170	56	20
Waltham Forest	24.7	18.0	29.2	220	47	27
Wandsworth	29.7	18.4	9.2	266	23	31
Westminster	41.6	32.2	42.9	189	19	30

*Based on 1993 Labour Force Survey.

and therefore rates for these areas could only be calculated for all ethnic groups combined.

Results

SURVEY POPULATIONS

Of the 2748 notifications during the 1988 survey period, 340 were excluded.⁴ There remained 2408 newly notified patients, of whom

245 had had previous treatment for tuberculosis (the corresponding figures for 1983 were 3302 and 300, respectively); 54% were of white origin, 38% of ISC origin, and 8% of "other" ethnic groups. During the 1993 survey period there were 3298 notifications of which 592 were excluded.⁵ The main reasons for exclusions were inappropriate notifications (chemoprophylaxis) and denotification (non-tuberculous mycobacteria and other diseases). There remained 2706 newly notified patients, of whom 248 had had previous treatment for tuberculosis 44% were of white origin, 41% were of ISC origin, and 14% of "other" ethnic groups.

The data are presented for all types of disease combined, although there were differences in the disease characteristics between the ethnic groups.

STANDARD REGIONS OF ENGLAND

Table 1 shows the estimated annual notification rates per 100 000 for 1988 and 1993, respectively, for all ethnic groups combined and the main ethnic groups for the standard regions with Greater London shown separately from the rest of the South East. There were wide regional variations with low rates in Anglia, South West, and the South East (excluding Greater London), and the highest rates in Greater London in both years. The rates for the white ethnic group varied from 3.0 to 8.8 per 100 000 in 1988, with a wider range from 2.5 to 10.4 per 100 000 in 1993. Increases in the notification rate for the white population between 1988 and 1993 were seen

Table 3 Local authorities: estimated annual notification rates/100 000, excluding Greater London, and restricted to those authorities with 15 cases or more and with rates of 15.0/100 000 or more in either 1988 or 1993, with 1993 population data

Local authority	Estimated annual rate/100 000		1993	
	1988	1993	Population* (1000s)	% notified ISC
Leicester	51.8	52.8	289	83
Pendle	34.4	22.0	86	90
Preston	34.0	37.1	132	54
Bolton	32.0	40.6	265	81
Slough	30.3	29.2	103	88
Blackburn	29.9	40.6	140	73
Bradford	28.6	31.4	480	80
Wolverhampton	24.2	21.4	246	36
Birmingham	23.6	32.4	1012	64
Stoke-on-Trent	23.1	13.4	253	22
Manchester	22.5	16.6	432	55
Luton	21.8	25.3	179	63
Derby	20.2	20.6	229	80
Rochdale	18.1	16.4	207	72
Walsall	17.0	19.2	265	63
Sandwell	17.0	4.5	294	71
Newcastle on Tyne	16.5	12.6	285	5
Coventry	16.3	25.4	304	68
Nottingham	14.7	25.4	283	26
Middlesborough	10.6	28.5	146	41
Oldham	7.1	23.1	221	70
Reading	8.5	21.9	138	50
Liverpool	8.4	17.0	477	5
Tameside	8.1	16.2	222	32
Kirklees	13.3	15.6	386	75

*Based on the 1993 Labour Force Survey.

Details of local authorities with estimated rates under 15/100 000 in both 1988 and 1993 are available from the authors.

Table 4 Patients notified in the six month survey periods by area and main ethnic group and estimated annual rates for 1988 and 1993

Local authority group	Population* (1000s)		Tuberculosis notifications		Estimated annual rate	Tuberculosis notifications								
	n	%	n	%		White			Indian		Pakistani & Bangladeshi			
						n	%	%†	n	%	n	%		
A. 1988														
Metropolitan areas:														
33 boroughs in Greater London	6770	13	680	28	19.4	246	19	36	264	44	74	23		
36 authorities in other metropolitan areas	11121	22	736	31	12.8	365	28	50	161	27	150	47		
Non-metropolitan areas														
63 authorities with a population of 125 000 or more	10343	20	464	19	8.7	272	21	59	119	20	53	17		
174 authorities with a population of 75 000–125 000	16814	33	394	16	4.5	295	23	75	45	7	41	13		
97 authorities with a population of less than 75 000	5439	11	134	6	4.8	116	9	87	12	2	3	1		
B. 1993														
Metropolitan areas:														
33 boroughs in Greater London	6933	14	924	34	25.1	290	24	31	252	42	118	23		
36 authorities in other metropolitan areas	11199	22	833	31	14.0	341	28	41	155	26	272	53		
Non-metropolitan areas:														
63 authorities with a population of 125 000 or more	10816	21	479	18	8.4	231	19	48	136	23	73	14		
174 authorities with a population of 75 000–125 000	17303	34	362	13	3.9	251	21	69	47	8	47	9		
97 authorities with a population of less than 75 000	5188	10	108	4	3.9	89	7	82	11	2	3	1		

*Based on the 1988 and 1993 Labour Force Surveys.
 †As a percentage of all notifications in the area grouping.

only in the North (9%) and Greater London (18%). The rate in the West Midlands stayed unchanged, and decreases were seen in the other regions ranging from 14% in the North West and South West to 31% in Yorkshire. The rates for the ISC ethnic group were high in all regions (59–192 per 100 000 in 1988 and 67–180 per 100 000 in 1993).

GREATER LONDON LOCAL AUTHORITY AREAS (BOROUGHES)

The rates for the London boroughs are given in table 2. All but six of the 33 boroughs showed an increase between 1988 and 1993 with five boroughs (Lambeth, Hackney, Enfield, Bromley and Merton) all showing increases of more than 100%. The rate for Greater London as a whole increased by 30% from 19.9 to 25.9 per 100 000 with the number of notifications rising from 680 to 924 with only a 2% increase in population. Whereas in 1988 seven of the boroughs had rates more than three times that of England and Wales, this had increased to 10 boroughs by 1993.

OTHER ETHNIC GROUPS

There have been changes in the distribution of the ethnic populations in England and Wales since 1983,³ and particularly since 1988. In

1988 non-ISC non-white ethnic groups contributed 8% of notifications⁴ which increased to 14% by 1993,⁵ this increase being particularly seen in Black Africans in whom notifications increased from 1.7% to 7.1% between 1988 and 1993. The contribution of the non-ISC ethnic groups is seen most in the London boroughs (table 2). There has been little change in the contributions of the Black Caribbean and Chinese ethnic groups.

In 1988 the possibility of an association between the overall rates of tuberculosis in the white population in the 33 London boroughs and the number of infectious tuberculosis cases in the ISC population in the same boroughs was examined but no association was found (Spearman's correlation coefficient 0.06). The same analysis was repeated in 1993 and again no evidence of an association was apparent (Spearman's correlation coefficient 0.45).

In the light of the increasing proportion of cases occurring in the non-white non-ISC group in 1993, an analysis was carried out using the number of cases of infectious tuberculosis in this group and the rates in the white population of the London boroughs (fig 1). No association was apparent in 1988 (Spearman's correlation coefficient 0.35) but the possibility of an association was apparent in 1993 (Spearman's correlation coefficient 0.69).

LOCAL AUTHORITY AREAS

The rates for those local authority areas outside Greater London with 15 or more cases and rates of 15 per 100 000 or more are presented in table 3. The number of such areas increased from 18 in 1988 to 21 in 1993. Of the 18 areas in 1988, 10 showed increases in 1993 and eight showed decreases—three of which (Stoke-on-Trent, Sandwell and Newcastle) had fallen below 15 per 100 000 by 1993. Liverpool, Reading, Oldham, and Middlesbrough all had

Table 5 Comparison of estimated annual rates in the 403 local authority areas of England and Wales in the 1983, 1988, and 1993 surveys

	1983	1988	1993
Estimated annual rate/100 000 for England and Wales	13.0	9.4	10.1
Ratio to England and Wales rate (n)			
Rate 4 times or greater	5	4	10
Rate 3.0–3.99 times	2	12	7
Rate 2.0–2.99 times	20	16	23
Rate 1.0–1.99 times	66	56	43
Rate 0.50–0.99 times	120	99	72
Rate up to 0.49 times	148	139	125
No cases	42	77	123

Table 6 Rates of tuberculosis notifications (per 100 000 population) in the six month survey periods of 1988 and 1993 by birthplace (UK or abroad) and number of years since first entry and region of residence (only regions with more than 25 notifications of Indian or Pakistani/Bangladeshi ethnic origin have been presented)

Ethnic group and region	Population* (1000s)	Overall		Born UK		Five or more years since first entry into UK		Less than five years since first entry into UK	
		1988	1993	1988	1993	1988	1993	1988	1993
Indian									
East Midlands	110.6	213	160	95	50	243	167	501	883
Greater London	370.9	138	128	12	41	135	114	706	809
Rest of South East	109.1	111	81	38	26	92	98	633	387
West Midlands	171.4	146	121	54	73	195	122	851	1057
North West	55.1	211	216	77	87	258	263	1901	1334
Pakistani & Bangladeshi									
Yorkshire	123.6	128	166	49	76	143	182	496	852
Greater London	186.7	102	119	27	42	127	118	239	242
Rest of South East	98.3	90	67	28	12	135	87	167	309
West Midlands	142.2	109	125	31	44	148	135	290	952
North West	128.8	131	164	19	69	188	210	723	362

*Based on 1993 Labour Force Survey.

substantially higher rates in 1993 with increases of over 100% since 1988. Although there were increases in notification rates in some areas, the general trend for 1988 and 1993 was for a decrease in most local authority areas.

NOTIFICATIONS BY LOCAL AUTHORITY GROUPS

Table 4 shows the 403 local authorities grouped according to type, area and population size for 1988 and 1993. The 13% of the population living in Greater London provided 28% of the total notifications in 1988 and 34% in 1993. The 22% of the population in the metropolitan areas outside London provided 31% of the total notifications in both years. Thus in 1988 the 65% of the population in non-metropolitan areas provided the remaining 41% of the notifications, falling to 35% by 1993. In 1988 the percentage of notifications in people of white ethnic origin was 87% in the local authority areas with populations of less than 75 000 compared with 36% and 50% in Greater London and in the other metropolitan counties, respectively. By 1993 the proportion of notifications in people of white ethnic origin in the local authorities with populations of under 75 000 had declined to 82%, and in Greater London and other metropolitan counties to 31% and 41%, respectively. In both 1988 and 1993 the highest proportions of Indian patients (44% in 1988, 42% in 1993) were resident in Greater London, whereas the greatest proportion of Pakistani and Bangladeshi patients were resident in the other metropolitan counties (47% in 1988, 53% in 1993). In 1983, 1988 and 1993, 74%, 70% and 72% of notifications, respectively, in people of ISC ethnic origin occurred in residents of either Greater London or the other metropolitan counties. The comparative figures for people of white ethnic origin were 50%, 47%, and 52%.

The varying trends in geographical distribution can be seen more clearly if all 403 local authority areas are stratified by comparison with the national rate (table 5). The number of districts with below national average notification rates have increased over the three surveys, and those districts with no notifications have trebled since 1983. Those areas with rates four times the national average, however, have dou-

bled between 1983 and 1993, with a lesser rise in those with rates between two and four times the national average notification rate.

DURATION OF STAY IN THE UK BY REGION FOR THE ISC ETHNIC GROUP

In both 1988 and 1993 there were considerable differences in the notification rates for the population of the ISC ethnic group between those born in the UK and those born abroad, with much higher rates for those recently arrived in the UK than for those who had been resident in the country for more than five years.^{4,5} Table 6 shows the notification rates in those born and those not born in the UK and the time since first entry for those regions with more than 25 notifications in the Indian or Pakistani and Bangladeshi ethnic groups. The overall rates generally decreased between the surveys in the Indian group but generally increased in the Pakistani and Bangladeshi group.

Although the rates amongst those born in the UK are relatively low, they are increasing in some areas—for example, the rates have more than doubled among the Indian group between the surveys in Greater London and for the Pakistani and Bangladeshi group in the North West.

The highest rates are seen in recent immigrants into the UK among whom the rates have generally increased between 1988 and 1993, especially within the Pakistani and Bangladeshi group. As most of these rates are essentially based on low numbers, changes in any group within one region should be interpreted with caution.

Discussion

The marked variation in tuberculosis notification rates in different parts of England and Wales reported in previous surveys^{3,7} was also seen in both 1988 and 1993, and in some respects has become more pronounced. The 1988 survey⁴ took place shortly after the nadir of notifications in 1987⁶ and that of 1993⁵ after the increase in notifications seen between 1987 and 1990 had levelled off. Analysis by region shows that the rate of tuberculosis declined in all regions between 1983 and 1988, but that whilst the decline continued in most regions in

1993, there were increases in some regions including the West Midlands (10%), North West (9%), Northern (11%), and within the South East region, particularly in Greater London (30%).

More detailed analysis of the 403 local authority areas, however, shows that most of the areas are still experiencing a decline. Those districts which are predominantly small, rural, or are in non-metropolitan areas generally have falling rates with an increasing number of areas reporting either no notifications or rates below national average. However, the number of districts with rates at least three times the national average increased, particularly in 1993. Thus, the distribution of tuberculosis is becoming increasingly uneven, with a small number of areas with very high rates and below average rates for most of the areas. The number of areas with either no cases or below average rates has increased progressively and, by 1993, 320 (79%) were experiencing rates below the national average of which 123 (31%) reported no cases in the survey period.

This analysis also highlights the continuing important contribution of patients belonging to the ethnic minority groups to tuberculosis notification rates. Those of non-white ethnic origin, who constitute 6.3% of the total population (ranging from 1.1% (Northern) to 21.6% (Greater London) in 1993), contributed 56% of the tuberculosis notifications. All regions showed an increase in the percentage of notifications in persons of non-white ethnic origin in 1993 compared with 1988, ranging from 13.6% of total notifications in Wales to 68.6% in Greater London.

The tuberculosis rates in the ISC ethnic population remain very strongly associated with place of birth and duration of residence in the UK, as well as to the age distribution of the population.⁴ Those born in the UK have the lowest rates, and those most recently arrived (less than five years) have the highest rates. This association was first reported over 20 years ago⁸ but remains important despite the overall fall in notification rates in ISC ethnic groups. Thus, the age distribution of the ISC ethnic group in a local authority area, the proportions born in the UK, and the time since first arrival of those born abroad will influence the overall rate of tuberculosis seen in that area.

The rates in the white population varied up to fourfold between regions in 1993 with some showing rising rates and others falling rates. The rates in the white population of the London boroughs did not show a correlation with the proportion of ISC smear and culture positive rate per 1000 000 in either 1988 or 1993, consistent with the findings in 1983. In

1988 there was no correlation between white population rates and non-white non-ISC smear and culture positive rate per 100 000, but there was a possible association in 1993. This possible association is difficult to interpret in terms of establishing causality because potentially important confounding factors such as recent immigration and poverty have not been taken into account. Although this report was not designed to consider deprivation, evidence from Liverpool,⁹ Leeds,¹⁰ London,¹¹ and England and Wales¹² suggest that socioeconomic deprivation is a substantial factor in the rates of tuberculosis of all ethnic groups.

In view of the marked variation in tuberculosis rates seen across the country, the tuberculosis services in a district will need to reflect not only absolute numbers and rates, but the proportions from the different ethnic groups, the proportions of those born abroad, and the ways in which these various factors are changing.

The success of the 1988 and 1993 surveys was due to the co-operation of the more than 150 Medical Officers in Environmental Health (1988) and Consultants in Communicable Disease Control (1993), all physicians with an interest in thoracic medicine (over 400), and many other clinicians who also provided information; to tuberculosis nurses and health visitors, clinic staff, records officers and clerks in many chest clinics and hospitals; and to infectious disease clerks and other local authority staff. The 1988 survey was funded by the Medical Research Council and that of 1993 by the Department of Health and the Welsh Office.

- 1 Hayward AC, Watson JM. Tuberculosis in England and Wales 1982-93: notifications exceeded predictions. *Commun Dis Rep* 1995;5:R29-33.
- 2 Medical Research Council Tuberculosis and Chest Diseases Unit. The geographical distribution of tuberculosis notifications in a national survey of England and Wales (1978-9). *Tubercle* 1982;63:75-88.
- 3 Medical Research Council Tuberculosis and Chest Diseases Unit. The geographical distribution of tuberculosis notifications in a national survey of England and Wales in 1983. *Tubercle* 1986;67:163-78.
- 4 Medical Research Council Cardiothoracic Epidemiology Group. National Survey of notifications of tuberculosis in England and Wales in 1988. *Thorax* 1992;47:770-5.
- 5 Kumar D, Watson JM, Charlett A, et al. Tuberculosis in England and Wales in 1993: results of a national survey. *Thorax* 1997;52:1060-7.
- 6 Medical Research Council Tuberculosis and Chest Diseases Unit. National survey of notifications of tuberculosis in England and Wales in 1983. *BMJ* 1985;291:658-61.
- 7 Medical Research Council Tuberculosis and Chest Diseases Unit. National survey of tuberculosis notifications in England and Wales 1978-9. *BMJ* 1980;281:895-8.
- 8 British Thoracic and Tuberculosis Association. Tuberculosis among immigrants related to length of residence in England and Wales. A report from the research committee of the British Thoracic and Tuberculosis Association. *BMJ* 1975;ii:698-9.
- 9 Spence DPS, Hotchkiss J, Williams CSD, et al. Tuberculosis and poverty. *BMJ* 1993;307:759-61.
- 10 Goldman JM, Teale C, Cundall DB, et al. Childhood tuberculosis in Leeds, 1982-90: social and ethnic factors and the role of the contact clinic in diagnosis. *Thorax* 1994;49:184-5.
- 11 Mangtani P, Jolley DJ, Watson JM, et al. Socio-economic deprivation and notification rates for tuberculosis rates in London during 1987-91. *BMJ* 1995;310:963-6.
- 12 Bhatti N, Law MR, Morris JK, et al. Increasing incidence of tuberculosis in England and Wales: a study of the likely causes. *BMJ* 1995;310:967-9.