

Alpha₁-antitrypsin deficiency, cirrhosis and emphysema

Ravi Mahadeva, David A Lomas

Emphysema is a chronic progressive lung disease characterised by abnormal permanent enlargement of airspaces as a result of destruction of alveolar walls.¹ Most patients develop emphysema as a consequence of smoking but 1-2% of patients with emphysema develop the condition as a result of a genetic deficiency of the plasma proteinase inhibitor α_1 -antitrypsin. The two common deficiency variants of α_1 -antitrypsin, S and Z, result from point mutations in the α_1 -antitrypsin gene²⁻⁴ and are named on the basis of their slower electrophoretic mobility on isoelectric focusing analysis compared with the normal M allele.⁵ S α_1 -antitrypsin (²⁶⁴Glu→Val) is found in up to 28% of Southern Europeans and, although it results in plasma α_1 -antitrypsin levels that are 60% of the M allele, it is not associated with any pulmonary sequelae. The Z variant (³⁴²Glu→Lys) results in a more severe deficiency that is characterised, in the homozygote, by plasma α_1 -antitrypsin levels of 10% of the normal M allele and by levels of 60% in the MZ heterozygote (50% from the M allele and 10% from the Z allele). The Z mutation results in the accumulation of α_1 -antitrypsin in the rough endoplasmic reticulum of the liver (fig 1A) and predisposes the homozygote to juvenile hepatitis, cirrhosis,⁶ and hepatocellular carcinoma.⁷ Z α_1 -antitrypsin inclusions are associated with abnormal liver function tests in over 90% of Z homozygotes in the first year of life but only 10-15% of these develop the prolonged cholestatic jaundice that can progress to cirrhosis and the requirement for hepatic transplantation.^{6, 8}

The role of α_1 -antitrypsin is to protect the tissues against enzymatic digestion by neutrophil elastase.⁹ The low circulating levels are unable to inhibit this proteinase and predispose the Z homozygote to early onset panlobular emphysema,¹⁰ bronchiectasis,¹¹ and vasculitis.¹² Alpha₁-antitrypsin deficiency related emphysema is predominantly panlobular and basal compared with the centrilobular upper lobe disease seen in smokers. Patients usually present with increasing dyspnoea and weight loss, with cor pulmonale and polycythaemia occurring late in the course of the disease. Chest radiographs and high resolution CT scans typically show bilateral basal emphysema with paucity and pruning of the basal pulmo-

nary vessels (fig 1B). Upper lobe vascularisation is relatively normal and ventilation perfusion radioisotope scans and angiography also show abnormalities with a lower zone distribution.¹³ Lung function tests are typical for emphysema with a reduced FEV₁/FVC ratio, evidence of air trapping and a low gas transfer factor.

Structure and function of α_1 -antitrypsin

Alpha₁-antitrypsin is the archetypal member of the serine proteinase inhibitor or serpin superfamily.^{14, 15} Members of the family have





Figure 1 (A) Electron micrograph of a hepatocyte from the liver of a child with Z α_1 -antitrypsin deficiency. The arrow shows the accumulation of α_1 -antitrypsin within the rough endoplasmic reticulum and the distortion of the hepatocellular architecture that predisposes to cell death and cirrhosis (reproduced from ref 26 with permission). (B) High resolution CT scan from a patient with Z α_1 -antitrypsin deficiency showing the characteristic basal panlobular emphysema rather than the apical centrilobular disease seen in smokers who have normal levels of α_1 -antitrypsin.

Respiratory Medicine
Unit, Department of
Medicine and
Department of
Haematology
D A Lomas
R Mahadeva

University of
Cambridge, MRC
Centre, Hills Road,
Cambridge CB2 2QH,
UK

Correspondence to:
Dr R Mahadeva.



more than 30% sequence homology with α_1 -antitrypsin and share a similar molecular architecture¹⁴ based on a dominant A β -pleated sheet and nine α -helices. This structure supports an exposed, mobile, reactive loop peptide that acts as a pseudosubstrate to inhibit the target proteinase. In the case of α_1 -antitrypsin the loop presents a methionine residue at the P_1 site to interact with the substrate binding pocket of neutrophil elastase. After binding the proteinase is inactivated by complex formation and cleared from the circulation.¹⁶ Crystallographic analysis of the serpins has yielded a series of structures which illustrate the striking mobility of the reactive centre loop, allowing its function to be likened to that of a mousetrap (fig 2A). More recently our crystal structures of intact α_1 -antitrypsin^{17, 18} have revealed the conformation of the loop that is ideal for docking with neutrophil elastase (fig 2B) before it is locked into a stable complex by the mousetrap mechanism. This mobility of the loop is essential for




Figure 3 (A) Loop-sheet polymers of α_1 -antitrypsin isolated from the plasma of a Siiyama α_1 -antitrypsin homozygote. This point mutation favours the formation of long chains of polymers as well as circlets (inset). These polymers tangle in the liver to form insoluble inclusions that predispose to liver damage (reproduced from ref 29 with permission). (B) Our recent crystal structure of α_1 -antitrypsin provides strong support for the loop-sheet linkage being between the β -pleated reactive centre loop of one molecule and the A β -pleated sheet of the next. These dimers extend to form long chains shown here as alternating yellow, blue and red α_1 -antitrypsin molecules (reproduced from ref 17 with permission). The prevention of this linkage would block polymerisation and so attenuate the liver disease associated with the Z mutation.

inhibitory function but may allow aberrant conformations by insertion into the A β -sheet of a second α_1 -antitrypsin molecule to form a dimer which then extends to form a chain of loop-sheet polymers.^{19–21} Such polymers form following treatment with mild denaturants²² and upon heating α_1 -antitrypsin²¹ and other serpins²³ at high temperatures.

Loop-sheet polymers and liver disease

The Z mutation of α_1 -antitrypsin is located at residue P₁₇ (17 residues proximal to the P₁ reactive centre) at the head of strand 5 of the A β -sheet and the base of the mobile reactive centre loop (fig 2). We predicted that the mutation would act to open the A β -sheet thereby favouring spontaneous reactive loop into β -sheet polymerisation. Indeed, functional studies showed that Z α_1 -antitrypsin was more unstable^{24 25} and able to polymerise at 37°C under physiological conditions.²⁶ The rate was accelerated by raising the temperature to 41°C and could be blocked by exogenous reactive loop peptides that compete with the loop for A β -sheet annealing. The role of this phenomenon in vivo was clarified when it was found that polymers with identical microscopic appearance could be isolated from the liver of a Z α_1 -antitrypsin homozygote. Although many α_1 -antitrypsin deficiency variants have been

described, only two other mutants of α_1 -antitrypsin have similarly been associated with plasma deficiency and hepatic inclusions: α_1 -antitrypsin Siiyama (⁵³Ser→Phe)²⁷ and α_1 -antitrypsin Mmalton (⁵²Phe deleted).²⁸ Both of these mutants also formed spontaneous loop-sheet polymers in vivo (fig 3A).^{29 30} The temperature and concentration dependence of polymerisation, along with genetic factors,^{31 32} may account for the heterogeneity in clinical features. As α_1 -antitrypsin is an acute phase protein the concentration will rise during episodes of inflammation. At these times the formation of polymers is likely to overwhelm the degradative pathway, thereby exacerbating hepatic inclusions and the associated hepatocellular damage.

The phenomenon of loop-sheet polymerisation is not restricted to α_1 -antitrypsin and has now been reported to account for the deficiency of mutants of C1 inhibitor³³ and antithrombin^{34 35} in patients with angioedema and thrombosis, respectively, and has been linked to a deficiency of α_1 -antichymotrypsin in patients with liver disease and emphysema.^{36 37} It also underlies the plasma deficiency associated with common S variant of α_1 -antitrypsin.³⁸ The rate of polymerisation of S α_1 -antitrypsin is slower than that associated with the Z variant which is compatible with the milder clinical phenotype. The precise linkage underlying the formation of polymers in vivo has been probed by site directed mutagenesis and protein expression in the *Xenopus* oocyte system. A point mutation in the region of opening of the A sheet (⁵¹Phe→Leu) stabilises the α_1 -antitrypsin molecule³⁹ and attenuates the effect of the Z mutation in vitro⁴⁰ and in vivo.⁴¹ Moreover, this mutation completely prevents the accumulation of α_1 -antitrypsin Siiyama in a *Xenopus* oocyte expression system.⁴¹ Recombinant α_1 -antitrypsin with and without the ⁵¹Phe→Leu mutation was expressed in *E coli* and purified to homogeneity. The proteins function normally as inhibitors of neutrophil elastase¹⁷ and were used for crystallographic analysis. Crystals of recombinant protein were grown and two structures of α_1 -antitrypsin solved to 2.9 Å. These structures,^{17 18} and that of Ryu *et al.*,⁴² provide an explanation for the ready formation of loop-sheet polymers as the reactive loop strand configuration fits easily into the A β -pleated sheet of α_1 -antitrypsin to form a dimer that can then extend as trimers, tetramers, and so on, to as many as 20 molecules in length (fig 3B). It is these filaments which then tangle within the endoplasmic reticulum of the hepatocyte to form the insoluble inclusions that are characteristic of Z α_1 -antitrypsin deficiency.

Loop-sheet polymers and emphysema

Alpha₁-antitrypsin deficiency related emphysema is believed to result from the reduced levels of circulating proteinase inhibitor being unable to protect the lungs against proteolytic attack.^{10 43} The Z α_1 -antitrypsin that does escape from the liver into the circulation is less efficient at protecting the tissues from enzyme

damage^{21 44 45} and, like M α_1 -antitrypsin,^{46 47} may be inactivated by oxidation of the P₁ methionine residue. The demonstration that Z α_1 -antitrypsin can undergo a spontaneous conformational transition in association with liver disease raised the possibility that this may also occur within the lung. Indeed, we have detected polymers in bronchoalveolar lavage fluid in two out of five patients with Z α_1 -antitrypsin deficiency but not in lavage fluid from 13 MZ, MS, or M α_1 -antitrypsin controls.⁴⁸ This may have important implications for the pathogenesis of disease as polymerisation obscures the reactive loop of α_1 -antitrypsin, rendering the protein inactive as an inhibitor of proteolytic enzymes.²¹ Thus, the spontaneous polymerisation of α_1 -antitrypsin within the lung will exacerbate the already reduced antiprotease screen, thereby increasing the susceptibility of the tissues to proteolytic attack. The relationship of α_1 -antitrypsin polymers to smoking, infection, and rate of decline in lung function remains to be established and other studies are required to determine their effects on neutrophils which are found in increased numbers in the lungs of patients with α_1 -antitrypsin deficiency.⁴⁹

Treatment

This new understanding of the structural basis of α_1 -antitrypsin deficiency provides a platform for rational drug design to block polymerisation *in vivo* and so attenuate the associated liver disease.¹⁸ Any treatment that improves secretion from the liver will raise the circulating levels of α_1 -antitrypsin and so enhance the antiprotease protection within the lung. Until that time the prevention of emphysema is better than cure and there is good evidence that many Z α_1 -antitrypsin homozygotes would develop only mild lung disease if they abstained from smoking.⁵⁰⁻⁵² The genetic deficiency in the anti-elastase screen may be rectified biochemically by intravenous infusions of α_1 -antitrypsin⁵³ but the role of this treatment in the prevention of emphysema is unproven. A theoretical option is the intravenous administration of genetically engineered α_1 -antitrypsin with the P₁ methionine mutated to a valine residue.⁵⁴ This has little effect on the function of the protein but makes it resistant to oxidation, thereby improving its anti-elastase activity. Other treatments at an early stage of development include gene therapy and retinoic acid. Vectors carrying the α_1 -antitrypsin gene have been targeted to liver^{55 56} and lung⁵⁷ but there is currently insufficient gene expression for this to be a useful therapeutic modality. Similarly, although the effects of retinoic acid on alveolar regeneration in the rat look promising,⁵⁸ it has yet to be tested in man.

In the meantime patients with α_1 -antitrypsin deficiency related emphysema should receive conventional therapy with trials of bronchodilators, inhaled corticosteroids and, where appropriate, assessment for long term oxygen therapy and single lung transplantation. The role of lung volume reduction surgery in this group is even more contentious than in smok-

ing related centrilobular emphysema as the disease is basal rather than apical and typically lacks the target areas which are most suitable for resection.

Conclusion

Z α_1 -antitrypsin deficiency results from the accumulation of loop-sheet polymers within the endoplasmic reticulum of the hepatocyte. These polymers also form within the lung where they inactivate α_1 -antitrypsin, thereby further reducing the antiprotease screen that protects the tissues against proteolytic attack. Recent advances in understanding the transition from monomeric to polymeric protein raises the prospect of rational drug design to treat the deficiency and provides a useful paradigm for other conformational diseases such as the prion diseases, amyloidosis and Alzheimer's disease.⁵⁹

- 1 American Thoracic Society. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease (COPD) and asthma. *Am Rev Respir Dis* 1986;136:225-44.
- 2 Owen MC, Carrell RW. Alpha-1-antitrypsin: molecular abnormality of S variant. *BMJ* 1976;1:130-1.
- 3 Jeppsson J-O. Amino acid substitution Glu→Lys in α_1 -antitrypsin PiZ. *FEBS Lett* 1976;65:195-7.
- 4 Yoshida A, Lieberman J, Gaidulis L, et al. Molecular abnormality of human alpha-antitrypsin variant (Pi-ZZ) associated with plasma activity deficiency. *Proc Natl Acad Sci USA* 1976;73:1324-8.
- 5 Brantly M, Nukiwa T, Crystal RG. Molecular basis of alpha-1-antitrypsin deficiency. *Am J Med* 1988;84(Suppl 6A):13-31.
- 6 Sveger T. Liver disease in alpha-antitrypsin deficiency detected by screening of 200,000 infants. *N Engl J Med* 1976;294:1316-21.
- 7 Eriksson S, Carlson J, Velez R. Risk of cirrhosis and primary liver cancer in alpha-antitrypsin deficiency. *N Engl J Med* 1986;314:736-9.
- 8 Sveger T. The natural history of liver disease in α_1 -antitrypsin deficient children. *Acta Paediatr Scand* 1988;77:847-51.
- 9 Carrell RW, Jeppsson J-O, Laurell C-B, et al. Structure and variation of human α_1 -antitrypsin. *Nature* 1982;298:329-34.
- 10 Eriksson S. Studies in α_1 -antitrypsin deficiency. *Acta Med Scand* 1965;Suppl 432:1-85.
- 11 King MA, Stone JA, Diaz PT, et al. α_1 -antitrypsin deficiency: evaluation of bronchiectasis with CT. *Radiology* 1996;199:137-41.
- 12 Griffith ME, Lovegrove JU, Gaskin G, et al. C-antineutrophil cytoplasmic antibody positivity in vasculitis patients is associated with the Z allele of alpha-1-antitrypsin, and P-antineutrophil cytoplasmic antibody positivity with the S allele. *Nephrol Dial Transplant* 1996;11:438-43.
- 13 Stein PD, Leu JD, Welsh MH, et al. Pathophysiology of the pulmonary circulation in emphysema associated with alpha-1 antitrypsin deficiency. *Circulation* 1971;43:227-39.
- 14 Huber R, Carrell RW. Implications of the three-dimensional structure of α_1 -antitrypsin for structure and function of serpins. *Biochemistry* 1989;28:8951-66.
- 15 Potempa J, Korzus E, Travis J. The serpin superfamily of proteinase inhibitors: structure, function, and regulation. *J Biol Chem* 1994;269:15957-60.
- 16 Mast AE, Enghild JJ, Pizzo SV, et al. Analysis of the plasma elimination kinetics and conformational stabilities of native, proteinase-complexed, and reactive site cleaved serpins: comparison of α_1 -proteinase inhibitor, α_1 -antichymotrypsin, antithrombin III, α_2 -antiplasmin, angiotensinogen, and ovalbumin. *Biochemistry* 1991;30:1723-30.
- 17 Elliott PR, Lomas DA, Carrell RW, et al. Inhibitory conformation of the reactive loop of α_1 -antitrypsin. *Nature Struct Biol* 1996;3:676-81.
- 18 Elliott PR, Abrahams J-P, Lomas DA. Wildtype α_1 -antitrypsin is in the canonical inhibitory conformation. *J Mol Biol* 1998;275:419-425.
- 19 Schulze AJ, Baumann U, Knof S, et al. Structural transition of α_1 -antitrypsin by a peptide sequentially similar to β -strand s4A. *Eur J Biochem* 1990;194:51-6.
- 20 Mast AE, Enghild JJ, Salvesen G. Conformation of the reactive site loop of α_1 -proteinase inhibitor probed by limited proteolysis. *Biochemistry* 1992;31:2720-8.
- 21 Lomas DA, Evans DL, Stone SR, et al. Effect of the Z mutation on the physical and inhibitory properties of α_1 -antitrypsin. *Biochemistry* 1993;32:500-8.
- 22 Lomas DA, Elliott PR, Chang W-SW, et al. Preparation and characterisation of latent α_1 -antitrypsin. *J Biol Chem* 1995;270:5282-88.

- 23 Patston PA, Hauert J, Michaud M, *et al.* Formation and properties of C1-inhibitor polymers. *FEBS Lett* 1995;**368**:401–4.
- 24 Le A, Ferrell GA, Dishon DS, *et al.* Soluble aggregates of the human PiZ α_1 -antitrypsin variant are degraded within the endoplasmic reticulum by a mechanism sensitive to inhibitors of protein synthesis. *J Biol Chem* 1992;**267**:1072–80.
- 25 Yu M-H, Lee KN, Kim J. The Z type variation of human α_1 -antitrypsin causes a protein folding defect. *Nature Struct Biol* 1995;**2**:363–7.
- 26 Lomas DA, Evans DL, Finch JT, *et al.* The mechanism of Z α_1 -antitrypsin accumulation in the liver. *Nature* 1992;**357**:605–7.
- 27 Seyama K, Nukiwa T, Takabe K, *et al.* Siiyama (serine 53 (TCC) to phenylalanine 53 (TTC)). A new α_1 -antitrypsin-deficient variant with mutation on a predicted conserved residue of the serpin backbone. *J Biol Chem* 1991;**266**:12627–32.
- 28 Roberts EA, Cox DW, Medline A, *et al.* Occurrence of alpha-1-antitrypsin deficiency in 155 patients with alcoholic liver disease. *Am J Clin Pathol* 1984;**82**:424–7.
- 29 Lomas DA, Finch JT, Seyama K, *et al.* α_1 -antitrypsin S_{Siiyama} (Ser⁵³→Phe); further evidence for intracellular loop-sheet polymerisation. *J Biol Chem* 1993;**268**:15333–5.
- 30 Lomas DA, Elliott PR, Sidhar SK, *et al.* Alpha₁-antitrypsin Mmalton (³²Phe deleted) forms loop-sheet polymers in vivo: evidence for the C sheet mechanism of polymerisation. *J Biol Chem* 1995;**270**:16864–70.
- 31 Wu Y, Whitman I, Molmenti E, *et al.* A lag in intracellular degradation of mutant α_1 -antitrypsin correlates with liver disease phenotype in homozygous PiZZ α_1 -antitrypsin deficiency. *Proc Natl Acad Sci USA* 1994;**91**:9014–8.
- 32 Teckman JH, Perlmutter DH. The endoplasmic reticulum degradation pathway for mutant secretory proteins α_1 -antitrypsin Z and S is distinct from that for an unassembled membrane protein. *J Biol Chem* 1996;**271**:13215–20.
- 33 Eldering E, Verpy E, Roem D, *et al.* COOH-terminal substitutions in the serpin C1 inhibitor that cause loop overinsertion and subsequent multimerization. *J Biol Chem* 1995;**270**:2579–87.
- 34 Bruce D, Perry DJ, Borg J-Y, *et al.* Thromboembolic disease due to thermolabile conformational changes of anti-thrombin Rouen VI (187 Asn→Asp). *J Clin Invest* 1994;**94**:2265–74.
- 35 Lindo VS, Kakkar VV, Learmonth M, *et al.* Antithrombin-TRI (Ala382 to Thr) causing severe thromboembolic tendency undergoes the S-to-R transition and is associated with a plasma-inactive high-molecular-weight complex of aggregated antithrombin. *Br J Haematol* 1995;**89**:589–601.
- 36 Faber J-P, Poller W, Olek K, *et al.* The molecular basis of α_1 -antichymotrypsin deficiency in a heterozygote with liver and lung disease. *J Hepatol* 1993;**18**:313–21.
- 37 Poller W, Faber J-P, Weidinger S, *et al.* A leucine-to-proline substitution causes a defective α_1 -antichymotrypsin allele associated with familial obstructive lung disease. *Genomics* 1993;**17**:740–3.
- 38 Elliott PR, Stein PE, Bilton D, *et al.* Structural explanation for the dysfunction of S α_1 -antitrypsin. *Nature Struct Biol* 1996;**3**:910–11.
- 39 Kwon K-S, Kim J, Shin HS, *et al.* Single amino acid substitutions of α_1 -antitrypsin that confer enhancement in thermal stability. *J Biol Chem* 1994;**269**:9627–31.
- 40 Kim J, Lee KN, Yi G-S, *et al.* A thermostable mutation located at the hydrophobic core of α_1 -antitrypsin suppresses the folding defect of the Z-type variant. *J Biol Chem* 1995;**270**:8597–601.
- 41 Sidhar SK, Lomas DA, Carrell RW, *et al.* Mutations which impede loop/sheet polymerisation enhance the secretion of human α_1 -antitrypsin deficiency variants. *J Biol Chem* 1995;**270**:8393–6.
- 42 Ryu S-E, Choi H-J, Kwon K-S, *et al.* The native strains in the hydrophobic core and flexible reactive loop of a serine protease inhibitor: crystal structure of an uncleaved α_1 -antitrypsin at 2.7 Å. *Structure* 1996;**4**:1181–92.
- 43 Snider GL. Emphysema: the first two centuries—and beyond. A historical overview, with suggestions for future research. *Am Rev Respir Dis* 1992;**146**:1615–22.
- 44 Ogushi F, Fells GA, Hubbard RC, *et al.* Z-type α_1 -antitrypsin is less competent than M1-type α_1 -antitrypsin as an inhibitor of neutrophil elastase. *J Clin Invest* 1987;**80**:1366–74.
- 45 Llewellyn-Jones CG, Lomas DA, Carrell RW, *et al.* The effect of the Z mutation on the ability of α_1 -antitrypsin to prevent neutrophil mediated tissue damage. *Biochim Biophys Acta* 1994;**1227**:155–60.
- 46 Beatty K, Bieth J, Travis J. Kinetics of association of serine proteinases with native and oxidized α_1 -proteinase inhibitor and α_1 -antichymotrypsin. *J Biol Chem* 1980;**255**:3931–4.
- 47 Carp H, Miller F, Hoidal JR, *et al.* Potential mechanism of emphysema: α_1 -proteinase inhibitor recovered from lungs of cigarette smokers contains oxidised methionine and has decreased elastase inhibitory capacity. *Proc Natl Acad Sci USA* 1982;**79**:2041–5.
- 48 Elliott PR, Bilton D, Lomas DA. Lung polymers in Z α_1 -antitrypsin related emphysema. *Am J Respir Cell Mol Biol* (in press).
- 49 Morrison HM, Kramps JA, Burnett D, *et al.* Lung lavage fluid from patients with α_1 -proteinase inhibitor deficiency or chronic obstructive bronchitis: anti-elastase function and cell profile. *Clin Sci* 1987;**72**:373–81.
- 50 Larsson C. Natural history and life expectancy in severe alpha₁-antitrypsin deficiency, PiZ. *Acta Med Scand* 1978;**204**:345–51.
- 51 Janus ED, Phillips NT, Carrell RW. Smoking, lung function, and α_1 -antitrypsin deficiency. *Lancet* 1985;**i**:152–4.
- 52 Hutchison DCS. Epidemiology of alpha₁-protease inhibitor deficiency. *Eur Respir J* 1990;**3**(Suppl 9):29s–34s.
- 53 Wewers MD, Casolaro MA, Sellers SE, *et al.* Replacement therapy for alpha₁-antitrypsin deficiency associated with emphysema. *N Engl J Med* 1987;**316**:1055–62.
- 54 Courtney M, Jallat S, Tessier LH, *et al.* Synthesis in *E coli* of α_1 -antitrypsin variants of therapeutic potential for emphysema and thrombosis. *Nature* 1985;**313**:149–51.
- 55 Kay MA, Baley P, Rothenberg S, *et al.* Expression of human α_1 -antitrypsin in dogs after autologous transplantation of retroviral transduced hepatocytes. *Proc Natl Acad Sci USA* 1992;**89**:89–93.
- 56 Jaffe HA, Danel C, Longenecker G, *et al.* Adenovirus-mediated in vivo gene transfer and expression in normal rat liver. *Nature Genetics* 1992;**1**:372–8.
- 57 Rosenfeld MA, Siegfried W, Yoshimura K, *et al.* Adenovirus-mediated transfer of a recombinant α_1 -antitrypsin gene to the lung epithelium in vivo. *Science* 1991;**252**:431–4.
- 58 Massaro GD, Massaro D. Retinoic acid treatment abrogates elastase-induced pulmonary emphysema in rats. *Nature Med* 1997;**3**:675–7.
- 59 Carrell RW, Lomas DA. Conformational diseases. *Lancet* 1997;**350**:134–8.
- 60 Carrell RW, Stein PE, Fermi G, *et al.* Biological implications of a 3 Å structure of dimeric antithrombin. *Structure* 1994;**2**:257–70.
- 61 Schreuder HA, de Boer B, Dijkema R, *et al.* The intact and cleaved human antithrombin III complex as a model for serpin-proteinase interactions. *Nature Struct Biol* 1994;**1**:48–54.