

SHORT PAPER

Wheeze not current asthma affects quality of life in young adults with asthma

M Matheson, J Raven, R K Woods, F Thien, E H Walters, M Abramson

See end of article for authors' affiliations

Thorax 2002;**57**:165–167

Correspondence to:
A/Prof Michael Abramson,
Department of
Epidemiology and
Preventive Medicine,
Central & Eastern Clinical
School, Monash University,
Prahran, Victoria 3181,
Australia;
michael.abramson@
med.monash.edu.au

Revised version received
17 September 2001
Accepted for publication
10 October 2001

Background: A study was undertaken to investigate quality of life in asthma, defined by differing criteria, to see which may be most appropriate in epidemiological studies.

Methods: The 426 adults were participants in the follow up phase of the European Community Respiratory Health Survey (ECRHS) in Melbourne. As part of the laboratory visit, participants completed the SF-36 quality of life questionnaire, a detailed respiratory questionnaire, and underwent lung function testing.

Results: Both the physical component summary and the mental component summary scores were significantly worse in those with wheeze in the previous 12 months than in those without wheeze. Only the mental component summary score was significantly worse in those with current asthma than in those without. In contrast, in those with current asthma or bronchial hyperreactivity only, neither of the summary scales was significantly different between cases and controls.

Conclusions: Quality of life is severely impaired in individuals with wheeze in the previous 12 months while individuals with current asthma or bronchial hyperreactivity alone did not appear to have significantly reduced quality of life.

Bronchial hyperreactivity (BHR) to histamine or methacholine has been used as an objective physiological marker of asthma and, in combination with wheeze in the previous 12 months, has been used to define "current asthma" in epidemiological studies. It has been claimed that this definition discriminates a group with more severe asthma than subjective definitions based on self-reported asthma symptoms alone.^{1,2} However, subjective measures of asthma severity have been found to correlate much better with measures of quality of life (QoL) than objective physiological measures such as BHR and forced expiratory volume in 1 second (FEV₁).³ The purpose of this study was to examine the relationship between QoL and symptom based and physiological definitions of asthma in a community setting to determine which definition might be most appropriate.

METHODS

The subjects were participants in the follow up phase of the ECRHS conducted in Melbourne in 1998/9. Full details of the original sampling protocol have been described elsewhere.⁴ Participants completed the detailed ECRHS questionnaire, spirometric tests, and a methacholine challenge. QoL was evaluated by the short form (SF-36) health survey which was completed by participants upon arrival at the laboratory. All questionnaires were checked for missing data by one of the trained interviewers after completion. Only the physical component summary (PCS) score and the mental component summary (MCS) score are reported in this analysis which were calculated using the three step procedure recommended by the developer.⁵ A total of 426 participants completed the methacholine challenge and were included in this analysis.

Wheeze only was defined as a positive response to the question: "Have you had wheezing or whistling in your chest at any time in the last 12 months?".¹ The ECRHS defined asthma as a positive response to any of the following questions: "Have you had an attack of asthma in the last 12 months?", "Have you been woken by an attack of shortness of breath at any time in last 12 months?", and "Are you currently taking any medicine for asthma?".⁴ BHR was defined as a provocative

dose of <2 mg methacholine causing a 20% fall in FEV₁ (PD₂₀). Current asthma was defined as a positive response to the question: "Have you had wheezing or whistling in your chest at any time in the last 12 months?" and measured BHR.¹ Doctor diagnosed asthma was defined as a positive response to the question: "Have you ever had asthma?" and then to the question: "Was this confirmed by a doctor?". BHR alone and FEV₁ % predicted were used as purely physiological definitions of asthma.

Comparisons in QoL scores were made between subjects meeting the various definitions of asthma and the remaining subjects who did not meet the criteria. The Mann-Whitney U test was used to test if the distribution of the SF-36 scores was the same across the different definitions and to assess whether SF-36 scores were associated with sex, age, and smoking status. A multivariate regression model was used to test for significance between cases and controls after adjusting for age, sex, and current smoking status. Spearman's rank correlation was used to assess the association between FEV₁ % predicted and SF-36 scores. A p value of <0.05 was considered significant. All analyses were conducted using Stata (Stata for Windows, Stata Corporation 1997, Texas, US). Post hoc power calculations indicated that there was a greater than 80% power to detect a three point difference in PCS and MCS scores between groups.

RESULTS

Participants had a mean (SD) age of 39.7 (6.4) years (range 26–50), 50.0% were women, and 81.5% were Australian born. The mean (SD) FEV₁ was 109.3 (14.5)% predicted. Eighty five participants (25%) were current smokers and 214 (50.2%) reported ever smoking. Doctor diagnosed asthma was reported by 120 participants (28.2%) and an attack of asthma in the previous 12 months was reported by 71 (16.7%).

The key asthma symptom of wheeze in the preceding 12 months was reported by 177 participants (41.5%). Nocturnal shortness of breath (SOB) and spontaneous SOB were reported by 43 (10.1%) and 53 (12.4%) participants, respectively. BHR was demonstrated by 106 participants

Table 1 Mean (SD) scores for SF-36 summary measures for the different definitions of asthma

	Physical component summary	Adjusted p value	Mental component summary	Adjusted p value
Wheeze only				
Cases (n=177)	52.94 (7.14)	0.006	48.87 (9.30)	0.002
Controls (n=249)	54.69 (6.04)		51.66 (7.49)	
ECRHS				
Cases (n=108)	52.27 (7.64)	0.001	49.08 (8.64)	0.042
Controls (n=318)	54.54 (6.07)		50.99 (8.27)	
Doctor diagnosed asthma				
Cases (n=120)	52.29 (7.53)	0.001	49.55 (8.64)	0.173
Controls (n=306)	54.62 (6.04)		50.88 (8.28)	
Current asthma				
Cases (n=80)	52.90 (7.86)	0.088	48.89 (8.94)	0.064
Controls (n=346)	54.21 (6.22)		50.88 (8.23)	
BHR				
Cases (n=106)	53.46 (7.24)	0.368	49.58 (8.42)	0.272
Controls (n=320)	54.13 (6.33)		50.81 (8.37)	

p values were calculated after adjusting for age, sex, and smoking status by multiple linear regression.

(24.9%). The number of participants who reported that they were currently on medication for their asthma was 145 (34.0%). While 38 (8.9%) reported seeing a GP specifically because of their breathing, only eight (1.9%) reported seeing a specialist physician.

In this population of young adults, sex, age, and current smoking status were not important determinants of QoL scores. SF-36 summary scores for the alternative definitions of asthma are presented in table 1. The PCS and MCS scores were both significantly worse in those with wheeze in the preceding 12 months than in those without wheeze. Similarly, for the ECRHS definition of asthma both the PCS and MCS scores were significantly different between the cases and controls. For doctor diagnosed asthma only the PCS score was significantly worse in adults with self-reported doctor diagnosed asthma than in those without. In contrast to these results, for current asthma neither the PCS nor MCS scores were significantly worse in those with current asthma than in those without. For those with BHR only neither PCS nor MCS scores were significantly different between cases and controls. The rank correlations between FEV₁, % predicted and the PCS and MCS scores were not significant (PCS $\rho=0.3$, $p=0.06$; MCS $\rho=0.2$, $p=0.4$).

DISCUSSION

This is the first study of which we are aware to investigate specifically the relationships between different epidemiological definitions of asthma and reported QoL using the SF-36 questionnaire in a community based sample of young adults. There is some controversy over which symptom questions are the most valid in diagnosing asthma for epidemiological studies.² We have found that subjective definitions of asthma such as wheeze in the previous 12 months, the ECRHS definition of asthma, and self-reported doctor diagnosed asthma define individuals with significantly worse QoL. On the other hand, the epidemiological definition of current asthma and purely physiologically based definitions of asthma such as BHR alone or FEV₁ were not associated with significantly worse QoL. These results would suggest that merely having respiratory symptoms of any type is sufficient to adversely affect QoL and, in the context of this particular population, the underlying problem was likely to be asthma.

Data from the French centres of the ECRHS found that subjects with asthma had significantly lower PCS and MCS scores than control subjects using a combination of self-reported symptoms and BHR to define cases.⁶ Several other studies have reported strong correlations between poor QoL and subjective measures of asthma severity such as diary

obtained symptom scores and β_2 agonist use for the relief of symptoms.^{3,7} On the other hand, correlations between objective measures such as FEV₁ have been found to be poor.⁸ Our results are consistent with these studies and show that, even for individuals with mostly mild asthma from a community setting, QoL is more strongly associated with self reported "subjective" than objective measures of asthma.

Clinical observations suggest that patients' concerns with regard to their asthma tend to focus on symptom frequency, activity limitation, and avoidance of irritants.⁹ For subjects with mild asthma who experience few symptoms QoL is equivalent to or better than population norms but when they experience symptoms their QoL is significantly affected,¹⁰ while individuals with more severe asthma and more frequent respiratory symptoms who have accommodated their lifestyle to this chronic condition might perceive less impact on QoL. This would suggest that most people tend to focus on subjective rather than objective measures of their asthma and, given that QoL is a self-assessed measure, it is not surprising that subjective definitions of asthma correlate more closely than do objective measures of asthma with poor QoL.

In conclusion, we have confirmed that symptom based definitions are more closely related to QoL than are physiological definitions of asthma. We suggest that, for epidemiological studies investigating the social, psychological, and economic costs of asthma, it might be more appropriate to use symptom based definitions when defining subjects. While our conclusions are valid for epidemiological studies of asthma in the community, they should not be generalised to clinical trials of patients with more severe disease. However, we do suggest that in clinical trials QoL measures would provide clinicians with valuable further information.

ACKNOWLEDGEMENTS

The authors are grateful to Dean McKenzie and Rory Wolfe for statistical advice and Trina Vincent for assistance with data collection. This project was supported by the Victorian Health Promotion Foundation and the National Health & Medical Research Council of Australia.

Authors' affiliations

M Matheson, R K Woods, M Abramson, Department of Epidemiology and Preventive Medicine, Central & Eastern Clinical School, Monash University, Prahran, Victoria 3181, Australia

J Raven, Department of Respiratory Medicine, The Alfred Hospital, Prahran, Victoria 3181, Australia

F Thien, Department of Allergy, Asthma and Clinical Immunology, The Alfred Hospital, Prahran, Victoria 3181, Australia

E H Walters, Department of Clinical Sciences, The University of Tasmania Medical School, Hobart, Tasmania 7001, Australia

REFERENCES

- 1 **Toelle BG**, Peat JK, Salome CM, *et al.* Towards a definition of asthma for epidemiology. *Am Rev Respir Dis* 1992;**146**:633–7.
- 2 **Jenkins MA**, Clarke JR, Carlin JB, *et al.* Validation of questionnaire and bronchial hyperresponsiveness against respiratory physician assessment in the diagnosis of asthma. *Int J Epidemiol* 1996;**25**:609–16.
- 3 **van der Molen T**, Postma DS, Schreurs AJ, *et al.* Discriminative aspects of two generic and two asthma-specific instruments: relation with symptoms, bronchodilator use and lung function in patients with mild asthma. *Qual Life Res* 1997;**6**:353–61.
- 4 **Burney PGJ**, Luczynska C, Chinn S, *et al.* The European Community Respiratory health survey. *Eur Respir J* 1994;**7**:954–60.
- 5 **Ware JE**, Kosinski M, Keller SD. *SF-36 physical and mental health summary scales: a user's manual*. 3rd ed. Boston: The Health Institute, New England Medical Centre, 1994.
- 6 **Leynaert B**, Neukirch C, Liard R, *et al.* Quality of life in allergic rhinitis and asthma: A population based study of young adults. *Am J Respir Crit Care Med* 2000;**162**:1391–6.
- 7 **Bousquet J**, Knani J, Dhivert H, *et al.* Quality of life in asthma: internal consistency and validity of the SF-36 questionnaire. *Am J Respir Crit Care Med* 1994;**149**:371–5.
- 8 **Ware JE**, Kemp JP, Buchner DA, *et al.* The responsiveness of disease-specific and generic health measures to changes in the severity of asthma among adults. *Qual Life Res* 1998;**7**:235–44.
- 9 **Schmier JK**, Chan KS, Leidy NK. The impact of asthma on health related quality of life. *J Asthma* 1998;**35**:585–97.
- 10 **Osman LM**, Calder C, Robertson R, *et al.* Symptoms, quality of life, and health service contact among young adults with mild asthma. *Am J Respir Crit Care Med* 2000;**161**:498–503.

SHORT PAPER

Patients' and carers' preferences in two models of care for acute exacerbations of COPD: results of a randomised controlled trial

J C Ojoo, T Moon, S McGlone, K Martin, E D Gardiner, M A Greenstone, A H Morice

Thorax 2002;**57**:167–169

See end of article for authors' affiliations

Correspondence to: Professor A H Morice, Academic Department of Medicine, University of Hull, Castle Hill Hospital, Cottingham, Hull HU16 5JQ, UK; A.H.Morice@hull.ac.uk

Revised version received 24 September 2001
Accepted for publication 5 October 2001

Background: Patients with an acute exacerbation of chronic obstructive pulmonary disease (COPD) were randomised to either hospital at home (HaH) or inpatient management, and patient and carer preferred site of management and satisfaction with care received in the two arms was determined.

Methods: Emergency admissions with an acute exacerbation of COPD were randomised to inpatient care or HaH care. After discharge an independent observer administered a questionnaire to both patients and carers on the preferred site of care and scored satisfaction with the care received.

Results: Of 60 patients recruited, 30 were randomised to receive HaH care. Retrospective patient preference for HaH care was 96.3% in the domiciliary arm and 59.3% in the conventional arm; carer preference figures were 85.7% and 42.9%, respectively. There was a higher preference for domiciliary care by both patients and carers in the HaH arm than in the inpatient arm ($p=0.001$ and $p=0.01$, respectively). Patients recorded equal satisfaction with care in the two arms (88.1% in the conventional arm, 91.7% in the domiciliary arm); carer scores were 91.3% and 91.9%, respectively.

Conclusions: The results of this study show that both patients and carers were significantly more likely to prefer domiciliary care if they were in the HaH arm. Since patients had to be willing to be looked after at home, both patients' and carers' perceptions of the benefits of HaH care were reinforced by their experience. HaH care of acute exacerbations of COPD is the preferred option in suitable patients.

Recent studies have shown that hospital at home (HaH) management of selected patients with acute exacerbations of chronic obstructive pulmonary disease (COPD) is a safe^{1,2} and cost effective option to inpatient care.³ As HaH schemes have become increasingly popular in recent years in hospitals across the UK, it is important to know their acceptability to patients and carers.

METHODS

Patients with an acute exacerbation of COPD were admitted to the Medical Chest Unit, Castle Hill Hospital and clinical management was instituted according to the British Thoracic Society guidelines.³ They were reviewed the following morning for possible inclusion in the trial. Both patients and carers gave informed consent for the study. A carer was defined as a provider of emotional or physical support to the

patient during his or her illness. Patients were randomised using sealed envelopes to receive either conventional inpatient care or HaH care under a team of two respiratory outreach nurses (RONs). Recruitment into the study was carried out from Monday to Thursday. The RONs were accessible by telephone between 09.00 and 17.00 hours daily. Outside these times patients could obtain advice from the Medical Chest Unit through a direct line.

Subjects

Inclusion criteria:

- Both sexes
- >18 years
- FEV₁/FVC ratio <70%
- FEV₁ reversibility to salbutamol <15% (obtained on a previous admission or clinic visit)