- 7 Polosa R, Puddicombe SM, Krishna MT, et al. Expression of c-erbB receptors and ligands in the bronchial epithelium of asthmatic subjects. J Allergy Clin Immunol 2002;109:75–81.
- 8 Giembycz MA. Development status of second generation PDE4 inhibitors for asthma and COPD: the story so far. *Monaldi Arch Chest Dis* 2002;57:48–64.
- 9 Kanehiro A, Ikemura T, Makela MJ, et al. Inhibition of phosphodiesterase 4 attenuates airway hyperresponsiveness and airway inflammation in a model of secondary allergen challenge. Am J Respir Crit Care Med 2001;163:173–84.
- 10 Toward TJ, Broadley KJ. Goblet cell hyperplasia, airway function, and leukocyte infiltration after chronic lipopolysaccharide exposure in conscious guinea pigs: effects of rolipram and dexamethasone. J Pharmacol Exp Ther 2002;302:814–21.
- 11 Dent G, White SR, Tenor H, et al. Cyclic nucleotide phosphodiesterase in human bronchial epithelial cells: characterization of isoenzymes and functional effects of PDE inhibitors. Pulm Pharmacol Ther 1998;11:47–56.
- 12 Fuhrmann M, Jahn H-U, Seybold J, et al. Identification and function of cyclic nucleotide phosphodiesterase isoenzymes in airway epithelial cells. Am J Respir Cell Mol Biol 1999;20:292–302.
- 13 Sevetson BR, Kong X, Lawrence JC Jr. Increasing cAMP attenuates activation of mitogen-activated protein kinase. Proc Natl Acad Sci USA 1993:90:10305–9.
- 14 Berger JT, Voynow JA, Peters KW, et al. Respiratory carcinoma cell lines. MUC genes and glycoconjugates. Am J Respir Cell Mol Biol 1999:20:500–10.
- 15 Sarria B, Naline E, Zhang Y, et al. Muscarinic M₂ receptors in acetylcholineisoproterenol functional antagonism in human isolated bronchus. Am J Physiol 2002;283:L1125–32.
- 16 Labat C, Bara J, Gascard JP, et al. M1/MUC5AC mucin released by human airways in vitro. Eur Respir J 1999;14:390–5.
- 17 Takeyama K, Dabbagh K, Shim JJ, et al. Oxidative stress causes mucin synthesis via transactivation of epidermal growth factor receptor: role of neutrophils. J Immunol 2000;164:1546–52.
- 18 Khori K, Ueki IF, Shim J-J, et al. Pseudomona aeruginosa induces MUC5AC production via epidermal growth factor receptor. Eur Respir J 2002;20:1263–70.
- 19 Mata A, Ruíz A, Cerdá M, et al. Oral N-acetylcysteine reduces bleomycininduced lung damage and mucin Muc5ac expression in rats. Eur Respir J 2003;22:900–5.
- 20 Bradford M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248–54.

- 21 Cortijo J, Villagrasa V, Pons R, et al. Bronchodilator and anti-inflammatory activities of glaucine: in vitro studies in human airway smooth muscle and polymorphonuclear leukocytes. Br J Pharmacol 1999;127:1641–51.
- Crúmp CM, Williams JL, Stephens DJ, et al. Inhibition of the interaction between tyrosine-based motifs and the medium chain subunit of the AP-2 adaptor complex by specific tyrphostins. J Biol Chem 1998;273:28073–7.
 Kawaguchi M, Kokubu F, Matsukura S, et al. Induction of C-X-C chemokines.
- 23 Kawaguchi M, Kokubu F, Matsukura S, et al. Induction of C-X-C chemokines, growth-related oncogene α expression, and epithelial cell-derived neutrophilactivating protein-78 by ML-1 (interleukin-17F) involves activation of raf1mitogen-activated protein kinase kinase-extracellular signal-regulated kinase 1/2 pathway. J Pharmacol Exp Ther 2003;**307**:1213–20.
- 24 Chijiwa T, Mishima A, Hagiwara M, et al. Inhibition of forskolin-induced outgrowth and portein phosphorylation by a newly synthesized selective inhibitor of cyclic AMP-dependent protein kinase, N-[2-(pbromocinnamylamino)ethyl]-5-isoquinolinesulfonamide (H-89), of PC 12D Pheochromocytoma cells. J Biol Chem 1990;265:5267–72.
- 25 Pedrosa R, Gomes P, Soares-da-Silva P. Distinct signalling cascades downstream to G_sα coupled dopamine D₁-like NHE3 inhibition in rat and opossum renal epithelial cells. *Cell Physiol Biochem* 2004;14:91–100.
- opossum renal epithelial cells. *Cell Physiol Biochem* 2004;14:91–100.
 Sandberg M, Butt E, Nolte C, et al. Characterization of Sp-5,6-dichloro-1beta-D-ribofuranosylbenzimidazole-3',5'-monophosphorothioate (Sp-5,6-DCl-cBiMPS) as a potent and specific activator of cyclic-AMP-dependent protein kinase in cell extracts and intact cells. *Biochem J* 1991;279:521–7.
- 27 Basbaum C, Li D, Gensch E, et al. Mechanisms by which Gram-positive bacteria and tobacco smoke stimulate mucin induction through the epidermal growth factor receptor (EGFR). In: Chadwick DJ, Goode JA, eds. Mucus hypersecretion in respiratory disease. Chichester: John Wiley & Sons, 2002:171–6.
- 28 Hauck CR, Sieg DJ, Hsia DA, et al. Inhibition of focal adhesion kinase expression or activity disrupts epidermal growth factor-stimulated signaling promoting the migration of invasive human carcinoma cells. *Cancer Res* 2001;61:7079–90.
- 29 Wetzker R, Böhmer F-D. Transactivation joins multiple tracks to the ERK/ MAPK cascade. Nature Rev Mol Cell Biol 2003;4:651–7.
- 30 Piiper A, Gebhardt R, Kronenberger B, et al. Pertussis toxin inhibits cholecystokinin- and epidermal growth factor-induced mitogen-activated protein kinase activation by disinhibition of the cAMP signaling pathway and inhibition of c-Raf-1. Mol Pharmacol 2000;58:608–13.
- Baillie GS, MacKenzie SJ, McPhee I, et al. Sub-family selective actions in the ability of Erk2 MAP kinase to phosphorylate and regulate the activity of PDE4 cyclic AMP-specific phosphodiesterases. Br J Pharmacol 2000;131:811–9.
 Hatzelmann A, Schudt C. Anti-inflammatory and immunomodulatory
- 32 Hatzelmann A, Schudt C. Anti-inflammatory and immunomodulatory potential of the novel PDE4 inhibitor roflumilast in vitro. J Pharmacol Exp Ther 2001;297:267–79.

Correction

It has been brought to our attention that there is an error in figure 3 on page ii55 of the Pleural Disease Guideline available at www.brit-thoracic.org.uk/docs/PleuralDiseaseChestDrain/pdf. Below is a corrected diagram illustrating the "safe triangle" for a chest drain. The publishers apologise for this error.

