

## OCCASIONAL REVIEW

## Focus on antioxidant enzymes and antioxidant strategies in smoking related airway diseases

V L Kinnula

*Thorax* 2005;60:693–700. doi: 10.1136/thx.2004.037473

Cigarette smoke causes significant oxidant stress which is further enhanced by recruitment and activation of inflammatory cells to the lung. Polymorphisms in some detoxification enzymes are thought to increase the risk of developing chronic obstructive pulmonary disease (COPD), but the ultimate role of genetic variability in antioxidant and/or detoxification enzymes in COPD remains obscure. Some antioxidant enzymes are induced, but the extent of induction is insufficient to protect the lung/alveolar epithelium against cigarette smoke. Exogenous antioxidants such as vitamins do not seem to protect against cigarette smoke related lung injury. Glutathione related synthetic drugs such as N-acetylcysteine have shown some benefits, but they may have pro-oxidant side effects. Synthetic compounds with superoxide dismutase and catalase activities have shown promising results in animal models against a variety of oxidant exposures including cigarette smoke in the lung. These results are in agreement with studies highlighting the importance of alveolar antioxidant protection mechanisms in oxidant stress and their inducibility. These new drugs need to be tested in cigarette smoking related lung injury/inflammation since inflammation/oxidant stress can continue after discontinuation of smoking.

antioxidants will be able to prevent the progression of cigarette smoke mediated injury in the human lung.

The lung is a unique tissue compared with many other vital organs since it is directly exposed to high levels of oxygen. It therefore has to have efficient antioxidant mechanisms for the primary protection of the airways against ambient oxygen and exogenous oxidants. In cigarette smokers the oxidant burden is further increased for a number of reasons. Firstly, one puff of cigarette smoke contains millions of oxygen free radicals. Cigarette smoke contains high concentrations of nitric oxide (NO). Tar includes organic compounds such as quinones, which react with oxygen and increase the generation of superoxide radicals in intracellular redox cycling reactions. Free radicals activate inflammatory cells which, in turn, generate high levels of reactive oxygen (ROS) and nitrogen species (RNS) and other toxic metabolites. The most important of these mechanisms include myeloperoxidase, NADPH oxidase, and nitric oxide synthases (NOSs) (such as inducible NOS) and, in certain inflammatory conditions, eosinophil peroxidase. Typical features of smoker's lung are inflammation with the recruitment of neutrophils and macrophages into the airways, small airway obstruction/fibrosis, and destructive processes of the lung parenchyma.<sup>2–3</sup> Some studies have suggested that oxidative stress/lipid peroxidation is associated with pulmonary airway narrowing in the general population.<sup>4–5</sup> Many studies have also shown that there is a significant oxidant burden both in the lung and systemic circulation in COPD.<sup>6–11</sup> The airways of patients with COPD, especially during disease exacerbations, contain high levels of activated neutrophils, activated macrophages, and also eosinophils, representing an even higher oxidant burden during the active phase of this disease.<sup>3–12</sup> A number of studies have detected increased levels of oxidant markers in COPD lung tissue as well as in sputum, exhaled air, and exhaled breath condensate. These markers include

The most important non-malignant lung disease caused by cigarette smoke is chronic obstructive lung disease (COPD), a globally escalating problem that mirrors the extent of cigarette smoking in the community. In addition to COPD, cigarette smoking is associated with respiratory bronchiolitis interstitial pneumonia (RB-ILD)/desquamative interstitial pneumonia (DIP), a disease entity classified as a histopathological group of interstitial pneumonias.<sup>1</sup> Both disease groups are underdiagnosed; this is a serious problem, especially for COPD because of its high prevalence, progressive nature, and cost to healthcare systems throughout the world. Smoking cessation is the only therapeutic intervention that has been shown to improve the prognosis of COPD with reduction of disease progression. This review will concentrate on the importance of the oxidant/antioxidant balance and the major antioxidant enzymes and detoxification mechanisms in smoker's lung and COPD, and the possibilities that new synthetic

**Abbreviations:** COPD, chronic obstructive pulmonary disease; CuZnSOD, copper zinc SOD; ECSOD, extracellular SOD; ELF, epithelial lining fluid; GCL, glutamate cysteine ligase; GPX, glutathione peroxidase; GR, glutathione reductase; GRX, glutaredoxin; GSH, glutathione; GST, glutathione-S-transferase;  $\gamma$ GT,  $\gamma$ -glutamyl transpeptidase; HO-1, heme oxygenase-1; MnSOD, manganese SOD; MRP, multidrug resistance protein; NAC, N-acetylcysteine; NO, nitric oxide; PRX, peroxiredoxin; ROS, reactive oxygen species; RNS, reactive nitrogen species; SOD, superoxide dismutase; TRX, thioredoxin

Correspondence to:  
Professor V L Kinnula,  
University of Helsinki,  
Department of Medicine,  
Pulmonary Division, P O  
Box 22, Haartmaninkatu  
4, Helsinki, FI-00014,  
Finland; vuokko.kinnula@  
helsinki.fi

Received  
25 November 2004  
Accepted  
17 February 2005

increased levels of myeloperoxidase, the neutrophil markers lipocalin and lactoferrin, and markers of lipid peroxidation such as 4-hydroxy-2-nonenal (4-HNE) in the tissue and sputum, increased levels of hydrogen peroxide, and markers of lipid peroxidation products (thiobarbituric acid reactive substances) in the exhaled air/breath condensate in COPD. These studies have been extensively reviewed previously.<sup>9 13–16</sup> Most studies have been conducted on chronic smokers. However, acute exposure to cigarette smoke has also been shown to lead to an immediate transient oxidant burden, although unchanged and even decreased levels of exhaled NO have also been reported (reviewed by van der Vaart<sup>8</sup>). Overall, there is a significant ROS/RNS production in COPD that apparently overwhelms lung antioxidant protection in this disease.

In contrast to the oxidant producing systems, relatively little is known about the primary antioxidant defence of human lung, and even less is known about this system in smokers. Furthermore, alterations in the primary antioxidant defence as well as the effects of exogenous antioxidants in COPD have been less extensively investigated. There are many questions that still remain unanswered, such as the importance of individual variability in the primary antioxidant defence and the risk for COPD, the significance of the induction of the primary antioxidant mechanisms in smoke related inflammation, the importance of specific lung cell types in lung protection, the effects of the ultrastructural localisation of these enzyme proteins and/or small molecular weight antioxidants in smoker's lung, and also whether antioxidant enzymes have any effects on the inflammatory process itself such as the survival of inflammatory cells or apoptosis. The balance between oxidants and antioxidants is complex and the interactions between these pathways have received little attention. However, the significance of the primary antioxidant defence mechanisms in the development of COPD as well as finding ways to improve lung antioxidant protection, especially in areas where the oxidant stress is highest, may be major questions in predicting COPD progression.

### ANTIOXIDANT DEFENCE OF HUMAN LUNG

The lung has an efficient protection capacity against exogenous oxygen and ROS. Studies on animal models have shown that mucins, which are cysteine rich glycoproteins present in the epithelial lining fluid, provide significant protection against oxidants, and that the levels of these glycoproteins are increased by oxidants.<sup>17 18</sup> Epithelial lining fluid contains a high glutathione (GSH) content,<sup>19</sup> a molecule that also has been considered as one of the primary antioxidants in the human lung. In addition to these compounds, lung tissue contains several proteins capable of binding free iron and other metals (albumin, transferrin, ferritin, ceruloplasmin, lactoferrin, metallothionein). These proteins may have fundamental roles in protecting lung tissue against oxidant attack since metals are potent generators of free radicals in the Fenton and superoxide derived Haber-Weiss reactions.<sup>20</sup> Human airways contain both lipid and water soluble vitamins, but their relative roles in the airways are poorly understood. One typical feature of the antioxidant enzymes and closely related thiol proteins of human lung is their highly cell specific arrangement and compartmentalisation, in that they are localised to different cell types, cell organelles (one example being mitochondria), and also different extracellular compartments.

### Superoxide dismutases

Superoxide dismutases (SODs) are the only enzyme family with activity against superoxide radicals. These enzymes include manganese SOD (MnSOD), copper zinc SOD

(CuZnSOD), and extracellular SOD (ECSOD).<sup>21–23</sup> All of them are widely expressed in human lung.<sup>24</sup> CuZnSOD, which is mainly a cytosolic enzyme, is evenly distributed intracellularly but is also found in the nucleus and lysosomes.<sup>25</sup> It is expressed in the bronchial epithelium of human lung,<sup>26</sup> and in the alveolar epithelium, mesenchymal cells, fibroblasts and vascular endothelial cells in rat lung.<sup>27</sup> MnSOD is localised to the mitochondria, a critical cell organelle both for cellular energy metabolism and cell survival. MnSOD is mainly expressed in alveolar type II epithelial cells and alveolar macrophages in the human lung.<sup>26 28</sup> ECSOD is primarily localised to the extracellular matrix, especially in areas containing high amounts of type I collagen fibres and around pulmonary and systemic vessels.<sup>29</sup> It has also been detected in the bronchial epithelium, alveolar epithelium, and alveolar macrophages. Overall, CuZnSOD and MnSOD are generally thought to act as bulk scavengers of superoxide radicals. However, the relatively high ECSOD level in the lung with its specific binding to the extracellular matrix components may represent a fundamental component of lung matrix protection. All SODs also have functional polymorphisms with potential consequences to the individual variability for the risk of lung diseases.<sup>30</sup> SODs are also induced by cytokines. In this respect, MnSOD is unique since it is highly responsive to multiple extracellular stimuli including cytokines, oxidants and radiation, and this induction is thought to have physiological significance, especially in the protection of alveolar epithelium.<sup>24</sup>

### Destruction of hydrogen peroxide

The major enzymes associated with hydrogen peroxide decomposition include catalase and enzymes associated with GSH metabolism. The latter group of enzymes includes glutathione peroxidases (GPXs), glutathione reductase, glutamate cysteine ligase (GCL)—that is, gammaglutamyl cysteine synthase ( $\gamma$ GCS) and glutathione synthase. Catalase, a peroxisomal hydrogen peroxide consuming enzyme, is expressed intracellularly mainly in alveolar macrophages and neutrophils.<sup>31 32</sup> Due to the high level of GSH (a tripeptide containing one cysteine amino acid—that is, a thiol group)<sup>19 33</sup> in the epithelial lining fluid (ELF), enzymes associated with GSH synthesis are thought to have central roles in lung protection. In contrast to catalase, GPXs, a selenium containing family of enzymes, can break down not only hydrogen peroxide but also lipid peroxides. Five GPX gene products have been characterised: GPX1 is the classic GPX; GPX2 is gastrointestinal; GPX3 is the plasma GPX that has also been detected in the epithelial lining fluid of human lung; GPX4 is the phospholipid hydroperoxide GPX; and GPX5 is a secreted protein synthesised by the caput epididymis. Of these, GPX3 has been most widely investigated in human lung.<sup>34</sup> “Oxidised GSH” (that is, glutathione disulphide (GSSG)) is reduced back to GSH by glutathione reductase; the level of GSSG in normal tissues is very low (below 1%) due to the immediate reaction by glutathione reductase. Another important pathway for GSH replenishment in the lung is, however, its synthesis by GCL and glutathione synthase. The rate limiting enzyme in GSH synthesis, GCL, has been suggested to play a fundamental role in the regulation of lung GSH homeostasis. Mammalian GCL is a heterodimer that contains two subunits—the catalytically active heavy subunit (GCLh) and a light subunit (GCLl) that modulates the affinity of the heavy subunit for substrates and inhibitors. Both subunits are localised to the cytosol and expressed especially in human bronchial epithelium and occasionally in alveolar macrophages.<sup>35 36</sup> The regulation of GCLh is still somewhat controversial but it does appear to be induced by oxidants and cytokines and downregulated by transforming growth factor  $\beta$  in vitro.<sup>37–39</sup>


GCL is cleaved during the apoptotic process of human cells,<sup>40-42</sup> but it is not known whether this cleavage also occurs in human lung and lung diseases.

In addition, disposal of hydrogen peroxide is closely associated with several thiol-containing proteins—namely, thioredoxins (TRX1 and TRX2), thioredoxin reductases (TRR), peroxiredoxins (PRX) (which are thioredoxin peroxidases), and glutaredoxins (GRX).<sup>43-46</sup> All these enzymes are considered to be powerful redox modulators; they also have antioxidant properties, protect cells against high oxygen tension, and participate in cell proliferation and survival. Two TRXs and TRRs have been characterised in human cells, existing in both cytosol and mitochondria. In human lung, TRX and TRR are expressed in bronchial and alveolar epithelium and macrophages.<sup>47</sup> Six different PRXs have been found in human cells, differing in their ultrastructural compartmentalisation. Experimental studies have revealed the importance of PRX VI in the protection of alveolar epithelium. Human lung expresses all PRXs in bronchial epithelium, alveolar epithelium, and macrophages.<sup>48</sup> PRX V has recently been found to function as a peroxynitrite reductase,<sup>49</sup> which means that it may function as a potential protective compound in the development of ROS/RNS mediated lung injury. GRXs (I-II) are also localised in both the cytosol and mitochondria.<sup>43</sup> GRX1 has recently been found especially in human alveolar macrophages and, to a lesser degree, in bronchial epithelium. It is downregulated by growth factors and possibly induced transiently by oxidant stress *in vitro*.<sup>50</sup> Thus, there are several redox modulatory thiol containing proteins in human lung that have recently been shown to participate not only in cell protection against oxidants but also in the regulation of cell growth and survival.

**Other detoxification pathways**

Glutathione-S-transferases (GSTs) and multidrug resistance proteins (MRPs) belong to a group of detoxification enzymes that require intracellular GSH for their catalytic activity. GSTs constitute a protein family with high genetic variability, having potential effects on the development of smoking related non-malignant and malignant diseases. Three

mammalian GST families—namely cytosolic, mitochondrial and microsomal—have been characterised.<sup>51</sup> The relative importance of GSTs and MRPs in human lung is still unknown. For example, there are nine related MRPs. MRP1, at least, is located in the epithelium at the intersection of many thiol-based protection mechanisms that modulate the extrusion of GSSG from the cells (reviewed by Kruh<sup>52</sup>). Both these families of detoxification enzymes are often considered to be important in malignant diseases and the resistance of cancer cells to oxidant generating drugs. However, they are also expressed in normal lung, mainly in the airways.<sup>53</sup> They protect cells against numerous oxidant generating compounds and drugs, including pollutants, quinones, hydroperoxides, and other end products of oxidative metabolism. MRP substrates include structurally diverse GSH, glucuronate and sulphate conjugates, but the precise mechanism by which GSH participates in MRP mediated efflux of the natural toxic products is currently unknown.<sup>52</sup> Gamma-glutamyltranspeptidase ( $\gamma$ GT) is a plasma membrane enzyme which is also expressed in lung epithelial cells. The released GSH/GSSG is not freely diffusible and needs first to be decomposed to the amino acids where  $\gamma$ GT decomposes the  $\gamma$ -glutamyl bond of GSH. Experimental studies have shown that  $\gamma$ GT is induced by oxidant stress,<sup>54</sup> and also that  $\gamma$ GT deficiency results in increased oxidative stress in normoxia.<sup>55</sup> Heme oxygenase-1 is a stress response protein with significant functions in cell protection and homeostasis; this is also an enzyme which can be induced by oxidants and cytokines.<sup>56</sup> Thus, in addition to the classical antioxidant enzymes and thiol proteins, human lung contains stress inducible enzymes and detoxification systems which seem to confer primary protection for the human airways. The reactions of the major antioxidant and detoxification enzymes are summarised in fig 1. Overall, there are numerous enzyme systems that participate in the scavenging of ROS in human lung with tight interaction; these enzymes are localised to specific cell types and organelles, mostly in bronchial epithelium, alveolar epithelial type II cells and alveolar macrophages. Some of the enzymes are also bound to the extracellular matrix, providing potential protection to the lung interstitium (fig 2).



**Figure 1** Oxidant generation by cigarette smoke and the major antioxidant defence systems in their disposal. O<sub>2</sub><sup>•-</sup>, superoxide; H<sub>2</sub>O<sub>2</sub>, hydrogen peroxide; •OH, hydroxyl radical; NO, nitric oxide; NOS, nitric oxide synthase; HOCl, hypochlorous acid; ONOO<sup>-</sup>, peroxynitrite; RNS, reactive nitrogen species; MMP, matrix metalloproteinase; TIMP, tissue inhibitor of matrix metalloproteinase. SOD, superoxide dismutase; CuZnSOD, copper zinc SOD; MnSOD, manganese SOD; ECSOD, extracellular SOD; GPX, glutathione peroxidase; GR, glutathione reductase; GSH, glutathione; GCL, glutamate cysteine ligase; GST, glutathione-S-transferase; GSSG, oxidised glutathione (glutathione disulphide); GS, glutathione synthase;  $\gamma$ GT,  $\gamma$ -glutamyltranspeptidase; MRP, multidrug resistance protein; TRX, thioredoxin; PRX, peroxiredoxin; GRX, glutaredoxin; CAT, catalase; HO-1, heme oxygenase-1.

## ANTIOXIDANT ENZYMES IN SMOKERS AND SMOKING RELATED LUNG DISEASES

### Polymorphisms of antioxidant and detoxification enzymes in COPD

Many antioxidant enzymes such as all three SODs exhibit functional polymorphisms which can impact on enzyme activity/localisation. No studies are available on the polymorphisms of MnSOD or CuZnSOD in the development/progression of COPD. Their polymorphisms appear to have no effect on asthma development.<sup>57</sup> In ECSOD, a single nucleotide substitution resulting in an amino acid change from Arg to Gly at 213 (in the middle of the positive charged sequence of the heparin binding domain of ECSOD) leads to increased circulating ECSOD activity (10-fold) and decreased tissue binding of ECSOD.<sup>58</sup> One preliminary report on this polymorphism suggests that ECSOD polymorphism might be an additional risk for the development of chronic lung disease.<sup>59</sup> However, it is likely that polymorphism of ECSOD, which is very rare, cannot alone explain the early development of common lung diseases like COPD. Several studies have suggested that polymorphisms of the GSH associated detoxification enzymes may be associated with COPD. In one study, three GST polymorphisms (GST T1, GST P1 and GST M1) but not hemeoxygenase-1 showed a significant association with the rapid decline in lung function parameters in smokers.<sup>60</sup> In some, but not all, studies there were also significant associations between GST M1 null genotypes, GST P1, microsomal epoxide hydrolase, and COPD.<sup>61–63</sup> Overall, it appears that GST genotypes are risk factors for a rapid decline or low function in smokers with mild to moderate airflow obstruction.<sup>64</sup>

### Regulation of antioxidant enzymes in smoker's lung, COPD, and smoking related desquamative interstitial pneumonia


A recent microarray study on the bronchial brushings assessed 44 antioxidant related genes in four categories (catalase, SOD/GSH metabolism/redox balance, and pentose phosphate cycle) in non-smokers and smokers (average smoking history 21 years, no lung disease).<sup>65</sup> There was some variation in the gene expression levels within the same individual, but a remarkable interindividual variability. Altogether, 16 of the 44 antioxidant related genes were increased in the epithelium of smokers compared with

non-smokers.<sup>65</sup> Neither SODs nor catalase were altered in the epithelium of smokers, while 33% (7 of 21) of the genes encoding enzymes of the GSH pathway were upregulated, the most prominent being GPX2, although others with minor induction included GCL (both subunits), glutathione reductase, GPX3, GSTs and—among the redox balance regulating enzymes—TRR 1. Another recent study that used microarray technology characterised the gene expression profile of lung tissue samples from healthy subjects and advanced emphysema.<sup>66</sup> A typical feature of severe emphysema, both in “usual emphysema” and alpha-1-antitrypsin deficiency, was a global decrease in gene expression but an increased expression of the transcripts encoding proteins involved in inflammation and proteolysis. The major finding among the classical antioxidant enzymes was a significant decline in GPX3. Overall, GSH associated enzymes represent a group of enzymes that can be induced by cigarette smoke in human lung but may probably also decline during the progression of COPD.

MnSOD is significantly induced by cigarette smoke in rat lungs,<sup>67</sup> which is consistent with the high inducibility of MnSOD by cytokines and oxidant exposure *in vivo*.<sup>24</sup> Many studies have, however, also observed that the induction is transient, and that antioxidant enzymes may also be down-regulated during severe and/or chronic oxidant exposure.<sup>68</sup> Superoxide dismutase and catalase activities have been reported to be increased in human alveolar macrophages in chronic smokers,<sup>69</sup> whereas several antioxidant enzymes such as CuZnSOD, GST, and GPX seem to decline in elderly smokers with long smoking histories.<sup>70</sup> A recent human study evaluated MnSOD in smokers and in subjects with mild/moderate COPD and found MnSOD to be increased especially in the alveolar pneumocyte II cells of smokers and patients with COPD.<sup>71</sup> In conclusion, it seems that MnSOD can be induced in smoker's lung, especially in the alveolar epithelium, and that this is probably a consequence of cytokine/oxidant mediated induction. Furthermore, it can be viewed as one indicator of the oxidant burden, especially at the alveolar level of smokers.

Cigarette smoke causes depletion of intracellular GSH in cultured airway epithelial cells with a transient decrease of GPX and glucose-6-phosphatase activities,<sup>72</sup> but also an increase in GPX 3 expression.<sup>73</sup> Some *in vitro* studies also suggest that, after initial GSH depletion, there is an increase in the GSH levels apparently due to GCL induction (reviewed by Rahman and MacNee<sup>74</sup>). In general, the results obtained from various animal species and cell cultures are, however, variable, which emphasises the importance of not generalising results from *in vitro* or animal experiments to human disease states. In humans, these enzymes have been investigated both after acute exposure to smoke and in chronic smokers. Acute smoking, which evidently causes remarkable oxidant stress, does not increase GSH levels in ELF, but several studies have shown that GSH levels in the BAL fluid and ELF of chronic smokers are raised.<sup>74–76</sup> On the other hand, GSH is not increased intracellularly—that is, in the BAL fluid cells (mainly alveolar macrophages) of smokers or patients with COPD; thus, extracellular GSH clearly does not reflect intracellular antioxidant enzymes in the lungs of smokers.<sup>76</sup> The regulation of ELF GSH levels is still poorly understood; it may be related both to the epithelial injury and also to complex mechanisms of GSH transport and synthesis.

Several studies indicate that GPX 3 is increased during the oxidative stress of human airways and is increased both in the bronchial epithelium and ELF of smokers<sup>77</sup> and in asthmatic airways.<sup>34</sup> The results with GCL are variable, partly due to the many experimental designs used in animal models and cells.<sup>38</sup> Two studies on smokers and/or patients with COPD have shown increased gene expression of GCL in the



**Figure 2** Localisation of major antioxidant defence systems in human lung. ELF, epithelial lining fluid. For abbreviations of antioxidant enzymes see fig 1.

### Box 1 Activity and/or expression of antioxidant enzymes, redox modulatory proteins, and detoxification enzymes in smoking related non-malignant lung diseases

#### Manganese superoxide dismutase (MnSOD)

- Unchanged gene expression in the airway epithelium of smokers<sup>65</sup>
- Increased activity and immunoreactivity in the alveolar epithelium of smokers<sup>71</sup>
- Increased immunoreactivity in the inflammatory areas of interstitial lung disorders (DIP)<sup>28</sup>

#### CuZn superoxide dismutase (CuZnSOD)

- Unchanged gene expression, protein and activity in the airway epithelium and lung homogenates of smokers<sup>65 71</sup>
- Decreased activity in the alveolar macrophages of elderly smokers<sup>70</sup>

#### Extracellular superoxide dismutase (ECSOD)

- Unchanged gene expression in the airway epithelium of smokers<sup>65</sup>
- Unchanged immunoreactivity in the lung homogenates of smokers<sup>71</sup>

#### Catalase

- Unchanged gene expression in the airway epithelium of smokers<sup>65</sup>
- Increased catalase activity in the alveolar macrophages of chronic smokers<sup>69</sup>

#### Enzymes associated with glutathione metabolism

- Increased glutamate cysteine ligase (GCL) (gene expression) or decreased (immunoreactive protein) in airway epithelium of smokers<sup>65 78 79</sup>
- Increased GCL immunoreactivity in metaplastic and dysplastic epithelium of smokers<sup>79 80</sup>
- Increased GCL immunoreactivity in the inflammatory areas of DIP<sup>35</sup>
- Increased GPX2, GPX3 and glutathione reductase gene expressions in the bronchial epithelium of smokers<sup>65</sup>
- Increased GPX3 gene expression in the bronchial epithelium of smokers<sup>113</sup>
- Decreased GPX3 gene expression in severe emphysema<sup>66</sup>
- Increased GSTs (gene expressions) in airway epithelium of smokers<sup>65</sup>
- Decreased GST and GPX activities in the alveolar macrophages of elderly smokers<sup>70</sup>

#### Thioredoxin and thioredoxin peroxidases (peroxiredoxins)

- Unchanged (thioredoxin gene expression) in the airways of smokers<sup>65</sup>
- Increased (thioredoxin reductase gene expression) in the airways of smokers<sup>65</sup>
- Increased thioredoxin immunoreactive protein in the inflammatory areas of DIP<sup>47</sup>

#### Heme oxygenase-1

- Increased immunoreactivity in the alveolar walls of smokers<sup>83</sup>
- Decreased immunoreactivity in the lung of severe COPD<sup>82</sup>
- Increased immunoreactivity in the inflammatory areas of DIP<sup>85</sup>

bronchial epithelium,<sup>65 78</sup> while others have reported lowered GCL immunoreactive protein in the bronchial epithelium and alveolar macrophages of subjects with smoker's lung and COPD.<sup>79 76</sup> On the other hand, the extent of GCL immunoreactivity was increased in the metaplastic/dysplastic bronchial lesions of COPD,<sup>79 80</sup> as was also the case with some other protective enzymes such as TRX, pointing to the importance of oxidant stress in lung/bronchial epithelial cell carcinogenesis.<sup>80</sup> The variable results in GCL expression in smoker's lung and COPD may therefore be related to the cell type present in the epithelium (either normal or metaplastic), and to differences in the smoking history and disease severity or its activity. Induction of antioxidant/detoxification enzymes protects not only normal bronchial epithelium against oxidative insults, but possibly also against cell death/apoptosis, which then might maintain cell proliferation even though the cells contain oxidised DNA alterations.

In addition to the classical antioxidant enzymes, GSTs and heme oxygenase-1 have been investigated in the airways of smokers since it is known that both these enzymes can protect lung cells against oxidants. In one study on a

relatively small number of patients, smoking had no effect on GST levels,<sup>81</sup> while the transcripts of several GSTs were increased in the bronchial epithelium of smokers without COPD.<sup>65</sup> Heme oxygenase-1, an enzyme that is highly inducible by oxidants, is increased in the alveolar macrophages of smokers but decreased in severe COPD.<sup>82 83</sup> Moreover, heme oxygenase-1 can be induced, especially in the alveolar macrophages of acute asthmatics.<sup>84</sup> There are remarkable differences in the individual variability of the detoxification enzymes (GSTs) and heme oxygenase-1 in human airways, but several studies suggest their induction during the oxidative stress caused by cigarette smoke. Recent results also suggest that, during the ongoing progression of COPD, there is some decline in the efficacy of protective mechanisms in human airways.

Cigarette smoking is also associated with airway/interstitial lung diseases, RB-ILD/DIP as smokers account for 90% of all cases. Several antioxidant enzymes and related mechanisms have been found to be increased in these diseases, probably due to oxidant stress and inflammation. Enzymes that have been found to be increased in these disorders include MnSOD, GCL, TRX, and heme oxygenase-1 (box 1).<sup>28 35 47 85</sup>

### Antioxidants and antioxidant enzymes of inflammatory cells and lung inflammation

A typical feature of cigarette smoking related lung diseases such as COPD is the recruitment of macrophages and neutrophils to the lung which increases oxidant burden also in the lung interstitium. Exogenous antioxidants and new synthetic molecules with SOD-like activity decrease lung inflammation in several experimental models of lung injury including cigarette smoke (see next section). However, little is known about the effects of antioxidant enzymes of inflammatory cells themselves on the survival/apoptosis or persistence of inflammation in human lung. Antioxidant enzymes of the inflammatory cells can have a pivotal role in the inflammatory reaction of the lung by offering significant protection against oxidants both intracellularly and extracellularly. They can prolong the survival of inflammatory cells, decrease their apoptosis, but some of them are also released into the interstitium with consequent protection of the extracellular matrix (GSH and extracellular antioxidant enzymes such as ECSOD). Alveolar macrophages have a long life span (several months) and there are suggestions that the survival of macrophages is prolonged in the lungs of smokers and subjects with COPD.<sup>86</sup> Macrophages express several antioxidant enzymes and related compounds including at least MnSOD, ECSOD, catalase, enzymes associated with the TRX-PRX system, and especially GRX and HO-1.<sup>26 32 47 48 50 87 88</sup> MnSOD and HO-1 appear to be increased in the alveolar macrophages of subjects with smoker's lung, mild COPD, and acute asthmatic inflammation.<sup>83 84</sup> In contrast to alveolar macrophages, the life span of neutrophils is very short. The physiological half life of neutrophils in circulation is only 6 hours, and there is evidence that neutrophil apoptosis is further enhanced by oxidants.<sup>89</sup> Very few studies have investigated antioxidant mechanisms of human neutrophils. These studies have shown that typical features of the antioxidant profile of neutrophils are their high catalase content, low levels of GSH-dependent enzymes, and nearly non-detectable levels of TRX and HO-1 by immunohistochemistry.<sup>32 87 90</sup> Antioxidant enzymes in the inflammatory cells have been shown to decrease during the progression of several systemic diseases (reviewed by Kinnula<sup>90</sup>), but whether this also occurs in smoking related human lung diseases or has clinical significance is unclear.

### POTENTIAL EXOGENOUS ANTIOXIDANTS AND ANTIOXIDANT MIMETICS IN SMOKING RELATED LUNG DISEASES

Several studies have investigated the systemic oxidant stress which occurs in COPD such as observations of a decline in lung and plasma levels of antioxidants, but variable results have been obtained by administering exogenous antioxidants (mainly vitamins such as vitamin E) to patients with COPD.<sup>91</sup> The disappointing results can be explained, at least in part, by the complex cell specific antioxidant systems present in human lung and the fact that vitamins taken orally do not reach the critical areas of the lung where the oxidant stress is highest. Since GSH is considered as one of the most important antioxidants in human lung, numerous studies have tried to improve GSH homeostasis in oxidant mediated lung injury. However, GSH has a poor penetration capability. Neither inhalation nor oral ingestion of GSH have improved the GSH balance in oxidative stress; in fact, it has caused side effects such as bronchoconstriction.<sup>74</sup> GSH is readily transported to the extracellular space by several mechanisms (including GSTs and MRPs), but is not directly transported from the extracellular milieu into the cells. N-acetylcysteine (NAC) is a widely used drug that might improve GSH homeostasis. In animal models NAC has attenuated elastase induced emphysema.<sup>92</sup> It is relatively safe also in humans and

has already been tested in a variety of conditions and treatment schedules, being used orally, by inhalation, and intravenously (reviewed by Rahman<sup>74</sup>). On the other hand, NAC may also have some pro-oxidant effects at the cellular level. In rats, NAC attenuated pulmonary inflammation caused by various antigens; it also decreased neutrophil recruitment and P-selectin upregulation.<sup>93</sup> NAC also decreased H<sub>2</sub>O<sub>2</sub> levels in the exhaled air in COPD and COPD exacerbations.<sup>94-97</sup> On the other hand, it had relatively minor effects on lung GSH homeostasis.<sup>98-100</sup> Both controlled studies and a systematic review with NAC in COPD have shown a small reduction in exacerbations.<sup>101 102</sup> Other compounds with NAC-like activity have been tested in many laboratories including N-acetylcysteinyl, GSH esters and GSH precursors, but there are no studies on these compounds in human smokers with COPD. Antioxidants that improve GSH homeostasis have potential possibilities in protecting the lung during oxidative stress, but these need to be investigated in future studies in combination with other new drugs with antioxidant activity.

Another group of antioxidants that have had beneficial results in several animal models are SODs. In the initial experiments, exogenous CuZnSOD or recombinant MnSOD were administered by inhalation or instillation, often encapsulated to liposomes or in combination with other antioxidant enzymes such as catalase (reviewed by Kinnula and Crapo<sup>24</sup>). However, antioxidant enzymes have several drawbacks such as high immunogenicity. In another approach, synthetic SOD mimetics have been developed to avoid the problems associated with endogenous antioxidant enzymes and to improve their penetration capability and efficacy. They have been shown to have at least four properties—namely, scavenging of superoxide, hydrogen peroxide, peroxynitrite, and lipid peroxides. These agents include salen compounds,<sup>103</sup> macrocyclics,<sup>104</sup> and metalloporphyrins.<sup>105</sup> Salen compounds (such as EUKs) can attenuate oxidant injury in numerous organs including lung injury in ARDS.<sup>106</sup> The macrocyclic agents (such as M40403) have been reported to decrease lung inflammation and pleurisy by inhibiting ROS/RNS formation, neutrophil infiltration, and release of pro-inflammatory cytokines.<sup>104</sup> Metalloporphyrins such as manganese (III) meso-tetrakis (4-benzoic acid) porphyrin (MnTBAP) have been tested in a number of models of lung injury including paraquat and carrageenan-induced pleurisy, exposure to bleomycin, asbestos and radiation, in airway inflammation, and models of bronchopulmonary dysplasia and lung haemorrhage.<sup>107-111</sup> In all these models, these synthetic compounds have had very positive results including a significant effect on lung protection. MnTBAP has also been found to reduce smoking related lung injury in rat lungs.<sup>112</sup> In that study, one of the compounds, AEOL 10150, significantly decreased the BAL cell number within the first weeks of exposure. Importantly, squamous metaplasia following 8 weeks of tobacco smoke exposure was only 2% in animals treated with the drug compared with 12% in animals exposed to cigarette smoke without AEOL 10150. This study clearly showed that new synthetic SOD mimetics can significantly reduce the adverse effects of exposure to tobacco smoke and suggest that they may even have beneficial effects in preventing malignant conversion of epithelial cells. These new synthetic antioxidants may prove also to have significant importance in human lung inflammation caused by cigarette smoke.

### CONCLUSIONS

The lung is exposed to high levels of oxidants, not only in normal conditions but especially in inflammatory states such as cigarette smoking related disorders. The lung has an efficient but complicated antioxidant defence system, where

antioxidant enzymes and related detoxification enzymes have a cell specific expression and compartmentalisation and variable induction capacities. Cigarette smoke leads to an induction of some of these enzymes, especially in the bronchial and alveolar epithelium, possibly reflecting high oxidant burden in these locations. Although there is a systemic oxidant stress in COPD, one key determinant in preventing COPD and emphysema progression might be provision of treatment that is specifically directed to the target areas where the oxidant stress is high and the cells are especially vulnerable to injury.

Funded by Helsinki University Hospital, Sigrid Juselius Foundation, Finnish Antituberculosis Association Foundation, and the Cancer Society of Finland.

## REFERENCES

- American Thoracic Society and European Respiratory Society.** American Thoracic Society/European Respiratory Society: International multidisciplinary consensus classification of the idiopathic interstitial pneumonias. *Am J Respir Crit Care Med* 2002;**165**:277–304.
- Gan WQ, Man SF, Senthilselvan A, et al.** Association between chronic obstructive pulmonary disease and systemic inflammation: a systematic review and a meta-analysis. *Thorax* 2004;**59**:574–80.
- Saetta M, Baraldo S, Zuin R.** Neutrophil chemokines in severe exacerbations of chronic obstructive pulmonary disease: fatal chemo-attraction? *Am J Respir Crit Care Med* 2003;**168**:911–3.
- Schunemann HJ, Muti P, Freudenheim JL, et al.** Oxidative stress and lung function. *Am J Epidemiol* 1997;**146**:939–48.
- Schunemann HJ, Grant BJ, Freudenheim JL, et al.** The relation of serum levels of antioxidant vitamins C and E, retinol and carotenoids with pulmonary function in the general population. *Am J Respir Crit Care Med* 2001;**163**:1246–55.
- Barnes PJ, Shapiro SD, Pauwels RA.** Chronic obstructive pulmonary disease: molecular and cellular mechanisms. *Eur Respir J* 2003;**22**:672–88.
- Rahman I.** Oxidative stress and gene transcription in asthma and chronic obstructive pulmonary disease: antioxidant therapeutic targets. *Curr Drug Targets Inflamm Allergy* 2002;**1**:291–315.
- van der Vaart, Postma DS, Timens W, et al.** Acute effects of cigarette smoke on inflammation and oxidative stress: a review. *Thorax* 2004;**59**:713–21.
- Langen RC, Korn SH, Wouters EF.** ROS in the local and systemic pathogenesis of COPD. *Free Radic Biol Med* 2003;**35**:226–35.
- MacNee W.** Oxidative stress and lung inflammation in airways disease. *Eur J Pharmacol* 2001;**429**:195–207.
- Repine JE, Bast A, Lankhorst I.** Oxidative stress in chronic obstructive pulmonary disease. Oxidative Stress Study Group. *Am J Respir Crit Care Med* 1997;**156**:341–57.
- Baraldo S, Turato G, Badin C, et al.** Neutrophilic infiltration within the airway smooth muscle in patients with COPD. *Thorax* 2004;**59**:308–12.
- Rahman I, MacNee W.** Role of oxidants/antioxidants in smoking-induced lung diseases. *Free Radic Biol Med* 1996;**21**:669–81.
- Rahman I, Skwarska E, MacNee W.** Attenuation of oxidant/antioxidant imbalance during treatment of exacerbations of chronic obstructive pulmonary disease. *Thorax* 1997;**52**:565–8.
- MacNee W.** Oxidants/antioxidants and COPD. *Chest* 2000;**117**:303–17S.
- Kharitonov SA, Barnes PJ.** Exhaled markers of pulmonary disease. *Am J Respir Crit Care Med* 2001;**163**:1693–722.
- Adler KB, Li Y.** Airway epithelium and mucus: intracellular signaling pathways for gene expression and secretion. *Am J Respir Cell Mol Biol* 2001;**25**:397–400.
- Cross CE, Halliwell B, Allen A.** Antioxidant protection: a function of tracheobronchial and gastrointestinal mucus. *Lancet* 1984;**1**:1328–30.
- Cantin AM, North SL, Hubbard RC, et al.** Normal alveolar epithelial lining fluid contains high levels of glutathione. *J Appl Physiol* 1987;**63**:152–7.
- Halliwell B, Gutteridge JM.** *Free radicals in biology and medicine*. Oxford: Clarendon, 1996.
- McCord JM, Fridovich I.** The utility of superoxide dismutase in studying free radical reactions. I. Radicals generated by the interaction of sulfite, dimethyl sulfoxide, and oxygen. *J Biol Chem* 1969;**244**:6056–63.
- Marklund SL.** Extracellular superoxide dismutase in human tissues and human cell lines. *J Clin Invest* 1984;**74**:1398–403.
- Oury TD, Day BJ, Crapo JD.** Extracellular superoxide dismutase: a regulator of nitric oxide bioavailability. *Lab Invest* 1996;**75**:617–36.
- Kinnula VL, Crapo JD.** Superoxide dismutases in the lung and human lung diseases. *Am J Respir Crit Care Med* 2003;**167**:1600–19.
- Slot JW, Geuze HJ, Freeman BA, et al.** Intracellular localization of the copper-zinc and manganese superoxide dismutases in rat liver parenchymal cells. *Lab Invest* 1986;**55**:363–71.
- Lakari E, Paakko P, Kinnula VL.** Manganese superoxide dismutase, but not CuZn superoxide dismutase, is highly expressed in the granulomas of pulmonary sarcoidosis and extrinsic allergic alveolitis. *Am J Respir Crit Care Med* 1998;**158**:589–96.
- Coursin DB, Cihla HP, Oberley TD, et al.** Immunolocalization of antioxidant enzymes and isozymes of glutathione S-transferase in normal rat lung. *Am J Physiol* 1992;**263**:L679–91.
- Lakari E, Paakko P, Pietarinen-Runtti P, et al.** Manganese superoxide dismutase and catalase are coordinately expressed in the alveolar region in chronic interstitial pneumonias and granulomatous diseases of the lung. *Am J Respir Crit Care Med* 2000;**161**:615–21.
- Oury TD, Chang LY, Marklund SL, et al.** Immunocytochemical localization of extracellular superoxide dismutase in human lung. *Lab Invest* 1994;**70**:889–98.
- Zelko IN, Mariani TJ, Folz RJ.** Superoxide dismutase multigene family: a comparison of the CuZn-SOD (SOD1), Mn-SOD (SOD2), and EC-SOD (SOD3) gene structures, evolution, and expression. *Free Radic Biol Med* 2002;**33**:337–49.
- Carter AB, Tephly LA, Venkataraman S, et al.** High levels of catalase and glutathione peroxidase activity dampen H<sub>2</sub>O<sub>2</sub> signaling in human alveolar macrophages. *Am J Respir Cell Mol Biol* 2004;**31**:43–53.
- Pietarinen P, Raivio K, Devlin RB, et al.** Catalase and glutathione reductase protection of human alveolar macrophages during oxidant exposure in vitro. *Am J Respir Cell Mol Biol* 1995;**13**:434–41.
- Cantin AM, North SL, Fells GA, et al.** Oxidant-mediated epithelial cell injury in idiopathic pulmonary fibrosis. *J Clin Invest* 1987;**79**:1665–73.
- Comhair SA, Bhatena PR, Farver C, et al.** Extracellular glutathione peroxidase induction in asthmatic lungs: evidence for redox regulation of expression in human airway epithelial cells. *FASEB J* 2001;**15**:70–8.
- Tiitto LH, Peltoniemi MJ, Kaarteenaho-Wiik RL, et al.** Cell-specific regulation of gamma-glutamylcysteine synthetase in human interstitial lung diseases. *Hum Pathol* 2004;**35**:832–9.
- Soini Y, Napankangas U, Jarvinen K, et al.** Expression of gamma-glutamyl cysteine synthetase in nonsmall cell lung carcinoma. *Cancer* 2001;**92**:2911–9.
- Liu RM, Hu H, Robison TW, et al.** Increased gamma-glutamylcysteine synthetase and gamma-glutamyl transpeptidase activities enhance resistance of rat lung epithelial L2 cells to quinone toxicity. *Am J Respir Cell Mol Biol* 1996;**14**:192–7.
- Dickinson DA, Forman HJ.** Cellular glutathione and thiols metabolism. *Biochem Pharmacol* 2002;**64**:1019–26.
- Arsalane K, Dubois CM, Muanza T, et al.** Transforming growth factor- $\beta$ 1 is a potent inhibitor of glutathione synthesis in the lung epithelial cell line A549: transcriptional effect on the GSH rate-limiting enzyme gamma-glutamylcysteine synthetase. *Am J Respir Cell Mol Biol* 1997;**17**:599–607.
- Siitonen T, Alarukka P, Mantymaa P, et al.** Protection of acute myeloblastic leukemia cells against apoptotic cell death by high glutathione and gamma-glutamylcysteine synthetase levels during etoposide-induced oxidative stress. *Ann Oncol* 1999;**10**:1361–7.
- Franklin CC, Rosenfeld-Franklin ME, White C, et al.** TGF $\beta$ 1-induced suppression of glutathione peroxidase defenses in hepatocytes: caspase-dependent post-translational and caspase-independent transcriptional regulatory mechanisms. *FASEB J* 2003;**17**:1535–7.
- Boito D, Franklin CC, White CC, et al.** Glutamate-cysteine ligase attenuates TNF-induced mitochondrial injury and apoptosis. *Free Radic Biol Med* 2004;**37**:632–42.
- Holmgren A.** Antioxidant function of thioredoxin and glutaredoxin systems. *Antioxid Redox Signal* 2000;**2**:811–20.
- Powis G, Mustacich D, Coon A.** The role of the redox protein thioredoxin in cell growth and cancer. *Free Radic Biol Med* 2000;**29**:312–22.
- Rhee SG, Kang SW, Netto LE, et al.** A family of novel peroxidases, peroxiredoxins. *Biofactors* 1999;**10**:207–9.
- Gromer S, Urig S, Becker K.** The thioredoxin system—from science to clinic. *Med Res Rev* 2004;**24**:40–89.
- Tiitto L, Kaarteenaho-Wiik R, Sormunen R, et al.** Expression of the thioredoxin system in interstitial lung disease. *J Pathol* 2003;**201**:363–70.
- Kinnula VL, Lehtonen S, Kaarteenaho-Wiik R, et al.** Cell specific expression of peroxiredoxins in human lung and pulmonary sarcoidosis. *Thorax* 2002;**57**:157–64.
- Dubuisson M, Vander SD, Clippe A, et al.** Human peroxiredoxin 5 is a peroxynitrite reductase. *FEBS Lett* 2004;**571**:161–5.
- Peltoniemi M, Kaarteenaho-Wiik R, Saily M, et al.** Expression of glutaredoxin is highly cell specific in human lung and is decreased by transforming growth factor-beta in vitro and in interstitial lung diseases in vivo. *Hum Pathol* 2004;**35**:1000–7.
- Hayes JD, Flanagan JU, Jowsey IR.** Glutathione transferases. *Annu Rev Pharmacol Toxicol* 2005;**45**:51–88.
- Kruh GD, Belinsky MG.** The MRP family of drug efflux pumps. *Oncogene* 2003;**22**:7537–52.
- Anttila S, Hirvonen A, Vainio H, et al.** Immunohistochemical localization of glutathione S-transferases in human lung. *Cancer Res* 1993;**53**:5643–8.
- Kugelman A, Choy HA, Liu R, et al.**  $\gamma$ -Glutamyl transpeptidase is increased by oxidative stress in rat alveolar L2 epithelial cells. *Am J Respir Cell Mol Biol* 1994;**11**:586–92.
- Jean JC, Liu Y, Brown LA, et al.** Gamma-glutamyl transferase deficiency results in lung oxidant stress in normoxia. *Am J Physiol Lung Cell Mol Physiol* 2002;**283**:L766–76.
- Choi AM, Alam J.** Heme oxygenase-1: function, regulation, and implication of a novel stress-inducible protein in oxidant-induced lung injury. *Am J Respir Cell Mol Biol* 1996;**15**:9–19.
- Kinnula VL, Lehtonen S, Koistinen P, et al.** Two functional variants of the superoxide dismutase genes in Finnish families with asthma. *Thorax* 2004;**59**:116–9.
- Sandstrom J, Nilsson P, Karlsson K, et al.** 10-fold increase in human plasma extracellular superoxide dismutase content caused by a mutation in heparin-binding domain. *J Biol Chem* 1994;**269**:19163–6.

- 59 **Folz RJ**, Crapo JD, Peno-Green LA. Elevated levels of extracellular superoxide dismutase in chronic lung disease and characterization of genetic variants. *Chest* 1997;**111**:745.
- 60 **He JQ**, Ruan J, Connett JE, et al. Antioxidant gene polymorphisms and susceptibility to a rapid decline in lung function in smokers. *Am J Respir Crit Care Med* 2002;**166**:323–8.
- 61 **Smith CA**, Harrison DJ. Association between polymorphism in gene for microsomal epoxide hydrolase and susceptibility to emphysema. *Lancet* 1997;**350**:630–3.
- 62 **Cheng SL**, Yu CJ, Chen CJ, et al. Genetic polymorphism of epoxide hydrolase and glutathione S-transferase in COPD. *Eur Respir J* 2004;**23**:818–24.
- 63 **Ishii T**, Matsuse T, Teramoto S, et al. Glutathione S-transferase P1 (GSTP1) polymorphism in patients with chronic obstructive pulmonary disease. *Thorax* 1999;**54**:693–6.
- 64 **He JQ**, Connett JE, Anthonisen NR, et al. Glutathione S-transferase variants and their interaction with smoking on lung function. *Am J Respir Crit Care Med* 2004;**170**:388–94.
- 65 **Hackett NR**, Heguy A, Harvey BG, et al. Variability of antioxidant-related gene expression in the airway epithelium of cigarette smokers. *Am J Respir Cell Mol Biol* 2003;**29**:331–43.
- 66 **Golpon HA**, Coldren CD, Zamora MR, et al. Emphysema lung tissue gene expression profiling. *Am J Respir Cell Mol Biol* 2004;**31**:595–600.
- 67 **Gilks CB**, Price K, Wright JL, et al. Antioxidant gene expression in rat lung after exposure to cigarette smoke. *Am J Pathol* 1998;**152**:269–78.
- 68 **Clerch LB**. Post-transcriptional regulation of lung antioxidant enzyme gene expression. *Ann N Y Acad Sci* 2000;**899**:103–11.
- 69 **McCusker K**, Hoidal J. Selective increase of antioxidant enzyme activity in the alveolar macrophages from cigarette smokers and smoke-exposed hamsters. *Am Rev Respir Dis* 1990;**141**:678–82.
- 70 **Kondo T**, Tagami S, Yoshioka A, et al. Current smoking of elderly men reduces antioxidants in alveolar macrophages. *Am J Respir Crit Care Med* 1994;**149**:178–82.
- 71 **Harju T**, Kaarteenaho-Wiik R, Sirvio R, et al. Manganese superoxide dismutase is increased in the airways of smokers' lungs. *Eur Respir J* 2004;**24**:765–71.
- 72 **Rahman I**, Li XY, Donaldson K, et al. Glutathione homeostasis in alveolar epithelial cells in vitro and lung in vivo under oxidative stress. *Am J Physiol* 1995;**269**:L285–92.
- 73 **Rahman I**, Bel A, Mulier B, et al. Differential regulation of glutathione by oxidants and dexamethasone in alveolar epithelial cells. *Am J Physiol* 1998;**275**:L80–6.
- 74 **Rahman I**, MacNee W. Lung glutathione and oxidative stress: implications in cigarette smoke-induced airway disease. *Am J Physiol* 1999;**277**:L1067–88.
- 75 **Linden M**, Hakansson L, Ohlsson K, et al. Glutathione in bronchoalveolar lavage fluid from smokers is related to humoral markers of inflammatory cell activity. *Inflammation* 1989;**13**:651–8.
- 76 **Neurohr C**, Lenz AG, Ding I, et al. Glutamate-cysteine ligase modulatory subunit in BAL alveolar macrophages of healthy smokers. *Eur Respir J* 2003;**22**:82–7.
- 77 **Comhair SA**, Lewis MJ, Bhatena PR, et al. Increased glutathione and glutathione peroxidase in lungs of individuals with chronic beryllium disease. *Am J Respir Crit Care Med* 1999;**159**:1824–9.
- 78 **Rahman I**, van Schadewijk AA, Hiemstra PS, et al. Localization of gamma-glutamylcysteine synthetase messenger rna expression in lungs of smokers and patients with chronic obstructive pulmonary disease. *Free Radic Biol Med* 2000;**28**:920–5.
- 79 **Harju T**, Kaarteenaho-Wiik R, Soini Y, et al. Diminished immunoreactivity of gamma-glutamylcysteine synthetase in the airways of smokers' lung. *Am J Respir Crit Care Med* 2002;**166**:754–9.
- 80 **Soini Y**, Kaarteenaho-Wiik R, Paakko P, et al. Expression of antioxidant enzymes in bronchial metaplastic and dysplastic epithelium. *Lung Cancer* 2003;**39**:15–22.
- 81 **Nakajima T**, Elovaara E, Anttila S, et al. Expression and polymorphism of glutathione S-transferase in human lungs: risk factors in smoking-related lung cancer. *Carcinogenesis* 1995;**16**:707–11.
- 82 **Maestrelli P**, Paska C, Saetta M, et al. Decreased haem oxygenase-1 and increased inducible nitric oxide synthase in the lung of severe COPD patients. *Eur Respir J* 2003;**21**:971–6.
- 83 **Maestrelli P**, El Messlemanni AH, De Fina O, et al. Increased expression of heme oxygenase (HO)-1 in alveolar spaces and HO-2 in alveolar walls of smokers. *Am J Respir Crit Care Med* 2001;**164**:1508–13.
- 84 **Harju T**, Soini Y, Paakko R, et al. Up-regulation of heme oxygenase-I in alveolar macrophages of newly diagnosed asthmatics. *Respir Med* 2002;**96**:418–23.
- 85 **Lakari E**, Pylkas P, Pietarinen-Runtti P, et al. Expression and regulation of hemeoxygenase 1 in healthy human lung and interstitial lung disorders. *Hum Pathol* 2001;**32**:1257–63.
- 86 **Barnes PJ**. Alveolar macrophages as orchestrators of COPD. *J COPD* 2004;**1**:59–70.
- 87 **Pietarinen-Runtti P**, Lakari E, Raivio KO, et al. Expression of antioxidant enzymes in human inflammatory cells. *Am J Physiol Cell Physiol* 2000;**278**:C118–25.
- 88 **Fattman CL**, Schaefer LM, Oury TD. Extracellular superoxide dismutase in biology and medicine. *Free Radic Biol Med* 2003;**35**:236–56.
- 89 **Haslett C**. Granulocyte apoptosis and its role in the resolution and control of lung inflammation. *Am J Respir Crit Care Med* 1999;**160**:S5–11.
- 90 **Kinnula VL**, Soini Y, Kvist-Makela K, et al. Antioxidant defense mechanisms in human neutrophils. *Antioxid Redox Signal* 2002;**4**:27–34.
- 91 **Boots AW**, Haenen GR, Bast A. Oxidant metabolism in chronic obstructive pulmonary disease. *Eur Respir J Suppl* 2003;**46**:14–27.
- 92 **Rubio ML**, Martin-Mosquera MC, Ortega M, et al. Oral N-acetylcysteine attenuates elastase-induced pulmonary emphysema in rats. *Chest* 2004;**125**:1500–6.
- 93 **Serrano-Mollar A**, Closa D, Cortijo J, et al. P-selectin upregulation in bleomycin induced lung injury in rats: effect of N-acetyl-L-cysteine. *Thorax* 2002;**57**:629–34.
- 94 **Kasielski M**, Nowak D. Long-term administration of N-acetylcysteine decreases hydrogen peroxide exhalation in subjects with chronic obstructive pulmonary disease. *Respir Med* 2001;**95**:448–56.
- 95 **Pela R**, Calcagni AM, Subiaco S, et al. N-acetylcysteine reduces the exacerbation rate in patients with moderate to severe COPD. *Respiration* 1999;**66**:495–500.
- 96 **Gerrits CM**, Herings RM, Leufkens HG, et al. N-acetylcysteine reduces the risk of re-hospitalisation among patients with chronic obstructive pulmonary disease. *Eur Respir J* 2003;**21**:795–8.
- 97 **Decramer M**, Dekhuijzen PN, Troosters T, et al. The Bronchitis Randomized On NAC Cost-Utility Study (BRONCUS): hypothesis and design. BRONCUS Trial Committee. *Eur Respir J* 2001;**17**:329–36.
- 98 **Behr J**, Degenkolb B, Krombach F, et al. Intracellular glutathione and bronchoalveolar cells in fibrosing alveolitis: effects of N-acetylcysteine. *Eur Respir J* 2002;**19**:906–11.
- 99 **Dueholm M**, Nielsen C, Thorshauge H, et al. N-acetylcysteine by metered dose inhaler in the treatment of chronic bronchitis: a multi-centre study. *Respir Med* 1992;**86**:89–92.
- 100 **Bridgeman MM**, Marsden M, Selby C, et al. Effect of N-acetyl cysteine on the concentrations of thiols in plasma, bronchoalveolar lavage fluid, and lung tissue. *Thorax* 1994;**49**:670–5.
- 101 **Poole PJ**, Black PN. Oral mucolytic drugs for exacerbations of chronic obstructive pulmonary disease: systematic review. *BMJ* 2001;**322**:1271–4.
- 102 **Poole PJ**, Black PN. Mucolytic agents for chronic bronchitis or chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2000;CD001287.
- 103 **Baudry M**, Etienne S, Bruce A, et al. Salen-manganese complexes are superoxide dismutase-mimics. *Biochem Biophys Res Commun* 1993;**192**:964–8.
- 104 **Salvemini D**, Mazzon E, Dugo L, et al. Pharmacological manipulation of the inflammatory cascade by the superoxide dismutase mimetic, M40403. *Br J Pharmacol* 2001;**132**:815–27.
- 105 **Szabo C**, Day BJ, Salzman AL. Evaluation of the relative contribution of nitric oxide and peroxynitrite to the suppression of mitochondrial respiration in immunostimulated macrophages using a manganese mesoporphyrin superoxide dismutase mimetic and peroxynitrite scavenger. *FEBS Lett* 1996;**381**:82–6.
- 106 **Gonzalez PK**, Zhuang J, Doctrow SR, et al. EUK-8, a synthetic superoxide dismutase and catalase mimetic, ameliorates acute lung injury in endotoxemic swine. *J Pharmacol Exp Ther* 1995;**275**:798–806.
- 107 **Chang LY**, Subramaniam M, Yoder BA, et al. A catalytic antioxidant attenuates alveolar structural remodeling in bronchopulmonary dysplasia. *Am J Respir Crit Care Med* 2003;**167**:57–64.
- 108 **Cuzzocrea S**, Zingarelli B, Costantino G, et al. Beneficial effects of Mn(III)tetrakis (4-benzoic acid) porphyrin (MnTBAP), a superoxide dismutase mimetic, in carrageenan-induced pleurisy. *Free Radic Biol Med* 1999;**26**:25–33.
- 109 **Cuzzocrea S**, Thiemermann C, Salvemini D. Potential therapeutic effect of antioxidant therapy in shock and inflammation. *Curr Med Chem* 2004;**11**:1147–62.
- 110 **Day BJ**, Shawen S, Liochev SI, et al. A metalloporphyrin superoxide dismutase mimetic protects against paraquat-induced endothelial cell injury, in vitro. *J Pharmacol Exp Ther* 1995;**275**:1227–32.
- 111 **Oury TD**, Thakker K, Menache M, et al. Attenuation of bleomycin-induced pulmonary fibrosis by a catalytic antioxidant metalloporphyrin. *Am J Respir Cell Mol Biol* 2001;**25**:164–9.
- 112 **Smith KR**, Uyemini DL, Kodavanti UP, et al. Inhibition of tobacco smoke-induced lung inflammation by a catalytic antioxidant. *Free Radic Biol Med* 2002;**33**:1106–14.
- 113 **Comhair SA**, Thomassen MJ, Erzurum SC. Differential induction of extracellular glutathione peroxidase and nitric oxide synthase 2 in airways of healthy individuals exposed to 100% O<sub>2</sub> or cigarette smoke. *Am J Respir Cell Mol Biol* 2000;**23**:350–4.