

ASTHMA

Moderate dose inhaled corticosteroids plus salmeterol versus higher doses of inhaled corticosteroids in symptomatic asthma

M Masoli, M Weatherall, S Holt, R Beasley

Thorax 2005;60:730–734. doi: 10.1136/thx.2004.039180

See end of article for authors' affiliations

Correspondence to: Professor R Beasley, Medical Research Institute of New Zealand, P O Box 10055, Wellington, New Zealand; richard.beasley@mrnz.ac.nz

Received 12 December 2004
Accepted 5 May 2005

Background: There is uncertainty as to the dose of inhaled corticosteroids (ICS) at which to start concomitant long acting β agonist (LABA) treatment in patients with asthma not adequately controlled by ICS alone.

Methods: A meta-analysis was carried out of randomised, double blind clinical trials that compared the efficacy of adding salmeterol to moderate doses of ICS (fluticasone propionate 200 μ g/day or equivalent) with increasing the ICS dose by at least twofold in symptomatic adult patients with asthma. The main outcome measures were the number of subjects withdrawn from the study due to asthma and the number of subjects with at least one moderate or severe exacerbation.

Results: Twelve studies with a total of 4576 subjects met the inclusion criteria for the analyses. The number of subjects withdrawn due to asthma and with at least one moderate or severe exacerbation was higher in the high dose ICS group (odds ratios 1.58, 95% CI 1.12 to 2.24 and 1.35, 95% CI 1.10 to 1.66, respectively). For the secondary outcome variables (forced expiratory volume in 1 second, morning and evening peak expiratory flow, and daytime β agonist use) there was significantly greater benefit in the salmeterol group.

Conclusions: This meta-analysis shows that the addition of salmeterol to moderate doses of ICS (fluticasone 200 μ g/day or equivalent) in patients with asthma symptomatic at that dose results in significantly greater clinical benefit than increasing the dose of ICS by twofold or more.

The recent British consensus guidelines on asthma management recommend a long acting β agonist (LABA) drug as first line add on therapy at step 3 in patients not controlled on inhaled corticosteroids (ICS) alone.¹ However, this recommendation is made over a fourfold ICS dose range (200–800 μ g/day beclometasone dipropionate (BDP) or equivalent). The dilemma for clinicians is to know at what dose of ICS within this range to start concomitant LABA treatment in a patient inadequately controlled on ICS alone.

This dilemma is partly due to the heterogeneity of ICS doses used in clinical studies which have compared the efficacy of adding a LABA with increasing the dose of ICS. For example, in the original MIASMA meta-analysis in which the addition of salmeterol was significantly more effective than increasing the dose of ICS, the baseline daily dose of ICS ranged from 400 to 1000 μ g BDP or equivalent.² It was not therefore possible to determine whether there was a differential response across the ICS therapeutic range. In addition, a number of studies were included that used doses of ICS already near the peak of the dose-response curve—that is, 500 μ g/day fluticasone or equivalent.^{3–5} While recognising that there is considerable individual variability in the response to ICS in asthma,⁶ it would be expected that these inclusion criteria would have favoured the salmeterol group because further increasing the dose of ICS above 500 μ g/day fluticasone or equivalent would have produced minimal further benefit. In contrast, most of the studies comparing the effect of the addition of formoterol to increasing the dose of ICS have examined low baseline doses of ICS at which most of the therapeutic response has not been obtained.^{7–10} Likewise, this approach is not favoured due to the clinical preference to obtain most of the therapeutic benefit from ICS before adding an additional agent.

In this meta-analysis we have compared the clinical benefit of adding salmeterol in patients not controlled on moderate doses of ICS (200 μ g/day fluticasone or equivalent) with increasing the dose of ICS by at least twofold. In this way we have attempted to compare more accurately the relative benefits of both therapeutic options and thereby provide some guidance on what is a common dilemma facing clinicians in daily practice.

METHODS

Search strategy

A search was conducted of Medline from January 1966 to August 2003, of EMBASE from 1980 to August 2003, and of the Cochrane Controlled Trials Register. On Medline, a search of studies containing the keyword “salmeterol” using the MeSH subject heading “asthma” and “chemical and pharmacologic phenomena” (MeSH) and “inhaled corticosteroid” was undertaken. When limited to English the total number of studies was 629. GlaxoSmithKline, the manufacturer of salmeterol, was approached for details of all relevant studies; no additional studies were identified. There were no additional relevant studies published in other languages on Medline or EMBASE. Finally, the reference lists of relevant studies were examined but no other studies were found. The search strategy as recommended by the QUOROM statement is shown in fig 1.

Inclusion criteria

Two people independently examined the title and abstract of each paper, then the full paper if necessary. To be included in

Abbreviations: BDP, beclometasone dipropionate; FEV₁, forced expiratory flow in 1 second; ICS, inhaled corticosteroid; LABA, long acting beta agonist; PEF, peak expiratory flow




Figure 1 Results of search of Medline, EMBASE and GlaxoSmithKline databases for clinical studies eligible for inclusion in meta-analysis.

the meta-analysis, studies had to meet all of the following criteria: a double blind, randomised trial of adolescents (≥ 12 years of age) or adults with asthma symptomatic on moderate doses of ICS (200 $\mu\text{g}/\text{day}$ fluticasone or equivalent); direct comparison between moderate doses of ICS (200 $\mu\text{g}/\text{day}$ fluticasone or equivalent) and salmeterol versus at least a twofold higher dose of ICS (≥ 400 $\mu\text{g}/\text{day}$ fluticasone or equivalent) administered twice daily; studies of at least 12 weeks duration; and data on measures of clinical efficacy. Studies in which participants were dependent on oral steroids or involved in oral steroid reduction regimes were excluded.

Data extraction

Data extraction was based on reported summary statistics (means, SD and SE). The primary outcome measures were the number of subjects withdrawn due to asthma and the number of subjects with at least one moderate or severe exacerbation. Secondary outcome measures included morning and evening peak expiratory flow (PEF), forced expiratory volume in 1 second (FEV_1), night awakenings, and daytime and night time β agonist use. Data were primarily obtained from the original publications. If the data were not available in the standardised format in the original publications, they were either obtained from the MIASMA publication or were provided by GlaxoSmithKline (the manufacturer of salmeterol) in the format required. To assess the validity of data, two investigators correlated the data from the different sources. If discrepancies between sources were found, the data were preferentially obtained from the original publication, followed by MIASMA, and then the data provided by GlaxoSmithKline.

In the individual studies, moderate or severe exacerbations were defined as any event requiring treatment with oral/parenteral corticosteroids and/or emergency hospital treatment and/or any asthma medication not allowed as concurrent treatment during the study. Symptom scores could not be analysed because of differences in assessment of

symptoms between studies or because the information was not available in the format required. For withdrawals due to asthma, a continuity correction was used in the analysis because the data sets in the papers by Van Noord *et al*¹¹ and Kalberg *et al*¹² had “zero” cell counts.

Data analysis

For continuous variables a standard method of pooling means based on standardised mean differences using an inverse variance weighting approach¹³ was used. A weighted mean standard deviation was used to translate the standardised mean difference back into the original units. In the forest plots the area of the box is proportional to the variance of the estimate for the individual trials.

For the categorical variables the log odds ratio was pooled using the inverse variance weighting approach.¹³ The forest plots are presented as the log of the odds ratio and the area of the box is, where possible, proportional to the variance of the estimate for the individual trials. If the variance of individual trial estimates was too large to fit the scale of the plot, the area of the box is reduced to a quarter of the variance; this was required for the two primary outcome variables. Heterogeneity statistics and inconsistency measurements, the latter according to the method of Higgins and Thompson,¹⁴ were calculated for all the comparisons. SAS Version 8.2 was used.

RESULTS

Twelve studies with a total of 4576 subjects met the inclusion criteria for the analysis (table 1).^{11–12, 15–23} The mean FEV_1 was 64% of predicted normal values (range 40–85%), indicating that subjects had moderate to severe asthma. By virtue of the inclusion criteria used, the methodological quality of included studies was high with all studies achieving a Jadad score of 4 or more.²⁴ In two studies^{20–21} subjects were only eligible to participate if they had been shown to be unstable on 200 $\mu\text{g}/\text{day}$ fluticasone. They were then treated with fluticasone 500 $\mu\text{g}/\text{day}$ in the run in phase, before randomisation to fluticasone 200 $\mu\text{g}/\text{day}$ and salmeterol or fluticasone 500 $\mu\text{g}/\text{day}$. Comparisons were made between the addition of salmeterol to BDP 400 *v* 1000 $\mu\text{g}/\text{day}$ in one study, BDP 400 *v* 800 $\mu\text{g}/\text{day}$ in three studies, fluticasone 200 *v* 500 $\mu\text{g}/\text{day}$ in six studies, fluticasone 200 *v* 400 $\mu\text{g}/\text{day}$ in one study, and fluticasone 200 $\mu\text{g}/\text{day}$ *v* budesonide 800 $\mu\text{g}/\text{day}$ in one study. The study by Baraniuk *et al*²² was the combined publication of two separate studies (FLTA 4021 and FLTA 4022). The combination salmeterol/fluticasone (Advair/Seretide) inhaler was used in three studies.

Evidence of heterogeneity was found for several of the outcome variables including morning PEF (where the heterogeneous studies were in the same direction as the others), FEV_1 (where one study was more positive and another more negative), daytime β agonist use (two studies showing less difference), and night time β agonist use (with two studies having more effect). For all these the confidence intervals became wider under the random effects models, and for FEV_1 and night time β agonist use this led to the lower confidence limit being close to or at zero, consistent with no difference in effectiveness between the two strategies. For the other secondary outcome variables there was no substantive difference to the conclusions.

Main outcome measures

There was a significant reduction in the number of subjects withdrawn due to asthma in the low dose ICS/salmeterol group (59/2036) compared with high dose ICS treatment (86/1992) with an odds ratio of 1.58 (95% CI 1.12 to 2.24) (table 2, fig 2). There was a significant reduction in the number of subjects with one or more moderate or severe exacerbation in

Table 1 Summary of studies included in the meta-analysis

Reference	N	Study duration (weeks)	Doses of FP/BDP/Sal ($\mu\text{g}/\text{day}$)	Device	Comparison dose ($\mu\text{g}/\text{day}$)	Range (mean) FEV ₁ % predicted
Greening ¹⁵	426	26	BDP 400 Sal 100	MDI Diskhaler	BDP 1000	≥ 50 (PEF 74)
Kelsen ¹⁶	483	24	BDP 400 Sal 100	MDI	BDP 800	45–80 (65)
Murray ¹⁷	514	24	BDP 400 Sal 100	MDI	BDP 800	50–80 (66)
Kalberg ¹²	488	24	FP 200 Sal 100	–	FP 500	40–80
Condemi ¹⁸	437	24	FP 200 Sal 100	MDI	FP 500	40–80 (61)
Van Noord ¹¹	60	12	FP 200 Sal 100	Diskhaler	FP 400	≥ 50 (72)
Vermetten ¹⁹	233	12	BDP 400 Sal 100	Diskhaler	BDP 800	PEF >60 (84)
Bloom ²⁰	574	12	FP 200 Sal 100	Diskus (comb)	FP 500	65–95 (78)
Busse ²¹	558	12	FP 200 Sal 100	Diskus (comb)	FP 500	≥ 65 (81)
FLTA 4021 ²²	232	12	FP 200 Sal 100	MDI	FP 500	40–85 (63)
FLTA 4022 ²²	222	12	FP 200 Sal 100	MDI	FP 500	40–85 (63)
Johansson ²³	349	12	FP 200 Sal 100	Diskus (comb)	BUD 800	65–85 (77)

FP, fluticasone propionate; ICS, inhaled corticosteroid; BDP, beclometasone dipropionate; Sal, salmeterol; BUD, budesonide; Comb, combination fluticasone/salmeterol; FEV₁, forced expiratory volume in 1 second; PEF, peak expiratory flow; MDI, metered dose inhaler.

the low dose ICS/salmeterol group (184/2312) compared with high dose ICS treatment (243/2264) with an odds ratio of 1.35 (95% CI 1.10 to 1.66) (table 2, fig 3).

Secondary outcome measures

Mean morning PEF and evening PEF were significantly greater and daytime β agonist use was reduced in those who received low dose ICS/salmeterol compared with high dose ICS, with differences of 23 l/min, 19 l/min, and -0.6 puffs per day, respectively (table 3). Significantly greater improvements were also observed for FEV₁, although the lower confidence limit was closer to no difference. For night awakenings and night time β agonist use, the lower confidence limits included no difference in effect although the point estimates favoured the ICS/salmeterol group.

DISCUSSION

In this meta-analysis we have shown that, in patients symptomatic on ICS at a dose of 200 μg fluticasone or equivalent, the addition of salmeterol is superior to increasing the dose of ICS by at least twofold for all major clinical outcome measures. For the main outcome variables of withdrawals due to asthma and at least one moderate or severe exacerbation, subjects receiving salmeterol had a 35–50% reduced risk compared with higher dose ICS treatment. The secondary outcome variables of lung function and β agonist use were also significantly improved compared with the higher ICS dose.

There are issues relating to the design of the study that are relevant to the interpretation of the results. The most important was the decision to restrict the analysis to studies that used a baseline dose of ICS of 200 $\mu\text{g}/\text{day}$ fluticasone or




Figure 2 Odds ratio for proportion of withdrawals due to asthma. Forest plot of fixed effects. ICS, inhaled corticosteroids; LABA, long acting β agonist.




Figure 3 Odds ratio for number of subjects with one or more moderate or severe exacerbation. Forest plot of fixed effects. ICS, inhaled corticosteroids; LABA, long acting β agonist.

Table 2 Numbers of subjects withdrawn due to asthma and number of subjects with at least one moderate or severe exacerbation, and associated odds ratios (95% CI) between treatment with salmeterol and increased dose of inhaled corticosteroid

Study	Withdrawals due to asthma			Moderate or severe exacerbations*		
	Sal	ICS	OR (95% CI)	Sal	ICS	OR (95% CI)
Greening van Noord (low dose)	7/220	14/206	2.22 (0.88 to 5.61)	19/220 0/30	19/206 2/30	1.07 (0.56 to 2.08) 5.35 (0.25 to 116.32)
Vermetten	6/113	10/120	1.62 (0.57 to 4.62)	9/113	17/120	1.86 (0.81 to 4.29)
Busse	14/281	20/277	1.48 (0.73 to 3.00)	8/281	6/277	0.77 (0.27 to 2.17)
Bloom	15/295	22/279	1.60 (0.81 to 3.15)	6/295	9/279	1.56 (0.57 to 4.30)
Kelsen	3/239	6/244	1.98 (0.49 to 8.02)	38/239	44/244	1.16 (0.72 to 1.72)
Murray	5/260	5/254	1.02 (0.29 to 3.58)	43/260	45/254	1.09 (0.69 to 1.72)
Kalberg	—	—	—	20/246	32/242	1.71 (0.95 to 3.06)
Condemi	2/221	6/216	3.13 (0.62 to 15.68)	21/221	31/216	1.58 (0.88 to 2.84)
FLTA4021	1/118	1/114	1.04 (0.06 to 16.75)	7/118	8/114	1.19 (0.43 to 3.28)
FLTA4022	1/113	1/109	1.04 (0.06 to 16.79)	4/113	20/109	5.57 (1.94 to 16.05)
Johansson	3/176	1/173	0.34 (0.03 to 3.26)	9/176	10/173	1.13 (0.46 to 2.80)
Pooled estimates of effect (95% CI)						
Fixed effect				1.58 (1.12 to 2.24)		
Random effect				1.35 (1.10 to 1.66)		
Inconsistency measurements				1.37 (1.10 to 1.70)		
				10.4 (0 to 49.7)		

OR, Odds ratio for withdrawal (greater than 1 means ICS more likely to withdraw and favours salmeterol). Note that a continuity correction was used for this analysis because the van Noord data set had a “zero” cell count.

Sal, group receiving added salmeterol; ICS, group receiving increased dose of inhaled steroid.

*Number of participants with one or more moderate or severe exacerbations of asthma.

equivalent. While recognising the marked individual variability in response to ICS⁵ and the requirement for higher doses in certain circumstances,^{25, 26} this ICS dose was chosen as its use results in 80–90% of the maximum obtainable benefit for all clinical outcome measures including exacerbations.^{3–5}

Consideration was given to broadening the inclusion criteria to include similar studies of the LABA formoterol. This was not undertaken due to the paucity of such studies which used an equivalent baseline daily dose of ICS and the lack of data presented in the same standardised format.^{7–10} However, the findings are complemented by the FACET study in which budesonide at 800 µg/day resulted in significantly fewer severe exacerbations than low dose budesonide (200 µg/day) plus formoterol.⁷ This study illustrates the importance of obtaining most of the therapeutic benefit from ICS therapy before considering add-on therapy in patients with symptomatic asthma.

By limiting the analysis to studies of salmeterol, we were able to include 12 studies involving over 4500 patients with asthma including five studies of 1935 subjects not included in the original MIASMA publication.² We are confident that we obtained all eligible studies as a result of the comprehensive search strategy undertaken. However, even with the strict inclusion criteria employed, our meta-analysis did include studies in which salmeterol was added to different ICS (BDP

and fluticasone), different inhaler devices (Diskus and MDI), and both combination and separate inhalers. We were provided with data in a standardised format from GlaxoSmithKline which, in addition to the data published in the original manuscripts and MIASMA, allowed data to be used in a standardised format across the studies included in the meta-analysis.

Another issue was the choice of the main outcome variables—withdrawals due to asthma and one or more moderate or severe exacerbations. This analysis differed from the original MIASMA study² in that we did not include mild exacerbations which incorporated an increase in β agonist use as this would be expected to be reduced with concomitant LABA treatment. However, even with the use of withdrawals due to asthma, the results must be interpreted with some caution as the number of withdrawals due to asthma was relatively few in comparison with total withdrawals. With regard to moderate to severe exacerbations, this may have included some episodes that reflected “loss of asthma control” rather than clinically severe exacerbations, and the paucity of hospital admissions prevented this specific outcome measure being used. Furthermore, it is likely that there was a close correlation between withdrawals due to asthma and moderate or severe exacerbations. However, despite these limitations we consider that withdrawals and moderate or severe exacerbations

Table 3 Mean difference (95% CI) for all secondary outcome measures between the group receiving added salmeterol and the group receiving increased dose of ICS

Outcome measure	Difference in original units†		
	Fixed effects	Random effects	Inconsistency measures
Morning PEF (l/min)	23 (19 to 28)	23 (15 to 30)	63.7 (25.5 to 82.3)
Evening PEF (l/min)	19 (15 to 23)	19 (13 to 24)	29.9 (0.00 to 82.3)
FEV ₁ (l)	0.12 (0.09 to 0.15)	0.12 (0.06 to 0.18)	71.0 (36.7 to 86.7)
Night awakenings (no/week)	−0.03 (0.00 to −0.07)	−0.03 (0.01 to −0.07)	20.5 (0.00 to 65.1)
Day time β agonist use (puffs/day)	−0.58 (−0.44 to −0.72)	−0.60 (−0.35 to −0.84)	70.3 (30.5 to 87.3)
Night time β agonist use (puffs/night)	−0.08 (−0.02 to −0.13)	−0.08 (−0.00 to −0.16)	58.0 (0.00 to 83.0)

†Mean difference represents the mean outcome measure for the group receiving added salmeterol minus the mean outcome measure for the group receiving increased dose of inhaled corticosteroid.

are of greater clinical relevance than lung function measures, particularly with comparisons between ICS and LABA treatment. Consistent with the findings for these main outcome measures, the secondary outcome measures including lung function measures such as clinic FEV₁, morning and evening PEF, and β agonist use also favoured the salmeterol group.

Finally, it is worthy of comment that, although the studies in this meta-analysis were restricted to salmeterol as the LABA, the evidence that salmeterol and formoterol have similar efficacy in the long term treatment of asthma^{27, 28} would suggest that the results are probably applicable to the class of LABAs.

In conclusion, recent meta-analyses of the dose-response of ICS demonstrate a relatively flat dose-response relationship with 80–90% of the maximum therapeutic benefit achieved with a daily dose of around 200 μ g fluticasone or 400 μ g budesonide.^{3–5} Notwithstanding the marked individual variability in response to ICS treatment,⁶ these studies would suggest that the 200 μ g/day dose of fluticasone or equivalent is likely to represent the preferable level at which to consider the addition of salmeterol in an asthmatic patient not well controlled on ICS. The results of this meta-analysis support this view by showing that the addition of salmeterol to moderate doses of ICS (200 μ g/day fluticasone or equivalent) is more effective than increasing the dose of ICS by twofold or more for clinical outcome measures including exacerbations and withdrawals due to asthma. These findings should provide clinicians with greater confidence when deciding the dose of ICS at which to consider adding salmeterol at step 3 in the asthma guidelines.

Authors' affiliations

M Masoli, R Beasley, Medical Research Institute of New Zealand, Wellington, New Zealand

M Weatherall, Wellington School of Medicine & Health Sciences, Wellington, New Zealand

S Holt, P3 Research, Wellington, New Zealand

R Beasley, University of Southampton, Southampton, UK

Competing interests: The Medical Research Institute of New Zealand, the Wellington School of Medicine & Health Sciences, and P3 Research have all received research grants from Astra Draco, GlaxoSmithKline, and Novartis. R Beasley has received fees for consulting and speaking and reimbursement for attending symposia from Astra Draco, and GlaxoSmithKline and Novartis. M Masoli and S Holt have received reimbursement for attending symposia from Astra Draco and Novartis, and S Holt also from GlaxoSmithKline.

MM developed the protocol, organised the search, undertook data extraction and wrote the paper. MW was responsible for the data analysis, statistical methodology and manuscript preparation. SH contributed to the study concept and manuscript preparation. RB provided senior oversight for the study and was responsible for the manuscript preparation. MM will act as guarantor for this paper.

REFERENCES

- 1 **British Thoracic Society/Scottish Guidelines Intercollegiate Network**. British guideline on the management of asthma. *Thorax* 2003;**58**(Suppl 1):i1–94.
- 2 **Shrewsbury S**, Pyke S, Britton M. Meta-analysis of increased dose of inhaled steroid or addition of salmeterol in symptomatic asthma (MIASMA). *BMJ* 2000;**320**:1368–73.
- 3 **Holt S**, Suder A, Weatherall M, et al. Dose-response relation of inhaled fluticasone propionate in adolescents and adults with asthma: meta-analysis. *BMJ* 2001;**323**:253–6.
- 4 **Masoli M**, Holt S, Weatherall M, et al. Dose-response relationship of inhaled budesonide in adult asthma: a meta-analysis. *Eur Respir J* 2004;**23**:1–7.
- 5 **Masoli M**, Weatherall M, Holt S, et al. Clinical dose-response relationship of fluticasone propionate in adults with asthma. *Thorax* 2004;**59**:16–20.
- 6 **Szeferl SJ**, Martin RJ, King TS, et al. Significant variability in response to inhaled corticosteroids for persistent asthma. *J Allergy Clin Immunol* 2002;**109**:410–8.
- 7 **Pauwels RA**, Lofdahl C-G, Postma DS, et al. Effect of inhaled formoterol and budesonide on exacerbations of asthma. *N Engl J Med* 1997;**337**:1405–11.
- 8 **O'Byrne P**, Barnes P, Rodriguez-Roisin R, et al. Low dose inhaled budesonide and formoterol in mild persistent asthma (OPTIMA). *Am J Respir Crit Care Med* 2001;**164**:1392–7.
- 9 **Lalloo U**, Malolepszy J, Kozma D, et al. Budesonide and formoterol in a single inhaler improves asthma control compared with increasing the dose of corticosteroid in adults with mild to moderate asthma. *Chest* 2003;**123**:1480–7.
- 10 **Bouras D**, Bachlitzanakis N, Kottakis J, et al. Formoterol and beclomethasone versus higher dose beclomethasone as maintenance therapy in adult asthma. *Eur Respir J* 1999;**14**:627–32.
- 11 **van Noord JA**, Schreurs AJM, Mol SJM, et al. Addition of salmeterol versus doubling the dose of fluticasone propionate in patients with mild to moderate asthma. *Thorax* 1999;**54**:207–12.
- 12 **Kalberg CJ**, Nelson H, Yancey S, et al. A comparison of added salmeterol versus increased-dose fluticasone in patients symptomatic on low-dose fluticasone. *J Allergy Clin Immunol* 1998;**101**:S6.
- 13 **Whitehead A**. *Meta-analysis of controlled clinical trials*. Chichester: Wiley, 2002.
- 14 **Higgins JPT**, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;**21**:1539–58.
- 15 **Greening A**, Ind P, Northfield M, et al. Added salmeterol versus higher dose corticosteroid in asthma patients with symptoms on existing inhaled corticosteroid. *Lancet* 1994;**344**:219–24.
- 16 **Kelsen S**, Church N, Gillman S, et al. Salmeterol added to inhaled corticosteroid therapy is superior to doubling the dose of inhaled corticosteroids: a randomized clinical trial. *J Asthma* 1999;**36**:703–15.
- 17 **Murray J**, Church N, Anderson W, et al. Concurrent use of salmeterol with inhaled corticosteroids is more effective than inhaled corticosteroid dose increases. *Allergy Asthma Proc* 1999;**20**:173–80.
- 18 **Condemi J**, Goldenstein S, Kalberg C, et al. The addition of salmeterol to fluticasone propionate versus increasing the dose of fluticasone propionate in patients with persistent asthma. *Ann Allergy Asthma Immunol* 1999;**82**:383–9.
- 19 **Vermetten FA**, Boermans AJ, Luiten WD, et al. Comparison of salmeterol with beclomethasone in adult patients with mild persistent asthma who are already on low dose inhaled steroids. *J Asthma* 1999;**36**:97–106.
- 20 **Bloom J**, Calhoun W, Koenig S, et al. Fluticasone propionate/salmeterol 100/50 μ g is inhaled steroid sparing in patients who require fluticasone propionate 250 μ g for asthma stability. *Am J Respir Crit Care Med* 2003;**167**:A891.
- 21 **Busse W**, Koenig S, Oppenheimer J, et al. Steroid-sparing effects of fluticasone propionate 100 μ g and salmeterol 50 μ g administered twice daily in a single product in patients previously controlled with fluticasone propionate 250 μ g administered twice daily. *J Allergy Clin Immunol* 2003;**111**:57–65.
- 22 **Baraniuk J**, Murray J, Nathan R, et al. Fluticasone alone or in combination with salmeterol vs triamcinolone in asthma. *Chest* 1999;**116**:625–32.
- 23 **Johansson G**, McIvor A, D'Ambrosio F, et al. Comparison of salmeterol/fluticasone propionate combination with budesonide in patients with mild to moderate asthma. *Clin Drug Invest* 2001;**21**:633–42.
- 24 **Jadad AR**, Moore RA, Carrol D, et al. Assessing the quality of reports of randomized clinical trials: Is blinding necessary? *Control Clin Trials* 1996;**17**:1–12.
- 25 **Nelson HS**, Busse WW, deBoisblanc BP, et al. Fluticasone propionate powder: oral corticosteroid-sparing effect and improved lung function and quality of life in patients with severe chronic asthma. *J Allergy Clin Immunol* 1999;**103**:267–75.
- 26 **Noonan M**, Chervinsky P, Busse WW, et al. Fluticasone propionate reduces oral prednisone use while it improves asthma control and quality of life. *Am J Respir Crit Care Med* 1995;**152**:1467–73.
- 27 **Vervloet D**, Ekström T, Pela R, et al. A 6-month comparison between formoterol and salmeterol in patients with reversible obstructive airways disease. *Respir Med* 1998;**92**:836–42.
- 28 **Nightingale JA**, Rogers DF, Barnes PJ. Comparison of the effects of salmeterol and formoterol in patients with severe asthma. *Chest* 2002;**121**:1401–6.