

# Mechanism of allosteric regulation of transglutaminase 2 by GTP

Gillian E. Begg\*, Lyle Carrington†, Philippa H. Stokes‡, Jacqueline M. Matthews‡, Merridee A. Wouters\*, Ahsan Husain§, Laszlo Lorand¶, Siiri E. Iismaa\*, and Robert M. Graham\*||

\*Victor Chang Cardiac Research Institute, University of New South Wales, 384 Victoria Street, Darlinghurst NSW 2010, Australia; †University of Queensland, Brisbane QLD 4072, Australia; ‡University of Sydney, Sydney NSW 2006, Australia; §University of Alabama at Birmingham, Birmingham, AL 35294; and ¶Northwestern University Medical School, Chicago, IL 60611

Contributed by Laszlo Lorand, October 23, 2006 (sent for review October 3, 2006)

**Allosteric regulation is a fundamental mechanism of biological control. Here, we investigated the allosteric mechanism by which GTP inhibits cross-linking activity of transglutaminase 2 (TG2), a multifunctional protein, with postulated roles in receptor signaling, extracellular matrix assembly, and apoptosis. Our findings indicate that at least two components are involved in functionally coupling the allosteric site and active center of TG2, namely (i) GTP binding to mask a conformationally destabilizing switch residue, Arg-579, and to facilitate interdomain interactions that promote adoption of a compact, catalytically inactive conformation and (ii) stabilization of the inactive conformation by an uncommon H bond between a cysteine (Cys-277, an active center residue) and a tyrosine (Tyr-516, a residue located on a loop of the  $\beta$ -barrel 1 domain that harbors the GTP-binding site). Although not essential for GTP-mediated inhibition of cross-linking, this H bond enhances the rate of formation of the inactive conformer.**

protein conformation | GTP inhibition | transamidase activity

**A**llosteric regulation of enzymes by conformational change is an important means of biological control, involving residues that functionally couple ligand binding at the allosteric site to modification of the catalytic site. Transglutaminase type 2 (TG2), also known as tissue TG or  $G_{II}$  (high molecular weight GTP-binding protein), is a multifunctional protein that is allosterically regulated by calcium and GTP (1). TG2 catalyzes calcium-dependent transamidation reactions, resulting in posttranslational amine modification of proteins or cross-linking of interchain glutamine and lysine residues to form  $N^\epsilon(\gamma\text{-glutamyl})$ lysine isopeptide bonds, which confer rigidity and protease resistance on protein complexes (2). TG2 is also a GTPase (3) and mediates intracellular signaling by various G protein-coupled receptors (4–6).

GDP-bound human TG2 (7) is comprised of four domains: an N-terminal  $\beta$ -sandwich, a core domain in which the transamidase active site catalytic triad (Cys-277, His-335, and Asp-358) and transition-state stabilizing residue (Trp-241) (8) are buried and inaccessible to substrate, and two  $\beta$ -barrels. Nucleotide binds mainly to residues from the first and last strands (amino acids 476–482 and 580–583) of  $\beta$ -barrel 1 and to two core domain residues (Lys-173 and Phe-174) that protrude on a loop to meet  $\beta$ -barrel 1 (7, 9, 10). This is postulated to stabilize two  $\beta$ -barrel 1 loops that block access to the catalytic site (7). One of these loops protrudes into the core domain localizing Tyr-516 within hydrogen-bonding distance of Cys-277 (7). This is postulated to prevent Cys-277 interaction with the substrate (7, 11, 12). Calcium-activated TG2 has unique conformational epitopes (13) and is less compact (14–16) and less resistant to protease digestion (1, 10, 14) than GTP-bound TG2. Allosteric mechanisms governing the conformational switch between transamidase and GTPase functions have yet to be elucidated.

Mutation of Arg-579 in rat TG2 (Arg-580 in human TG2) to Ala reduces GTP affinity by 100-fold and uncouples GTP-dependent allosteric inhibition of transamidase activity, resulting in dysregulated intracellular cross-linking (10). R579A is also more resistant

to  $\mu$ -calpain digestion than wild-type TG2 (WT) in the absence of GTP (10), suggesting an important conformational role for this residue. Here, we show that Arg-579 is a conformationally destabilizing residue that is masked by GTP binding. Further, we provide evidence that Tyr-516 is not directly involved in transamidase catalysis but, by forming a H bond with Cys-277, stabilizes and enhances the rate of conversion to the compact, transamidase-inactive conformation. Thus, Arg-579 and Tyr-516 form an important part of the allosteric mechanism that functionally couples GTP binding to inhibition of cross-linking at the distal TG2 active site.

## Results

**GTP Stabilizes a Compact Conformation of TG2.** As shown by agarose gel electrophoresis (17), the addition of GTP caused a shift in WT mobility in native PAGE (nPAGE) to a faster-migrating species (Fig. 1*B*) that could be specifically labeled with [ $\alpha$ - $^{32}$ P]GTP (data not shown). Point mutants of core (Ser-171 and Phe-174) and barrel 1 (Arg-476, Arg-478, and Arg-579) residues (Fig. 1*A*) showed GTP-dependent shifts in mobility (Fig. 1*B* and *C*) in order from most to least sensitive: WT  $\approx$  F174W > R476A  $\approx$  R478A > R476A/R478A  $\approx$  F174A > R579A  $\approx$  R579K > S171E. This order is consistent with their relative reduction in GTP photolabeling and GTP-dependent inhibition of transamidase activity relative to WT (9, 10).

**Arg-579 Destabilizes the Compact TG2 Conformation.** Independent of GTP addition, R579A migrated as a fast-migrating species (Fig. 1*B* and *C*) and had a dominant effect on migration when combined with R476A or R478A in the double mutants, R476A/R579A and R478A/R579A (Fig. 1*C*). A similar migration pattern was observed for R579K, which retains the charge and much of the bulk of the arginine side chain, but like R579A is impaired in GTP binding (10). Of note, the fast-migrating species observed for Arg-579 mutants migrated slightly more slowly than GTP-bound WT or other mutants.

The homogeneity and approximate shape of GTP-bound WT and R579A was evaluated in sedimentation velocity experiments at 4°C [see supporting information (SI) Fig. 6]. The accuracy of the molar mass estimate (Table 1) of the major GTP-bound WT species (Fig. 1*D*) (within 4% of the calculated molar mass of 78.0 kDa for rat TG2 with one GTP bound) indicates GTP-bound TG2 adopts a stable conformation. Similar results, i.e.,  $s_{20,w}$  of 4.94 S and a molar mass of 74.8 kDa, were obtained for WT in 500  $\mu$ M GTP at 24°C, indicating the GTP-bound conformation is stable at higher temperatures. R579A consistently sedimented as a single species in the


Author contributions: G.E.B., S.E.I., and R.M.G. designed research; G.E.B., L.C., P.H.S., M.A.W., and S.E.I. performed research; G.E.B., P.H.S., J.M.M., M.A.W., A.H., L.L., S.E.I., and R.M.G. analyzed data; and G.E.B., S.E.I., and R.M.G. wrote the paper.

The authors declare no conflict of interest.

Abbreviation: nPAGE, native PAGE.

||To whom correspondence may be addressed. E-mail: l.lorand@northwestern.edu, s.iismaa@victorchang.unsw.edu.au, or b.graham@victorchang.unsw.edu.au.

© 2006 by The National Academy of Sciences of the USA



**Fig. 1.** Mutations of residues involved in GTP-binding or the Cys-277–Tyr-516 H bond affect TG2 conformation. (A) Model of GDP-bound human TG2 showing the transamidase active center (Cys-277, His-335, and Asp-358) and other key residues [(Protein Data Bank ID code 1KV3) (7); core domain backbone, pale pink;  $\beta$ -barrel 1 domain, green]. Carbon atoms are gray, except those of active site residues (cyan), and GDP (orange). Other atoms include oxygen, red; nitrogen, deep blue; sulfur, yellow; phosphorous, bright pink. Arg-580 in human TG2 is equivalent to Arg-579 in rat TG2. (B and C) Samples (3  $\mu$ g) of WT and mutant TG2 were incubated (3 h at 25°C) in buffer (50 mM Tris-HCl, pH 7.5/50 mM NaCl/0.5 mM EDTA/1.0 mM MgCl<sub>2</sub>/5 mM DTT/GTP as indicated). nPAGE was performed with 0.5 mM MgCl<sub>2</sub>, and the indicated GTP concentrations were included in gel solutions and running buffer. Data are representative of two to eight experiments. (D) Sedimentation coefficient distributions,  $c_M(s)$ , for 13  $\mu$ M WT (solid line), Y516F (dashed line), and R579A (dotted line) in 10 mM Tris-HCl pH 7.5/100 mM NaCl/0.5 mM EDTA/1 mM DTT/500  $\mu$ M GTP at 4°C. Molar mass was fixed at 78.0 kDa for 2.5–3.8 S species. Data are representative of three to four experiments. (E) Disulfide trapping of Cys-277 with a cysteine substituted for Tyr-516. Samples (3  $\mu$ g) of WT and mutant TG2 were oxidized (2 h at 25°C) in buffer (50 mM Tris-HCl, pH 7.5/50 mM NaCl/200:800  $\mu$ M Cu<sup>2+</sup>:phenanthroline/50  $\mu$ M GTP), then incubated (1 h at 25°C) in 0.5 mM MgCl<sub>2</sub>, without or with 20 mM DTT, as indicated. nPAGE was performed as in B. Data are representative of two experiments.

presence of GTP and at a slightly slower rate than GTP-bound WT (Fig. 1D and Table 1). This difference was not due to incomplete saturation of R579A with GTP as a 10-fold increase in GTP concentration did not affect the R579A  $s$  value (Table 1). The compact R579A conformation, therefore, is slightly more asymmetric or expanded than GTP-bound WT (dimensions of equivalent prolate ellipsoids are compared in Table 1). The fast-migrating species displayed by Arg-579 mutants (Fig. 1B and C), thus, is not due to charge but most likely to GTP-independent isomerization to a semicompact conformation. This indicates that Arg-579 destabilizes the compact TG2 conformation in the absence of GTP.

**The Cys-277–Tyr-516 H Bond Stabilizes the Compact TG2 Conformation.** The electronegative group required for the Cys-277–Tyr-516 H bond was removed by substitution of Cys-277 with Ala or Ser or Tyr-516 with Phe. Transamidase-inactive (18) C277A (Fig. 1C) and C277S (data not shown) showed markedly weaker GTP-dependent shifts to fast-migrating species and weaker [ $\alpha$ -<sup>32</sup>P]GTP photolabeling (Fig. 2A) compared with WT, consistent with our previous finding (19) that their affinity for 2'-(or 3')-O-(*N*-methylanthraniloyl) GTP is greatly reduced. Y516F showed an impaired GTP-dependent shift to fast-migrating species (Fig. 1B and C) and eliminated the fast-migrating species

**Table 1. Hydrodynamic size of TG2 at 4°C**

TG2	$s_{20,w}$ , S	Molar mass,* kDa	Frictional ratio†	Prolate ellipsoid,‡ nm
WT (500 $\mu$ M GTP)	4.93	81.1 (80.5, 82.0)	1.33	15.5 $\times$ 4.3
Y516F (500 $\mu$ M GTP)	4.90	76.5 (76.0, 77.0)	1.34	15.9 $\times$ 4.3
R579A (500 $\mu$ M GTP)	4.82	80.1 (79.0, 82.0)	1.36	16.6 $\times$ 4.2
R579A (5 mM GTP)	4.78	82.0 (78.0, 88.0)	1.37	17.1 $\times$ 4.1

\*Data fitted with hybrid discrete/continuous model. Confidence limits (parentheses) were calculated by using  $F$  statistics at confidence level of 1 SD.

†Calculated from sequence molar mass and  $s$  by using SEDNTERP.

‡Two-fold major and minor axis, calculated assuming hydration of 0.402 g/g, as predicted from amino acid composition (38) by using SEDNTERP.

seen with R579A mutants when combined with R579A in the double mutant, Y516F/R579A (Fig. 1C). Surprisingly, Y516F photolabeled (Fig. 2A) and bound GTP $\gamma$ S (Fig. 2B;  $K_d = 0.8 \pm 0.08 \mu$ M), equivalent to WT ( $K_d = 1.6 \pm 0.09 \mu$ M; ref. 10). Thus, unlike Cys-277 mutants, decreased stability of the compact Y516F form is not due to weaker GTP binding. Isothermal titration calorimetry of WT, F174A, and R579A showed sharp peaks indicating rapid binding (10). However, GTP $\gamma$ S titration of Y516F (Fig. 2B) indicated a slow interaction process, with heat liberated after each injection taking >30 min to reach baseline. This indicates either slow interconversion of Y516F to a conformation competent to bind GTP, or a rapid, low-affinity interaction with GTP followed by a slow conformational change to high-affinity binding (20).

GTP-bound Y516F sedimented as a major species (Fig. 1D) with a very slight decrease in  $s$  value relative to WT (Table 1), indicating that in contrast to R579A, compact GTP-bound Y516F is not measurably different from WT. This is consistent with equal migration of GTP-bound forms of WT and Y516F (Fig. 1B and C). The minor peak at  $\approx 2.5$  S (Fig. 1D) suggests an expanded Y516F form, consistent with decreased stability of the compact form (Fig. 1B and C).

**The Cys-277–Tyr-516 Interaction Enhances Allosteric Inhibition of TG2 by GTP.** Y516F showed similar dose-dependent calcium-stimulated transamidase activity to WT (see SI Fig. 7), and the same  $K_m$  for the glutamyl substrate, dimethyl casein (0.095% versus 0.093% wt/vol for WT), indicating that Tyr-516 is not involved in substrate binding or catalysis. GTP $\gamma$ S inhibition of transamidase activity, evaluated at  $\approx 20\%$  maximal activity (Fig. 2C), resulted in a 20-fold higher  $IC_{50}$  for Y516F (255  $\mu$ M compared with 13  $\mu$ M for WT), demonstrating that the loss of the Cys-277–Tyr-516 H bond substantially weakens, but does not prevent, GTP inhibition of transamidase activity. Together, this data indicate that transition to the compact, transamidase-inactive form is enhanced by GTP binding and that the Cys-277–Tyr-516 H bond is important for conformer stability.

**A Disulfide Bond Between Cys-277 and a Cys Substituted for Tyr-516 Stabilizes the Compact TG2 Conformation.** Cys was substituted for Tyr-516 to evaluate whether oxidative trapping of Y516C in a disulfide bond with Cys-277 can mimic the native Cys–Tyr H bond to stabilize the compact form. Cys-336, which is within bonding distance of Cys-277 (Fig. 1A), was mutated to Val (the equivalent residue in TG3), either alone (control) or in combination with Y516C, to eliminate the possibility of a competing Cys-277–Cys-336 disulfide bond (12). Maximal calcium-stimulated transamidase activities of Y516C, C336V, and Y516C/C336V were 76%, 83% and 71% of WT, respectively.

To maximize disulfide bond formation between Cys-277 and Cys-substituted Tyr-516, proteins were oxidized in the presence of GTP. After GTP removal by nPAGE (Fig. 1E Upper Left), Y516C and Y516C/C336V, but not WT, C336V, or Y516F,

displayed a range of fast-migrating species that were eliminated by DTT reduction before nPAGE (Fig. 1E Upper Right), demonstrating stabilization of these species by a disulfide bond, likely between Cys-277 and Cys-516. Oxidation, followed by nPAGE in the presence of GTP (Fig. 1E Lower Left), showed a marked increase in compact GTP-bound Y516C and Y516C/C336V relative to WT and C336V, indicating Cys-277–Cys-516-mediated stabilization of the fully compact GTP-bound conformation. Oxidation followed by DTT reduction before nPAGE in the presence of GTP (Fig. 1E Lower Right) completely reversed the effects of oxidation (compare to untreated mutants in Fig. 1C). The increase in the relative proportion of compact WT, C336V, and Y516F forms (Fig. 1E, Lower Right compared with Lower Left) indicates that this form is more stable under reducing conditions. In contrast, stability of the compact form of Y516C mutants already was maximized under oxidizing conditions, likely due to the Cys-277–Cys-516 disulfide bond.

**Transition Between GTP-Free and GTP-Bound Forms Is Faster for R579A.** Undiluted samples of WT, R579A, and Y516F (Fig. 3) showed a major GTP-bound species of  $\approx 3.16$  S for WT and Y516F (equivalent to  $s_{20,w} \approx 4.9$  S) and  $\approx 3.1$  S for R579A (equivalent to  $s_{20,w} \approx 4.8$  S). The 3.9 S R579A species is likely a dimer. Upon sample dilution, the proportion of GTP-bound species decreased and species at 0.5–1.5 S and 1.5–2.5 S increased. Free GTP sediments predominantly at  $<0.25$  S (data not shown). Species at 1.5–2.5 S have a frictional ratio of  $\approx 1.9$  and are equivalent to a prolate ellipsoid with a long axis (36 nm) approximately double that of GTP-bound TG2 (Table 1). These expanded form(s) may be similar to calcium-bound forms described in a Monte Carlo simulation of small-angle x-ray scattering data (16), where the  $\beta$ -barrels rotated up to  $130^\circ$  away from the core domain. The 0.5–1.5 S species are extremely expanded/asymmetric, equivalent to a prolate ellipsoid with an axial ratio of  $>30:1$  or a cylinder with a length-to-diameter ratio  $>10.2$ , and may be partially unfolded. This may point to an important role for GTP in stabilizing TG2 tertiary structure.

The absence of a concentration-dependent shift of the  $\approx 3.2$  S peak upon WT and Y516F dilution (Fig. 3) indicates a very slow exchange between this and slower sedimenting species at 4°C ( $k_{off} \approx 10^{-4}$ /s or slower) (21). This is consistent with a significant GTP-dependent conformational change requiring a relatively large activation energy. In contrast, R579A dilution showed progressive decrease of the main boundary from 3.1 S to an endpoint (observed with 33- and 100-fold dilution) of 2.8 S. This indicates a reaction on the time scale of the experiment and likely reflects faster GTP dissociation from R579A, consistent with its lower GTP affinity (10). Moreover, GTP dissociation is accompanied by conformational change, because reduction from 3.1 to 2.8 S is much greater than expected from loss of one GTP molecule, using the 2/3 rule ( $s \approx M^{2/3}$ ). The semicompact form of R579A mutants (Fig. 1B and C) may correspond to a rapidly





It is tempting to speculate that Arg-587 also may be destabilizing in TG3 and that nucleotide or  $\beta$ -octyl glucoside binding to mask Arg-587 promotes crystallization in the compact conformation.

Robust transamidase activity of Y516F and Y516C (SI Fig. 7 and Results) is contrary to recent molecular modeling data that suggest Tyr-516 functions in transition-state stabilization (25) but is entirely consistent with our earlier work demonstrating that Trp-241 is a transition-state stabilizing residue (8). In addition, calcium dependence of Y516F transamidase activity shows that loss of the Cys-277–Tyr-516 bond, and destabilization of the compact form alone, is not sufficient to activate transamidase activity; rather, calcium binding is required. Molecular details of this activation are as yet unknown, however, active site access to substrate would disrupt  $\approx 3,000 \text{ \AA}^2$  of buried surface area between the core and  $\beta$ -barrel 1 (26). Cis/trans isomerization of three energetically unfavorable cis-peptide bonds (two of which are nonproline), found in all transglutaminases crystallized to date, may reduce the estimated 11 kcal/mol of binding energy (26). A schematic HERA diagram (27) of TG2 secondary structure (data not shown) indicates that two of the cis-peptide bonds (Lys-273–Tyr-274 and Gly-372–Pro-373) are components of  $\beta$ -bulges (28), important determinants of protein stability and function (29) that have been associated with conformational isomerization (30). Gly-372–Pro-373 is in the  $\beta$ -sheet containing the active site Asp-358 and is close to the nonproline cis-peptide bond Lys-387–Tyr-388 (Fig. 5). Lys-273–Tyr-274 is in a small parallel  $\beta$ -sheet immediately N-terminal to the  $\alpha$ -helix containing Cys-277. Abrogation of the Cys-277–Tyr-516 H bond may promote the initial step(s) in a sequence of cis/trans-peptide bond isomerizations of Lys-273–Tyr-274, Gly-372–Pro-373, and Lys-387–Tyr-388, leading to destabilization of the compact TG2 form.

In conclusion, GTP-mediated conformational transition of TG2 to its compact, transamidase-inactive form involves masking of a destabilizing Arg-579 residue by GTP binding and formation of a stabilizing Cys-277–Tyr-516 H bond.

## Materials and Methods

**Protein Preparation, Enzyme, and Binding Assays.** Site-directed mutants were constructed in rat TG2 cDNA (19) by using GST-

TG2/pGEX2T (31) as a template. *Escherichia coli*-expressed, thrombin-cleaved TG2-GST fusion proteins were purified by using affinity and ion exchange chromatography (10). Transamidase activity assays (9), GTP photolabeling, and isothermal titration calorimetry (10) were performed as described. Approximately 30% of Y516F (0.3 Y516F:1 GTP $\gamma$ S binding stoichiometry) was available to bind GTP, likely due to residual nucleotide binding as observed for WT and other mutants (10).

**nPAGE.** This was performed on 8–10% resolving gels (100 V at 4°C) by using Laemmli buffers (32) without SDS, with pH adjusted at 4°C. Loading buffer (0.5 M Tris/20% glycerol/bromophenol blue/20 mM DTT, pH 6.8 at 4°C) was added to ice-chilled samples in a 3:1 sample:buffer ratio. Gels were stained with 0.5% Coomassie brilliant blue R-250 in 40% methanol/10% acetic acid and destained in 10% methanol/10% acetic acid.

**Disulfide Bond Formation.** The redox catalyst Cu(II)(1,10-phenanthroline)<sub>3</sub> (made by diluting 20 mM CuSO<sub>4</sub>/20 mM 1,10-phenanthroline in buffer) was used to trigger oxidation by ambient dissolved oxygen (33).

**Sedimentation Velocity Experiments.** TG2 was dialyzed against buffer containing GTP or GTP $\gamma$ S at indicated concentrations. Interference optical fringe displacement data were acquired (30- to 45-sec intervals) at 201,600  $\times$  g by using a Beckman-Coulter (Fullerton, CA) Optima XL1 ultracentrifuge fitted with an AnTi60 rotor. Temperature-equilibrated cells were loaded with sample (350  $\mu$ l) and dialysate (350  $\mu$ l). Data were analyzed by the *c(s)* method (and variations thereof) with SEDFIT (34) and SEDPHAT (35, 36) programs. For further details, see *SI Materials and Methods*.

We thank Sara Holman for protein purification. This work was supported by Australian National Health and Medical Research Council Grant 142000, Australian Research Council Grant DP0665422, and a National Heart Foundation Postdoctoral Fellowship (to G.E.B.).

- Achyuthan KE, Greenberg CS (1987) *J Biol Chem* 262:1901–1906.
- Lorand L, Graham RM (2003) *Nat Rev Mol Cell Biol* 4:140–157.
- Lee KN, Birckbichler PJ, Patterson MK, Jr (1989) *Biochem Biophys Res Commun* 162:1370–1375.
- Nakaoka H, Perez DM, Baek KJ, Das T, Husain A, Misono K, Im MJ, Graham RM (1994) *Science* 264:1593–1596.
- Park ES, Won JH, Han KJ, Suh PG, Ryu SH, Lee HS, Yun HY, Kwon NS, Baek KJ (1998) *Biochem J* 331:283–289.
- Veza R, Habib A, FitzGerald GA (1999) *J Biol Chem* 274:12774–12779.
- Liu S, Cerione RA, Clardy J (2002) *Proc Natl Acad Sci USA* 99:2743–2747.
- Iismaa SE, Holman S, Wouters MA, Lorand L, Graham RM, Husain A (2003) *Proc Natl Acad Sci USA* 100:12636–12641.
- Iismaa SE, Wu MJ, Nanda N, Church WB, Graham RM (2000) *J Biol Chem* 275:18259–18265.
- Begg GE, Holman SR, Stokes PH, Matthews JM, Graham RM, Iismaa SE (2006) *J Biol Chem* 281:12603–12609.
- Yee VC, Pedersen LC, Le Trong I, Bishop PD, Stenkamp RE, Teller DC (1994) *Proc Natl Acad Sci USA* 91:7296–7300.
- Noguchi K, Ishikawa K, Yokoyama K, Ohtsuka T, Nio N, Suzuki E (2001) *J Biol Chem* 276:12055–12059.
- Fesüs L, Laki K (1977) *Biochemistry* 16:4061–4066.
- Casadio R, Polverini E, Mariani P, Spinozzi F, Carsughi F, Fontana A, Polverino de Lauro P, Matteucci G, Bergamini CM (1999) *Eur J Biochem* 262:672–679.
- Di Venere A, Rossi A, De Matteis F, Rosato N, Agro AF, Mei G (2000) *J Biol Chem* 275:3915–3921.
- Mariani P, Carsughi F, Spinozzi F, Romanzetti S, Meier G, Casadio R, Bergamini CM (2000) *Biophys J* 78:3240–3251.
- Prasanna Murthy SN, Lomasney JW, Mak EC, Lorand L (1999) *Proc Natl Acad Sci USA* 96:11815–11819.
- Lee KN, Arnold SA, Birckbichler PJ, Patterson MK, Jr, Fraij BM, Takeuchi Y, Carter HA (1993) *Biochim Biophys Acta* 1202:1–6.
- Prasanna Murthy SN, Iismaa SE, Begg G, Freymann DM, Graham RM, Lorand L (2002) *Proc Natl Acad Sci USA* 99:2738–2742.
- Thompson G, Owen D, Chalk PA, Lowe PN (1998) *Biochemistry* 37:7885–7891.
- Dam J, Velikovskiy CA, Mariuzza RA, Urbanke C, Schuck P (2005) *Biophys J* 89:619–634.
- Ahvazi B, Boeshans KM, Idler W, Baxa U, Steinert PM, Rastinejad F (2004) *J Biol Chem* 279:7180–7192.
- Ahvazi B, Boeshans KM, Steinert PM (2004) *J Biol Chem* 279:26716–26725.
- Ahvazi B, Kim HC, Kee S-H, Nemes Z, Steinert PM (2002) *EMBO J* 21:2055–2067.
- Chica RA, Gagnon P, Keillor JW, Pelletier JN (2004) *Protein Sci* 13:979–991.
- Weiss MS, Metzner HJ, Hilgenfeld R (1998) *FEBS Lett* 423:291–296.
- Hutchinson EG, Thornton JM (1990) *Proteins* 8:203–212.
- Richardson JS, Getzoff ED, Richardson DC (1978) *Proc Natl Acad Sci USA* 75:2574–2578.
- Axe DD, Foster NW, Fersht AR (1999) *J Mol Biol* 286:1471–1485.
- Steadman DJ, Zhao PS, Spencer HT, Dunlap RB, Berger SH (1998) *Biochemistry* 37:7089–7095.
- Iismaa SE, Chung L, Wu MJ, Teller DC, Yee VC, Graham RM (1997) *Biochemistry* 36:11655–11664.
- Laemmli UK (1970) *Nature* 227:680–685.
- Kobashi K (1968) *Biochim Biophys Acta* 158:239–245.
- Schuck P (2000) *Biophys J* 78:1606–1619.
- Schuck P (2005) in *Analytical Ultracentrifugation: Techniques and Methods*, eds Scott DJ, Harding SE, Rowe AJ (R Soc Chem, Cambridge, UK), pp 26–49.
- Boukari H, Nossal R, Sackett DL, Schuck P *Phys Rev Lett* 93:098106, 2004.
- Chan AW, Hutchinson EG, Harris D, Thornton JM (1993) *Protein Sci* 2:1574–1590.
- Kunz ID (1971) *J Amer Chem Soc* 93:516–518.